

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Effect of dietary inclusion of different lipid supplements on quality and oxidative susceptibility of beef meat

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1689142> since 2020-01-14T12:01:24Z

Published version:

DOI:10.1080/1828051X.2018.1485517

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



Volume 15
Number 1
January 2016

italian
journal
of
animal
science



ISSN 1528-0845

Effect of dietary inclusion of different lipid supplements on quality and oxidative susceptibility of beef meat

Journal:	<i>Italian Journal of Animal Science</i>
Manuscript ID	TJAS-2016-0320.R4
Manuscript Type:	Short Communication
Date Submitted by the Author:	n/a
Complete List of Authors:	Mordenti, Attilio; Universita degli Studi di Bologna, DIMEVET Brogna, Nico; Universita degli Studi di Bologna, DIMEVET Merendi, Flavia; Universita degli Studi di Bologna, DIMEVET Formigoni, Andrea; Universita degli Studi di Bologna, DIMEVET Sardi, Luca; Universita degli Studi di Bologna Dipartimento di Scienze Mediche Veterinarie Cardenia, Vladimiro; Universita degli Studi di Bologna, DISTAL Rodriguez Estrada, Maria; Universita degli Studi di Bologna, DISTAL
Keywords:	beef quality, flaxseed, CLA, vitamin E, lipid fatty acid composition and oxidation
Abstract:	The aim of this study was to evaluate the impact of dietary extruded flaxseed (EF) and/or rumen-protected (rp) lipids on beef meat quality and oxidation. Sixty-three crossbred heifers (Charolais X Limousin) were evenly distributed into seven experimental groups, balanced in terms of age and BW. The feeding groups differed in both, the dietary lipid source (EF and/or rp conjugated linoleic acid (CLA), supplemented with vitamin E (VE)) and the supplementation length (90 or 180 days before slaughter); the same total amount of lipids was administered to the animals. With the respect of the control group, the α -linolenic acid content significantly increased and the n-6/n-3 ratio decreased in cattle groups fed with EF. The results also

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

	<p>show that providing CLA and/or EF did not lead to an increase in CLA isomers (CLAs) in the lipid fraction of the meat. Our finding show that administering EF for a shorter period with respect to longer, moderately enhance the nutritional value of the meat with regard to fat composition. No significant effects of dietary supplementation on TBA-reactive substances (TBARs) levels of fresh meat were observed; however, the protection provided by vitamin supplementation might become evident during meat storage under commercial retail conditions.</p>

SCHOLARONE™
Manuscripts

For Peer Review Only

1
2
3 1 **Effect of dietary inclusion of different lipid supplements on quality and oxidative**
4
5 2 **susceptibility of beef meat**
6
7 3

8
9 4 A.L. Mordenti^{1*}, N. Brogna¹, F. Merendi¹, A. Formigoni¹, L. Sardi¹, V. Cardenia², M.T.
10
11 5 Rodriguez-Estrada²
12
13 6

14
15 7 ¹ *Dipartimento di Scienze Mediche Veterinarie, Università di Bologna (DIMEVET; Via Tolara di*
16
17 *sopra 50; 40064, Ozzano Emilia, Bologna), Italy.*
18
19 8

20 9 ² *Dipartimento di Scienze e Tecnologie Agro-Alimentari, Università di Bologna (DISTAL; Viale G.*
21
22 *Fanin 40; 40127, Bologna), Italy.*
23
24 10
25 11

26 12 *Corresponding author: Attilio Luigi Mordenti; <http://orcid.org/0000-0002-4162-6726>; Phone:
27
28 13 +390512097381; Fax: +390512097373; email: attilio.mordenti@unibo.it
29
30 14

31
32
33 15 **Keywords:** beef quality; flaxseed; CLA; vitamin E; lipid fatty acid composition and oxidation.
34
35 16

36
37 17 **Abstract**
38

39 18 The aim of this study was to evaluate the impact of dietary extruded flaxseed (EF) and/or rumen-
40
41 19 protected (rp) lipids on beef meat quality and oxidation. Sixty-three crossbred heifers (Charolais X
42
43 20 Limousin) were evenly distributed into seven experimental groups, balanced in terms of age and
44
45 21 BW. The feeding groups differed in both, the dietary lipid source (EF and/or rp conjugated linoleic
46
47 22 acid (CLA), supplemented with vitamin E (VE)) and the supplementation length (90 or 180 days
48
49 23 before slaughter); the same total amount of lipids was administered to the animals. With the respect
50
51 24 of the control group, the α -linolenic acid content significantly increased and the n-6/n-3 ratio
52
53 25 decreased in cattle groups fed with EF. The results also show that providing CLA and/or EF did not
54
55 26 lead to an increase in CLA isomers (CLAs) in the lipid fraction of the meat. Our finding show that
56
57
58
59
60

1
2
3 27 administering EF for a shorter period with respect to longer, moderately enhance the nutritional
4
5 28 value of the meat with regard to fat composition. No significant effects of dietary supplementation
6
7 29 on TBA-reactive substances (TBARs) levels of fresh meat were observed; however, the protection
8
9 30 provided by vitamin supplementation might become evident during meat storage under commercial
10
11 31 retail conditions.
12
13
14 32

15 33 **Introduction**

16
17
18 34 In the common consumer perception of beef meat is considered a high-fat food and its consumption
19
20 35 may increase the risk of cardiovascular disease and colon cancer in humans (McAfee et al., 2010;
21
22 36 McNeill, 2014). Recent nutritional guidelines (EFSA, 2005; U.S. Department of Health and Human
23
24 37 Services, 2010) recommend that people enhance their intake with foods rich in polyunsaturated
25
26 38 fatty acids (PUFA). Particularly the intake of PUFA n-3 should be increased, so that the western
27
28 39 diet n-6/n-3 FA ratio can decrease from 16/1 to 5/1, thus promoting health conditions and
29
30 40 preventing the risk of diseases (Moloney et al, 2008). Human health also seems to significantly
31
32 41 benefit from the intake of several CLAs, two of which are commonly found in beef and dairy
33
34 42 products and are known to possess biological activity: the anti-carcinogenic and anti-atherogenic
35
36 43 effects of cis-9, trans-11 CLA and the anti-obesity effects of trans-10, cis-12 CLA have been
37
38 44 documented in animal experiments (McAfee et al., 2010; Salter, 2013). With the aim to positively
39
40 45 influence the FA composition of beef intramuscular fat, pasture-based farming is the better strategy
41
42 46 (Priolo et al., 2001). The possibility to influence the FA content can be achieved also in intensive
43
44 47 farming conditions, with cereals and corn silage based diets, by modifying the composition of the
45
46 48 lipid fraction of these diets (Pouzo et al., 2016; Scollan et al., 2014; Juarez et al., 2012). However, a
47
48 49 higher content of PUFA makes meat more prone to oxidation, which can impact its overall quality
49
50 50 and sensory traits (such as colour, flavour, aroma) (Priolo et al., 2001). This may also compromise
51
52 51 animal health and could result in the development of unhealthy oxidation products in beef, together
53
54 52 with decreased colour stability with a consequent reduction in shelf life (Moloney et al. 2008).
55
56
57
58
59
60

1
2
3 53 Therefore, the addition of suitable dietary protections, such as vitamin E (VE), may be important to
4
5 54 control oxidation, especially during storage (Juarez et al., 2012). In fact, VE supplementation can
6
7 55 lengthen the retail shelf-life of beef by 1.6 to 5 days (Nassu et al., 2011b). The aim of this study was
8
9 56 to improve the meat nutritional properties by evaluating the impact of these dietary lipid treatments
10
11 57 on beef meat quality (total fat content, FA composition and oxidation susceptibility).
12
13
14 58

15 16 59 **Materials and methods**

17
18 60 The Scientific Ethics Committee on Animal Experimentation of the University of Bologna
19
20 61 approved the experimental protocol of this study (n.: 71674-X/6 - All.: 63). Sixty-three crossbred
21
22 62 heifers (Charolais X Limousin), approximately 11 months old and 380 kg weight, were randomly
23
24 63 distributed into seven groups. Each group received a basal diet included (kg/head/d): corn silage (8),
25
26 64 beet pulp silage (5), corn meal (2.5), straw (0.8) and a concentrate (2). Feeding was enriched with
27
28 65 various experimental supplements (EF, D α -tocopheril acetate (VE), CLAs (LUTA-CLA 60[®]
29
30 66 (BASF) guarantees a minimum of 56% of conjugated linoleic acid methyl ester (28% cis-9, trans-11
31
32 67 CLA and 28% trans-10, cis-12 CLA)) at different doses, leading to 7 independent diet groups
33
34 68 (Table 1). The diets of the experimental groups differed with respect to the lipid source and to the
35
36 69 length of administration but, considering their entire experimental period (180 days), the same total
37
38 70 amount of lipids was administered to the animals. When animals reached an average BW of 675 kg,
39
40 71 they were transported to a slaughtering facility (Unipeg, Reggio Emilia). Twenty-four hours after
41
42 72 slaughtering, a three-rib sample (20 cm of thickness from the 7th-9th thoracic vertebrae) of
43
44 73 *Longissimus thoracis et lumborum* muscle (LT), was taken from the half right carcass of each of the
45
46 74 63 heifers. The muscle samples were vacuum-packaged and wet-aged for 14 days in a refrigerator at
47
48 75 2-4 °C. At the end of the aging period, each sample was cut into three subsamples, then vacuum-
49
50 76 packaged, rapidly frozen and stored at -20 °C. At the time of analysis, the 63 beef samples were
51
52 77 thawed in 3 different work sessions. Each first subsample was deboned and the first and last slices
53
54 78 were cut and discarded. After 30 min of exposure to air, colour (L*, a*, b*) was measured using a
55
56
57
58
59
60

1
2
3 79 Minolta Chromameter CR-200 (Illuminant D65, aperture size 8 mm; Minolta Camera, Osaka,
4
5 80 Japan). The LT muscle was then divided into a 50 x 50 x 10 mm thick sample to measure cooking
6
7 81 loss; thereafter, in the same cooked sample, Warner-Bratzler Shear Force (WBSF) was evaluated.
8
9 82 The WBSF method was assessed also on raw meat. Cooking loss (%) was determined using the
10
11 83 Honikel gravimetric method (Honikel, 1998). The WBSF test was carried out on the cooked meat
12
13 84 samples, after the cooking loss assessment (AMSA, 1995); the same procedure was carried out on
14
15 85 raw meat samples. The second subsample (weight 200 g) was used for proximate analysis; moisture
16
17 86 and ash content were determined in ground freeze-dried samples. Crude protein was determined
18
19 87 using a Kjeldahl Nitrogen/Protein Analyzer (Gerhardt Vapodest 50, Gerhardt. GMBH; Germany).
20
21 88 The ash content was determined by incinerating samples in a muffle furnace at 500 °C. The LT fat
22
23 89 content was determined with the Soxhlet method (2055 Soxtec Avanti, Foss Tecator AB, Höganäs,
24
25 90 Sweden), according to the AOAC (1992). Intramuscular fat was extracted from LT muscle
26
27 91 according to Folch et al. (1957). Fatty acid methyl esters were prepared according to the method
28
29 92 proposed by Christopherson et al. (1969). The upper organic phase was collected and filtered with
30
31 93 anhydrous sodium sulphate and 1 µL was injected into a Fisons HRGC 8560 series MEGA2 gas
32
33 94 chromatograph (Thermo Electron Corporation, Milan, Italy) equipped with a flame ionization
34
35 95 detector and an AS 2000 automatic injection system (Thermo Electron Corporation, Milan, Italy).
36
37 96 The column was a Varian (Varian Inc. Ass 1985, USA) CP-SIL 88 WCOT Fused Silica capillary
38
39 97 column (100 m x 0.25 mm i.d. x 0.20 µm film thickness). FAME identification was performed by
40
41 98 comparing their retention times with those of known standard FAME mixtures (SUPELCO, Sigma
42
43 99 Aldrich Inc., St. Louis, MO). Each FAME was expressed as a percentage of the total areas of all
44
45 100 FAME peaks. Peroxide value (PV) was determined in the lipid extract (50 mg) from the third
46
47 101 subsample, as suggested by Shantha and Decker (1994). This method is based on the ability of
48
49 102 peroxides to oxidize ferrous ions to ferric ones. Ammonium thiocyanate then reacts with ferric ions,
50
51 103 resulting in a coloured complex that is measured at 500 nm with a spectrophotometer (double-beam
52
53 104 UV-visible Jasco model V-550, Jasco International, Tokyo, Japan). PV was calculated from the
54
55
56
57
58
59
60

1
2
3 105 absorbance at 500 nm. For the quantitative determination of PV, an Fe(III) standard calibration
4
5 106 curve was used with a concentration range of 0.1-5 $\mu\text{g/mL}$ ($y = 0.0282x - 0.0003$; $r^2 = 0.999$). PV
6
7 107 was expressed as meq of O_2 per kg of fat. Three replicates were run per sample. TBA-reactive
8
9 108 substances (TBARS) were determined in a 2 g sample (ground meat), according to the Witte (1970)
10
11 109 method. This method is based on the reaction between thiobarbituric acid and the aldehydes that
12
13 110 derive from secondary oxidation of the lipids present in the sample, resulting in a coloured complex
14
15 111 that can be measured at 530 nm. TBARS were evaluated at such wavelength with a double-beam
16
17 112 UV-visible spectrophotometer (Jasco model V-550, Jasco International, Tokyo, Japan), and they
18
19 113 were calculated from the absorbance. For the quantitative determination of TBARS, a 1,1,3,3-
20
21 114 tetramethoxypropane standard calibration curve was used with a concentration range of 0.045-0.113
22
23 115 $\mu\text{g/mL}$ ($y = 0.0087x - 0.0051$; $r^2 = 0.999$). The TBARS values were expressed as mg of
24
25 116 malonylaldehyde (MDA) per kg of sample. Three replicates were run per sample. All data were
26
27 117 analysed by one-way factorial ANOVA with repeated measures, using Tukey's HSD as *post-hoc*
28
29 118 test. The dietary treatment was used as the source of variation to determine whether there were any
30
31 119 significant differences ($P \leq 0.05$). Pearson's correlation coefficient ($\alpha = 0.05$) was used to separate
32
33 120 the means of statistically different parameters and interactions and examine the possible
34
35 121 relationships between the degree of unsaturation of the meat and the main oxidative indices. The
36
37 122 Statistica version 10 software (StatSoft Inc., Padova, Italy) was used.
38
39
40
41
42
43

44 124 **Results and discussion**

45
46 125 Table 2 shows the chemical composition of the LT muscle: no significant increase in the lipid
47
48 126 content of meat from heifers that had received flaxseed, or CLA, was noted. This result is in
49
50 127 agreement with Juarez et al. (2012), who administered 10% of ground flaxseed to feedlot steers for
51
52 128 129 days. WBSF data here obtained, can be classified as "very tender", according to the Destefanis
53
54 129 et al. (2008) WBSF scale (category n. 5; <3.36 kg). As observed by Juarez et al. (2012) and Nassu
55
56 130 et al. (2011b), meat tenderness was not affected by lipid supplementation. Regarding the colour

1
2
3 131 measurements performed in the aged meat, L value (lightness) of group C was significantly less
4
5 132 bright than those of B, F and G groups. This finding might be related to those presented by Priolo et
6
7 133 al. (2001) who have noted by the linear regression between objective L value and time on pasture
8
9 134 that was significant ($P < 0.001$; $r^2 = 0.74$). The cause of this effect is extremely difficult to evaluate
10
11 135 because several factors play an important role, such as: carcass fatness, meat ultimate pH, animal
12
13 136 age, carcass weight and intramuscular fat content. Table 3 shows LT muscle fatty acid profile, in
14
15 137 particular for α -linolenic acid content, we observed a dose effect in groups supplemented with EF
16
17 138 compared with the A diet (C, D and E, average 0.31%). Moreover the α -linolenic acid content in a
18
19 139 short term EF supplementation (90 days, C, average 0.71%) is to prefer than a long term
20
21 140 supplementation (180 days, B, F and G, average 0.54%). Furthermore, Pearson test ($\alpha=0.05$),
22
23 141 between EF increasing doses (0, 250 and 500 g/head/day) and α -linolenic acid level in lipid meat,
24
25 142 was statistically significant ($r = 0.74825$). Priolo et al. (2001) reported the variation of α -linolenic
26
27 143 acid in intramuscular fat of cattle: grazing for six months on pasture has an effect of increasing
28
29 144 intramuscular α -linolenic acid by 50%. However, considering the lipid meat content shown in Table
30
31 145 3, these differences were relatively low from a quantitative point of view. Considering these results,
32
33 146 administering the same amount of flaxseed in a shorter time may be the best strategy (Nassu et al.,
34
35 147 2011a) to reduce partially the n-6/n-3 ratio (calculated on all FA detected) as noticed in previous
36
37 148 studies (Mordenti et al., 2005; Mordenti et al., 2013). Regarding the two main CLAs contents, our
38
39 149 results indicate that administering rp CLA, with or without EF, do not have had effectiveness on
40
41 150 both CLAs contents as compared to A group; obviously the CLAs contents were not influenced by
42
43 151 the treatment length. This finding is in agreement with what has been reported by other authors
44
45 152 (Schiavon et al., 2011) and might have been due to an CLA utilization directly by animals, reducing
46
47 153 body mass (trans-10, cis-12 CLA) and improving feed efficiency (cis-9, trans-11 CLA). Among
48
49 154 PUFA, arachidonic acid content was significantly higher ($P \leq 0.05$) in group A than in groups B and
50
51 155 D. The content of eicosatrienoic acid (C20:3 n-3 and n-6) was significantly higher ($P \leq 0.05$) in
52
53 156 group A than in groups B, D and G, and the A group amount of n-3 docosapentaenoic acid (DPA)

1
2
3 157 was significantly higher compared to the B. The n-6/n-3 ratio was significantly higher in group A
4
5 158 than in groups B, C, F and G, by 27%, 37%, 25% and 30%, respectively. Regarding the oxidative
6
7 159 parameters (Table 3), PV varied from 0.57 to 4.92 meq O₂/kg lipids, being significantly lower in the
8
9 160 A diet with respect to the B diet. TBARs values ranged from 0.19 to 0.74 mg MDA/kg meat,
10
11 161 remaining below the acceptance threshold of 2 mg MDA/kg beef meat (Campo et al., 2006). No
12
13 162 significant effects of dietary supplementation on TBARs levels of fresh meat were observed. No
14
15 163 correlations (Pearson test, $\alpha=0.05$) were found between the results obtained for lipid composition
16
17 164 and the oxidative parameters resulting from the seven types of feeding conditions. In general, a low
18
19 165 oxidation level was found in all samples, similar to that reported by Boselli et al. (2009), but both
20
21 166 PV and TBARs were greater than those reported by other authors (Insani et al., 2008). This could be
22
23 167 due to Italian carcass processing practices, which provide for a storage period, aimed at improving
24
25 168 meat tenderness and promoting the formation of aroma compounds, which develop during cooking
26
27 169 (Rodriguez-Estrada et al., 1997; Boselli et al., 2009). The long-term supplementation of EF seems
28
29 170 to favour primary oxidation, even in the presence of vitamin E. Although the dietary
30
31 171 supplementation did not have any significant impact on TBARs levels of fresh meat, the protective
32
33 172 action of vitamin addition might become evident during meat storage under commercial retail
34
35 173 conditions (Cardenia et al., 2011). On the other hand, it might be pointed out that TBARs describe
36
37 174 an overall oxidative behaviour, which is useful if compared with PV; however, other specific
38
39 175 antioxidant parameters (such as antioxidant enzyme activities and antioxidant capacity) could
40
41 176 provide additional information (Mahecha et al., 2011).

42 177

43 178 **Conclusions**

44 179 The results obtained in the present study confirm that the diets that included EF supplementation
45
46 180 proved to be effective at improving the n-3 FA and lowering the n-6/n-3 ratio in lipid fraction of
47
48 181 beef meat. For this aspect, a shorter period of supplementation (90 days) was a more successful
49
50 182 strategy than a longer one (180 days). However, considering the lipid meat content, these quantities

1
2
3 183 were relatively low for humans, from a quantitative point of view. Although oxidation parameters
4
5 184 did not reflect the antioxidant effect of the latter, this effect might become evident during meat
6
7 185 storage (Cardenia et al., 2011).
8

9 186

11 187 **Disclosure statement**

13 188 The authors report no conflicts of interest.
14
15

16 189

18 190 **References**

20 191 AOAC (Association of Official Analytical Chemists). 1992. (991.36) Fat (Crude) in Meat and Meat
21
22 192 Products. Association of Official Analytical Chemists. Arlington, VA, USA.

24 193 AMSA. 1995. Research guidelines for cookery, sensory evaluation and instrumental tenderness
25
26 194 measurements of fresh meat. In: American Meat Science Association and National Live Stock
27
28 195 and Meat Board. Chicago, IL, USA.

31 196 Boselli E, Rodriguez-Estrada MT, Fedrizzi G, Caboni MF. 2009. Cholesterol photosensitised
32
33 197 oxidation of beef meat under standard and modified atmosphere at retail conditions. *Meat Sci.*
34
35 198 81, 224-229.

37 199 Campo MN, Nute GR, Hughes SI, Enser M, Wood JD, Richardson RI. 2006. Flavour perception of
38
39 200 oxidation in beef. *Meat Sci.* 72, 303–311.

41 201 Cardenia V, Rodriguez-Estrada MT, Cumella F, Sardi L, Della Casa G, Lercker G. 2011. Oxidative
42
43 202 stability of pork meat lipids as related to high-oleic sunflower oil and vitamin E diet
44
45 203 supplementation and storage conditions. *Meat Sci.* 88, 271-279.

48 204 Christopherson S, Glass RL. 1969. Preparation of milk fat methyl esters by alcoholysis in an
49
50 205 essential nonalcolic solution. *J. Dairy Sci.* 52, 1289-1290.

52 206 Destefanis G, Brugiapaglia A, Barge MT, Dal Molin E. 2008. Relationship between beef consumer
53
54 207 tenderness perception and Warner–Bratzler shear force. *Meat Sci.* 78, 153–156.
55
56
57

- 1
2
3 208 EFSA 2005. Opinion of the scientific panel on dietetic products, nutrition and allergies on a request
4
5 209 from the commission related to nutrition claims concerning omega-3 fatty acids,
6
7 210 monounsaturated fat, polyunsaturated fat and unsaturated fat. The EFSA Journal 253, 1-29.
8
9 211 Folch J, Lees M, Sloane-Stanley GH. 1957. A simple method for the isolation and purification of
10
11 212 total lipids from animal tissues. Journal of Biological Chemistry 226, 497–509.
12
13 213 Honikel KO. 1998. Reference methods for the assessment of physical characteristics of meat. Meat
14
15 214 Sci. 49, 447–457.
16
17 215 Juarez M, Dugan MER, Aldai N, Basarab JA, Baron VS, McAllister TA, Aalhus JL. 2012. Beef
18
19 216 quality attributes as affected by increasing the intramuscular levels of vitamin E and omega-3
20
21 217 fatty acids Meat Sci. 90, 764–769.
22
23 218 Insani EM, Eyherabide A, Grigioni G, Sancho AM, Pensel NA, Descalzo AM. 2008. Oxidative
24
25 219 stability and its relationship with natural antioxidants during refrigerated retail display of beef
26
27 220 produced in Argentina. Meat Sci. 79, 444-452.
28
29 221 Mahecha L, Nuernberg K, Nuernberg G, Martin J, Hubbermann EM, Knoeller S, Claeys E, De
30
31 222 Smet S, Dannenberger D. 2011. Antioxidant enzyme activities an antioxidant capacity in
32
33 223 *longissimus* muscle from bulls fed diets rich in polyunsaturated fatty acids. Food Chemistry 127,
34
35 224 379-386.
36
37 225 McAfee AJ, McSorley EM, Cuskelly GJ, Moss BW, Wallace JMW, Bonham MP, Fearon AM.
38
39 226 2010. Red meat consumption: An overview of the risks and benefits. Meat Sci. 84,1–13.
40
41 227 McNeill SH. 2014. Inclusion of red meat in healthful dietary patterns. Meat Sci. 98(3), 452-460
42
43 228 Moloney AP, Scollan ND, Miles L. 2008. Enrichment of n-3 fatty acids and conjugated linoleic
44
45 229 acid in beef: ProSafeBeef. Nutrition Bulletin 33, 374–381.
46
47 230 Mordenti AL, Gatta PP, Lambertini L, Tassinari M. 2005. Effect of dietary supplementation with
48
49 231 LCPUFA of vegetable origin in lipid content of muscular and subcutaneous tissues in heifers.
50
51 232 Progress in Nutrition 7, 201-210.
52
53
54
55
56
57
58
59
60

- 1
2
3 233 Mordenti AL, Merendi F, Vecchia P, Panciroli A, Tassinari M, Brogna N, 2013. Effect of different
4
5 234 lipid supplements on steer performance, meat quality and fatty acid composition of Longissimus
6
7 235 dorsi muscle. ASPA 20th Congress Bologna, June 11-13. Ital. J. Anim. Sci. 12, C-036.
8
9 236 Nassu RT, Dugan MER, He ML, McAllister TA, Aalhus JL, Aldai JL, Kramer JKG. 2011a. The
10
11 237 effects of feeding flaxseed to beef cows given forage based diets on fatty acids of longissimus
12
13 238 thoracis muscle and backfat. Meat Sci. 89, 469-477.
14
15 239 Nassu RT, Dugan MER, Juarez M, Basarab JA, Baron VS, Aalhus JL, 2011b. Effect of α -
16
17 240 tocopherol tissue levels on beef quality. Animal 5, 2010-2018.
18
19 241 Priolo A, Micol D, Agabriel J. 2001. Effects of grass feeding systems on ruminant meat colour and
20
21 242 flavour. A review. Animal Research 50, 185-200.
22
23 243 Pouzo LB, Descalzo AM, Zaritzky NE, Rossetti L, Pavan E. 2016. Antioxidant status, lipid and
24
25 244 color stability of aged beef from grazing steers supplemented with corn grain and increasing
26
27 245 levels of flaxseed. Meat Sci. 111, 1-8.
28
29 246 Rodriguez-Estrada MT, Penazzi G, Caboni MF, Bertacco G, Lercker G. 1997. Effect of different
30
31 247 cooking methods on some lipid and protein components of hamburgers. Meat Sci. 45, 365-375.
32
33 248 Salter AM. 2013. Dietary fatty acids and cardiovascular disease. Animal 7, 163-171.
34
35 249 Schiavon S, Tagliapietra F, Dal Maso M, Bailoni L, Bittante G. 2011. Effects of low-protein diets
36
37 250 and rumen-protected conjugated linoleic acid on production and carcass traits of growing
38
39 251 double-muscled Piemontese bulls. Journal of Animal Science 88, 3372-3383.
40
41 252 Scollan ND, Dannenberger D, Nuenberg K, Richardson I, MacKintosh S, Hocquette JF, Moloney
42
43 253 AP. 2014. Enhancing the nutritional and health value of beef lipids and their relationship with
44
45 254 quality. Meat Sci. 97, 384-394.
46
47 255 Shantha NC, Decker EA. 1994. Rapid, sensitive, iron-based spectrophotometric method for
48
49 256 determination of peroxide values of food lipids. Journal of AOAC International 77, 421-424.
50
51 257 U.S. Department of Health and Human Services. 2010. Dietary guidelines for Americans. Retrieved
52
53 258 September 17, 2014, from. <http://health.gov/dietaryguidelines/2010.asp>

1
2
3 259 Van Soest PJ, Robertson JB, Lewis BA. 1991. Methods for dietary fiber neutral detergent fiber, and
4
5 260 non-starch polysaccharides in relation to animal nutrition. *J. Dairy Sci.* 74, 583-597.

6
7 261 Witte VC, Krause GF, Bailet ME. 1970. A new extraction method for determining 2-thiobarbituric
8
9 262 acid values of pork and beef during storage. *J. Food Sci.* 35, 582–585.
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only

1
2
3
4 **1 Tables**5
6 **2 Table 1: Description of the seven dietary treatments administered to heifers (g as feed)**
7

8 Groups	Supplementation	Lenght	Description
9 A (control)		180	Basal diet supplemented with Megalac® and soybean meal (protein 14% and energy 1762 kcal/kg)
10 B	EF/VE	180	Basal diet plus 250 g EF head ⁻¹ day ⁻¹ and 1 g VE head ⁻¹ day ⁻¹
11 C	Control + EF/VE	90 + 90	Basal diet (first 90 days) followed by basal diet supplemented with 500 g EF head ⁻¹ day ⁻¹ and 2 g VE head ⁻¹ day ⁻¹
12 D	CLA/VE	180	Basal diet supplemented with 25 g CLA head ⁻¹ day ⁻¹ and 1 g VE head ⁻¹ day ⁻¹
13 E	Control + CLA/VE	90 + 90	Basal diet (first 90 days) followed by basal diet supplemented with 50 g CLA head ⁻¹ day ⁻¹ and 2 g VE head ⁻¹ day ⁻¹
14 F	EF/CLA/VE	180	Basal diet supplemented with 250 g EF head ⁻¹ day ⁻¹ , 25 g CLA head ⁻¹ day ⁻¹ and 1 g VE head ⁻¹ day ⁻¹
15 G	EF/CLA	180	Basal diet supplemented with 250 g EF head ⁻¹ day ⁻¹ and 25 g CLA head ⁻¹ day ⁻¹

16
17
18
19
20
21
22
23
24
25 **3** EF, extruded flaxseed; VE, DL- α tocopheril acetate; CLA, conjugated linoleic acid isomers.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Table 2: Composition (% of DM) and physical properties of the *Longissimus* muscle in heifers

Item	Experimental groups							SEM	P
	A	B	C	D	E	F	G		
Dry Matter	26.9	28.1	26.8	27.2	26.6	27.8	27.2	0.15	ns
Protein	79.0	77.6	78.2	79.3	80.2	78.3	80.3	0.57	ns
Ether extract	13.6	15.3	14.6	12.7	11.9	14.1	12.7	0.56	ns
Cooking loss (%)	30.1	28.9	30.3	30.2	30.2	28.3	29.1	0.37	ns
WBSF fresh (kg)	1.84	1.75	2.10	2.08	2.08	1.89	1.76	0.06	ns
WBSF cooked (kg)	3.38	3.52	3.55	3.23	3.77	4.29	3.56	0.10	ns
Color at day 14									
L value	42.7 ^{ab}	44.6 ^a	40.9 ^b	42.9 ^{ab}	43.3 ^{ab}	44.5 ^a	45.4 ^a	0.35	≤ 0.05
a* value	20.9	21.8	21.3	20.5	20.7	20.2	20.0	0.23	ns
b* value	11.0	11.8	10.5	10.4	10.8	10.6	11.2	0.19	ns

Values in the same row with different superscripts are different (a and b, $P \leq 0.05$; A and B $P \leq 0.01$).

SEM=standard error of the mean; WBSF = Warner-Bratzler Shear Force

1 Table 3: Fatty acid content, fatty acid classes and oxidative parameters of *Longissimus* muscle in
 2 heifers

Fatty acids (% of total FA)	Experimental groups							SEM	P
	A	B	C	D	E	F	G		
C14:0	2.95	3.05	3.18	3.33	3.24	2.95	2.80	0.08	ns
C16:0	28.34	28.61	28.22	29.32	28.51	28.54	27.76	0.25	ns
C16:1 n-7	2.67	2.55	2.45	2.57	2.47	2.74	2.45	0.05	ns
C18:0	17.43	19.22	19.67	19.54	19.51	17.44	19.21	0.26	ns
C18:1 n-9	35.92	36.17	33.59	34.83	34.45	35.97	37.48	0.36	ns
C18:1 n-7	1.94 ^{ab}	1.59 ^{ab}	1.99 ^a	1.65 ^{ab}	1.84 ^{ab}	1.68 ^{ab}	1.43 ^b	0.14	≤ 0.05
C18:2 n-6	6.20	4.91	6.04	5.00	5.59	6.29	4.86	0.21	ns
C18:3 n-3	0.29 ^{Bb}	0.54 ^{Ac}	0.71 ^{Aa}	0.33 ^{Bb}	0.30 ^{Bb}	0.56 ^{Ac}	0.52 ^{Ac}	0.02	≤ 0.05
CLA cis9,trans11	0.34	0.41	0.40	0.33	0.35	0.34	0.36	0.03	ns
CLA trans10,cis12	0.16	0.15	0.14	0.13	0.12	0.15	0.16	0.01	ns
C20:4 n-6	1.10 ^a	0.48 ^b	0.93 ^{ab}	0.57 ^b	0.88 ^{ab}	0.83 ^{ab}	0.58 ^b	0.05	≤ 0.05
C20:5 n-3	0.10 ^{ab}	0.06 ^b	0.12 ^a	0.07 ^b	0.08 ^{ab}	0.11 ^a	0.06 ^b	0.01	≤ 0.05
C22:5 n-3	0.14 ^A	0.06 ^B	0.11 ^{AB}	0.09 ^{AB}	0.11 ^{AB}	0.11 ^{AB}	0.09 ^{AB}	0.01	≤ 0.01
C22:6 n-3	0.29	0.16	0.29	0.18	0.23	0.29	0.19	0.02	ns
SFA	50.01 ^B	52.26 ^{AB}	52.51 ^A	53.61 ^A	52.83 ^A	50.21 ^B	51.14 ^B	0.83	≤ 0.01
MUFA	41.08	40.80	38.53	39.53	39.29	40.89	41.88	0.89	ns
PUFA	8.91	6.94	8.96	6.86	7.88	8.89	6.98	0.74	ns
PUFA/SFA	1.00 ^A	0.91 ^{AB}	0.91 ^{AB}	0.87 ^B	0.90 ^{AB}	1.00 ^{AB}	0.96 ^{AB}	0.03	≤ 0.01
n-6/n-3	8.99 ^a	6.54 ^b	5.70 ^b	8.63 ^a	8.98 ^a	6.76 ^b	6.26 ^b	0.46	≤ 0.05
TBARs	0.70	0.26	0.19	0.44	0.21	0.32	0.73	0.23	ns
PV	0.54 ^b	4.11 ^a	0.85 ^b	2.80 ^{ab}	1.07 ^{ab}	4.28 ^a	2.59 ^{ab}	1.45	≤ 0.05

3 Values in the same row with different superscripts are different (a and b, P≤0.05; A and B P≤0.01).

1 SEM= standard error of the mean; SFA = saturated fatty acids; MUFA = monounsaturated fatty
2 acids; PUFA = polyunsaturated fatty acids; TBARs = TBA-reactive substances (mg MDA/kg
3 meat); PV = peroxide value (meq O₂/kg lipids)
4

For Peer Review Only