

This is the author's manuscript



### AperTO - Archivio Istituzionale Open Access dell'Università di Torino

## Targeting the extracellular signature of metastatic colon cancers

Original Citation:	
Availability:	
This version is available http://hdl.handle.net/2318/1695411	since 2019-03-25T14:38:22Z
Terms of use:	
Open Access	
Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright	
protection by the applicable law.	, , , ,

(Article begins on next page)



Advertisement

# **Cancer Research**

Search... Q

**Advanced Search** 

Home

**About** 

**Articles** 

**For Authors** 

**Alerts** 

News

Experimental and Molecular Therapeutics

# Targeting the extracellular signature of metastatic colon cancers

Serena Marchio, Marco Soster, Fabio Pastorino, Jessica Sun, Maria Monti, Piero Pucci, Davide Dovetta, Marco Botta, Mirco Ponzoni, Wadih Arap, Renata Pasqualini, and Federico Bussolino

DOI: Published May 2008

**Article** 

Info & Metrics


AACR Annual Meeting-- Apr 12-16, 2008; San Diego, CA

### **Abstract**

### 4094

Colon cancer is a leading cause of mortality, mainly as a consequence of malignant cell spreading to liver, brain and lungs. Despite the possibility of a certain range of intervention at least for liver metastases, the success of surgery is related to the very early detection of few lesions, which is rarely achieved with our present knowledge.

Epithelial and endothelial cells in metastatic sites express peculiar proteins, to accomplish for their survival in the host environment. Investigating the complex of these characteristics (the "extracellular signature of metastasis") is pivotal to understand the molecular mechanisms leading to metastases. This extracellular signature is dependent both on the intrinsic properties of tumor cells and on the tissue microenvironment. Consequently, the search for metastasis-specific players has to be conducted with selective processing of tumor tissues. We set up a



protocol for the isolation of heterogeneous cell populations by tissue fractionation of liver metastasis biopsies from colon cancer patients immediately after surgical removal. We screened these tissues with phage-displayed peptide libraries, obtaining more than 200 single peptides binding to liver metastasis cells, 7 of which were further validated.

We next followed two mirror approaches. From the ligand side, we investigated homologies between the 7 metastasis-validated peptides and known human extracellular proteins. This first part of the metastatic signature revealed several proteins potentially involved in tumor progression, angiogenesis and metastasis, among which angiopoietins, cadherins, heregulins, integrins and matrix proteins. From the receptor side, we set up pull-down and proteomics experiments, in which we fished potential receptors with one of the selected peptides. This second part of the metastatic signature revealed two potential players, E-cadherin and α6 integrin, possibly interacting in a molecular complex.

We propose one of more of these ligand/receptor pairs be involved in tumor angiogenesis and/or, more specifically, in the process of liver metastatization. On verification of the molecular mechanism(s), we are presently developing innovative diagnostic (imaging with nanopartilcles) and therapeutic (targeted liposomes) approaches.

### **Footnotes**

• 99th AACR Annual Meeting-- Apr 12-16, 2008; San Diego, CA

American Association for Cancer Research

Previous

▲ Back to top

**Articles** 

Online First

Past Issues

Current Issue

No related articles found.

- ▶ Cited By...
- ▶ More in this TOC Section

Info for **About Cancer** Research Authors

> About the Journal

**Editorial Board** 

Feedback

Home

Alerts

Subscribers

Advertisers