

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Pentachlorophenol Removal from Water by Soybean Peroxidase and Iron(II) Salts Concerted Action

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1705060> since 2019-06-28T11:29:01Z

Published version:

DOI:10.1007/s11270-019-4189-7

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Pentachlorophenol removal from water by Soybean peroxidase and iron(II) salts concerted action

Valentina Tolardo ^{a,1}, Sara García-Ballesteros ^b, Lucas Santos-Juanes ^b, Rosa Vercher ^b, Ana M. Amat ^b, Antonio Arques ^b, Enzo Laurenti ^{a,*}

^a Department of Chemistry, Università di Torino, Via P. Giuria 7, 10125 Torino, Italy.

^b Grupo Procesos de Oxidación Avanzada, Campus de Alcoy, Universitat Politècnica de València, Spain.

¹ Present address: Smart Materials, Istituto Italiano di Tecnologia, Via Morego, 30, 16163 Genova, Italy.

* Corresponding author: enzo.laurenti@unito.it, phone: +390116707951, fax: +39 0116707855, ORCID: 0000-0001-7363-4002

Abstract

Soybean peroxidase (SBP) has been employed for the treatment of aqueous solutions containing pentachlorophenol (PCP) in the presence of hydrogen peroxide at pH range 5-7. Reaction carried out with 1 mg/L of PCP, 4 mg/L of H₂O₂ and 1.3 x 10⁻⁹ M of SBP, showed a fast initial elimination of PCP (ca. 30% in 20 min), but the reaction does not go beyond the removal of 50% of the initial concentration of PCP.

Modification in SBP and PCP amounts did not change the reaction profile and higher amounts of H₂O₂ were detrimental for the reaction. Addition of Fe(II) to the system resulted in an acceleration of the process to reach nearly complete PCP removal at pH 5 or 6; this is more probably due to a synergetic effect of the enzymatic process and Fenton reaction. However, experiments developed in tap water resulted in a lower PCP elimination, but this inconvenience can be partly overcome by leaving the tap water overnight in an open vessel before reaction.

Keywords: soybean peroxidase; Fenton; pentachlorophenol; hydrogen peroxide; wastewater; iron

Acknowledgments

The authors want to thank the financial support of the European Union (PIRSES-GA-2010-269128, EnvironBOS) and Spanish Ministerio de Educación y Ciencia (CTQ2015-69832-C4-4-R). Sara García-Ballesteros would like to thank Spanish Ministerio de Economía y Competitividad for her fellowship (BES-

2013-066201). We also want to acknowledge Davide Mainero from Acea Pinerolese for his collaboration in this research.

Introduction

Chlorophenols are a group of pollutants that constitute a serious environmental and health concern (Li 2018; Zheng et al. 2012). Despite their natural formation in soil (Hoekstra et al. 1999) or in marine environment (Ballschmiter 2003), the main contribution to the presence of these substances in the environment is due to anthropogenic activities, since they are widely used as bactericides, insecticides, herbicides, fungicides or wood preservatives, and as intermediates in the production of dyes and pharmaceuticals (Czaplicka 2004; Olaniran and Igbinosa 2011; Tsai 2013; Verbrugge et al. 2018). In addition, natural microbial degradation of herbicides and pesticides results in a worldwide release of highly toxic derivatives in the environment.

Although chlorophenols can be in turn be degraded by aerobic and anaerobic microorganism, their persistence and the toxicity of degradation products led to the development of new methods to obtain more effective and rapid removal of these substances from the environment.

Since chlorophenols can also inhibit the efficiency of activated sludge bacteria (Essam et al. 2007), the development of more efficient wastewater treatment processes is continuously proposed. Some of these new methods are based on the use of isolated enzymes. According to their high specificity, tailored processes can be designed for the removal of groups of pollutants (Durán and Esposito 2000). Moreover, the use of isolated enzymes instead of whole microorganisms maintains enzyme concentration independent from bacterial growth rates and permits an easier storage (Caza et al. 1999).

Some oxidoreductases (i.e. ligninases, laccases, peroxidases, or tyrosinases) isolated from plants and fungi were tested for this kind of treatment as they are able to transform phenols (including chlorophenols) into less hazardous compounds (Naghdi et al. 2018; Qayyum et al. 2009). In particular, the use of peroxidases is justified by their high oxidative potential and relatively low selectivity, which permits to catalyse the peroxide-mediated oxidation of a wide number of compounds (Passardi et al. 2005). In this context, a stable and efficient peroxidase isolated from soybean (*Glycine max*, L.) seed hulls was proposed for the application in wastewater remediation, showing its ability to degrade both aromatic compounds and synthetic dyes (Caza et al. 1999; Marchis et al. 2011; Steevensz et al. 2014).

On the other hand chemical processes like electrochemical reduction (Sun et al. 2015), hydrodechlorination by hydrogen peroxide (Muñoz et al. 2013) and advanced oxidation processes (AOPs) (Karci 2014; Pera-Titus et al. 2004; Sharma et al. 2013) have been employed for the removal of chlorophenols. In particular,

emerging approaches for wastewater treatment involve, in most cases, the coupling of biological methods with AOPs (Oller et al. 2011; Sarria et al. 2002). For instance, a combined use of ozonation and microbiological degradation has been reported for the removal of 4-chloro and 2,4-dichlorophenol (Garcia-Peña et al. 2012).

The Fenton and photo-Fenton process are among the most widely used AOPs (Babuponnusami and Muthukumar 2014). They use the ability of iron salts to decompose hydrogen peroxide into more reactive species. However, acidic pH (ca. 2.8) is required to avoid iron inactivation and strategies are now being developed to work at mild conditions (Lucas Santos-Juanes et al. 2017); some of them involve coupling treatments, such as combining the reducing ability of zero-valent iron (ZVI) to induce changes in the structure of the molecules and, at the same time, using ZVI as a source of iron for a Fenton process (Donadelli et al. 2018; L. Santos-Juanes et al. 2017). Starting from these data, the possibility of coupling Fenton with SBP seems particularly interesting, as they both employ H_2O_2 as oxidising agent and this strategy might combine the mild pH optimal values of SBP activity, in the range 5-7 (Caza et al. 1999; Marchis et al. 2012), with the higher oxidative efficiency of Fenton, even at low amounts of iron. As far as we know, this approach remains unexplored; therefore in this paper we investigated the potential use of the system SBP/hydrogen peroxide in the presence of iron(II) as a method for wastewater treatment, using pentachlorophenol (PCP) as model compound.

Material and methods

Chemical reagents

Pentachlorophenol (PCP) was purchased from Sigma Aldrich. Hydrogen peroxide (30% v/v) and ferrous sulphate were supplied by Panreac. Water employed in the experiments was Milli-Q grade, except for those run with tap water. All reagents were at high purity grade and used as received.

Soybean peroxidase isolation and purification

Soybean peroxidase (SBP) was extracted from the hulls of soybean seeds, kindly furnished by Prof. D. Sacco (Department of Agricultural, Forest and Food Sciences, Università di Torino, Italy). After the peeling of seeds, the hulls had been stored at -12 °C until use, then SBP was extracted and purified by a process based

on a previously published method (Calza et al. 2016). One hundred grams of soybean hulls were ground in a mortar, added to 600 mL of phosphate buffer (0.025 M, pH 7) and left under stirring for two hours at room temperature. Then, the hulls were separated from the solution by filtration with cotton gauze and the filtrate was centrifuged for 15 min at 4000 rpm. After the centrifugation, the supernatant was separated and tested for peroxidase activity by the H_2O_2 /DMAB-MBTH system: 10 μL of SBP containing supernatant were added to 3 mL of a solution containing 3-(dimethylamino)benzoic acid (DMAB, 5×10^{-4} M), 3-methyl-2-benzothiazolinonehydrazone (MBTH, 2×10^{-3} M) and H_2O_2 (1×10^{-2} M) in acetate buffer 0.1 M pH 5.4. The enzymatic activity was measured by following the increase of absorbance at 590 nm of the reaction product (Ngo and Lenhoff 1980).

The treatment of the hulls with phosphate buffer was repeated five times with decreasing buffer volume (600 mL, 500 mL, 400 mL) until the resulting solution gave a negative response to the activity test. Then, SBP containing solutions were collected with a Vivaspin 20 (Sartorius, 10000 MWCO) tangential filter in a centrifuge (4000 rpm, 30 min). Successively, the proteins were precipitated by addition of ammonium sulphate until saturation (53 g in 100 mL), and the mixture was left under stirring for one night at 4 °C. The precipitate was centrifuged for 20 min at 4000 rpm and dissolved in 250 mL of phosphate buffer (0.025 M, pH 7). The resulting solution was then dialyzed for 24 hours at 4 °C in cellulose tubes (Sigma, 12000 MWCO) against several aliquots of the same buffer.

The dialyzed fraction was loaded onto a column (4 cm \times 20 cm) containing DEAE-Sepharose CL-6B (Sigma-Aldrich) ionic exchange resin, washed with three volumes of phosphate buffer 0.025 M pH 7 and eluted with a KCl gradient 0–0.5 M (500 mL) in the same buffer. The fractions were collected and analysed by means of UV–visible spectroscopy. The selected fractions were pooled and concentrated by ultrafiltration on Vivaspin 20 (Sartorius, 10000 MWCO). The final SBP sample was then stored at –12 °C until use.

Pentachlorophenol removal studies

All the experiments were conducted at room temperature (30 ± 1 °C) in an open glass beaker. SBP (1.3×10^{-9} – 2.6×10^{-9} M) and, when needed, hydrogen peroxide (4–40 mg/L) and iron(II) (4 mg/L, added as sulphate salt) were added to 250 mL of a PCP solution (1 mg/L), kept in the dark and under stirring until use. The pH was adjusted to the desired value by dropwise addition of HCl or NaOH diluted solutions. A sample was

withdrawn as reference before adding H_2O_2 to the PCP/SBP mixture. After the start of the reaction, 1 mL of reaction mixture was collected every 3 minutes up to 30 minutes and then every 30 minutes. Immediately, 1 mL of methanol was added to each sample in order to stop the reaction. The pH of the reaction mixture was measured before the start of the reaction and at the end.

The PCP concentration was determined by HPLC-UV analysis in a Perkin Elmer model Flexiar UPLC FX-10). A Bronwnlee Analytical column (DB-C18) was employed as stationary phase. The elution was carried out with acetonitrile and formic acid 10 mM (85:15 % v/v) in a 0.3 mL/min flow rate. The detector was set up at 215 nm and the results were expressed as A/A_0 vs reaction time, where A_0 is the initial area of the PCP chromatographic peak and A is the corresponding area at each reaction time.

Results and Discussion

3.1. SBP-based enzymatic process

A first series of experiments were performed at pH 5 to study the possibility of using the SBP/ H_2O_2 system to remove PCP from aqueous solutions. For this purpose, the behavior of a solution containing 1 mg/L of PCP (3.75×10^{-6} M) and hydrogen peroxide 4 mg/L (1.17×10^{-4} M) was followed, both in absence and in the presence of SBP (1.3×10^{-9} M). Figure 1 shows as H_2O_2 alone was not able to degrade PCP, while in the presence of the enzyme the reaction occurred, reaching a 35% percent of PCP degradation in 30 min of reaction; beyond this point the reaction slowed down and in the subsequent 60 minutes only a further 5% of PCP degradation occurred. Similar results were obtained in the past for the degradation of PCP with free and immobilized horseradish peroxidase (J. Zhang et al. 2007).




Fig 1. Degradation profiles of PCP (1 mg/L) vs. time under the following experimental conditions: (■) 4 mg/L of H₂O₂ without SBP, (▲) 4 mg/L of H₂O₂ and 1.3·10⁻⁹ M of SBP and (●) 40 mg/L of H₂O₂ and 1.3·10⁻⁹ M of SBP

Moreover, the presence of higher amounts of H₂O₂, 40 mg/L (1.17 x 10⁻³ M), did not result in an enhancement of the process. On the contrary, the results showed a similar time profile, but a slightly lower percentage of PCP degradation was reached (Figure 1). In fact, a detrimental effect of the excess of hydrogen peroxide, with a decrease in the enzyme activity, has been already observed for SBP (Wright and Nicell 1999) and attributed to an irreversible inhibition mechanism typical of heme-peroxidases, as extensively reviewed (Valderrama et al. 2002).

Some experiments were successively conducted in order to better understand the reasons of the incomplete degradation of PCP, even at low H₂O₂ concentration, and the extent of hydrogen peroxide inhibition. In these experiments the hydrogen peroxide concentration was halved (2mg/L) or doubled (8 mg/L), but in the latter case H₂O₂ was added in two aliquots: the first at the start of the reaction and the second after 10 minutes (Figure 2). Moreover, also the concentrations of SBP and PCP were changed, and the corresponding results are shown in Figure 3.




Fig 2. Removal of PCP (1 mg/L) vs. time under the following experimental conditions: (■) 4 mg/L of H₂O₂ and 1.3x10⁻⁹ M of SBP – same data of Figure 1, reported for comparison, (▲) two successive additions of 4 mg/L of H₂O₂ in the presence of 1.3·10⁻⁹ M of SBP and (●) 2 mg/L of H₂O₂ and 1.3x10⁻⁹ M of SBP

The strict similarity of the results obtained with 2, 4 and 8 mg/L of H₂O₂ concentrations (Figure 2) indicates that the inhibition due to hydrogen peroxide is relative unimportant in this range of concentrations and that it does not significantly affect the difficulty of achieving complete PCP degradation. On the other hand, the higher initial rate and percent of PCP removal obtained when the PCP concentration was reduced to 0.25 mg/L (Figure 3) seems to highlight a role for PCP or its reaction products in the inhibition of the enzymatic action.




Fig 3. Degradation profiles of PCP vs. time under the following experimental conditions: A) 1 mg/L of PCP, 4 mg/L of H₂O₂ and two concentrations of SBP: 1.3×10^{-9} M (■) and 2.6×10^{-9} M (●); B) 4 mg/L of H₂O₂, 1.3×10^{-9} M of SBP and two concentrations of PCP: 1 mg/L (■) and 0.25 mg/L (●)

Hence, the loss of efficiency of the reaction in these range of H₂O₂ concentrations could be instead attributed to the persistence of the reaction products into the catalytic site of SBP, which make progressively more difficult the access of new substrate molecules and slow down the catalytic process. This hypothesis is supported by the recognition, in similar experimental conditions, of both 2,3,5,6-tetrachloro-1,4-benzoquinone and 2,2',3,3',5,5',6,6'-octachloro-1,1'-biphenyl-4,4'-diol as product of the PCP reaction catalyzed by horseradish peroxidase (Samokyszyn et al. 1995; G. Zhang and Nicell 2000).

3.2 Evaluating the synergetic effect of SBP and Fenton reaction

Since in none of the previous experiments PCP removal was complete, an alternative approach was used. In particular, iron(II) ions were added to the reaction mixture in order to drive a Fenton process, induced by the presence of hydrogen peroxide, able to enhance the PCP removal. 4 mg/L of iron(II) were added as FeSO₄ to

250 mL of solution containing 1 mg/L of PCP, 2.68×10^{-9} M of SBP and 4 mg/L of hydrogen peroxide. The experiments were carried out at pH 5 and 6 and a control experiment without SBP was also performed at pH = 5.

As shown in Figure 4, the Fenton reaction (in the absence of SBP) was able to remove 60-70% of PCP in the early stages of the process, but also in this case the PCP removal was not complete since iron inactivation by the formation of iron oxides or hydroxides stopped the reaction. Otherwise, in the presence of SBP, a clear synergetic effect was observed and a nearly complete elimination of PCP was reached in almost 1 hour when both SBP and iron(II) were present, even at pH as high as 6. This might be due to a complementary effect of both processes: while the enzymatic reaction is efficient to degrade PCP, the presence of photo-Fenton prevents inhibitory effects due to the presence of reaction intermediates or H_2O_2 in excess.




Fig 4. Progressive removal of PCP vs time in the presence of 4 mg/L of Fe(II), 1 mg/L of PCP and 4 mg/L of H_2O_2 : (●) 2.68×10^{-9} M of SBP at pH = 5, (▲) 2.68×10^{-9} M of SBP at pH = 6, (■) without SBP at pH = 5

Finally, the reaction was also carried out in tap water in order to test the reaction system in conditions closer to natural matrices. The reaction was studied at pH 5, 6 and 7 and results are shown in Figure 5. At pH 7, PCP removal did not occur, while at pH 6 it was very limited (about 10%). On the contrary, at pH 5 more than 50% of PCP was removed in 30 minutes, but the reaction did not proceed further and the PCP concentration remained the same even after 2 hours of reaction.




Fig 5. Removal of PCP vs time in tap water in the presence of 4 mg/L of Fe(II), 1 mg/L of PCP and 4 mg/L of H₂O₂, 2.68 x 10⁻⁹ M of SBP: (▲) at pH = 5, (●) at pH = 6, (■) at pH = 7, (▼) at pH = 5 in tap water stirred overnight in an open vessel

We hypothesized that this result could be due to the scavenging effect of some ions (namely chloride and hydrogen carbonate) towards the reactive species involved in the Fenton reaction (Soler et al. 2009) and to the influence on the enzymatic process of the chlorine present in tap water. In order to clarify this point, tap water was left in an open vessel under magnetic stirring for one day to allow chlorine elimination. After this treatment, actually the reaction did not stop after 30 minutes, reaching 60% of PCP removal after 2 hours (Figure 5) and 70% after 3 hours (data not shown).

Conclusions

The experimental results showed as SBP could be able to remove PCP from aqueous solutions at slightly acidic pH. But the progressive slowing down of the reaction rate, probably due to the interference of intermediates or reaction products, did not allowed the complete removal of PCP.

The presence of iron salts allowed overcoming this inconvenience, most probably due to the existence of a synergetic effect between the Fenton and the enzymatic reactions, which results in a fast removal of the formed by-products. Nonetheless, this is not so efficient in tap water.

Hence, further research is needed in this field in order to determine with other pollutants the real applicability of this approach and to remove the process inconveniences towards real application, in particular, in real aqueous matrices.

References

- Babuponnusami, A., & Muthukumar, K. (2014). A review on Fenton and improvements to the Fenton process for wastewater treatment. *Journal of Environmental Chemical Engineering*, 2(1), 557–572. doi:10.1016/j.jece.2013.10.011
- Ballschmiter, K. (2003). Pattern and sources of naturally produced organohalogens in the marine environment: biogenic formation of organohalogens. *Chemosphere*, 52(2), 313–324. doi:10.1016/S0045-6535(03)00211-X
- Calza, P., Zacchigna, D., & Laurenti, E. (2016). Degradation of orange dyes and carbamazepine by soybean peroxidase immobilized on silica monoliths and titanium dioxide. *Environmental Science and Pollution Research*, 23(23), 23742–23749. doi:10.1007/s11356-016-7399-1
- Caza, N., Bewtra, J. ., Biswas, N., & Taylor, K. . (1999). Removal of phenolic compounds from synthetic wastewater using soybean peroxidase. *Water Research*, 33(13), 3012–3018. doi:10.1016/S0043-1354(98)00525-9
- Czaplicka, M. (2004). Sources and transformations of chlorophenols in the natural environment. *Science of The Total Environment*, 322(1–3), 21–39. doi:10.1016/j.scitotenv.2003.09.015
- Donadelli, J. A., Carlos, L., Arques, A., & García Einschlag, F. S. (2018). Kinetic and mechanistic analysis of azo dyes decolorization by ZVI-assisted Fenton systems: pH-dependent shift in the contributions of reductive and oxidative transformation pathways. *Applied Catalysis B: Environmental*, 231, 51–61. doi:10.1016/j.apcatb.2018.02.057
- Durán, N., & Esposito, E. (2000). Potential applications of oxidative enzymes and phenoloxidase-like compounds in wastewater and soil treatment: a review. *Applied Catalysis B: Environmental*, 28(2), 83–99. doi:10.1016/S0926-3373(00)00168-5
- Essam, T., Amin, M. A., El Tayeb, O., Mattiasson, B., & Guieysse, B. (2007). Sequential photochemical–biological degradation of chlorophenols. *Chemosphere*, 66(11), 2201–2209. doi:10.1016/j.chemosphere.2006.08.036
- Garcia-Peña, E. I., Zarate-Segura, P., Guerra-Blanco, P., Poznyak, T., & Chairez, I. (2012). Enhanced phenol and chlorinated phenols removal by combining ozonation and biodegradation. *Water, Air, and Soil Pollution*, 223(7), 4047–4064. doi:10.1007/s11270-012-1172-y

- Hoekstra, E. J., De Weerd, H., De Leer, E. W. B., & Brinkman, U. A. T. (1999). Natural formation of chlorinated phenols, dibenzo-p-dioxins, and dibenzofurans in soil of a Douglas fir forest. *Environmental Science and Technology*, 33(15), 2543–2549. doi:10.1021/es9900104
- Karci, A. (2014). Degradation of chlorophenols and alkylphenol ethoxylates, two representative textile chemicals, in water by advanced oxidation processes: The state of the art on transformation products and toxicity. *Chemosphere*, 99, 1–18. doi:10.1016/j.chemosphere.2013.10.034
- Li, Z. (2018). Health risk characterization of maximum legal exposures for persistent organic pollutant (POP) pesticides in residential soil: An analysis. *Journal of Environmental Management*, 205, 163–173. doi:10.1016/j.jenvman.2017.09.070
- Marchis, T., Avetta, P., Bianco-Prevot, A., Fabbri, D., Viscardi, G., & Laurenti, E. (2011). Oxidative degradation of Remazol Turquoise Blue G 133 by soybean peroxidase. *Journal of Inorganic Biochemistry*, 105(2), 321–327. doi:10.1016/j.jinorgbio.2010.11.009
- Marchis, T., Cerrato, G., Magnacca, G., Crocellà, V., & Laurenti, E. (2012). Immobilization of soybean peroxidase on aminopropyl glass beads: Structural and kinetic studies. *Biochemical Engineering Journal*, 67, 28–34. doi:10.1016/j.bej.2012.05.002
- Muñoz, M., de Pedro, Z. M., Casas, J. A., & Rodriguez, J. J. (2013). Chlorophenols breakdown by a sequential hydrodechlorination-oxidation treatment with a magnetic Pd-Fe⁰-Al₂O₃ catalyst. *Water Research*, 47(9), 3070–3080. doi:10.1016/j.watres.2013.03.024
- Naghdi, M., Taheran, M., Brar, S. K., Kermanshahi-pour, A., Verma, M., & Surampalli, R. Y. (2018, March 1). Removal of pharmaceutical compounds in water and wastewater using fungal oxidoreductase enzymes. *Environmental Pollution*. Elsevier. doi:10.1016/j.envpol.2017.11.060
- Ngo, T. T., & Lenhoff, H. M. (1980). A sensitive and versatile chromogenic assay for peroxidase and peroxidase-coupled reactions. *Analytical Biochemistry*, 105(1), 389–397. doi:10.1016/0003-2697(80)90475-3
- Olaniran, A. O., & Igbinsola, E. O. (2011). Chlorophenols and other related derivatives of environmental concern: Properties, distribution and microbial degradation processes. *Chemosphere*, 83(10), 1297–1306. doi:10.1016/j.chemosphere.2011.04.009
- Oller, I., Malato, S., & Sánchez-Pérez, J. A. (2011). Combination of Advanced Oxidation Processes and

- biological treatments for wastewater decontamination—A review. *Science of The Total Environment*, 409(20), 4141–4166. doi:10.1016/j.scitotenv.2010.08.061
- Passardi, F., Cosio, C., Penel, C., & Dunand, C. (2005, July 22). Peroxidases have more functions than a Swiss army knife. *Plant Cell Reports*. Springer-Verlag. doi:10.1007/s00299-005-0972-6
- Pera-Titus, M., García-Molina, V., Baños, M. A., Giménez, J., & Esplugas, S. (2004). Degradation of chlorophenols by means of advanced oxidation processes: a general review. *Applied Catalysis B: Environmental*, 47(4), 219–256. doi:10.1016/j.apcatb.2003.09.010
- Qayyum, H., Maroof, H., & Yasha, K. (2009). Remediation and treatment of organopollutants mediated by peroxidases: a review. *Critical Reviews in Biotechnology*, 29(2), 94–119. doi:10.1080/07388550802685306
- Samokyszyn, V. M., Freeman, J. P., Rao Maddipati, K., & Lloyd, R. V. (1995). Peroxidase-Catalyzed Oxidation of Pentachlorophenol. *Chem. Res. Toxicol*, 8, 349–355. <http://pubs.acs.org/doi/pdf/10.1021/tx00045a005>. Accessed 23 June 2017
- Santos-Juanes, L., Amat, A. M., & Arques, A. (2017). Strategies to Drive Photo-Fenton Process at Mild Conditions for the Removal of Xenobiotics from Aqueous Systems. *Current Organic Chemistry*, 21(12), 1074–1083. doi:10.1136/adc.2010.199901
- Santos-Juanes, L., García Einschlag, F. S., Amat, A. M., & Arques, A. (2017). Combining ZVI reduction with photo-Fenton process for the removal of persistent pollutants. *Chemical Engineering Journal*, 310, 484–490. doi:10.1016/j.cej.2016.04.114
- Sarria, V., Parra, S., Adler, N., Péringer, P., Benitez, N., & Pulgarin, C. (2002). Recent developments in the coupling of photoassisted and aerobic biological processes for the treatment of biorecalcitrant compounds. *Catalysis Today*, 76(2–4), 301–315. doi:10.1016/S0920-5861(02)00228-6
- Sharma, S., Mukhopadhyay, M., & Murthy, Z. V. P. (2013). Treatment of chlorophenols from wastewaters by Advanced Oxidation Processes. *Separation & Purification Reviews*, 42(May 2015), 37–41. doi:10.1080/15422119.2012.669804
- Soler, J., García-Ripoll, A., Hayek, N., Miró, P., Vicente, R., Arques, A., & Amat, A. M. (2009). Effect of inorganic ions on the solar detoxification of water polluted with pesticides. *Water Research*, 43(18), 4441–4450. doi:10.1016/j.watres.2009.07.011

- Steevensz, A., Cordova Villegas, L. G., Feng, W., Taylor, K. E., Bewtra, J. K., & Biswas, N. (2014). Soybean peroxidase for industrial wastewater treatment: a mini review. *Journal of Environmental Engineering and Science*, 9(3), 181–186. doi:10.1680/jees.13.00013
- Sun, Z., Wei, X., Zhang, H., & Hu, X. (2015). Dechlorination of pentachlorophenol (PCP) in aqueous solution on novel Pd-loaded electrode modified with PPy-SDBS composite film. *Environmental Science and Pollution Research*, 22(5), 3828–3837. doi:10.1007/s11356-014-3641-x
- Tsai, W.-T. (2013). A review on environmental distributions and risk management of phenols pertaining to the endocrine disrupting chemicals in Taiwan. *Toxicological & Environmental Chemistry*, 95(5), 723–736. doi:10.1080/02772248.2013.818150
- Valderrama, B., Ayala, M., & Vazquez-Duhalt, R. (2002, May 1). Suicide inactivation of peroxidases and the challenge of engineering more robust enzymes. *Chemistry and Biology*. Cell Press. doi:10.1016/S1074-5521(02)00149-7
- Verbrugge, L. A., Kahn, L., & Morton, J. M. (2018). Pentachlorophenol, polychlorinated dibenzo-p-dioxins and polychlorinated dibenzo furans in surface soil surrounding pentachlorophenol-treated utility poles on the Kenai National Wildlife Refuge, Alaska USA. *Environmental Science and Pollution Research*, 25(19), 19187–19195. doi:10.1007/s11356-018-2269-7
- Wright, H., & Nicell, J. A. (1999). Characterization of soybean peroxidase for the treatment of aqueous phenols. *Bioresource Technology*, 70(1), 69–79. doi:10.1016/S0960-8524(99)00007-3
- Zhang, G., & Nicell, J. A. (2000). Treatment of aqueous pentachlorophenol by horseradish peroxidase and hydrogen peroxide. *Water Research*, 34(5), 1629–1637. doi:10.1016/S0043-1354(99)00326-7
- Zhang, J., Ye, P., Chen, S., & Wang, W. (2007). Removal of pentachlorophenol by immobilized horseradish peroxidase. *International Biodeterioration & Biodegradation*, 59, 307–314. doi:10.1016/j.ibiod.2006.09.003
- Zheng, W., Yu, H., Wang, X., & Qu, W. (2012, July 1). Systematic review of pentachlorophenol occurrence in the environment and in humans in China: Not a negligible health risk due to the re-emergence of schistosomiasis. *Environment International*. Pergamon. doi:10.1016/j.envint.2011.04.014