

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Review: Peering through a keyhole: liquid biopsy in primary and metastatic central nervous system tumours**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1701022> since 2022-11-25T15:10:15Z

*Published version:*

DOI:10.1111/nan.12553

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

DR LUCA BERTERO (Orcid ID : 0000-0001-9887-7668)

Article type : Review

## Peering through a keyhole: Liquid biopsy in primary and metastatic central nervous system tumours

**Running title:** Liquid biopsy in CNS tumours

Luca Bertero<sup>1,2</sup>, Giulia Siravegna<sup>3,4</sup>, Roberta Rudà<sup>5,6</sup>, Riccardo Soffietti<sup>5,6</sup>, Alberto Bardelli<sup>3,4</sup>, Paola Cassoni<sup>1,2</sup>

<sup>1</sup> Pathology Unit, Department of Medical Sciences, University of Turin, Torino, Italy

<sup>2</sup> Pathology Unit, AOU Città della Salute e della Scienza di Torino, Torino, Italy

<sup>3</sup> Department of Oncology, University of Turin, Candiolo (TO), Italy

<sup>4</sup> Candiolo Cancer Institute, FPO-IRCCS, Candiolo (TO), Italy

<sup>5</sup> Neuro-oncology Unit, Department of Neurosciences, University of Turin, Torino, Italy

<sup>6</sup> Neuro-oncology Unit, AOU Città della Salute e della Scienza di Torino, Torino, Italy

Corresponding author

Dr. Luca Bertero

Pathology Unit, Department of Medical Sciences

University of Turin and AOU Città della Salute e della Scienza di Torino

Via Santena 7

10126 Torino

Italy

Email: luca.bertero@unito.it

Phone: +390116335466

Fax: +390116635267

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/nan.12553

This article is protected by copyright. All rights reserved.

## Keywords

Liquid biopsy; CNS tumours; CSF; molecular profiling; precision oncology; cfDNA; circulating tumour cells; tumour educated platelets; extracellular vesicles

## Abstract

Tumour molecular profiling by liquid biopsy is being investigated for a wide range of research and clinical purposes. The possibility of repeatedly interrogating the tumour profile using minimally invasive procedures is helping to understand spatial and temporal tumour heterogeneity, and to shed a light on mechanisms of resistance to targeted therapies. Moreover, this approach has been already implemented in clinical practice to address specific decisions regarding patients' follow up and therapeutic management. For central nervous system tumours (CNS), molecular profiling is particularly relevant for the proper characterization of primary neoplasms, while CNS metastases can significantly diverge from primary disease or extra-CNS metastases, thus compelling a dedicated assessment. Based on these considerations, effective liquid biopsy tools for CNS tumours are highly warranted and a significant amount of data has been accrued over the last few years. These results have shown that liquid biopsy can provide clinically-meaningful information about both primary and metastatic CNS tumours, but specific considerations must be taken into account, for example when choosing the source of liquid biopsy. Nevertheless, this approach is especially attractive for CNS tumours, since repeated tumour sampling is not feasible. The aim of our review is, therefore, to thoroughly report the state-of-the-art regarding the opportunities and challenges posed by liquid biopsy in both primary and secondary CNS tumours.

## List of abbreviations

ATRX: Transcriptional regulator ATRX

BBB: Blood brain barrier

BM: Brain metastasis

cfNA: Cell-free nucleic acids

CDKN2A: Cyclin Dependent Kinase Inhibitor 2A

CNS: Central nervous system

CSF: Cerebrospinal fluid

CTC: Circulating tumour cell

EGFR: Epidermal growth factor receptor

EpCAM: Epithelial cell adhesion molecule

EV: Extracellular vesicles

H3: Histone 3

IDH1/2: Isocitrate dehydrogenase 1/2

MGMT: O<sup>6</sup>-methylguanine DNA methyltransferase

MYD88: Myeloid differentiation primary response 88

miRNA: Micro RNA

mRNA: Messenger RNA

NM: Neoplastic meningitis

NSCLC: Non-small-cell lung cancer

PCNSL: Primary central nervous system lymphoma

SOX2: SRY (Sex Determining Region Y)-Box 2

TERT: Telomerase reverse transcriptase

TP53: Tumour protein p53

WHO: World Health Organization

## Introduction

The possibility of detecting circulating tumour biomarkers in body fluids, including cerebrospinal fluid (CSF), has been exploited in routine clinical practice for tumour diagnosis, treatment and follow up for a long time.[1-3] Protein markers are commonly used, but this approach does not provide information regarding the tumour's molecular profile.[3] Comprehensive molecular profiling of a neoplasm has now been made possible by the analysis of circulating tumour components, and specific applications of this approach, which is usually referred to as "liquid biopsy", are exponentially growing for both research and patient care.[4]

Central nervous system (CNS) tumours are an unmet need: outcome is often dismal, and therapies offer limited efficacy. Tumour molecular profiling is especially relevant in this setting as the CNS is a peculiar anatomic and functional compartment; even biopsy can carry significant risks. Thus liquid biopsy could fulfil its potential in this setting. However, CNS specificities apply to liquid biopsy too. For example, the blood brain barrier (BBB) hinders the circulation of tumour components from the CSF to the blood, thus potentially impairing CNS tumour sampling through the commonly used blood-based approaches.

Over the last few years, a significant amount of data has been accrued regarding liquid biopsy in both primary and secondary CNS tumours. The aim of our review is to thoroughly report the state-of-the-art regarding the opportunities and challenges posed by liquid biopsy in this setting.

## The role of blood brain barrier

Although blood is an ideal, easily accessible source for liquid biopsy sampling and is routinely used to identify many tumour types, data show that its sensitivity can be severely limited when dealing with CNS neoplasms. The main factor hampering the detection of CNS tumour components in blood is the BBB. The latter is a peculiar anatomic and functional structure aimed at regulating the traffic of molecules and cells into and outside of the CNS. Its functions rely on the synergic activity of a wide range of partners including endothelial cells, pericytes and astrocytes.[5] In particular, the widespread presence of tight junctions allows the passage of very small molecules only (less than a few nanometres) and this could explain the overall poor sensitivity observed by blood sampling when looking for components derived from CNS neoplasms.[6]

As will be discussed, however, in specific situations it seems to be possible to detect the molecular signatures of CNS tumours in blood. For example, larger tumour volumes and presence of contrast enhancement were correlated with an increased sensitivity for *IDH1* mutations.[7] This result can be understood if we consider that contrast enhancement is a sign of BBB leakage, which is usually associated with higher grade tumours.

To overcome this limitation, CSF seems the best alternative source for CNS neoplasms considering its direct contact with CNS structures. Present data suggest a higher sensitivity of CSF compared with blood (Figure 1). Nevertheless, CSF sampling is not free of risks especially for patients with CNS neoplasms, although it is usually a feasible and safe approach.

A different approach would be to alter the BBB permeability to allow the transfer of tumour-derived markers into the blood. Until recently, this research topic was mostly aimed at increasing drug penetration into the CNS[8]. However, it is now being investigated as a tool to increase the sensitivity of liquid biopsy for CNS tumours, for example, using focused ultrasound enabled blood-based liquid biopsy in animal models of glioblastoma.[9] The specific mechanisms through which the BBB hampers the transfer of specific tumour components deserve further study.

Last, it should be remembered that the BBB plays an active role in facilitating or preventing brain metastases (BM) from systemic tumours. Primary tumour cells can impair the BBB by releasing specific molecules such as nucleic acids or proteins, thus increasing the risk of BM development. Improved understanding of these mechanisms could pave the way not only to early detection of BM, but could also allow risk prediction or prevention.[10]

Technical considerations: what to analyse and how

The number of potential technical approaches is both extremely high and rapidly evolving.[4] The first choice is about which tumour component or marker would be the most informative in the specific clinical setting (Figure 2). Analysis of circulating cell-free nucleic acids (cfNA), including cfDNA and cfRNA, is the most used approach for the time being, potentially allowing the detection of somatic mutations, insertions, deletions, copy number variations and also enabling assessment of methylation status and of regulatory nucleic acids, such as microRNAs (miRNA) or long non-coding RNAs (lncRNA). These data are what usually matters the most to the pathologist and the clinician to achieve a diagnosis and molecular profiling for clinical management. Conversely, cfNA analysis does not allow reliable assessment of RNA expression, but this limitation can be overcome by analysing circulating tumour cells (CTCs) or extracellular vesicles (EVs) which also allow proteomic studies.[11, 12]

Another important choice regarding genomic studies is whether to use a candidate gene strategy or larger scale approaches like whole exome sequencing. In the diagnostic clinical setting, the first is usually advised: i) it allows reliable assessment of a specific set of genes with known diagnostic and/or predictive value; ii) it enables very low limits of detection; iii) it provides a faster, clinically-suitable, turnaround time with lower costs; iv) bioinformatic support is usually not required; v) germline DNA analysis and its ethical/legal implications can usually be avoided. Conversely, for exploratory/research studies, extensive approaches, also including deep sequencing, can be

considered. Among candidate gene analyses techniques, droplet digital PCR allows an extremely low limit of detection (0.001%) and provides precise DNA copy number quantification,[13] thus allowing monitoring of residual disease.

For CTCs and EVs analysis, different strategies have also to be used for primary and secondary CNS tumours since these tumour components have to be selectively captured and enriched; the proposed approaches will be discussed in the relevant paragraphs. A further informative component worthy of mention are circulating platelets. Analysis of tumour educated platelets by RNA sequencing has been proven to be highly informative in both primary and secondary CNS neoplasms.[14, 15]

#### Liquid biopsy in primary CNS tumours

Molecular profiling is of paramount importance for the characterization of primary CNS tumours. For this reason, the latest WHO classification of these tumours introduced the concept of “integrated diagnosis”: a diagnosis based upon the integration of both morphological and molecular findings. This approach is necessary to distinguish between entities with significant biological and clinical differences, but overlapping histological features.[16]

Even considering just the group of diffuse gliomas, a wide range of molecular alterations is relevant for their proper classification and treatment: *IDH1/2*, *ATRX*, *TP53*, *TERT* promoter and histone H3-coding genes mutations, 1p/19q chromosomal arms codeletion, *EGFR* alterations and *MGMT* promoter methylation.[16, 17] Overall, specific molecular hallmarks are relevant for diagnosis of most primary CNS tumours, while assessment of temporal molecular heterogeneity could help shed a light on mechanisms of resistance to treatments. Thus, liquid biopsy could be extremely useful for the diagnosis and follow up of primary CNS tumours.



## Circulating cell-free nucleic acids analysis

Analysis of the circulating cfDNA shed by tumour cells allows detection of gene alterations, including mutations or fusions, changes in methylation profile, copy number variations, and to quantify tumour burden; moreover, overall cfDNA (i.e. tumour and non-tumour derived) concentration has been suggested to play a prognostic role in primary CNS tumours.[18]

Boisselier *et al.* (2012) evaluated the detection of *IDH1* mutations in plasma cfDNA of patients with histologically-proven glioma with known tumour IDH-status:[7] sensitivity and specificity were 60% and 100%, respectively (Table 1). Sensitivity correlated with tumour volume and contrast enhancement: a higher contrast-enhancing tumour volume was associated with higher sensitivity. More recently, Bettegowda *et al.* (2014) investigated the possibility of detecting a wide range of alterations in a large series of tumour types by analysis of plasma cfDNA by digital PCR.[19] A cohort of 41 primary brain tumours (including gliomas and medulloblastomas) was also included and, unlike most extra-CNS tumours, the detection rates were limited (<50% for medulloblastoma and <10% for gliomas).

The possibility of using CSF as a source for cfDNA analysis in primary CNS tumours was suggested more than 20 years ago by Rhodes *et al.* who identified *EGFR* amplification and a *TP53* mutation in glioblastoma patients (Table 1).[1, 2] De Mattos-Arruda *et al.* (2015) further showed that CSF-derived cfDNA analysis detects mutations of both primary and secondary brain tumours.[20] Regarding the primary neoplasms, 4 glioblastomas and 2 medulloblastomas were investigated; at least one tumour mutation was detected in all cases by CSF cfDNA analysis, while blood-derived liquid biopsies were negative. Tumour burden, assessed by cfDNA quantification, mirrored the neuroimaging findings, thus supporting liquid biopsy as an effective tool to track tumour evolution over time and possibly anticipate recurrence/progression, although specific prospective studies are needed for clinical validation. Pentsova *et al.* (2016) confirmed the usefulness of CSF liquid biopsy,

but observed a low sensitivity (6/9 patients, 66,7%) in a series of primary CNS tumours, mostly including grade III-IV (WHO) diffuse gliomas.[21]

Tumour variables, such as histological type, grade and site, could influence the efficacy of cfDNA analysis. Medulloblastoma, for example, a malignant cerebellar tumour with high cellularity could be expected to more easily shed cfDNA into CSF compared to hemispheric diffuse low-grade gliomas with an infiltrative growth pattern. This hypothesis is supported by Wang *et al.* (2015) who found that tumours closer to CSF spaces have a higher probability of being detected by CSF cfDNA analysis compared to tumours not in contact with the ventricles.[22] This finding was true for different tumour types, including high-grade gliomas, ependymomas and medulloblastomas. Interestingly, Connolly *et al.* (2017) analysed matched CSF and blood samples of three cases of grade II intramedullary ependymomas without identifying any alteration in blood or in CSF[23]. In this regard, other variables such as CSF flow alterations may affect cfDNA circulation and liquid biopsy sensitivity.

A very recent study specifically investigated a large series of diffuse gliomas (n=85), including 46/85 (54%) glioblastomas and 39/85 (46%) lower grade gliomas (grade II-III), using a targeted next generation sequencing-based assay.[24] CSF was collected by lumbar puncture in most cases (82/85, 96%). Overall sensitivity was of 49.4% (42/85): 59% (27/46) in glioblastomas and 38% (15/39) in lower grade gliomas, although a significant association between tumour grade and CSF ctDNA detection was not found. Conversely, the presence of ctDNA was associated with tumour progression, a higher tumour burden, tumour spread in the ventricular system/subarachnoid space and a shorter survival time since CSF collection. This latter finding was found to be independent of tumour burden. Regarding the diagnostic efficacy of this approach, in a subset of 10 lower grade gliomas with available tissue sample, the genetic alterations currently used to define the glioma subtype according to WHO criteria were always consistent between CSF and tissue. Plasma was analysed in 19 cases with positive CSF ctDNA and no mutations were detected in most of them

(16/19, 84%), while in the remaining samples the mutant allele frequencies were very low. An important finding regarding tumour heterogeneity was that, as the time between tissue and CSF sampling increased, a greater genetic divergence was observed, in particular for genes involved in growth factor signalling pathways. These data highlight the importance of longitudinal molecular evaluation in diffuse gliomas and show that liquid biopsy may successfully fulfil this need.

*EGFR* alterations are another relevant molecular hallmark since *EGFR* amplification and/or EGFRvIII variant are present in many glioblastomas[25, 26] and have been targeted in many clinical trials employing a wide range of strategies (i.e. direct inhibition, vaccination, antibody-drug conjugate therapy). Unfortunately, the results have been unsatisfactory so far,[27] but new strategies are currently under investigation.[28, 29] Salkeni *et al.* (2013) were able to detect EGFRvIII in plasma-derived cfDNA of 3/3 patients with EGFRvIII-positive tumours (out of a cohort of 13 glioblastomas), making this marker potentially assessable through liquid biopsy.[30]

Regarding the diagnostic potential of CSF cfDNA analysis, evaluation of a selected panel of molecular alterations in CSF cfDNA allowed the identification of tumour molecular alterations in 17/20 (85%) cases of diffuse gliomas, allowing their classification according to the latest WHO criteria.[31] Also detection of mutations affecting histone H3 genes in a case series of paediatric brain tumours showed high sensitivity and specificity: 87.5% and 100%, respectively.[32] This finding is especially relevant as the H3 K27M mutation characterizes, although not exclusively, a subgroup of highly-malignant midline diffuse gliomas, usually occurring in children. Considering the location of these tumours, which often involve the brainstem, this approach could enable to achieve a specific diagnosis without the potential risks of a surgical biopsy.

Recent data further support the sensitivity of liquid biopsy for brainstem gliomas: Pan *et al.* (2018) evaluated a series of 57 patients and were able to detect at least one tumour-specific mutation in 82.5% (47/57) of cases.[33] Sensitivity reached 96.3% (36/37) when considering only patients with a detectable mutation in the tumour tissue sample, while in 83% (31/37), it was possible to achieve

the same results as tissue analysis by liquid biopsy. CSF and plasma cfDNA were compared in 8 patients producing sensitivities of 100% and 37.5%, respectively. Higher levels of cfDNA, tumour directly abutting the CSF spaces and higher tumour grade were associated with an effective liquid biopsy, while a trend was observed for higher tumour volumes. An important consideration is that in most cases (91.2%, 52/57), CSF was collected intraoperatively, thus results could differ from lumbar puncture sampling. Last, in 3 cases out of 10 with negative tissue analysis, mutations were detected in cfDNA. This result could support the capability of liquid biopsy to capture tumour heterogeneity, but it must be interpreted cautiously since sequencing depths differed significantly between the two assays. Another recent study by Panditharatna *et al.* (2018) focused on midline gliomas and evaluated H3 K27M mutations and typical partner alterations in cfDNA by droplet digital PCR.[34] CSF analysis sensitivity reached 87% (20/23) in H3 K27M-mutant cases (defined through tissue analysis), but, interestingly, plasma sensitivity was very high, ranging from 90% (18/20) in the original series to 80% (16/20) in a prospective cohort collected during a clinical trial. However, mutant allele fractions were lower compared to CSF. The relationship between sampling site and mutations detection was also evaluated: a higher mutant allele frequency was observed if CSF was collected adjacent to the tumour location, but mutant alleles could be potentially detected in all sample sites. Notably, longitudinal comparisons showed a correlation between clinical course, radiological findings and mutant allele frequencies.

A concluding remark regarding the use of liquid biopsy as a diagnostic tool for gliomas is that particular care should be applied when selecting the gene panel to be assessed, since dramatic differences exist between the molecular landscapes of adult and paediatric tumours. Moreover, reliance on single markers should be avoided. For instance, the H3 K27M mutation, which was initially considered as a potential pathognomonic alteration of malignant diffuse brainstem glioma, has been reported in a wider range of tumours.[35]

Liquid biopsy could be useful for diagnosis and follow up of non-glial primary brain tumours as well, although less data is available.[36] Hiemcke-Jiwa *et al.* (2018) demonstrated the possibility of detecting *MYD88* mutations, a common alteration of primary central nervous system lymphomas (PCNSL) (Table 1).[37] This is especially intriguing considered that PCNSL are usually extremely chemo/radiosensitive, thus surgery could be avoided.

Epigenetic alterations are also relevant biomarkers in primary brain tumours. The possibility of assessing *MGMT* promoter methylation in blood-derived cfDNA was first reported by Weaver *et al.* (2006) in a small case series of high-grade diffuse gliomas, achieving a 50% sensitivity.[38] Lavon *et al.* (2010) and further studies confirmed this possibility and also explored the correlations with outcome and longitudinal evolution during disease course.[39-41] cfDNA methylation analysis has also been proposed for diagnostic purposes in gliomas using *CDKN2A* promoter methylation assessment.[42]

Analysis of circulating micro RNAs (miRNAs) is another potential strategy. miRNAs are non-coding short RNA regulatory molecules, which can widely affect transcription and translation. Most available data relate to diffuse glioma, in particular glioblastoma, and a broad range of miRNAs have been investigated, alone or in specific combinations, as possible diagnostic or prognostic biomarkers.[43-52] Both CSF[44, 45, 49, 51] and blood[43, 46-48, 50, 52] have been evaluated as sources of genetic material. The possibility of distinguishing between primary and secondary CNS tumours by miRNA profiling, and of estimating patients' outcome based on their concentrations has been suggested.

Overall, the main limitation of this approach is the heterogeneity of the reported data. Although many different candidate miRNAs have been proposed, their validation in larger cohorts is lacking, hampering translation into clinical practice.

Last, the possibility of exploiting other nucleic acids such as long non-coding RNAs or mitochondrial DNA should also be taken into consideration.[53, 54]

### Circulating tumour cells

Circulating tumour cells (CTCs) analysis is another potential approach for liquid biopsy and allows the assessment of nucleic acids, including mRNAs, which are usually quickly degraded in the circulation, and other tumour cell components such as cytoplasmic proteins. Detection and capture of CTCs is challenging due to their extremely low concentration in blood, thus many different techniques have been investigated.[55] Moreover, primary CNS tumour cells do not express a common membrane marker, thus hampering their detection and collection.

The presence of CTCs from blood in primary CNS tumours was debated due to the extremely low incidence of extra-CNS metastases [56]. However, blood CTCs were successfully detected in a series of 33 glioblastomas using a combination of markers (SOX2, Tubulin beta-3, EGFR, A2B5, and c-MET) with a sensitivity (defined as the rate of patients with at least one positive sample) of 13/33 (39%).[57] Specific probes for molecular hallmarks of glioblastoma, such as telomerase activity have been also investigated. MacArthur *et al.* (2014) observed a sensitivity of 8/11 (72%) in a series of grade III and IV gliomas sampled before radiotherapy.[58] Interestingly, sampling after radiotherapy was successful in only 1/8 (12.5%). Müller *et al.* (2014) were able to detect circulating neoplastic cells in 21% (29/141) of patients with glioblastoma by GFAP immunostaining of mononuclear circulating cells.[59] Moreover, an integrated positive and negative selection approach was also effective in grade II diffuse gliomas reaching a sensitivity of 82% (9/11) in this specific subset of tumours. Using this assay, CTCs count resulted associated with tumour status (i.e. progression *versus* radiation necrosis) in an additional series of 5 cases.[60] The significant variability in terms of

detection rates is likely due to the differences in technical approaches and to the overall low sample sizes. Further larger studies are warranted to confirm the real efficacy of the different analytical methods.

Last, an interesting observation regarding tumour heterogeneity was reported by Sullivan *et al.* (2014): CTCs can show a different gene expression profile compared to the previously resected tumour sample. Specifically, increased mesenchymal differentiation and a reduced neural-like profile was observed. These data suggest that a single approach or a single set of markers may be unable to fully sample tumour heterogeneity.[57]

#### Tumour educated platelets

Analysis of circulating platelets is another intriguing approach to achieve liquid biopsy of CNS tumours.[61, 62] Many biomolecules are transferred between tumours and platelets and this process has been termed platelet “education”.[63, 64]. As initially shown by Nilsson *et al.* (2011), platelets can sequester tumour-shed extracellular vesicles which allow detecting of tumour-derived biomarkers, including proteins and nucleic acids like the EGFRvIII mRNA, with high sensitivity and specificity.[14] Different patterns of mRNA expression were also observed in platelets of patients with glioblastoma compared with healthy controls. The shaping of platelets mRNA splicing by the external environment can thus be exploited for tumour diagnostics and profiling. Indeed, Best *et al.* (2015), showed that analysis of RNA splicing profiles of tumour educated platelets was able to identify patients with tumours and further distinguish those with localized or metastatic disease. In particular, this approach correctly classified 85% (33/39) of patients as patients with a neoplasm in a series of 39 glioblastomas. Regarding the specific diagnostic potential, 78% of glioblastoma patients were correctly classified when compared with controls and five other tumour types in a training series of 175 samples. Similar results were observed in a validation set of 108 cases. Furthermore, in a series of 114 samples (62 patients) with brain lesions, tumour educated platelet analysis allowed

correct identification of cancer patients in 91%, discrimination between primary and secondary tumours in 93% and discerned between the mutational subtypes in 82%.[15] It has also to be remembered that blood components can actively promote glioma growth and malignancy, thus further studies investigating strategies to antagonize this cross-talk are warranted.[65]

#### Extracellular vesicles and other approaches

Extracellular vesicles (EVs) are a heterogeneous range of membrane-bound carriers secreted or shed by both normal and neoplastic cells. Many studies have investigated their physiological role in intercellular communication, which can occur between different tissues or compartments.[66]

Similar to CTC, liquid biopsy by EVs analysis offers some advantages over cfNAs, allowing mRNAs characterization and evaluation of cytoplasmic proteins.[67] In primary brain tumours, this approach has been mainly investigated in malignant gliomas, and both blood and CSF have been explored as potential sources of EVs. As initially shown by Skog *et al.* (2008),[11] EVs shed by glioblastoma contain a wide range of molecules, including nucleic acids and proteins, and can significantly affect distant cells, for instance by increasing proliferation or angiogenesis. Moreover, specific molecular hallmarks of gliomas can be detected in EVs: for example, Skog *et al.* (2008) were able to detect EGFRvIII in EVs of 7/25 (28%) patients with glioblastoma. [11]

More recently, Chen *et al.* (2013) detected *IDH1* mutations in CSF-derived EVs of patients with IDH-mutant gliomas using highly-sensitive techniques; conversely, serum-derived EVs of the same patients were negative.[68] Although based on significantly different approaches and techniques, these results somehow conflict with data reported by Boisselier *et al.* (2012)[7] and indicate the importance of further research evaluating how the interplay between biological and technical variabilities can affect the results. *EGFR* alterations can also be assessed in EVs. [11, 69]



Concerning immune system evasion, a recent study identified PD-L1 in glioblastoma-derived EVs and presence of this ligand was associated with impaired T cell activation.[70]

miRNAs are another cargo of EVs through which they can exert their regulatory functions on distant tissues.[71-73] As for circulating free miRNAs, many candidates have been investigated as potential diagnostic biomarkers and different subsets of EVs have been compared.[74, 75] A panel of three miRNAs (miR-21, miR-222 and miR-124-3p), detected in blood-derived exosomes, was recently shown to be differentially expressed in high grade gliomas compared with low grade gliomas or healthy controls.[76]

Last, circulating proteins and metabolites can be also exploited as biomarkers of neoplastic diseases, including brain tumours. The improved knowledge of tumour molecular profiles and their correlation with cellular metabolism could enable new avenues for diagnosis and monitoring of CNS neoplasms, as recently shown by Ballester *et al.* (2018).[77] This extensive topic, however, is outside the scope of the present review.

#### Liquid biopsy in secondary CNS tumours

Brain metastases are the most frequent intracranial tumours. They develop in 9-17% of patients with solid tumours (usually from lung and breast cancers and melanoma) and their incidence is increasing.[78, 79] Quality of life and outcome are severely affected by BMs[80], but encouraging results have been recently achieved thanks to targeted therapies and immunotherapeutic drugs.[81]

BMs from solid tumours often show a divergent molecular profile compared to the primary lesion. Spatial and temporal heterogeneity is an intrinsic characteristic of malignant tumours and this hallmark is strictly intertwined with the continuous evolutionary pressure which selects the neoplastic cells that best fit in a specific anatomic compartment or timepoint during the course of disease, also because of external variables, such as ongoing treatments. Brastianos *et al.* (2015), showed that more than 50% of BMs harbour private molecular alterations (compared to the primary

tumour or extra-CNS lesions) which are potentially actionable.[82] These data suggest that, to offer the optimal treatment to these patients, BM sampling is necessary to fully recapitulate tumour heterogeneity; since surgical resection of BMs is not indicated in most cases, liquid biopsy could help overcome this limitation.

Neoplastic meningitis (NM) is a specific type of secondary CNS involvement which is characterized by the spread of neoplastic cells across the leptomeninges and their circulation through the CSF. Definitive diagnosis is based upon cytological demonstration of malignant cells in the CSF, but sensitivity is limited and thus diagnosis usually relies on supporting clinical and neuroimaging findings.[83] Liquid biopsy could be a valuable diagnostic/profiling tool also in this setting.

#### Circulating cell-free nucleic acids analysis

The possibility of detecting tumour-derived cfDNA in patients with secondary CNS involvement has long been reported,[1] but only during the last few years this approach has received specific consideration as a potential tool for the routine care of patients.[20, 21, 36, 84-96] Most data have been obtained in NM from non-small-cell lung carcinoma (NSCLC), but studies on breast carcinoma[1], melanoma[89, 97] and secondary haematological malignancies[96] have also been reported (Table 2).

cfDNA analysis proved a quite high overall sensitivity as a diagnostic tool for BMs or NM from solid tumours, ranging from 40% to 100%, but sample sizes of the available studies are still limited. Nevertheless, if we compare this tool with traditional cytology (the gold-standard for NM diagnosis at present),[98] cfDNA appears to be superior (Table 2). Another intrinsic advantage of liquid biopsy compared with cytology is that it allows tumour molecular profiling, while cytology can only provide a diagnostic confirmation.

With respect to the variables potentially affecting liquid biopsy yield, the type of CNS involvement should be considered. BMs may have limited contact with CSF, especially when dealing with a single or few lesions in a supratentorial intraparenchymal location. Conversely, tumour cells circulation in the CSF is a hallmark of NM, and thus a higher sensitivity can be expected in the latter condition. The results of Yang *et al.* (2014)[87] and Pentsova *et al.* (2016)[21] support these considerations, although De Mattos-Arruda *et al.* (2015)[20] and Pan *et al.* (2015)[36] were able to identify tumour cfDNA in all BMs cases (Table 2).

As for primary CNS tumours, the most pressing question is choice of the source for cfNAs (or of other analytes). Blood or other easily accessible body fluids (like saliva or urine) would obviously be preferable, but they may not be representative of CNS-restricted lesions (Table 2). Some important considerations stem from the available data: i) a negative result may be obtained from blood-based liquid biopsy even when active disease is present within the CNS and this is an important concern when dealing with minimal residual disease monitoring; ii) even if we find alterations in blood-based liquid biopsy, these could be unrepresentative of the CNS disease.

Within the possible practical applications of liquid biopsy in this setting, CSF testing in patients with CNS involvement by non-small cell lung carcinoma (NSCLC) seems one of the most promising, mirroring the already widespread practice of blood-based testing to promptly detect emerging resistance-associated mutations in extra-CNS NSCLC. As recently shown by Nanjo *et al.*, the specific detection of the *EGFR* T790M mutation in CSF cfDNA was associated with clinical efficacy of treatment with osimertinib, a third-generation *EGFR* tyrosine kinase inhibitor developed to overcome T790M-induced resistance.[99] Conversely, in patients with positive CSF liquid biopsy, but without *EGFR* T790M detection, no response to treatment was observed.

Last, it is worth pointing out that the significant differences observed between studies may depend on the wide range of techniques that have been tested so far. This variability should be addressed in future studies.

## Circulating tumour cells and other approaches

The presence of common membrane markers, like EpCAM, significantly helps CTCs collection in secondary CNS tumours, but alternative strategies are also possible.[100] As a diagnostic tool, CSF CTCs proved to be superior to traditional cytology[101]; moreover, the possibility of using this approach also for molecular profiling has been demonstrated. Magbanua *et al.* (2013 and 2014)[85, 86] and Jiang *et al.* (2017)[91] were able to assess copy number and mutational profile of CSF CTCs in metastatic breast and lung cancer, respectively. Similar to cfNAs analysis, CTCs collection by blood-derived liquid biopsy was not possible or showed a low yield in cases with CNS restricted disease.

Best *et al.* (2015) showed that analysis of tumour educated platelets in patients with BMs allows identification of primary tumour type in 70% of samples.[15] Moreover, blood components indices (like white blood cells or platelets values and ratios) may harbour prognostic significance in patients with BMs[102]: this finding can be expected considering the role played by the systemic immune system in BM development.[103]

Finally, EVs analysis in secondary CNS tumours is especially intriguing not only as a diagnostic/profiling tool, but also as a prognostic assay. Multiple studies suggested that tumour-derived EVs may play an active role in creating a pro-metastatic niche in distant tissues[104], including the CNS.[10, 105] Thus, EVs characterization could allow to estimate the metastatic potential of a tumour and, possibly, to devise therapeutic strategies capable of hampering this process even before BMs are established.

## Future perspectives

The recent technological advancements paired with the growing importance of molecular profiling in CNS neoplasms can explain the far-reaching results achieved in just the last few years regarding liquid biopsy use in CNS tumours.[106, 107]

Accepted Article

The ever-increasing reliance on molecular traits for proper classification of primary CNS tumours will favour the clinical adoption of liquid biopsy as a routine diagnostic tool in selected cases, for example when surgical resection is not possible. Conversely, resection and histological examination will probably remain the cornerstone diagnostic approach if feasible, considered its therapeutic relevance and the risk of diagnostic pitfalls due to overlapping molecular features even between significantly different tumour entities [for example, a pilocytic astrocytoma (grade I) can rarely harbour the H3 K27M mutation which is characteristic of diffuse midline gliomas (grade IV)]. Moreover, a tissue sample is needed for whole genome or proteomic studies which are now increasingly warranted in translational/clinical research protocols.

Liquid biopsy could instead represent a game-changing development for disease follow up considering that repeated surgical sampling is not feasible in CNS tumours. Its first implementation, in this setting, could probably be as a companion tool for disease monitoring and tumour burden quantification: liquid biopsy data could help resolve conflicting clinical and radiological findings, for instance when dealing with a suspected pseudo-progression. The following step could be the longitudinal assessment of tumour heterogeneity: for example, patients with IDH-mutant diffuse gliomas can have very long disease courses characterized by a progressive increase of tumour malignancy. Liquid biopsy could enable prompt detection and molecular characterization of disease progression, allowing optimisation of clinical management. Nevertheless, the relevance of liquid biopsy in this setting will ultimately depend on the identification of significant, actionable, prognostic and/or predictive markers.

The same considerations apply to secondary neoplasms, but special caution should be applied when dealing with synchronous intra- and extra-CNS disease progression, since blood-derived results could be uninformative of the CNS disease. Nevertheless, the outcome of patients with BMs has dramatically changed in the last few years thanks to the newly available targeted treatments,[108] thus liquid biopsy is expected to become a mandatory assessment in an increasing number of

tumour types and disease settings for optimizing treatment decisions and prompt detection of resistance-associated mutations.

## Conclusion

In the coming years, liquid biopsy will probably become a common tool for the diagnosis and follow up of both primary and secondary CNS tumours. For research purposes, liquid biopsy will be a cornerstone to determine longitudinal changes in the molecular profile of tumours, thus improving our knowledge of tumour resistance/progression mechanisms. Further studies, evaluating larger, prospective series, are needed to evaluate technical variabilities, validate its use in specific disease settings and, most importantly, to assess which is the ultimate clinical benefit for patients with CNS tumours.

## Acknowledgements

We apologize to colleagues whose work could not be cited because of space limitations.

All authors contributed to the conception and design of the manuscript, collection, review and interpretation of data, manuscript drafting, writing and final approval.

## Funding

This work was supported by the Rete Oncologica del Piemonte e della Valle d'Aosta (no specific grant number applicable) and received funding specifically dedicated to the Department of Medical Sciences from Italian Ministry for Education, University and Research (Ministero dell'Istruzione, dell'Università e della Ricerca - MIUR) under the programme "Dipartimenti di Eccellenza 2018 – 2022". Project n° D15D18000410001. G.S. was supported by a 3-year FIRC-AIRC fellowship and by "Roche per la ricerca" - Grant 2017. A.B. was supported by IMI contract n. 115749 CANCER-ID.

## Disclosure

The authors declare they have no conflicts of interest.

Figure legend

**Fig 1 CSF sampling for liquid biopsy.** CSF, usually sampled by lumbar puncture, allows gathering of multiple tumour components which can be submitted to a wide range of molecular tests.

**Fig 2 Liquid biopsy possible analytes and relevant assays.** Different tumour components can be collected in liquid biopsy sources (blood or CSF) allowing a wide range of analyses.

## References

- 1 Rhodes CH, Honsinger C, Sorenson GD. Detection of tumor-derived DNA in cerebrospinal fluid. *J Neuropathol Exp Neurol* 1994; 53: 364-8
- 2 Rhodes CH, Honsinger C, Sorenson GD. PCR-detection of tumor-derived p53 DNA in cerebrospinal fluid. *Am J Clin Pathol* 1995; 103: 404-8
- 3 Henry NL, Hayes DF. Cancer biomarkers. *Mol Oncol* 2012; 6: 140-6
- 4 Siravegna G, Marsoni S, Siena S, Bardelli A. Integrating liquid biopsies into the management of cancer. *Nat Rev Clin Oncol* 2017; 14: 531-48
- 5 Sharif Y, Jumah F, Coplan L, Krosser A, Sharif K, Tubbs RS. Blood brain barrier: A review of its anatomy and physiology in health and disease. *Clin Anat* 2018; 31: 812-23
- 6 Van Itallie CM, Anderson JM. Claudins and epithelial paracellular transport. *Annu Rev Physiol* 2006; 68: 403-29
- 7 Boisselier B, Gallego Perez-Larraya J, Rossetto M, Labussiere M, Ciccarino P, Marie Y, Delattre JY, Sanson M. Detection of IDH1 mutation in the plasma of patients with glioma. *Neurology* 2012; 79: 1693-8
- 8 Banks WA. From blood-brain barrier to blood-brain interface: new opportunities for CNS drug delivery. *Nat Rev Drug Discov* 2016; 15: 275-92
- 9 Zhu L, Cheng G, Ye D, Nazeri A, Yue Y, Liu W, Wang X, Dunn GP, Petti AA, Leuthardt EC, Chen H. Focused Ultrasound-enabled Brain Tumor Liquid Biopsy. *Sci Rep* 2018; 8: 6553
- 10 Tominaga N, Kosaka N, Ono M, Katsuda T, Yoshioka Y, Tamura K, Lotvall J, Nakagama H, Ochiya T. Brain metastatic cancer cells release microRNA-181c-containing extracellular vesicles capable of destructing blood-brain barrier. *Nat Commun* 2015; 6: 6716
- 11 Skog J, Wurdinger T, van Rijn S, Meijer DH, Gainche L, Sena-Esteves M, Curry WT, Jr., Carter BS, Krichevsky AM, Breakefield XO. Glioblastoma microvesicles transport RNA and proteins that promote tumour growth and provide diagnostic biomarkers. *Nat Cell Biol* 2008; 10: 1470-6
- 12 Yu M, Bardia A, Wittner BS, Stott SL, Smas ME, Ting DT, Isakoff SJ, Ciciliano JC, Wells MN, Shah AM, Concanon KF, Donaldson MC, Sequist LV, Brachtel E, Sgroi D, Baselga J, Ramaswamy S, Toner M, Haber DA, Maheswaran S. Circulating breast tumor cells exhibit dynamic changes in epithelial and mesenchymal composition. *Science* 2013; 339: 580-4
- 13 Hindson BJ, Ness KD, Masquelier DA, Belgrader P, Heredia NJ, Makarewicz AJ, Bright IJ, Lucero MY, Hiddessen AL, Legler TC, Kitano TK, Hodel MR, Petersen JF, Wyatt PW, Steenblock ER, Shah PH, Bousse LJ, Troup CB, Mellen JC, Wittmann DK, Erndt NG, Cauley TH, Koehler RT, So AP, Dube S, Rose KA, Montesclaros L, Wang S, Stumbo DP, Hodges SP, Romine S, Milanovich FP, White HE, Regan JF, Karlin-Neumann GA, Hindson CM, Saxonov S, Colston BW. High-throughput droplet digital PCR system for absolute quantitation of DNA copy number. *Anal Chem* 2011; 83: 8604-10
- 14 Nilsson RJ, Balaj L, Hulleman E, van Rijn S, Pegtel DM, Walraven M, Widmark A, Gerritsen WR, Verheul HM, Vandertop WP, Noske DP, Skog J, Wurdinger T. Blood platelets contain tumor-derived RNA biomarkers. *Blood* 2011; 118: 3680-3
- 15 Best MG, Sol N, Kooi I, Tannous J, Westerman BA, Rustenburg F, Schellen P, Verschueren H, Post E, Koster J, Ylstra B, Ameziane N, Dorsman J, Smit EF, Verheul HM, Noske DP, Reijneveld JC, Nilsson RJA, Tannous BA, Wesseling P, Wurdinger T. RNA-Seq of Tumor-Educated Platelets Enables Blood-Based Pan-Cancer, Multiclass, and Molecular Pathway Cancer Diagnostics. *Cancer Cell* 2015; 28: 666-76
- 16 Louis DN, Ohgaki H, Wiestler OD, Cavenee WK, Ellison DW, Figarella-Branger D, Perry A, Reifenberger G, von Deimling A, International Agency for Research on Cancer. WHO classification of tumours of the central nervous system. Revised 4th edition. ed. Lyon: International Agency For Research On Cancer. 2016
- 17 NCCN Clinical Practice Guidelines in Oncology - Central Nervous Systems Cancers (Version 1.2017). 18/08/2017 [cited 2018 June 28]; Available from: [https://www.nccn.org/professionals/physician\\_gls/pdf/cns.pdf](https://www.nccn.org/professionals/physician_gls/pdf/cns.pdf)



- 18 Shi W, Lv C, Qi J, Zhao W, Wu X, Jing R, Wu X, Ju S, Chen J. Prognostic value of free DNA quantification in serum and cerebrospinal fluid in glioma patients. *J Mol Neurosci* 2012; 46: 470-5
- 19 Bettgowda C, Sausen M, Leary RJ, Kinde I, Wang Y, Agrawal N, Bartlett BR, Wang H, Lubner B, Alani RM, Antonarakis ES, Azad NS, Bardelli A, Brem H, Cameron JL, Lee CC, Fecher LA, Gallia GL, Gibbs P, Le D, Giuntoli RL, Goggins M, Hogarty MD, Holdhoff M, Hong SM, Jiao Y, Juhl HH, Kim JJ, Siravegna G, Laheru DA, Lauricella C, Lim M, Lipson EJ, Marie SK, Netto GJ, Oliner KS, Olivi A, Olsson L, Riggins GJ, Sartore-Bianchi A, Schmidt K, Shih I M, Oba-Shinjo SM, Siena S, Theodorescu D, Tie J, Harkins TT, Veronese S, Wang TL, Weingart JD, Wolfgang CL, Wood LD, Xing D, Hruban RH, Wu J, Allen PJ, Schmidt CM, Choti MA, Velculescu VE, Kinzler KW, Vogelstein B, Papadopoulos N, Diaz LA, Jr. Detection of circulating tumor DNA in early- and late-stage human malignancies. *Sci Transl Med* 2014; 6: 224ra24
- 20 De Mattos-Arruda L, Mayor R, Ng CK, Weigelt B, Martinez-Ricarte F, Torrejon D, Oliveira M, Arias A, Raventos C, Tang J, Guerini-Rocco E, Martinez-Saez E, Lois S, Marin O, de la Cruz X, Piscuoglio S, Towers R, Vivancos A, Peg V, Ramon y Cajal S, Carles J, Rodon J, Gonzalez-Cao M, Tabernero J, Felip E, Sahuquillo J, Berger MF, Cortes J, Reis-Filho JS, Seoane J. Cerebrospinal fluid-derived circulating tumour DNA better represents the genomic alterations of brain tumours than plasma. *Nat Commun* 2015; 6: 8839
- 21 Pentsova EI, Shah RH, Tang J, Boire A, You D, Briggs S, Omuro A, Lin X, Fleisher M, Grommes C, Panageas KS, Meng F, Selcuklu SD, Ogilvie S, Distefano N, Shagabayeva L, Rosenblum M, DeAngelis LM, Viale A, Mellinghoff IK, Berger MF. Evaluating Cancer of the Central Nervous System Through Next-Generation Sequencing of Cerebrospinal Fluid. *J Clin Oncol* 2016; 34: 2404-15
- 22 Wang Y, Springer S, Zhang M, McMahon KW, Kinde I, Dobbyn L, Ptak J, Brem H, Chaichana K, Gallia GL, Gokaslan ZL, Groves ML, Jallo GI, Lim M, Olivi A, Quinones-Hinojosa A, Rigamonti D, Riggins GJ, Sciubba DM, Weingart JD, Wolinsky JP, Ye X, Oba-Shinjo SM, Marie SK, Holdhoff M, Agrawal N, Diaz LA, Jr., Papadopoulos N, Kinzler KW, Vogelstein B, Bettgowda C. Detection of tumor-derived DNA in cerebrospinal fluid of patients with primary tumors of the brain and spinal cord. *Proc Natl Acad Sci U S A* 2015; 112: 9704-9
- 23 Connolly ID, Li Y, Pan W, Johnson E, You L, Vogel H, Ratliff J, Hayden Gephart M. A pilot study on the use of cerebrospinal fluid cell-free DNA in intramedullary spinal ependymoma. *J Neurooncol* 2017; 135: 29-36
- 24 Miller AM, Shah RH, Pentsova EI, Pourmaleki M, Briggs S, Distefano N, Zheng Y, Skakodub A, Mehta SA, Campos C, Hsieh WY, Selcuklu SD, Ling L, Meng F, Jing X, Samoila A, Bale TA, Tsui DWY, Grommes C, Viale A, Souweidane MM, Tabar V, Brennan CW, Reiner AS, Rosenblum M, Panageas KS, DeAngelis LM, Young RJ, Berger MF, Mellinghoff IK. Tracking tumour evolution in glioma through liquid biopsies of cerebrospinal fluid. *Nature* 2019; 565: 654-8
- 25 An Z, Aksoy O, Zheng T, Fan QW, Weiss WA. Epidermal growth factor receptor and EGFRvIII in glioblastoma: signaling pathways and targeted therapies. *Oncogene* 2018; 37: 1561-75
- 26 Koga T, Li B, Figueroa JM, Ren B, Chen CC, Carter BS, Furnari FB. Mapping of genomic EGFRvIII deletions in glioblastoma: insight into rearrangement mechanisms and biomarker development. *Neuro Oncol* 2018;
- 27 Weller M, Butowski N, Tran DD, Recht LD, Lim M, Hirte H, Ashby L, Mechtler L, Goldlust SA, Iwamoto F, Drappatz J, O'Rourke DM, Wong M, Hamilton MG, Finocchiaro G, Perry J, Wick W, Green J, He Y, Turner CD, Yellin MJ, Keler T, Davis TA, Stupp R, Sampson JH, investigators Alt. Rindopepimut with temozolomide for patients with newly diagnosed, EGFRvIII-expressing glioblastoma (ACT IV): a randomised, double-blind, international phase 3 trial. *Lancet Oncol* 2017; 18: 1373-85
- 28 van den Bent M, Gan HK, Lassman AB, Kumthekar P, Merrell R, Butowski N, Lwin Z, Mikkelsen T, Nabors LB, Papadopoulos KP, Penas-Prado M, Simes J, Wheeler H, Walbert T, Scott AM, Gomez E, Lee HJ, Roberts-Rapp L, Xiong H, Bain E, Ansell PJ, Holen KD, Maag D, Reardon DA. Efficacy of depatuxizumab mafodotin (ABT-414) monotherapy in patients with EGFR-amplified, recurrent glioblastoma: results from a multi-center, international study. *Cancer Chemother Pharmacol* 2017; 80: 1209-17

- 29 Reardon DA, Lassman AB, van den Bent M, Kumthekar P, Merrell R, Scott AM, Fichtel L, Sulman EP, Gomez E, Fischer J, Lee HJ, Munasinghe W, Xiong H, Mandich H, Roberts-Rapp L, Ansell P, Holen KD, Gan HK. Efficacy and safety results of ABT-414 in combination with radiation and temozolomide in newly diagnosed glioblastoma. *Neuro Oncol* 2017; 19: 965-75
- 30 Salkeni MA, Zarzour A, Ansay TY, McPherson CM, Warnick RE, Rixe O, Bahassi el M. Detection of EGFRvIII mutant DNA in the peripheral blood of brain tumor patients. *J Neurooncol* 2013; 115: 27-35
- 31 Martinez-Ricarte F, Mayor R, Martinez-Saez E, Rubio-Perez C, Pineda E, Cordero E, Cicuendez M, Poca MA, Lopez-Bigas N, Ramon YCS, Vieito M, Carles J, Tabernero J, Vivancos A, Gallego S, Graus F, Sahuquillo J, Seoane J. Molecular Diagnosis of Diffuse Gliomas through Sequencing of Cell-Free Circulating Tumor DNA from Cerebrospinal Fluid. *Clin Cancer Res* 2018; 24: 2812-9
- 32 Huang TY, Piunti A, Lulla RR, Qi J, Horbinski CM, Tomita T, James CD, Shilatfard A, Saratsis AM. Detection of Histone H3 mutations in cerebrospinal fluid-derived tumor DNA from children with diffuse midline glioma. *Acta Neuropathol Commun* 2017; 5: 28
- 33 Pan C, Diplas BH, Chen X, Wu Y, Xiao X, Jiang L, Geng Y, Xu C, Sun Y, Zhang P, Wu W, Wang Y, Wu Z, Zhang J, Jiao Y, Yan H, Zhang L. Molecular profiling of tumors of the brainstem by sequencing of CSF-derived circulating tumor DNA. *Acta Neuropathol* 2018:
- 34 Panditharatna E, Kilburn LB, Aboian MS, Kambhampati M, Gordish-Dressman H, Magge SN, Gupta N, Myseros JS, Hwang EI, Kline C, Crawford JR, Warren KE, Cha S, Liang WS, Berens ME, Packer RJ, Resnick AC, Prados M, Mueller S, Nazarian J. Clinically Relevant and Minimally Invasive Tumor Surveillance of Pediatric Diffuse Midline Gliomas Using Patient-Derived Liquid Biopsy. *Clin Cancer Res* 2018; 24: 5850-9
- 35 Louis DN, Giannini C, Capper D, Paulus W, Figarella-Branger D, Lopes MB, Batchelor TT, Cairncross JG, van den Bent M, Wick W, Wesseling P. cIMPACT-NOW update 2: diagnostic clarifications for diffuse midline glioma, H3 K27M-mutant and diffuse astrocytoma/anaplastic astrocytoma, IDH-mutant. *Acta Neuropathol* 2018; 135: 639-42
- 36 Pan W, Gu W, Nagpal S, Gephart MH, Quake SR. Brain tumor mutations detected in cerebral spinal fluid. *Clin Chem* 2015; 61: 514-22
- 37 Hiemcke-Jiwa LS, Minnema MC, Radersma-van Loon JH, Jiwa NM, de Boer M, Leguit RJ, de Weger RA, Huibers MMH. The use of droplet digital PCR in liquid biopsies: A highly sensitive technique for MYD88 p.(L265P) detection in cerebrospinal fluid. *Hematol Oncol* 2018; 36: 429-35
- 38 Weaver KD, Grossman SA, Herman JG. Methylated tumor-specific DNA as a plasma biomarker in patients with glioma. *Cancer Invest* 2006; 24: 35-40
- 39 Lavon I, Refael M, Zelikovitch B, Shalom E, Siegal T. Serum DNA can define tumor-specific genetic and epigenetic markers in gliomas of various grades. *Neuro Oncol* 2010; 12: 173-80
- 40 Majchrzak-Celinska A, Paluszczak J, Kleszcz R, Magiera M, Barciszewska AM, Nowak S, Baer-Dubowska W. Detection of MGMT, RASSF1A, p15INK4B, and p14ARF promoter methylation in circulating tumor-derived DNA of central nervous system cancer patients. *J Appl Genet* 2013; 54: 335-44
- 41 Fiano V, Trevisan M, Trevisan E, Senetta R, Castiglione A, Sacerdote C, Gillio-Tos A, De Marco L, Grasso C, Magistrello M, Tondat F, Ruda R, Cassoni P, Soffietti R, Merletti F. MGMT promoter methylation in plasma of glioma patients receiving temozolomide. *J Neurooncol* 2014; 117: 347-57
- 42 Wakabayashi T, Natsume A, Hatano H, Fujii M, Shimato S, Ito M, Ohno M, Ito S, Ogura M, Yoshida J. p16 promoter methylation in the serum as a basis for the molecular diagnosis of gliomas. *Neurosurgery* 2009; 64: 455-61; discussion 61-2
- 43 Roth P, Wischhusen J, Happold C, Chandran PA, Hofer S, Eisele G, Weller M, Keller A. A specific miRNA signature in the peripheral blood of glioblastoma patients. *J Neurochem* 2011; 118: 449-57
- 44 Teplyuk NM, Mollenhauer B, Gabriely G, Giese A, Kim E, Smolsky M, Kim RY, Saria MG, Pastorino S, Kesari S, Krichevsky AM. MicroRNAs in cerebrospinal fluid identify glioblastoma and metastatic brain cancers and reflect disease activity. *Neuro Oncol* 2012; 14: 689-700

- 45 Baraniskin A, Kuhnhen J, Schlegel U, Maghnouj A, Zollner H, Schmiegell W, Hahn S, Schroers R. Identification of microRNAs in the cerebrospinal fluid as biomarker for the diagnosis of glioma. *Neuro Oncol* 2012; 14: 29-33
- 46 Yang C, Wang C, Chen X, Chen S, Zhang Y, Zhi F, Wang J, Li L, Zhou X, Li N, Pan H, Zhang J, Zen K, Zhang CY, Zhang C. Identification of seven serum microRNAs from a genome-wide serum microRNA expression profile as potential noninvasive biomarkers for malignant astrocytomas. *Int J Cancer* 2013; 132: 116-27
- 47 Shao N, Wang L, Xue L, Wang R, Lan Q. Plasma miR-454-3p as a potential prognostic indicator in human glioma. *Neurol Sci* 2015; 36: 309-13
- 48 Lai NS, Wu DG, Fang XG, Lin YC, Chen SS, Li ZB, Xu SS. Serum microRNA-210 as a potential noninvasive biomarker for the diagnosis and prognosis of glioma. *Br J Cancer* 2015; 112: 1241-6
- 49 Drusco A, Bottoni A, Lagana A, Acunzo M, Fassan M, Cascione L, Antenucci A, Kumchala P, Vicentini C, Gardiman MP, Alder H, Carosi MA, Ammirati M, Gherardi S, Luscri M, Carapella C, Zanesi N, Croce CM. A differentially expressed set of microRNAs in cerebro-spinal fluid (CSF) can diagnose CNS malignancies. *Oncotarget* 2015; 6: 20829-39
- 50 Yue X, Lan F, Hu M, Pan Q, Wang Q, Wang J. Downregulation of serum microRNA-205 as a potential diagnostic and prognostic biomarker for human glioma. *J Neurosurg* 2016; 124: 122-8
- 51 Akers JC, Hua W, Li H, Ramakrishnan V, Yang Z, Quan K, Zhu W, Li J, Figueroa J, Hirshman BR, Miller B, Piccioni D, Ringel F, Komotar R, Messer K, Galasko DR, Hochberg F, Mao Y, Carter BS, Chen CC. A cerebrospinal fluid microRNA signature as biomarker for glioblastoma. *Oncotarget* 2017; 8: 68769-79
- 52 Lopez-Aguilar JE, Velazquez-Flores MA, Simon-Martinez LA, Avila-Miranda R, Rodriguez-Flrido MA, Ruiz-Esparza Garrido R. Circulating microRNAs as Biomarkers for Pediatric Astrocytomas. *Arch Med Res* 2017; 48: 323-32
- 53 Li R, Qian J, Wang YY, Zhang JX, You YP. Long noncoding RNA profiles reveal three molecular subtypes in glioma. *CNS Neurosci Ther* 2014; 20: 339-43
- 54 Wong LJ, Lueth M, Li XN, Lau CC, Vogel H. Detection of mitochondrial DNA mutations in the tumor and cerebrospinal fluid of medulloblastoma patients. *Cancer Res* 2003; 63: 3866-71
- 55 Rawal S, Yang YP, Cote R, Agarwal A. Identification and Quantitation of Circulating Tumor Cells. *Annu Rev Anal Chem (Palo Alto Calif)* 2017; 10: 321-43
- 56 Bohm C, Wassmann H, Paulus W. No evidence of tumour cells in blood of patients with glioma. *Mol Pathol* 2003; 56: 187-9
- 57 Sullivan JP, Nahed BV, Madden MW, Oliveira SM, Springer S, Bhare D, Chi AS, Wakimoto H, Rothenberg SM, Sequist LV, Kapur R, Shah K, Iafrate AJ, Curry WT, Loeffler JS, Batchelor TT, Louis DN, Toner M, Maheswaran S, Haber DA. Brain tumor cells in circulation are enriched for mesenchymal gene expression. *Cancer Discov* 2014; 4: 1299-309
- 58 MacArthur KM, Kao GD, Chandrasekaran S, Alonso-Basanta M, Chapman C, Lustig RA, Wileyto EP, Hahn SM, Dorsey JF. Detection of brain tumor cells in the peripheral blood by a telomerase promoter-based assay. *Cancer Res* 2014; 74: 2152-9
- 59 Muller C, Holtschmidt J, Auer M, Heitzer E, Lamszus K, Schulte A, Matschke J, Langer-Freitag S, Gasch C, Stoupiec M, Mauermann O, Peine S, Glatzel M, Speicher MR, Geigl JB, Westphal M, Pantel K, Riethdorf S. Hematogenous dissemination of glioblastoma multiforme. *Sci Transl Med* 2014; 6: 247ra101
- 60 Gao F, Cui Y, Jiang H, Sui D, Wang Y, Jiang Z, Zhao J, Lin S. Circulating tumor cell is a common property of brain glioma and promotes the monitoring system. *Oncotarget* 2016; 7: 71330-40
- 61 Best MG, Wesseling P, Wurdinger T. Tumor-Educated Platelets as a Noninvasive Biomarker Source for Cancer Detection and Progression Monitoring. *Cancer Res* 2018; 78: 3407-12
- 62 Sol N, Wurdinger T. Platelet RNA signatures for the detection of cancer. *Cancer Metastasis Rev* 2017; 36: 263-72

- 63 Klement GL, Yip TT, Cassiola F, Kikuchi L, Cervi D, Podust V, Italiano JE, Wheatley E, Abou-Slaybi A, Bender E, Almog N, Kieran MW, Folkman J. Platelets actively sequester angiogenesis regulators. *Blood* 2009; 113: 2835-42
- 64 Kuznetsov HS, Marsh T, Markens BA, Castano Z, Greene-Colozzi A, Hay SA, Brown VE, Richardson AL, Signoretti S, Battinelli EM, McAllister SS. Identification of luminal breast cancers that establish a tumor-supportive macroenvironment defined by proangiogenic platelets and bone marrow-derived cells. *Cancer Discov* 2012; 2: 1150-65
- 65 Wurdinger T, Deumelandt K, van der Vliet HJ, Wesseling P, de Gruijl TD. Mechanisms of intimate and long-distance cross-talk between glioma and myeloid cells: how to break a vicious cycle. *Biochim Biophys Acta* 2014; 1846: 560-75
- 66 van Niel G, D'Angelo G, Raposo G. Shedding light on the cell biology of extracellular vesicles. *Nat Rev Mol Cell Biol* 2018; 19: 213-28
- 67 Shao H, Chung J, Lee K, Balaj L, Min C, Carter BS, Hochberg FH, Breakefield XO, Lee H, Weissleder R. Chip-based analysis of exosomal mRNA mediating drug resistance in glioblastoma. *Nat Commun* 2015; 6: 6999
- 68 Chen WW, Balaj L, Liao LM, Samuels ML, Kotsopoulos SK, Maguire CA, Loguidice L, Soto H, Garrett M, Zhu LD, Sivaraman S, Chen C, Wong ET, Carter BS, Hochberg FH, Breakefield XO, Skog J. BEAMing and Droplet Digital PCR Analysis of Mutant IDH1 mRNA in Glioma Patient Serum and Cerebrospinal Fluid Extracellular Vesicles. *Mol Ther Nucleic Acids* 2013; 2: e109
- 69 Figueroa JM, Skog J, Akers J, Li H, Komotar R, Jensen R, Ringel F, Yang I, Kalkanis S, Thompson R, LoGuidice L, Berghoff E, Parsa A, Liao L, Curry W, Cahill D, Bettegowda C, Lang FF, Chiocca EA, Henson J, Kim R, Breakefield X, Chen C, Messer K, Hochberg F, Carter BS. Detection of wild-type EGFR amplification and EGFRvIII mutation in CSF-derived extracellular vesicles of glioblastoma patients. *Neuro Oncol* 2017; 19: 1494-502
- 70 Ricklefs FL, Alayo Q, Krenzlin H, Mahmoud AB, Speranza MC, Nakashima H, Hayes JL, Lee K, Balaj L, Passaro C, Roj AK, Krasemann S, Carter BS, Chen CC, Steed T, Treiber J, Rodig S, Yang K, Nakano I, Lee H, Weissleder R, Breakefield XO, Godlewski J, Westphal M, Lamszus K, Freeman GJ, Bronisz A, Lawler SE, Chiocca EA. Immune evasion mediated by PD-L1 on glioblastoma-derived extracellular vesicles. *Sci Adv* 2018; 4: eaar2766
- 71 Figueroa J, Phillips LM, Shahar T, Hossain A, Gumin J, Kim H, Bean AJ, Calin GA, Fueyo J, Walters ET, Kalluri R, Verhaak RG, Lang FF. Exosomes from Glioma-Associated Mesenchymal Stem Cells Increase the Tumorigenicity of Glioma Stem-like Cells via Transfer of miR-1587. *Cancer Res* 2017; 77: 5808-19
- 72 Sun X, Ma X, Wang J, Zhao Y, Wang Y, Bihl JC, Chen Y, Jiang C. Glioma stem cells-derived exosomes promote the angiogenic ability of endothelial cells through miR-21/VEGF signal. *Oncotarget* 2017; 8: 36137-48
- 73 Yang JK, Yang JP, Tong J, Jing SY, Fan B, Wang F, Sun GZ, Jiao BH. Exosomal miR-221 targets DNMT3 to induce tumor progression and temozolomide resistance in glioma. *J Neurooncol* 2017; 131: 255-65
- 74 Akers JC, Ramakrishnan V, Kim R, Skog J, Nakano I, Pingle S, Kalinina J, Hua W, Kesari S, Mao Y, Breakefield XO, Hochberg FH, Van Meir EG, Carter BS, Chen CC. MiR-21 in the extracellular vesicles (EVs) of cerebrospinal fluid (CSF): a platform for glioblastoma biomarker development. *PLoS One* 2013; 8: e78115
- 75 Akers JC, Ramakrishnan V, Kim R, Phillips S, Kaimal V, Mao Y, Hua W, Yang I, Fu CC, Nolan J, Nakano I, Yang Y, Beaulieu M, Carter BS, Chen CC. miRNA contents of cerebrospinal fluid extracellular vesicles in glioblastoma patients. *J Neurooncol* 2015; 123: 205-16
- 76 Santangelo A, Imbruce P, Gardenghi B, Belli L, Agushi R, Tamanini A, Munari S, Bossi AM, Scambi I, Benati D, Mariotti R, Di Gennaro G, Sbarbati A, Eccher A, Ricciardi GK, Cicceri EM, Sala F, Pinna G, Lippi G, Cabrini G, Dechecchi MC. A microRNA signature from serum exosomes of patients with glioma as complementary diagnostic biomarker. *J Neurooncol* 2018; 136: 51-62

- 77 Ballester LY, Lu G, Zorofchian S, Vantaku V, Putluri V, Yan Y, Arevalo O, Zhu P, Riascos RF, Sreekumar A, Esquenazi Y, Putluri N, Zhu JJ. Analysis of cerebrospinal fluid metabolites in patients with primary or metastatic central nervous system tumors. *Acta Neuropathol Commun* 2018; 6: 85
- 78 Nayak L, Lee EQ, Wen PY. Epidemiology of brain metastases. *Curr Oncol Rep* 2012; 14: 48-54
- 79 Barnholtz-Sloan JS, Sloan AE, Davis FG, Vignea FD, Lai P, Sawaya RE. Incidence proportions of brain metastases in patients diagnosed (1973 to 2001) in the Metropolitan Detroit Cancer Surveillance System. *J Clin Oncol* 2004; 22: 2865-72
- 80 Peters S, Bexelius C, Munk V, Leighl N. The impact of brain metastasis on quality of life, resource utilization and survival in patients with non-small-cell lung cancer. *Cancer Treat Rev* 2016; 45: 139-62
- 81 Dagogo-Jack I, Gill CM, Cahill DP, Santagata S, Brastianos PK. Treatment of brain metastases in the modern genomic era. *Pharmacol Ther* 2017; 170: 64-72
- 82 Brastianos PK, Carter SL, Santagata S, Cahill DP, Taylor-Weiner A, Jones RT, Van Allen EM, Lawrence MS, Horowitz PM, Cibulskis K, Ligon KL, Tabernero J, Seoane J, Martinez-Saez E, Curry WT, Dunn IF, Paek SH, Park SH, McKenna A, Chevalier A, Rosenberg M, Barker FG, 2nd, Gill CM, Van Hummelen P, Thorner AR, Johnson BE, Hoang MP, Choueiri TK, Signoretti S, Sougnez C, Rabin MS, Lin NU, Winer EP, Stemmer-Rachamimov A, Meyerson M, Garraway L, Gabriel S, Lander ES, Beroukheim R, Batchelor TT, Baselga J, Louis DN, Getz G, Hahn WC. Genomic Characterization of Brain Metastases Reveals Branched Evolution and Potential Therapeutic Targets. *Cancer Discov* 2015; 5: 1164-77
- 83 Le Rhun E, Weller M, Brandsma D, Van den Bent M, de Azambuja E, Henriksson R, Boulanger T, Peters S, Watts C, Wick W, Wesseling P, Ruda R, Preusser M, Board EE, Committee EG. EANO-ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up of patients with leptomeningeal metastasis from solid tumours. *Ann Oncol* 2017; 28: iv84-iv99
- 84 Shingyoji M, Kageyama H, Sakaida T, Nakajima T, Matsui Y, Itakura M, Iuchi T, Yokoi S, Kimura H, Iizasa T. Detection of epithelial growth factor receptor mutations in cerebrospinal fluid from patients with lung adenocarcinoma suspected of neoplastic meningitis. *J Thorac Oncol* 2011; 6: 1215-20
- 85 Magbanua MJ, Melisko M, Roy R, Sosa EV, Hauranieh L, Kablanian A, Eisenbud LE, Ryazantsev A, Au A, Scott JH, Park JW. Molecular profiling of tumor cells in cerebrospinal fluid and matched primary tumors from metastatic breast cancer patients with leptomeningeal carcinomatosis. *Cancer Res* 2013; 73: 7134-43
- 86 Magbanua MJ, Roy R, Sosa EV, Hauranieh L, Kablanian A, Eisenbud LE, Ryazantsev A, Au A, Scott JH, Melisko M, Park JW. Genome-wide copy number analysis of cerebrospinal fluid tumor cells and their corresponding archival primary tumors. *Genom Data* 2014; 2: 60-2
- 87 Yang H, Cai L, Zhang Y, Tan H, Deng Q, Zhao M, Xu X. Sensitive detection of EGFR mutations in cerebrospinal fluid from lung adenocarcinoma patients with brain metastases. *J Mol Diagn* 2014; 16: 558-63
- 88 Sasaki S, Yoshioka Y, Ko R, Katsura Y, Namba Y, Shukuya T, Kido K, Iwakami S, Tominaga S, Takahashi K. Diagnostic significance of cerebrospinal fluid EGFR mutation analysis for leptomeningeal metastasis in non-small-cell lung cancer patients harboring an active EGFR mutation following gefitinib therapy failure. *Respir Investig* 2016; 54: 14-9
- 89 Li Y, Pan W, Connolly ID, Reddy S, Nagpal S, Quake S, Gephart MH. Tumor DNA in cerebral spinal fluid reflects clinical course in a patient with melanoma leptomeningeal brain metastases. *J Neurooncol* 2016; 128: 93-100
- 90 Marchio C, Mariani S, Bertero L, Di Bello C, Francia Di Celle P, Papotti M, Ruda R, Soffietti R, Cassoni P. Liquoral liquid biopsy in neoplastic meningitis enables molecular diagnosis and mutation tracking: a proof of concept. *Neuro Oncol* 2017; 19: 451-3
- 91 Jiang BY, Li YS, Guo WB, Zhang XC, Chen ZH, Su J, Zhong WZ, Yang XN, Yang JJ, Shao Y, Huang B, Liu YH, Zhou Q, Tu HY, Chen HJ, Wang Z, Xu CR, Wang BC, Wu SY, Gao CY, Zhang X, Wu YL.



Detection of Driver and Resistance Mutations in Leptomeningeal Metastases of NSCLC by Next-Generation Sequencing of Cerebrospinal Fluid Circulating Tumor Cells. *Clin Cancer Res* 2017;

92 Siravegna G, Geuna E, Mussolin B, Crisafulli G, Bartolini A, Galizia D, Casorzo L, Sarotto I, Scaltriti M, Sapino A, Bardelli A, Montemurro F. Genotyping tumour DNA in cerebrospinal fluid and plasma of a HER2-positive breast cancer patient with brain metastases. *ESMO Open* 2017; 2: e000253

93 Huang WT, Lu NM, Hsu WY, Chang SE, Atkins A, Mei R, Javey M. CSF-ctDNA SMSEQ Analysis to Tailor the Treatment of a Patient with Brain Metastases: A Case Report. *Case Rep Oncol* 2018; 11: 68-74

94 Li YS, Jiang BY, Yang JJ, Zhang XC, Zhang Z, Ye JY, Zhong WZ, Tu HY, Chen HJ, Wang Z, Xu CR, Wang BC, Du HJ, Chuai S, Han-Zhang H, Su J, Zhou Q, Yang XN, Guo WB, Yan HH, Liu YH, Yan LX, Huang B, Zheng MM, Wu YL. Unique genetic profiles from cerebrospinal fluid cell-free DNA in leptomeningeal metastases of EGFR-mutant non-small-cell lung cancer: a new medium of liquid biopsy. *Ann Oncol* 2018; 29: 945-52

95 Zhao J, Ye X, Xu Y, Chen M, Zhong W, Sun Y, Yang Z, Zhu G, Gu Y, Wang M. EGFR mutation status of paired cerebrospinal fluid and plasma samples in EGFR mutant non-small cell lung cancer with leptomeningeal metastases. *Cancer Chemother Pharmacol* 2016; 78: 1305-10

96 Jang NE, Baek SK, Jeong J-h, Kim S-Y, Yoon H-J, Cho KS, Jeong K-h, Park TS. Early Detection of BCR-ABL Fusion Gene of Cerebrospinal Fluid (CSF) by RT-PCR in Relapsed Acute Lymphoblastic Leukemia with Philadelphia Chromosome. *Lab Med* 2012; 43: e33-e7

97 Ballester LY, Glitza Oliva IC, Douse DY, Chen MM, Lan C, Haydu LE, Huse JT, Roy-Chowdhuri S, Luthra R, Wistuba II, Davies MA. Evaluating Circulating Tumor DNA From the Cerebrospinal Fluid of Patients With Melanoma and Leptomeningeal Disease. *J Neuropathol Exp Neurol* 2018; 77: 628-35

98 Pellerino A, Bertero L, Ruda R, Soffietti R. Neoplastic meningitis in solid tumors: from diagnosis to personalized treatments. *Ther Adv Neurol Disord* 2018; 11: 1756286418759618

99 Nanjo S, Hata A, Okuda C, Kaji R, Okada H, Tamura D, Irie K, Okada H, Fukushima S, Katakami N. Standard-dose osimertinib for refractory leptomeningeal metastases in T790M-positive EGFR-mutant non-small cell lung cancer. *Br J Cancer* 2018; 118: 32-7

100 Gabriel MT, Calleja LR, Chalopin A, Ory B, Heymann D. Circulating Tumor Cells: A Review of Non-EpCAM-Based Approaches for Cell Enrichment and Isolation. *Clin Chem* 2016; 62: 571-81

101 Lin X, Fleisher M, Rosenblum M, Lin O, Boire A, Briggs S, Bensman Y, Hurtado B, Shagabayeva L, DeAngelis LM, Panageas KS, Omuro A, Pentsova EI. Cerebrospinal fluid circulating tumor cells: a novel tool to diagnose leptomeningeal metastases from epithelial tumors. *Neuro Oncol* 2017; 19: 1248-54

102 Wang W, Bian C, Xia D, He JX, Hai P, Zhao R, Wang YY. Combining Carcinoembryonic Antigen and Platelet to Lymphocyte Ratio to Predict Brain Metastasis of Resected Lung Adenocarcinoma Patients. *Biomed Res Int* 2017; 2017: 8076384

103 Hamilton A, Sibson NR. Role of the systemic immune system in brain metastasis. *Mol Cell Neurosci* 2013; 53: 42-51

104 Peinado H, Aleckovic M, Lavotshkin S, Matei I, Costa-Silva B, Moreno-Bueno G, Hergueta-Redondo M, Williams C, Garcia-Santos G, Ghajar C, Nitadori-Hoshino A, Hoffman C, Badal K, Garcia BA, Callahan MK, Yuan J, Martins VR, Skog J, Kaplan RN, Brady MS, Wolchok JD, Chapman PB, Kang Y, Bromberg J, Lyden D. Melanoma exosomes educate bone marrow progenitor cells toward a pro-metastatic phenotype through MET. *Nat Med* 2012; 18: 883-91

105 Boire A, Zou Y, Shieh J, Macalinao DG, Pentsova E, Massague J. Complement Component 3 Adapts the Cerebrospinal Fluid for Leptomeningeal Metastasis. *Cell* 2017; 168: 1101-13 e13

106 Nevel KS, Wilcox JA, Robell LJ, Umemura Y. The Utility of Liquid Biopsy in Central Nervous System Malignancies. *Curr Oncol Rep* 2018; 20: 60

107 Zachariah MA, Oliveira-Costa JP, Carter BS, Stott SL, Nahed BV. Blood-based biomarkers for the diagnosis and monitoring of gliomas. *Neuro Oncol* 2018; 20: 1155-61

108 Preusser M, Winkler F, Valiente M, Manegold C, Moyal E, Widhalm G, Tonn JC, Zielinski C.  
Recent advances in the biology and treatment of brain metastases of non-small cell lung cancer:  
summary of a multidisciplinary roundtable discussion. ESMO Open 2018; 3: e000262

**Table 1. Studies investigating circulating cfDNA in primary CNS tumours.**

	<b>Tumour</b>	<b>Positive CSF cytology</b>	<b>Positive CSF molecular profiling</b>	<b>Positive blood molecular profiling</b>
<b>Rhodes CH et al., J Neuropathol Exp Neurol 1994</b>	Glioblastoma	ND/NR	1/1 (100%)	ND/NR
<b>Rhodes CH et al., Am J Clin Pathol 1995</b>	Glioblastoma	ND/NR	1/1 (100%)	ND/NR
<b>Boisselier B et al., Neurology 2012</b>	Glioma (low grade=8, high grade=17)	ND/NR	ND/NR	Low grade: 3/8 (37.5%) High grade: 12/17 (70.6%)
<b>Salkeni MA et al., J Neurooncol 2013</b>	Glioblastoma	ND/NR	ND/NR	3/3 (100%)
<b>Bettegowda et al., Sci Transl Med 2014</b>	Glioma (n=27) Medulloblastoma (n=14)	ND/NR	ND/NR	Glioma: <10% Medulloblastoma : <50%
<b>Pan W et al., Clin Chem 2015</b>	Meningioma (n=1) Schwannoma (n=1)	ND/NR	Meningioma: 1/1 (100%) Schwannoma: 0/1 (0%)	Meningioma: 0/1 (0%) Schwannoma: 0/1 (0%)
<b>De Mattos-Arruda L et al., Nat Commun 2015</b>	Glioblastoma (n=4) Medulloblastoma (n=2)	ND/NR	Glioblastoma: 4/4 (100%) Medulloblastoma: 2/2 (100%)	Glioblastoma: 0/4 (0%) Medulloblastoma : 0/2 (0%)
<b>Wang Y et al., Proc Natl Acad Sci USA 2015</b>	Low grade glioma (n=8) High grade glioma (n=13) Ependymoma (n=7) Medulloblastoma (n=6) Other low grade tumour (n=1)	ND/NR	Low grade glioma: 6/8 (75%) High grade glioma: 13/13 (100%) Ependymoma: 5/7 (71%) Medulloblastoma: 5/6 (83%) Other low grade tumour: 1/1 (100%)	ND/NR
<b>Pentsova EI et al., J Clin</b>	Glioma (n=8) Ependymoma	Glioma: 0/8 Ependymoma: 0/1	Glioma: 6/8 (75%) Ependymoma: 0/1	ND/NR



<b>Oncol 2016</b>	(n=1)		(0%)	
<b>Connolly ID et al., J Neurooncol 2017</b>	Ependymoma (n=3)	ND/NR	0/3 (0%)	0/3 (0%)
<b>Huang TY et al., Acta Neuropathol Comm 2017</b>	Diffuse midline glioma (n=5)	ND/NR	4/5 (80%)	ND/NR
<b>Martinez-Ricarte F et al., Clin Cancer Res 2018</b>	High grade glioma (n=15) Low grade glioma (n=5)	ND/NR	High grade glioma: 15/15 (100%) Low grade glioma: 2/5 (40%)	ND/NR
<b>Pan C et al., Acta Neuropathol 2018</b>	Brainstem glioma	ND/NR	39/40 (98%)	3/8 (38%)
<b>Panditharatna E et al., Clin Cancer Res 2018</b>	Diffuse midline glioma	ND/NR	24/27 (89%)	34/40 (85%)
<b>Hiemcke-Jiwa LS et al., Hematol Oncol 2018</b>	Lymphoplasmacytic lymphoma (n=6) PCNSL (n=1)	Lymphoplasmacytic lymphoma: 2/6 (33%) PCNSL: 1/1 (100%)	Lymphoplasmacytic lymphoma: 5/6 (83%) PCNSL: 1/1 (100%)	ND/NR
<b>Miller AM et al., Nature 2019</b>	Diffuse glioma (grade II-III-IV)	7/80 (9%) CSF cytology not available in 5 cases	42/85 (49%)	3/19 (16%)

Notes: i) CSF molecular profiling was considered positive if at least one tumour mutation was detected in cfDNA, but in some cases only a subset of alterations was detected compared to tissue samples; ii) Cases with negative/not available tissue profiling and negative liquid profiling were excluded. Conversely, cases with positive CSF despite negative or untested primary were considered; iii) Atypical cells were considered positive in terms of CSF cytology evaluation. CNS: central nervous system; CSF: cerebrospinal fluid; ND/NR: not done/not reported, PCNSL: primary central nervous system lymphoma.

**Table 2. Studies investigating circulating cfDNA in secondary CNS tumours from solid neoplasms.**

	Primary tumour	Type of CNS involvement	Positive CSF cytology	Positive CSF molecular profiling	Positive blood molecular profiling
<b>Rhodes CH <i>et al.</i>, <i>J Neuropathol Exp Neurol</i> 1994</b>	Breast	NM	ND/NR	1/1 (100%)	ND/NR
<b>Swinkels DW <i>et al.</i>, <i>Clin Chem</i> 2000</b>	NSCLC	NM	0/2 (0%)	2/2 (100%)	ND/NR
<b>Shingyoji M <i>et al.</i>, <i>J Thorac Oncol</i> 2011</b>	NSCLC	NM	10/21 (47.6%)	13/21 (61.9%)	ND/NR
<b>Yang H <i>et al.</i>, <i>J Mol Diagn</i> 2014</b>	NSCLC	BM and NM	ND/NR	BM: 2/5 (40%) NM: 4/4 (100%)	ND/NR
<b>De Mattos-Arruda L <i>et al.</i>, <i>Nat Commun</i> 2015</b>	Breast and lung	BM and NM	BM: 13/24 (54.2%) NM: 1/3 (33%)	BM: 17/17 (100%) NM: 3/3 (100%)	0% in CNS restricted disease
<b>Pan W <i>et al.</i>, <i>Clin Chem</i> 2015</b>	Mixed	BM and NM	BM: ND/NR NM: 2/2 (100%)	BM: 5/5 (100%) NM: 1/1 (total DNA was analysed)	BM: 3/5, 2/3 with also extra-CNS progression NM: ND/NR
<b>Sasaki S <i>et al.</i>, <i>Respir Investig</i> 2016</b>	NSCLC	NM	2/7 (28.6%)	7/7 (100%)	0/3 (0%)
<b>Li Y <i>et al.</i>, <i>J Neurooncol</i> 2016</b>	Melanoma	NM	1/1 (100%)	1/1 (100%)	0/1 (0%)
<b>Pentsova EI <i>et al.</i>, <i>J Clin Oncol</i> 2016</b>	Mixed	BM and NM	BM: ND/NR NM: 3/3 (100%)	BM: 17/24 (70.8%) NM: 3/3 (100%)	ND/NR
<b>Zhao J <i>et al.</i>, <i>Cancer Chemother Pharmacol</i> 2016</b>	NSCLC	NM		7/7 (100%)	2/7 (28.6%), 1/2 with also extra-CNS progression
<b>Marchiò C <i>et al.</i>, <i>Neuro Oncol</i> 2017</b>	NSCLC	NM	2/2 (100%)	2/2 (100%)	0/2 (0%)

<b>Siravegna G <i>et al.</i>, ESMO Open 2017</b>	Breast	NM	ND/NR	1/1 (100%)	1/1 (100%), but showed decreasing mutant allele frequencies despite CNS disease progression
<b>Huang W <i>et al.</i>, Case Rep Oncol 2018</b>	CUP	NM	1/1 (100%)	1/1 (100%)	ND/NR
<b>Li <i>et al.</i>, Ann Oncol 2018</b>	NSCLC	NM	18/28 (64%)	28/28 (100%)	19/26 (73.1%)
<b>Ballester LY <i>et al.</i>, J Neuropathol Exp Neurol 2018</b>	Melanoma	NM	3/3 (100%)	2/3 (67%)	ND/NR
<b>Nanjo S <i>et al.</i>, Br J Cancer 2018</b>	NSCLC	NM	5/13 (38%)	5/13 (38%)	ND/NR

Notes: i) CSF molecular profiling was considered positive if at least one tumour mutation was detected in cfDNA, but in some cases only a subset of alterations was detected compared to tissue samples; ii) In many cases NM was present together with BM: these cases were considered together with NM only cases; iii) Cases with negative/not available tissue profiling and negative liquid profiling were excluded. Conversely, cases with positive CSF despite negative or untested primary were considered; iv) Atypical cells were considered positive in terms of CSF cytology evaluation. BM: brain metastasis; CNS: central nervous system; CSF: cerebrospinal fluid; CUP: cancer of unknown primary; NM: neoplastic meningitis; NSCLC: non-small cell lung cancer; ND/NR: not done/not reported.



