

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Power Doppler sonography to predict the risk of surgical recurrence of Crohn's disease

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1632798> since 2019-07-30T14:18:12Z

Published version:

DOI:10.1007/s40477-014-0101-x

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Journal of Ultrasound

Power Doppler sonography to predict the risk of surgical recurrence of Crohn's disease --Manuscript Draft--

Manuscript Number:	JUSD-D-14-00019R2
Full Title:	Power Doppler sonography to predict the risk of surgical recurrence of Crohn's disease
Article Type:	Original Paper
Abstract:	<p>Purpose. The aim of this work is to investigate the role of power Doppler sonography as an additional predictor of surgical recurrence in Crohn's disease.</p> <p>Methods. A sample of 33 patients, with ileal or ileocolonic Crohn's disease, that had underwent intestinal resection, were retrospectively enrolled. All patients had bowel ultrasonography 7-16 months after resection. Power Doppler sonography of the preanastomotic ileum was evaluated as a possible prognostication tool to assess the risk of long-term need for reoperation.</p> <p>Results. The absolute incidence of surgical recurrence in those who had a positive power Doppler was 42%, while that of those who had a negative power Doppler was 28.6%. Combining the power Doppler with bowel wall thickness, the surgical recurrence risk grew from 41.2% of those with a positive power Doppler and thickness >3 mm to 55.6% of those with a positive power Doppler and thickness >6 mm.</p> <p>Conclusions. Power Doppler look to be another useful prediction tool for the personalization of patient's care. It could be useful to perform power Doppler in all patients with a wall thickness greater than 5 mm: for those who have a positive power Doppler it may be indicated a more aggressive prophylactic therapy.</p>
Corresponding Author:	Davide Giuseppe Ribaldone ITALY
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	
Corresponding Author's Secondary Institution:	
First Author:	Davide Giuseppe Ribaldone
First Author Secondary Information:	
Order of Authors:	Davide Giuseppe Ribaldone Teresa Cammarota Andrea Resegotti Antonino Sarno Daniela Robotti Paola Debani Giovanni Bonenti Francesca Bresso Rinaldo Pellicano Alessandro Adriani Carlo Sguazzini Stefania Reggiani Marco Astegiano
Order of Authors Secondary Information:	

Author Comments:	This is the first serie in the literature in which is studied the predictive power of the color Doppler of surgical recurrence of Crohn's disease; we have found another useful factor for the personalization of patient's care: knowing that the 54.5% of patients who at bowel ultrasound performed within 1 year after operation have a bowel wall thickness > 5 mm and positive power Doppler will be reoperated could select the patients that should undergo colonoscopy for assessing endoscopic recurrence and authorize the use of a more aggressive medical therapy to try to change the natural history of their disease.
Response to Reviewers:	1) the color Doppler imagine has been removed 2) declaration about Conflict of interest, Informed Consent, Human & Animal Studies have been added

Power Doppler sonography to predict the risk of surgical recurrence of Crohn's disease.

Abstract

Purpose. The aim of this work is to investigate the role of power Doppler sonography as an additional predictor of surgical recurrence in Crohn's disease.

Methods. A sample of 33 patients, with ileal or ileocolonic Crohn's disease, that had underwent intestinal resection, were retrospectively enrolled. All patients had bowel ultrasonography 7–16 months after resection. Power Doppler sonography of the preanastomotic ileum was evaluated as a possible prognostication tool to assess the risk of long-term need for reoperation.

Results. The absolute incidence of surgical recurrence in those who had a positive power Doppler was 42%, while that of those who had a negative power Doppler was 28.6%. Combining the power Doppler with bowel wall thickness, the surgical recurrence risk grew from 41.2% of those with a positive power Doppler and thickness >3 mm to 55.6% of those with a positive power Doppler and thickness >6 mm.

Conclusions. Power Doppler look to be another useful prediction tool for the personalization of patient's care. It could be useful to perform power Doppler in all patients with a wall thickness greater than 5 mm: for those who have a positive power Doppler it may be indicated a more aggressive prophylactic therapy.

Keywords: Color Doppler; Inflammation; Small Bowel; Ultrasound.

Introduction

1
2
3
4
5
6 In a previous study [1], we found that bowel wall thickness at the anastomosis >3
7
8
9 mm, detected through bowel ultrasound performed 1 year after surgery, could be
10
11
12 added to the list of non-invasive predictors of early recurrence after ileo-colonic
13
14
15 resection besides those implying a double risk of recurrence, such as smoking,
16
17
18 previous bowel surgery, penetrating behavior of the disease, perianal location,
19
20
21 extensive small bowel resection, and absence of prophylactic treatment [2].
22
23

24 In recent years, power Doppler has been proposed as an additional
25
26
27 prognostication tool to increase the accuracy of bowel ultrasound in the assessment
28
29
30 of disease activity [3-5].
31
32

33 A wide deployment of these additional techniques is impaired by the intrinsic
34
35
36 variability. It has been also proposed that the morphological characteristics of the
37
38
39 intestinal wall correlate with the histological pattern [6, 7] and with the clinical activity
40
41
42 [8-11], although the correlation between the intestinal wall thickness and Crohn's
43
44
45 disease activity index (CDAI) is not close [12].
46
47
48
49
50

Methodology

51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 We performed a retrospective study on 196 patients that underwent ileal or ileo-
2 colonic resection with ileo-colonic anastomosis between December 1993 and March
3
4
5
6 2009 (all diagnosis of Crohn's disease was confirmed through histologic
7
8
9 examination). All patients underwent bowel ultrasound after at least 6 months from
10
11
12 surgery; in this study, we analyzed a total of 33 patients who were evaluated by
13
14
15 power Doppler.
16

17 The study was reviewed and authorized by the Local Ethical Committee.
18

19
20
21 Bowel ultrasound was performed using the following ultrasonographic
22
23
24 equipments: ESAOTE AU4, ESAOTE Technos, ESAOTE MyLab 70, and TOSHIBA
25
26
27 Aplio. A first evaluation of the bowel was made with a convex transducer (frequency
28
29
30 3.5 MHz) and then with a high frequency linear-array transducer (7.5–10 MHz);
31
32
33 power Doppler was performed at the preanastomotic ileum.
34
35

36 The four ultrasonographers who performed the examinations were all
37
38
39 radiologists with at least 3 years of experience in ultrasound scanning of patients with
40
41
42 intestinal bowel diseases and all working in the same department. Each investigation
43
44
45 was performed by one radiologist, C. T., S. A, R. D., D. P., who, respectively,
46
47
48 performed 11, 10, 7 and 5 investigations. During the procedure, the involved tract
49
50
51 wall was examined in a transverse section, from the central hyperechoic line of the
52
53
54 lumen to the outer hyperechoic margin of the wall with a linear-array transducer. Wall
55
56
57 thickness of the ileo-colonic anastomosis was calculated as the average of at least 3
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

measurements; at the end of the US investigation, measurements of intestinal wall and power Doppler results at the ileo-colonic anastomosis level were reported on a standardized form. Flow was considered to be present when colored pixels could be constantly reproduced in the area of interest or when flowmetric signals could be documented by spectral analysis of the pulsed color Doppler signal. Color Doppler flow was subjectively characterized either as positive (clearly visible, i.e. numerous color signals distributed within and/or around the inflammatory mass or clear identification of vessel paths or minimally visible, i.e. few and/or scattered color signals) or negative (no color signal obtained either inside the lesion or in the periphery).

Ileocolonoscopy was performed in a little percentage of patients and at a different time than bowel ultrasound (due to the invasiveness of this type of exam in patients with resection of the ileum for Crohn's disease), so it is impossible to prove any correlation.

The CDAI was calculated in the follow-up visits.

Statistical analysis

Statistical analysis was performed by using MedCalc software (version 9.2.1.0). The relative risk was calculated with a confidence interval of 95%.

Results

Clinical characteristics of the study population are reported in Table 1.

Table 1.

Other data: 28 patients were clinically inactive (CDAI < 150 at the time of the bowel ultrasound); bowel ultrasound was performed after a mean time of 13 months (range 7–16 months) after surgery.

The average follow-up was 95 months (range: 25–156 months) after surgery.

Major surgical complications occurred in 5 out of the 33 patients (15%).

Clearly, our final analysis refers to the reoperation for the disease recurrence and not to the reoperation for these immediate complications (Table 2).

Table 2.

The rate of surgical recurrence for the whole population was 36.4% (12 of 33 patients), at a mean distance of 50 months from previous surgery (range: 20-144 months).

Discussion

1
2
3
4
5
6 Although a sample of 33 patients is not very a large one, considering that power
7
8
9 doppler in real world is usually assessed only in a subgroup of patients that
10
11
12 underwent bowel ultrasound and the follow up of 95 months is long enough, this
13
14
15 looks to be the first case in the literature in which the predictive power of color
16
17
18 Doppler for the surgical recurrence of Crohn's disease is assessed.

19
20
21 Out of 33 patients, 19 of them had a positive power Doppler after surgery and
22
23
24 this seems to be linked to the fact that color Doppler in clinical practice is performed
25
26
27 mostly in patients with a thickened bowel wall and, thus, with a greater probability of
28
29
30 active disease.

31
32
33 The surgical recurrence was found to be 36.4%; this percentage is higher than
34
35
36 the 20.4% found in our previous study [1]; this seems linked to the fact that the
37
38
39 population in which the power Doppler is normally assessed usually has an
40
41
42 increased risk of recurrence (increased thickness of intestinal wall).

43
44
45 Patients with positive power Doppler at the anastomosis run a relative risk =
46
47
48 1.5 (95% CI = 0.6–3.9) of undergoing surgery compared to patients with negative
49
50
51 power Doppler.

52
53
54 The RR of surgical recurrence of those with positive color Doppler compared to those
55
56
57 with negative color Doppler does not reach the statistical significance, since there is
58
59
60
61
62
63
64
65

1 a selection bias of the population, as there are very few patients with negative power
2 Doppler and physiological thickness of the wall (those who have physiological
3 thickness of the wall generally will not be evaluated by power Doppler in real world,
4
5
6
7
8
9 which is the setting of our study).

10
11 The RR of surgical recurrence for patients with positive power Doppler and a bowel
12 wall thickness at the anastomosis greater than 3 mm (18 patients), 4 mm (16
13
14
15 patients), 5 mm (12 patients), 6 mm (10 patients), respectively, compared to patients
16
17
18
19
20
21 with negative power Doppler and lower bowel thickness is shown in Table 3.

22
23
24
25
26
27 Table 3.

28
29
30
31
32
33 The RR of patients with positive power Doppler and increased wall thickness
34
35
36 does not reach the statistical significance compared to those with negative power
37
38
39 Doppler and lower wall thickness because of the smaller sample size compared to
40
41
42 our previous study [1] (in fact, only a limited number of patients who underwent bowel
43
44
45 ultrasound were assessed using power Doppler).

46
47
48 The absolute incidence of new surgical intervention was: 28.6% in patients
49
50
51 with negative power Doppler, 42% with positive power Doppler, 41.2% with positive
52
53
54 power Doppler and thickness >3 mm, 46.7% with positive power Doppler and
55
56
57 thickness >4 mm, 54.5% with positive power Doppler and thickness >5 mm, and,
58
59
60
61
62
63
64
65

finally, 55.6% with positive power Doppler and thickness >6 mm (Figure 1).

Figure 1.

As far as the therapy, the percentage of patients not taking prophylactic therapy of post-surgical recurrence fell from 31.1% of the previous study [1] to 20%; only 12% of patients were receiving steroids and 3% immunosuppressive agents (Figure 2).

Figure 2.

The percentage of patients with positive power Doppler that had taken steroids or immunosuppressants or biologics at the time of bowel ultrasound (2 out of 19, 10.5%) is similar to that of the whole population (5 out of 33, 15%).

Some critical issues should be considered. The retrospective design of the study represents a limitation and this is the reason why the color Doppler has been assessed mainly in patients with thickened bowel wall, but this is exactly what happens in daily clinical practice; in addition, the examinations have been assessed by the same group of radiologists and all patients were treated along the years by the same doctor (A. M.).

1
2
3 In conclusion, it seems we have found another useful factor for the
4
5 personalization of patient's care: knowing that 54.5% of patients who performed
6
7 bowel ultrasound within 1 year after operation and showed both a bowel wall
8
9 thickness >5 mm and a positive power Doppler will be reoperated, this could help
10
11 selecting the patients that should undergo colonoscopy for assessing the endoscopic
12
13 recurrence and suggest the use of a more aggressive medical therapy to try to
14
15 change the natural history of their disease.
16
17
18
19
20

21 A prospective study would also be useful, where the population with bowel
22
23 wall thickness >5 mm within 1 year after surgery would be assessed by color
24
25 Doppler: those with positive color Doppler should undergo colonoscopy for assessing
26
27 endoscopic recurrence and should be treated with more aggressive drugs (topical or
28
29 systemic steroids, immunosuppressants, biological agents) to modify the disease
30
31 behavior and to extend their surgical recurrence-free interval.
32
33
34
35
36
37
38
39
40
41

42 Conflict of interest

43
44
45 We have no conflict of interest.
46
47
48
49
50

51 Informed Consent

52
53
54 All procedures followed were in accordance with the ethical standards of the
55
56 responsible committee on human experimentation (institutional and national) and
57
58
59
60
61
62
63
64
65

1
2
3 with the Helsinki Declaration of 1975, as revised in 2000. All patients provided written
4
5 informed consent to enrolment in the study and to the inclusion in this article of
6
7 information that could potentially lead to their identification.
8
9

10 11 12 Human & Animal Studies

13
14
15 The study was conducted in accordance with all institutional and national guidelines
16
17
18 for the care and use of laboratory animals.
19
20
21
22
23

24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65

- 1) Cammarota T, Ribaldone DG, Resegotti A, Repici A, Danese S, Fiorino G, et al (2013) Role of bowel ultrasound as a predictor of surgical recurrence of Crohn's disease. *Scand J Gastroenterol* 48:552-555
- 2) Bernstein C, Rawsthorne P, Cheang M, Blanchard JF (2006) A population-based case control study of potential risk factors for IBD. *Am J Gastroenterol* 101:993-1002
- 3) Esteban JM, Aleixandre A, Hurtado MJ, Maldonado L, Mora FJ, Nogues E (2003) Contrast-enhanced power Doppler ultrasound in the diagnosis and follow-up of inflammatory abdominal masses in Crohn's disease. *Eur J Gastroenterol Hepatol* 2003 15:253-259
- 4) Robotti D, Cammarota T, Deboni P, Sarno A, Astegiano M (2004) Activity of Crohn disease: value of Color-Power-Doppler and contrast-enhanced ultrasonography. *Abdom Imaging* 29:648-652
- 5) Bolondi L, Gaiani S, Brignola C, Campieri M, Rigamonti A, Zironi G, et al (1992) Changes in splanchnic hemodynamics in inflammatory bowel disease. Non-invasive assessment by Doppler ultrasound flowmetry. *Scand J Gastroenterol* 27:501-507
- 6) Parente F, Maconi G, Bollani S, Anderloni A, Sampietro G, Cristaldi M, et al (2002) Bowel ultrasound in assessment of Crohn's disease and detection of related small bowel strictures: a prospective

comparative study versus x ray and intraoperative findings. Gut 50:490-495

- 1
2 7) Maconi G, Carsana L, Fociani P, Sampietro GM, Ardizzone S, Cristaldi M, et al (2003) Small bowel stenosis
3
4 in Crohn's disease: clinical, biochemical and ultrasonographic evaluation of histological features.
5
6 Aliment Pharmacol Ther 18:749-756
7
- 8 8) Mayer D, Reinshagen M, Mason RA, Muche R, von Tirpitz C, Eckelt D, et al (2000) Sonographic
9
10 measurement of thickened bowel wall segments as a quantitative parameter for activity in inflammatory bowel
11
12 disease. Z Gastroenterol 38:295-300
13
14
- 15 9) Haber HP, Busch A, Ziebach R, Stern M (2000) Bowel wall thickness measured by ultrasound as a marker of
16
17 Crohn's disease activity in children. Lancet 355:1239-1240
18
19
- 20 10) Castiglione F, deSio I, Cozzolino A, Rispo A, Manguso F, Del Vecchio Blanco G (2004) Bowel wall
21
22 thickness at abdominal ultrasound and the one-year-risk of surgery in patients with Crohn's disease. Am J
23
24 Gastroenterol 99:1977-1983
25
26
- 27 11) Maconi G, Parente F, Bollani S, Cesana B, Bianchi Porro G (1996) Abdominal ultrasound in the assessment
28
29 of extent and activity of Crohn's disease: clinical significance and implication of bowel wall thickening. Am J
30
31 Gastroenterol 91:1604-1609
32
33
- 34 12) Best WR, Beckett JM, Singleton JW (1979) Rederived values of the eight coefficients of the Crohn's Disease
35
36 Activity Index (CDAI). Gastroenterology 77:843-846
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Fig. 1 Surgical recurrence rate

Fig. 2 Medications at the time of the bowel ultrasound

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65

Power Doppler sonography to predict the risk of surgical recurrence of Crohn's disease.

Davide Giuseppe Ribaldone ^a, Teresa Cammarota ^b, Andrea Resegotti ^c, Antonino Sarno ^b, Daniela Robotti ^b, Paola Debanì ^b, Giovanni Bonenti ^b, Francesca Bresso ^d, Rinaldo Pellicano ^a, Alessandro Adriani ^a, Carlo Sguazzini ^a, Stefania Reggiani ^a, Marco Astegiano

^a

^a Department of Gastro-Hepatology, Città della Salute e della Scienza, Turin, Italy

^b Radiologia 5, Department of Radiology, Città della Salute e della Scienza, Turin, Italy

^c 7th Department of Surgery, Città della Salute e della Scienza, Turin, Italy

^d Gastrocentrum medicin, Karolinska university hospital, Stockholm, Sweden

Davide Giuseppe Ribaldone: Department of Gastro-Hepatology, Città della Salute e della Scienza, Corso Bramante, n° 88, 10126 Turin, Italy; tel (0039)3476861126, fax (0039)0116333623, davrib_1998@yahoo.com

Running title: Doppler and Crohn's disease recurrence.

Table 1. Characteristics of the study population ($n = 33$, Crohn's disease patients).

Males/females	18/15
Median age, years (range)	44 (24–72)
Duration of Crohn's disease, years (range)	12 (0–54)
Smoking habit at surgery, n (%)	
Ever smoker	15 (45.5)
Never smoker	18 (54.5)
Previous surgery, n (%)	
Never	17 (51.5)
1	13 (39.4)
2	1 (3.0)
> 2	2 (6.1)
Total length of bowel resection (cm)	
< 50	20 (60.6%)
>= 50	13 (39.4%)
Indication for surgery, n (%)	
Strictures	20 (60.6)
Fistula	5 (15.2)
Refractoriness to medical therapy	6 (18.2)
Perforation	2 (6.1)

Table 2. Major surgical complications.

Patients with major surgical complications	5 (15%)
Complications	
Anastomotic dehiscence	2
Entero-cutaneous fistula	1
Hemorrhage	1
Venous thrombosis	1
Reoperation	2
Number of patients with anastomosis wall thickness >3 mm 1 year after operation	5 (100%)
Number of patients with surgical recurrence	3 (60%)

Table 3. RR of surgical recurrence.

Power Doppler positive and wall thickness at the anastomosis (mm)	RR
> 3	1.3 (95% CI = 0.5 ÷ 3.3)
> 4	1.7 (95% CI = 0.7 ÷ 4.2)
> 5	2.0 (95% CI = 0.8 ÷ 4.8)
> 6	1.9 (95% CI = 0.8 ÷ 4.5)





Figure 2

[Click here to download high resolution image](#)



Authorship disclosure form

[Click here to download Authorship disclosure form: Cover letter.doc](#)