

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

## Effects of astaxanthin and canthaxanthin on oxidative stress biomarkers in rainbow trout

### **This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1719480> since 2019-12-22T19:33:09Z

*Published version:*

DOI:10.1080/15287394.2019.1648346

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



### Effects of astaxanthin and canthaxanthin on oxidative stress biomarkers in rainbow trout

Journal:	<i>Journal of Toxicology and Environmental Health, Part A: Current Issues</i>
Manuscript ID	Draft
Manuscript Type:	Original Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Elia, Antonia; Universita degli Studi di Perugia, Prearo, Marino; Istituto Zooprofilattico Sperimentale del Piemonte Liguria e Valle d'Aosta Dörr, Ambrosius; Universita degli Studi di Perugia Pacini, Nicole; Universita degli Studi di Perugia Magara, Gabriele; Universita degli Studi di Perugia Brizio, Paola; Istituto Zooprofilattico Sperimentale del Piemonte Liguria e Valle d'Aosta Gasco, Laura; Department of Agricultural, Forest and Food Sciences, University of Torino, Largo P. Braccini 2, 10095 Grugliasco (TO), Italy Abete, Maria; Istituto Zooprofilattico Sperimentale del Piemonte Liguria e Valle d'Aosta
Keywords:	astaxanthin, canthaxanthin, oxidative stress biomarkers, rainbow trout, tissues

SCHOLARONE™  
Manuscripts

1  
2  
3 **Effects of astaxanthin and canthaxanthin on oxidative stress biomarkers in**  
4  
5  
6 **rainbow trout**  
7

8 *Antonia Concetta Elia<sup>a\*</sup>, Marino Prearo<sup>b</sup>, Ambrosius Josef Martin Dörr<sup>a</sup>, Nicole*  
9  
10  
11 *Pacini<sup>a</sup>, Gabriele Magara<sup>a</sup>, Paola Brizio<sup>b</sup>, Laura Gasco<sup>c</sup>, Maria Cesarina Abete<sup>b</sup>*  
12

13  
14  
15 *a Department of Chemistry, Biology and Biotechnology, via Elce di sotto 8, University of Perugia,*  
16  
17 *06123 Perugia, Italy*  
18

19  
20 *b Veterinary Medical Research Institute for Piedmont, Liguria and Aosta Valley, Via Bologna, 148,*  
21  
22 *10154 Torino, Italy;*  
23

24  
25  
26 *c Department of Agricultural, Forest and Food Sciences, University of Torino, Largo P. Braccini 2,*  
27  
28 *10095 Grugliasco (TO), Italy.*  
29

30  
31 Running title: Xanthophylls mediated-oxidative stress in *Oncorhynchus mykiss*  
32

33  
34  
35  
36  
37  
38 *\* Corresponding author: Dr. Antonia Concetta Elia*  
39

40  
41 *Department of Chemistry,*  
42

43  
44  
45 *Biology and Biotechnology,*  
46

47  
48 *University of Perugia, Italy*  
49

50  
51 *Email: antonia.elia@unipg.it*  
52  
53  
54  
55  
56  
57  
58  
59  
60

**ABSTRACT**

Farmed trout are commonly fed carotenoid-enriched diets during the finishing period to acquire the typical red-to-pink flesh color in salmonid muscle. The aim of this study was to examine the effects of the two xanthophylls, astaxanthin (Ax) or canthaxanthin (Cx), administered individually or in combination, on oxidative stress biomarkers in kidney and liver of rainbow trout. Specimens were fed Ax and/or Cx enriched diets with 75 or 25 mg of each compound/kg for eight weeks. Changes in concentration of oxidative stress biomarkers, such as glutathione, superoxide dismutase, catalase, glutathione peroxidase, glutathione reductase, and glutathione *S*-transferase, were recorded in Ax- and Cx-dosed trout. The two carotenoids, mainly Cx, impelled antioxidant defense in rainbow trout, while lipid peroxidation process was not recorded for all Ax or Cx-dosed trout. Furthermore, both combined xanthophylls did not exert remarkable combined effects in liver and kidney. Biomarker responses were generally altered in both tissues throughout both experimental endpoints, suggesting that different time-dependent pathways may be involved in strengthening the anti-oxidative defense in Ax and/or Cx-fed trout.

**Keywords:** astaxanthin, canthaxanthin, oxidative stress biomarkers, rainbow trout, tissues

## Introduction

Salmonid farming strongly depends on market demand. Excellent organoleptic fish grade can be very attractive for consumers, and industries are struggling to conceive efficient strategies to enhance the appeal of aquaculture products. The aspect of fish fillet mostly influences customer's choice, and therefore farmed salmonids are often nourished during the finishing period with carotenoid-enriched diets. Likewise, pigmentation degree of meat is an important criterion of quality for farmers and consumers.

Oxocarotenoids (xanthophylls), such as Astaxanthin (Ax, 3,3'-dihydroxy- $\beta,\beta$ -carotene-4,4'-dione) and canthaxanthin (Cx, 4,4-diketo-beta-carotene) have been widely used in order to develop the typical red-to-pink flesh color in salmonid muscle, due to fish ability to combine Ax and Cx with tissue actomyosinic complexes. Furthermore, a previous study reported that astaxanthin had a greater binding affinity for muscle fibers than canthaxanthin (Henmi et al., 1991). However, xanthophylls are considered as safe additives for animal and human consumption, and the maximum content in feed for Ax has been established by EU legislation (Commission implementing Regulation (EU) 2015/1415), while for Cx, also known as E161g, the maximum level as an additive in aquaculture feed was reduced from 100 to 25 mg/kg (Commission Directive 2003/7/EC).

Changes in oxidative stress biomarkers may be related with oxidative catabolism and nutritional factors (Martínez-Álvarez et al., 2005). In addition to its role as food additive, astaxanthin as well as other carotenoids are recognized as efficient physical and chemical scavengers of reactive oxygen species (ROS; hydrogen peroxide and hydroxyl radicals), quenchers of singlet oxygen ( $^1O_2$ ) and inhibitors of lipid peroxidation (Chew, 1995; Sies and Stahl, 1995; Chien et al., 2003; Shimidzu et al., 1996; Stahl and Sies, 2003).

In fact, the Ax double bonds conjugated with relatively unstable electronic orbitals can reduce the concentration of prooxidant molecules (Conn et al., 1991). Astaxanthin can also enhance total antioxidant status (TAS) (Chien et al., 2003) and modulate lipid peroxidation and antioxidant

1  
2  
3 enzymes activity (Shih et al., 2008). Therefore, xanthophyll may be involved in improving the  
4  
5 defense mechanisms against ROS, thus preventing oxidative stress impairment. However, although  
6  
7 Ax and Cx are widely used in fish industry, the potential oxidative stress modulating role of the two  
8  
9 xanthophylls on rainbow trout (*Oncorhynchus mykiss*) has been poorly investigated (Nakano et al.,  
10  
11 1999).

12  
13  
14 The aim of the study was to investigate the effects of Ax (100 mg/kg) or Cx (25 mg/kg) diets  
15  
16 administered alone or in combination for eight weeks on antioxidant stress biomarkers in liver and  
17  
18 kidney of rainbow trout *Oncorhynchus mykiss*. The chosen doses would mimic the regulatory limits  
19  
20 for dietary inclusion of astaxanthin and canthaxanthin in aquaculture industry (Commission  
21  
22 Directive 2003/7/EC).  
23  
24  
25  
26  
27

## 28 **Materials and Methods**

### 29 *Fish exposure*

30  
31  
32 Control and pigment diets were manufactured in the laboratory at the Experimental Station of the  
33  
34 Department of Agricultural, Forest and Food Sciences of the University of Turin. Briefly, in-house  
35  
36 pelleted feeding stuffs were prepared from the same basal composition (control diet) and  
37  
38 supplemented with Ax and/or Cx. Ingredients of the experimental diets (on weight basis) were: fish  
39  
40 meal (50%), corn gluten meal (10%), dehulled barley meal (18%), corn meal (9%), cod liver oil  
41  
42 (7%), binder (merigel, 3%), mineral mixture (1.5%) and vitamin mixture (1.5%). Astaxanthin and  
43  
44 canthaxanthin doses were chosen according to the previous study of Brizio et al. (2013). The three  
45  
46 pigment-diets were: 1) Ax (75 mg/kg feed); 2) Cx (25 mg/kg feed); 3) Ax + Cx (75+25 mg/kg feed,  
47  
48 respectively). Control diet was supplied to fish without carotenoids.  
49  
50  
51  
52  
53

54 Females of rainbow trout (*O. mykiss* n=80) were divided into 8 tanks (10 specimens each tank)  
55  
56 placed at same time (e.g 2 control groups, and 2 Ax, 2 Cx and 2 Ax+Cx groups) and then  
57  
58 acclimatized for 15 days in flow-through open system with artesian well water ( $13 \pm 1^\circ\text{C}$ ) with each  
59  
60 tank (1500 L) having a water inflow of 8 L/min. Dissolved oxygen was measured daily and ranged

1  
2  
3 between 7.6 and 8.7 mg/L. The trial lasted for 8 weeks and specimens were fed by hand twice a  
4 day, 7d per week and the daily quantity of feed distributed was set at 1% of the tank biomass. No  
5 feed refusals were recorded during the experiment. Mortality was checked every day.  
6  
7

8  
9  
10 At week four and eight, the specimens of each experimental group were sampled and killed by over  
11 anesthesia (MS-222; PHARMAQ Ltd., UK). Each fish was weighed and total length was measured  
12 from the tip of the snout to the tip of the longer lobe of the caudal fin. Liver and kidney were  
13 collected immediately and stored in dry ice and next at -80°C, until used for analyses of antioxidant  
14 enzymes and total glutathione in liver and kidney.  
15  
16

17  
18  
19 The experiments were conducted in accordance with the European and national guidelines for the  
20 protection of animals used for experimental and other scientific purposes (European Commission,  
21 Directive 86/609/EC and Italian Directive 116/1992, respectively).  
22  
23  
24  
25  
26  
27  
28  
29

### 30 ***Biochemical analyses***

31  
32  
33 The biochemical analyses were performed on the cytosolic fractions of liver and kidney as  
34 previously reported (Elia et al., 2017; Pacini et al., 2013).  
35  
36

37  
38 Levels of lipid peroxidation products (MDA) were analyzed in each liver or kidney (1:10 and 1:5,  
39 respectively) in 20 mM 2-amino-2-hydroxymethylpropane-1,3-diol (TRIS/HCl), pH 7.4, and 0.5 M  
40 butylated hydroxytoluene (BHT), centrifuged at 3000×g for 20 min at 4 °C. The supernatant was  
41 derivatized in 1-methyl-2-phenylindole (10.32 mM in acetonitrile/methanol 3:1), HCl, and dilution  
42 buffer (TRIS/HCl, pH 7.4), sample or MDA standard (0-4 µM of 1,1,3,3-tetramethoxypropane). All  
43 samples were incubated at 45 °C for 60 min, then centrifuged at 15000×g for 10 min and read  
44 spectrophotometrically at 586 nm.  
45  
46  
47  
48  
49  
50  
51  
52

53  
54 For antioxidant biomarkers, liver and kidney (0.2 g) were homogenized (1:10 w/v) in 100 mM  
55 TRIS buffer, pH 7.8, 100 µM phenylmethylsulphonyl fluoride (PMSF), 0.008 TIU mL<sup>-1</sup> and  
56 bacitracin 0.1mg mL<sup>-1</sup>, and then centrifuged at 100,000 g for 1 h at 4 °C.  
57  
58  
59  
60

1  
2  
3 Superoxide dismutase (SOD) activity was evaluated according to the original method of McCord  
4 and Fridovich (1969) and slightly modified (Pacini et al., 2012). The reduction of cytochrome C by  
5 the system xanthine/hypoxanthine was measured at 550 nm and a standard curve of SOD units was  
6 considered. One unit of SOD is defined as the amount of enzyme that inhibits by 50% the reduction  
7 of cytochrome C.  
8  
9

10  
11  
12 Catalase (CAT) activity was measured by following the decrease in absorbance at 240 nm due to  
13 H<sub>2</sub>O<sub>2</sub> consumption. The assay was carried out in Na-phosphate buffer pH 7 and 12 mM H<sub>2</sub>O<sub>2</sub>.  
14  
15

16  
17 Glutathione peroxidase (GPx) activity towards hydrogen peroxidase (H<sub>2</sub>O<sub>2</sub>) as substrate, was  
18 determined and the oxidation of NADPH was followed at 340. The assay condition was: 100 mM  
19 Na-phosphate buffer with pH 7.5, 1mM EDTA, 0.12 mM NADPH, 2 mM GSH, 1 mM NaN<sub>3</sub>, 1 U  
20 GR, 0.6 mM H<sub>2</sub>O<sub>2</sub>.  
21  
22

23  
24 Glutathione reductase (GR) activity was assayed following the decrease in absorbance at 340 nm  
25 due to the oxidation of NADPH. The assay condition was: 100 mM Na-phosphate buffer with pH 7,  
26 1 mM GSSG and 60 μM NADPH.  
27  
28

29  
30 Glutathione S-transferase (GST) activity was measured in 100 mM Na-phosphate buffer with pH  
31 6.5 with 1 mM CDNB that forming the conjugate with 1 mM GSH at 340 nm.  
32  
33

34  
35 Protein concentration was determined according to the method reported in Lowry et al. (1951).  
36  
37

38  
39 Total glutathione was measured in liver and kidney. Weighted tissues (0.5 g) were homogenized  
40 (1:5 w/v) in 5% sulphosalicylic acid with 4 mM ethylene diamine tetraacetic acid (EDTA), and  
41 centrifuged at 30,000 x g for 15 min. Total glutathione (GSH+2GSSG) content was measured in  
42 potassium phosphate (KP) buffer 100 mM pH 7, EDTA 1 mM, glutathione reductase (GR) 1U,  
43 NADPH 4 mg ml<sup>-1</sup> and 5,5'-dithiobis (2-nitrobenzoic acid) (DTNB) 1.5 mg ml<sup>-1</sup>, both dissolved in  
44 NaCO<sub>3</sub> 0.5%. The oxidized glutathione was used as the standard and the absorbance was read at  
45 412 nm.  
46  
47

48  
49 All biochemical analyses were performed in triplicate on each sample with a Varian  
50 spectrophotometer (Cary 50 Thermostat Cell Holder) at a constant temperature of 25°C.  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



### ***Statistical analyses***

Data are reported as mean and standard deviation (SD). Homogeneity of variance was checked by Bartlett test. To highlight the statistically significant differences among specimens fed carotenoid-supplied diets and controls, we used one-way ANOVA and the Tukey's Multiple Comparison Test. Statistical significance was checked at  $p < 0.05$ .

### **Results**

No statistically significant differences in growth and length were recorded between treated and control rainbow trout within the same experimental period (4 or 8 weeks). However, length of control and Cx groups and weight of Cx dosed trout were slightly higher at 8 weeks, when compared to those recorded at 4 weeks (Table 1).

Ax- and Cx- enriched feed administered to rainbow trout, either singly or in combination, did not produce changes of MDA levels in liver, whereas same metabolite was higher (75%) in kidney of Ax + Cx trout at 4 weeks, if compared to the last experimental time (Figure 1).

Elevated hepatic SOD activity was noted in fish dosed with Cx (65%) and Ax + Cx (95%), if compared with control group and the Ax and Ax+Cx one at 4 weeks. Moreover, SOD levels were also higher in Cx fish (2 fold) and Ax+Cx (3 fold), if compared with the same groups at 8 weeks (Figure 2). Therefore, SOD activity did not show a monotonic exposure response relationship, since at 8 weeks activity in all carotenoids treated fish was similar to control ones.

No significant changes of SOD levels were recorded in kidney for the whole sample (Figure 2).

Administration of carotenoids, Ax and/or Cx, to rainbow trout produced only slightly higher CAT activity in Ax groups at both experimental times. No significant changes of CAT activity were recorded in kidney (Figure 3) and for GPx in liver (Figure 4).

GPx activity was higher (about 95%) in kidney treated with Cx dose administered singly or combined with Ax (Ax+Cx), when compared with own control at 4 weeks and with same experimental groups at 8 weeks (about 2 fold) (Figure 4). Therefore, GPx activity similar to SOD in

1  
2  
3 liver did not exhibit a monotonic exposure response relationship, since at 8 weeks activity for all  
4  
5 experimental groups was not markedly different from controls.  
6

7  
8 GR activity (Figure 5) was enhanced (2.8-3.9 fold) in liver of fish fed with Ax, Cx and both  
9  
10 combined carotenoids at 8 weeks, whereas in kidney activity was slight higher in Ax- and Cx-  
11  
12 treated specimens, when compared to own controls.  
13

14  
15 GST activity (Figure 6) increased about 2-3 times in liver and kidney of fish exposed to both  
16  
17 carotenoids for 8 weeks. When the two compounds were co-administered, activity in liver was at  
18  
19 the same as when each molecule was administered singly. On the contrary, mainly at 8 weeks GST  
20  
21 activity in kidney of Ax+Cx fed trout was significantly higher than for Ax or Cx groups.  
22

23  
24 No significant changes of thiol concentrations (Figure 7) were recorded in liver among all  
25  
26 carotenoids treated groups. Higher total glutathione levels were recorded in kidney of Ax + Cx  
27  
28 group, if compared with control (30%) and Ax-trout (40%) at 4 weeks. All three carotenoid treated  
29  
30 groups showed increased thiol levels at 8 weeks, when compared own control (1.8-2.2-fold) and the  
31  
32 same experimental groups (1.5-2-fold) at 4 weeks, with the exception of the Ax+Cx one (Figure 7).  
33  
34  
35  
36

### 37 **Discussion**

38  
39  
40 In our study, Ax and Cx fed continuously for eight weeks to rainbow trout, individually or  
41  
42 concurrently, exerted no noticeable effects on length and weight. Our outcome agrees with previous  
43  
44 findings on *O. mykiss* (Řehulka 2000; Choi et al., 2016; Rahman et al., 2016) as well as on Atlantic  
45  
46 salmon, *Salmo salar* (Bell et al., 2000) or gilthead seabream, *Sparus aurata* (Gomes et al., 2002)  
47  
48 treated with astaxanthin. However, both Ax and Cx diets stimulated different responses of oxidative  
49  
50 stress biomarkers in liver and kidney of xanthophyll-fed trout.  
51  
52

53  
54 Carotenoids play an essential role in protection of cell membranes and lipoproteins from oxidative  
55  
56 damage, offsetting the effects of peroxy radicals (Angeles and Chien, 2016; Dose et al., 2016;  
57  
58 Esatbeyoglu and Rimbach, 2017). The improvement of total antioxidant ability of trout following  
59  
60 single administration of 100 mg/kg astaxanthin or cantaxanthin or combined (50 mg/kg Ax + 50

1  
2  
3 mg/kg Cx) for 60 days was previously reported by Cui et al. (2009). Moreover, rats fed with  
4  
5 canthaxanthin and  $\beta$ -carotene for 6 weeks showed higher GPx and catalase activity in erythrocytes,  
6  
7 when compared to rats fed diets with a high cholesterol and lipid content (CC) (Shih et al., 2008).  
8  
9 Conversely, Ax lowered SOD in juvenile tiger prawn *Penaeus monodon* or SOD and GPx activity  
10  
11 in *Hyphessobrycon callistus* (Chien et al., 2003; Wang et al., 2006).  
12  
13

14 In the present study, biomarkers response may corroborate the antioxidant property of xanthophylls,  
15  
16 mainly for Cx and Ax + Cx-dosed specimens. Indeed, although the unchanged hepatic and renal  
17  
18 MDA concentration during the eight weeks represent a reassuring outcome, detoxifying biomarker  
19  
20 levels in Ax and/or Cx fed rainbow trout suggest that different mechanisms may be involved in  
21  
22 strengthening the antioxidant shield in both tissues and through the two experimental endpoints. In  
23  
24 liver, the transient raise of SOD concentration was followed by the unchanged activity of both main  
25  
26  $H_2O_2$  scavengers, such as catalase and GPx. This outcome is surprising, since the high SOD levels  
27  
28 in xanthophylls-dosed rainbow trout suggest an increase of  $H_2O_2$  hepatic load. Hydrogen peroxide  
29  
30 is a strong oxidizer that can reduce cell viability due to membrane oxidative damage (Xie et al.,  
31  
32 2017). Therefore, the unaltered lipid peroxidation process contributes to outline a more intriguing  
33  
34 scenario. In particular, astaxanthin is recognized as effective molecule in preventing lipid  
35  
36 peroxidation (Dose et al., 2016; Xie et al., 2017) and in lowering MDA concentration in *O. mykiss*  
37  
38 (Brambilla et al., 2009). According to these results we measured a slight lowering of MDA levels in  
39  
40 Ax and Ax + Cx-treated rainbow trout liver, although our values did not reach the statistical  
41  
42 significant differences. Lack of lipid peroxidation process in both tissues of rainbow trout fed with  
43  
44 xanthophyll diets along with high levels of the main antioxidant biomarkers suggest an alternative  
45  
46 route for CAT and GPx in scavenging  $H_2O_2$ . Previous studies suggested that carotenoids, and in  
47  
48 particular canthaxanthin, can act as radical scavenger by quenching free radicals in unsaturated fatty  
49  
50 acids systems (Daubrawa et al., 2005; Mansour et al., 2006). A mechanistic study focused on the  
51  
52 antioxidant role of carotenoids against the hydroperoxides indicated that astaxanthin and  
53  
54 canthaxanthin are able to retard the hydroperoxide formation on azo-initiated lipid peroxidation.  
55  
56  
57  
58  
59  
60

1  
2  
3 Linkage of carbonyl groups at the 4 and 4' positions that is different from other carotenoids may  
4  
5 play a key role in increasing their efficiency to trap peroxy radicals. Indeed, Ax and Cx carbonyl  
6  
7 oxygen has the property to attract electrons, hence reducing the unpaired electron density  
8  
9 throughout the carbon skeleton of radical molecules. As a consequence, the carbon-centred radical  
10  
11 of hydroperoxides reduces its affinity for oxygen (Mortensen et al., 1997). Although H<sub>2</sub>O<sub>2</sub> may  
12  
13 trigger lipid peroxidation and weaken the total antioxidant capacity, xanthophylls can act as a direct  
14  
15 oxyradical scavenger, as previously reported in golden pompano (*Trachinotus ovatus*) exposed to  
16  
17 astaxanthin (Xie et al., 2017). Unexpected response was also observed in kidney, where increasing  
18  
19 GPx activity, mainly in Cx and Ax + Cx-fed rainbow trout, was not accompanied by a rising SOD  
20  
21 level. Moreover, transcription of several antioxidant enzymes, such as SOD, CAT and GPx, can be  
22  
23 regulated by the NF-E2-related nuclear factor 2 (Nrf2) in fish (Jiang et al., 2014). Furthermore, a  
24  
25 previous study showed that xanthophylls may modulate the mRNA expression of Nrf2 and  
26  
27 associated genes, and an optimal Nrf2-mediated oxidative defense can be achieved in HepG2 cell  
28  
29 line following exposure at low (12 µM) astaxanthin concentration (Saw et al., 2013). Therefore, it is  
30  
31 feasible that the enhancement of renal GPx activity was related to the ability of xanthophyll to  
32  
33 modulate the Nrf-2 mRNA expression, reinforcing the antioxidant shield of *O. mykiss*.  
34  
35  
36  
37  
38

39  
40 Astaxanthin and canthaxanthin have been documented as weak inducers of the xenobiotic-  
41  
42 metabolizing enzyme GST in liver or kidney of *O. mykiss* after short term treatment (3 weeks)  
43  
44 (Page and Davies, 2002). Indeed, when dietary substances are absorbed at a higher rate than  
45  
46 catabolism and excretion, astaxanthin and canthaxanthin can gradually accumulate in tissues. Brizio  
47  
48 et al. (2013) reported muscle concentration of 2.2 and 3.70 mg/kg for Ax or 0.4 and 1.21 mg/kg for  
49  
50 Cx, at 4 and 8 weeks, respectively. Moreover, both carotenoids are metabolized reductively, and  
51  
52 geometrical E/Z-isomers of astaxanthin are utilized differently in rainbow trout. Although, all-E-  
53  
54 astaxanthin accumulates in plasma and muscle, while 13Z-astaxanthin accumulates in liver (Ahmadi  
55  
56 et al., 2006; Østerlie et al., 1999). In our study, no changes of GST activity were measured in  
57  
58 xanthophyll-treated fish during the first 4 weeks of the trial, while enzymes level increased  
59  
60

1  
2  
3 noticeably mainly in liver after eight weeks. Although Ax and Cx concentrations were not measured  
4  
5 in all tissues of rainbow trout, we may assume that the activation of this important defense line  
6  
7 occurred along with higher accumulation levels of carotenoids or metabolites production.  
8

9  
10 Additionally, it is ascertained that antioxidant compounds may act as prooxidant factors when their  
11  
12 concentration exceeds the optimal threshold (Dörr et al., 2008, 2013; Elia et al., 2011, 2014; Pacini  
13  
14 et al., 2012). Accordingly, previous studies showed that high carotenoid concentrations may  
15  
16 produce oxidative effects, such as alteration of the biological membrane properties and permeability  
17  
18 to toxins, due to production of radical molecules (Stahl and Sies, 2003; El-Agamey et al., 2004).  
19  
20 Furthermore, the highly reactive carotenoid-peroxyl molecules can impair the activity of antioxidant  
21  
22 enzymes or cause lipid peroxidation (Lygren et al. 1999; El-Agamey et al., 2004). However,  
23  
24 astaxanthin can activate the extracellular signal-regulated protein kinase (ERK) signaling pathway,  
25  
26 following an up-regulation of expression of GST- $\alpha$ 1, strengthen the biological defense against  
27  
28 oxidative stress damages *in vitro* and *in vivo* (Fakhri et al., 2018). The induced GST activity in both  
29  
30 rainbow trout tissues in this study could thus suggest a regulatory mechanism preventing Ax and/or  
31  
32 Cx disturbance. Accordingly, unchanged activity of hepatic SOD and renal GPx during the last  
33  
34 endpoint might have been balanced by higher GST levels offsetting the potential adverse effects of  
35  
36 xanthophylls. Particularly, GST was differently supported in both analyzed tissues. While in liver  
37  
38 the suitable GSH amount was ensured by the increased GR activity, in kidney an *ex novo*  
39  
40 production of reduced thiol may be assumed, as suggested by the raised concentrations of total  
41  
42 glutathione and the unchanged GR levels. Astaxanthin has also been described as enhancing factor  
43  
44 of GR activity in giant freshwater prawn *Macrobrachium rosenbergii* following challenging  
45  
46 conditions (Angeles et al., 2009). The role of GSH as enzymatic co-factor or radical scavenger may  
47  
48 protect tissues from free radical damages (Saw et al., 2013). Furthermore, previous study showed  
49  
50 that Ax in combination with fish oil enhances the immune response by improving the glutathione-  
51  
52 based redox balance in rat plasma and neutrophils (Barros et al., 2012). This evidence could suggest  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 that in this study glutathione could act not only as antioxidant protector, but also reinforce promptly  
4  
5 the immune system defense.  
6  
7  
8  
9

## 10 **Conclusions**

11  
12 Astaxanthin and canthaxanthin stimulate the antioxidant defense in liver and kidney of rainbow  
13  
14 trout involving different time-dependent routes. Furthermore, both combined xanthophylls diets did  
15  
16 not exert additive effects in the investigated rainbow trout tissues throughout the trial. The current  
17  
18 findings may deepen knowledge on the prooxidant effects of the considered carotenoids widely  
19  
20 used in aquaculture, as safe additives, in order to sustain fish health and productivity.  
21  
22  
23  
24  
25

## 26 **Acknowledgements**

27  
28 This research was funded by the Italian Ministry of Health (grant n. IZS PLV 21/08 RC).  
29  
30  
31

## 32 **References**

33  
34 Ahmadi, M. R., A. A. Bazyar, S.Safi , T. Ytrestøyl, and B. Bjerkeng. 2006. Effects of dietary  
35  
36 astaxanthin supplementation on reproductive characteristics of rainbow trout (*Oncorhynchus*  
37  
38 *mykiss*). *Journal of Applied Ichthyology* 22: 388-394.  
39  
40  
41

42 Angeles, I.P., Y.H. Chien, and A.V. Yambot. 2009. Effect of injected astaxanthin on survival,  
43  
44 antioxidant capacity, and immune response of the giant freshwater prawn  
45  
46 *Macrobrachium rosenbergii* (de Man, 1879) challenged with *Lactococcus garvieae*. *Journal of*  
47  
48 *Shellfish Research* 28: 931–937.  
49  
50  
51

52 Angeles, I.P., and Y.H. Chien. 2016. Antioxidant capacity and metabolic response of red striped  
53  
54 snapper *Lutjanus erythropterus* fed diet with astaxanthin and/or oxidized oil. *Journal of Marine*  
55  
56 *Science and Technology* 24: 338-346.  
57  
58  
59  
60

- 1  
2  
3 Barros, M.P., D.P. Marin, A.P. Bolin, de P. Cassia Santos Macedo, T.R. Campoio, C.Jr. Fineto,  
4 B.A., Guerra, T.G. Polotow, C. Vardaris, R. Mattei, and R. Otton. 2012. Combined astaxanthin  
5 and fish oil supplementation improves glutathione-based redox balance in rat plasma and  
6 neutrophils. *Chemico-Biological Interaction* 197: 58-67.  
7  
8  
9  
10  
11  
12  
13 Bell, J.G., J. McEvoy, D.R. Tocher, and J.R. Sargent. 2000. Depletion of  $\alpha$ -tocopherol and  
14 astaxanthin in Atlantic salmon (*Salmo salar*) affects autoxidative defense and fatty acid  
15 metabolism. *Journal of Nutrition* 130: 1800-1808.  
16  
17  
18  
19  
20  
21 Brambilla, F., A. Forchino, M. Antonini, S. Rimoldi, G. Terova, and M. Saroglia. 2009. Effect of  
22 dietary Astaxanthin sources supplementation on muscle pigmentation and lipid peroxidation in  
23 rainbow trout (*Oncorhynchus mykiss*). Italian Journal of Animal Science 8(2), Proceedings of  
24 the 18th ASPA Congress, Palermo, June 9-12, 2009.  
25  
26  
27  
28  
29  
30  
31 Brizio, P., A. Benedetto, M. Righetti, M. Prearo, L. Gasco, S. Squadrone, and M.C. Abete. 2013.  
32 Astaxanthin and canthaxanthin (xanthophyll) as supplements in rainbow trout diet: in vivo  
33 assessment of residual levels and contributions to human health. *Journal of Agricultural and*  
34 *Food Chemistry* 61: 10954–1095.  
35  
36  
37  
38  
39  
40  
41 Chew. B.P., 1995. Antioxidant vitamins affect food animal immunity and health. *Journal of*  
42 *Nutrition* 125: 1804S-1808S.  
43  
44  
45  
46  
47 Chien Y.H., C.H. Pan , and B. Hunter. 2003. The resistance to physical stresses by *Penaeus*  
48 *monodon* juveniles fed diets supplemented with astaxanthin. *Aquaculture* 216: 177–191.  
49  
50  
51  
52 Choi, J., M.M. Rahman, S.Y. Lee, K.H. Chang, and S.M. Lee. 2016. Effects of Dietary Inclusion of  
53 Fermented Soybean Meal with Phaffiarhodozyma on Growth, Muscle Pigmentation, and  
54 Antioxidant Activity of Juvenile Rainbow Trout (*Oncorhynchus mykiss*). *Turkish Journal of*  
55 *Fisheries and Aquatic Sciences* 16: 91-101.  
56  
57  
58  
59  
60

- 1  
2  
3 Commission Implementing Regulation (EU) 2015/1415 of 20 August 2015 concerning the  
4 authorisation of astaxanthin as a feed additive for fish, crustaceans and ornamental fish (Text  
5 with EEA relevance). [http://data.europa.eu/eli/reg\\_impl/2015/1415/oj](http://data.europa.eu/eli/reg_impl/2015/1415/oj)  
6  
7  
8  
9  
10 Commission Directive (EC) No 2003/7 of 24 January 2003 amending the conditions for  
11 authorization of canthaxanthin in feeding stuffs in accordance with Council Directive  
12 70/524/EEC .  
13  
14  
15  
16  
17  
18  
19 Conn, P.F., W. Schalch, and T.G. Truscott. 1991. The singlet oxygen carotenoid interaction.  
20 *Journal of Photochemistry and Photobiology B* 11: 41–47.  
21  
22  
23  
24 Cui, W., X. Leng, X. Li, X. Li, and J. Xu. 2009. Effects of astaxanthin and canthaxanthin on  
25 pigmentation of muscle and total antioxidant capacity of liver in rainbow trout (*Oncorhynchus*  
26 *mykiss*). *Journal of Fisheries of China* 33: 987-995.  
27  
28  
29  
30  
31 Daubrawa, F., H. Sies, and W. Stahl. 2005. Astaxanthin diminishes gap junctional intercellular  
32 communication in primary human fibroblasts. *Journal of Nutrition* 135: 2507-2511.  
33  
34  
35  
36  
37 Dörr, A.J.M., N. Pacini, M.C. Abete, M. Prearo, and A.C. Elia. 2008. Effects of a selenium-  
38 enriched diet on antioxidant response in adult crayfish (*Procambarus clarkii*). *Chemosphere* 73:  
39 1090-1095.  
40  
41  
42  
43  
44  
45 Dörr, A.J.M., M.C. Abete, M. Prearo, N. Pacini, G. La Porta, M. Natali, and A.C. Elia. 2013.  
46 Effects of selenium supplemented diets on growth and condition indexes in juvenile red swamp  
47 crayfish, *Procambarus clarkii*. *Environmental Toxicology and Pharmacology* 36: 484-492.  
48  
49  
50  
51  
52  
53 Dose, J., S. Matsugo, H. Yokokawa, Y. Koshida, S. Okazaki, U. Seidel, M. Eggersdorfer, G.  
54 Rimbach, and T. Esatbeyoglu. 2016. Free radical scavenging and cellular antioxidant properties  
55 of astaxanthin. *International Journal of Molecular Sciences* 17: 103.  
56  
57  
58  
59  
60



- 1  
2  
3 El-Agamey, A., G.M. Lowe, D.J. McGarvey, A. Mortensen, D.M. Phillip, T.G. Truscott, and A.J.  
4  
5 Young. 2004. Carotenoid radical chemistry and antioxidant/pro-oxidant properties. *Archives of*  
6  
7 *Biochemistry and Biophysics* 430: 37–48.  
8  
9  
10 Elia, A.C., M. Prearo, N. Pacini, A.J.M. Dörr, and M.C. Abete. 2011. Effects of selenium diets on  
11  
12 growth, accumulation and antioxidant response in juvenile carp. *Ecotoxicology and*  
13  
14 *Environmental Safety* 74: 166-173.  
15  
16  
17 Elia, A.C., M.C. Abete, A.J.M. Dörr, T. Scanzio, and M. Prearo. 2014. Antioxidant biomarker  
18  
19 survey ensuing long-term selenium withdrawal in *Acipenser baeri* fed Se-cysteine diets.  
20  
21 *Environmental Toxicology and Pharmacology* 37: 1131-1139.  
22  
23  
24 Elia, A.C., G. Magara, M. Righetti, A.J.M. Dörr, T. Scanzio, N. Pacini, M.C. Abete, and M. Prearo.  
25  
26 2017. Oxidative stress and related biomarkers in cupric and cuprous chloride-treated rainbow  
27  
28 trout. *Environmental Science and Pollution Research* 24: 10205-10219.  
29  
30  
31  
32  
33 Esatbeyoglu, T., and G. Rimbach. 2017. Canthaxanthin: from molecule to function. *Molecular*  
34  
35 *Nutrition and Food Research* 61: 6.  
36  
37  
38  
39 Fakhri, S., F. Abbaszadeh, L. Dargahi, M. Jorjani. 2018. Astaxanthin: a mechanistic review on its  
40  
41 biological activities and health benefits. *Pharmacology Research* 136: 1-20.  
42  
43  
44  
45 Gomes, E., J. Dias, P. Silva, L. Valente, J. Empis, L. Gouveia, J. Bowen, and A. Young. 2002.  
46  
47 Utilization of natural and synthetic sources of carotenoids in the skin pigmentation of gilthead  
48  
49 seabream (*Sparus aurata*). *European Food Research and Technology* 214: 287-293.  
50  
51  
52  
53 Henmi, H., M. Hata, and M. Takeuchi. 1991. Studies on the carotenoids in the muscle of salmon-V.  
54  
55 Combination of astaxanthin and canthaxanthin with bovine serum albumin and egg albumin.  
56  
57 *Comparative Biochemistry and Physiology B* 99: 609-612.  
58  
59  
60

- 1  
2  
3 Jiang, W.D., Y. Liu, K. Hu, J. Jiang, S.H. Li, L. Feng, and X.Q. Zhou. 2014. Copper exposure  
4 induces oxidative injury, disturbs the antioxidant system and changes the Nrf2/ARE  
5 (CuZnSOD) signaling in the fish brain: protective effects of myo-inositol. *Aquatic Toxicology*  
6  
7 155: 301-313.  
8  
9  
10  
11  
12  
13 Lowry, O.H., N.L. Rosebrough, and R.I. Randall. 1951. Protein measurement with Folin phenol  
14 reagent. *Journal of Biological Chemistry* 193: 265- 275.  
15  
16  
17  
18 Lygren, B., K. Hamre, and R. Waagboe. 1999. Effects of dietary pro- and antioxidants on some  
19 protective mechanisms and health parameters in Atlantic salmon. *Journal of Aquatic Animal*  
20  
21 *Health* 11: 211 -221.  
22  
23  
24  
25  
26 Mansour, N., M.A. McNiven, and G.F. Richardson. 2006. The effects of dietary supplementation  
27 with blueberry, a-tocopherol or astaxanthin on oxidative stability of Arctic char (*Salvelinus*  
28 *alpinus*) semen. *Theriogenology* 66: 373-382.  
29  
30  
31  
32  
33  
34 Martínez-Álvarez, R.M., A.E. Morales, and A. Sanz. 2005. Antioxidant defenses in fish: biotic and  
35 abiotic factors. *Reviews in Fish Biology and Fisheries* 15: 75-88.  
36  
37  
38  
39 Mortensen, A., L.H. Skibsteda, J. Sampson, C. Rice-Evans, and S.A. Everet. 1997. Comparative  
40 mechanisms and rates of free radical scavenging by carotenoid antioxidants. *FEBS Letters* 418:  
41  
42 91-97.  
43  
44  
45  
46  
47 McCord, J.M., and I. Fridovich. 1969. Superoxide dismutase: an enzymatic function for  
48 erythrocyte hemocuprein (hemocuprein). *Journal of Biological Chemistry* 244: 6049-6055.  
49  
50  
51  
52 Nakano, T., T. Kanmuri, M. Sato, and M. Takeuchi. 1999. Effect of astaxanthin rich red yeast  
53 (*Phaffia rhodozyma*) on oxidative stress in rainbow trout. *Biochimica et Biophysica Acta* 1426:  
54  
55 119-125.  
56  
57  
58  
59  
60

- 1  
2  
3 Østerlie, M., B. Bjerkeng, and S. Liaaen-Jensen. 1999. Accumulation of astaxanthin all-E, 9Z and  
4 13Z geometrical isomers and 3 and 3 $\phi$  RS optical isomers in rainbow trout (*Oncorhynchus*  
5 *mykiss*) is selective. *Journal of Nutrition* 129: 391–398.  
6  
7  
8  
9  
10 Pacini, N., M.C. Abete, A.J.M. Dörr, M. Prearo, M. Natali, and A.C. Elia. 2012. Detoxifying  
11 response in juvenile tench fed by selenium diet. *Environmental Toxicology and Pharmacology*  
12 33: 46-52.  
13  
14  
15  
16  
17  
18 Pacini, N., M. Prearo, M.C. Abete, P. Brizio, A.J.M. Dörr, R. Reimschuessel, W. Andersen, L.  
19 Gasco, M. Righetti, and A.C. Elia. 2013. Antioxidant responses and renal crystal formation in  
20 rainbow trout treated with melamine administered individually or in combination with cyanuric  
21 acid. *Journal of Toxicology and Environmental Health, Part A* 76: 491-508.  
22  
23  
24  
25  
26  
27  
28 Page, G.I., and S.J. Davies. 2002. Astaxanthin and canthaxanthin do not induce liver or kidney  
29 xenobiotic-metabolizing enzymes in rainbow trout (*Oncorhynchus mykiss* Walbaum).  
30 *Comparative Biochemistry and Physiology C* 133: 443–451.  
31  
32  
33  
34  
35  
36 Rahman, M.M., S. Khosravi, K.H. Chang, and S.M. Lee. 2016. Effects of dietary inclusion of  
37 Astaxanthin on growth, muscle pigmentation and antioxidant capacity of juvenile Rainbow  
38 Trout (*Oncorhynchus mykiss*). *Preventive Nutrition and Food Science* 21: 281-288.  
39  
40  
41  
42  
43  
44 Řehulka, J. 2000. Influence of astaxanthin on growth rate, condition, and some blood indices of  
45 rainbow trout, *Oncorhynchus mykiss*. *Aquaculture* 190: 27-47.  
46  
47  
48  
49 Saw, C.L.L., A.Y. Yang, Y. Guo, and A.N.T. Kong. 2013. Astaxanthin and omega-3 fatty acids  
50 individually and in combination protect against oxidative stress via the Nrf2-ARE pathway.  
51 *Food Chemistry and Technology* 62: 869-875.  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 Shih, C.K., J.H. Chang, S.H. Yang, T.W. Chou, and H.H. Cheng. 2008.  $\beta$ -Carotene and  
4 canthaxanthin alter the pro-oxidation and antioxidation balance in rats fed a high-cholesterol  
5 and high-fat diet. *British Journal of Nutrition* 99: 59-66.  
6  
7  
8  
9  
10 Shimidzu, N., M. Goto, and W. Miki. 1996. Carotenoids as singlet oxygen quenchers in marine  
11 organisms. *Fisheries Science* 62: 134–137.  
12  
13  
14  
15  
16 Sies, H., and W. Stahl. 1995. Vitamins E and C, beta-carotene, and other carotenoids as  
17 antioxidants. *The American Journal of Clinical Nutrition* 62: 1315S–1321S.  
18  
19  
20  
21  
22 Stahl, W., and H. Sies. 2003. Antioxidant activity of carotenoids. *Molecular Aspects of Medicine*  
23 24: 345–351.  
24  
25  
26  
27 Wang, Y.J., Y.H. Chien, and C.H. Pan. 2006. Effects of dietary supplementation of carotenoids on  
28 survival, growth, pigmentation, and antioxidant capacity of characins, *Hyphessobrycon*  
29 *callistus*. *Aquaculture* 261: 641 – 648.  
30  
31  
32  
33  
34  
35 Xie, J.J., X. Chen, J. Niu., J. Wang, Y. Wang, and Q. Liu. 2017. Effects of astaxanthin on  
36 antioxidant capacity of golden pompano (*Trachinotus ovatus*) in vivo and in vitro. *Fisheries and*  
37 *Aquatic Sciences* 20: 6.  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Table 1- Length and weight of treated rainbow trout fed astaxanthin (Ax) and/or canthaxanthin (Cx) diets

	Length	Weight
C T1	31.05 ± 0.83 <sup>a*</sup>	306.5 ± 26.57 <sup>a</sup>
Ax T1	31.7 ± 1.00 <sup>a</sup>	329.5 ± 32.1 <sup>a</sup>
Cx T1	31.25 ± 1.11 <sup>a*</sup>	314.5 ± 22.66 <sup>a*</sup>
Ax + Cx T1	31.5 ± 0.88 <sup>a</sup>	315 ± 23.57 <sup>a</sup>
C T2	32.65 ± 1.44 <sup>a*</sup>	337.5 ± 48.95 <sup>a</sup>
Ax T2	33.1 ± 1.24 <sup>a</sup>	356 ± 45.39 <sup>a</sup>
Cx T2	33.7 ± 1.25 <sup>a*</sup>	367 ± 42.24 <sup>a*</sup>
Ax + Cx T2	33.2 ± 0.82 <sup>a</sup>	358.5 ± 32.83 <sup>a</sup>

The results are reported as mean and standard deviation. Statistical significant differences ( $p < 0.05$ ) are indicated by the following symbols: between the same group at both experimental times, asterisk (\*), and between Ax, Cx and Ax + Cx samples at the same endpoint, different letters (a, b, c). T1= 4 weeks; T2 = 8 weeks.

## Legend of Figures

**Figure 1.** Malondialdehyde concentration in rainbow trout fed Ax (astaxanthin), Cx (canthaxanthin), and Ax + Cx diets for 8 weeks. Liver (L) or kidney (K) of each group was analyzed individually ( $n = 10$ ). The results are reported as mean and standard deviation. Statistical significant differences ( $p < 0.05$ ) are indicated by the following symbols: between the same group at both experimental times, asterisk (\*), and between Ax, Cx and Ax + Cx samples at the same endpoint, different letters (a, b, c). T1= 4 weeks; T2 = 8 weeks.

**Figure 2.** Superoxide dismutase activity in rainbow trout fed Ax (astaxanthin), Cx (canthaxanthin), and Ax + Cx diets for 8 weeks. Liver (L) or kidney (K) of each group was analyzed individually ( $n = 10$ ). The results are reported as mean and standard deviation. Statistical significant differences ( $p < 0.05$ ) are indicated by the following symbols: between the same group of both experimental times, asterisk (\*), and between Ax, Cx and Ax + Cx samples at the same endpoint, different letters (a, b, c). T1= 4 weeks; T2 = 8 weeks.

**Figure 3.** Catalase activity in rainbow trout fed Ax (astaxanthin), Cx (canthaxanthin), and Ax + Cx diets for 8 weeks. Liver (L) or kidney (K) of each group was analyzed individually ( $n = 10$ ). The results are reported as mean and standard deviation. Statistical significant differences ( $p < 0.05$ ) are indicated by the following symbols: between the same group of both experimental times, asterisk (\*), and between Ax, Cx and Ax + Cx samples at the same endpoint, different letters (a, b, c). T1= 4 weeks; T2 = 8 weeks.

**Figure 4.** Glutathione peroxidase activity in rainbow trout fed Ax (astaxanthin), Cx (canthaxanthin), and Ax + Cx diets for 8 weeks. Liver (L) or kidney (K) of each group was analyzed individually ( $n = 10$ ). The results are reported as mean and standard deviation. Statistical significant differences ( $p < 0.05$ ) are indicated by the following symbols: between the same group of

1  
2  
3 both experimental times, asterisk (\*), and between Ax, Cx and Ax + Cx samples at the same  
4  
5 endpoint, different letters (a, b, c). T1= 4 weeks; T2 = 8 weeks.  
6  
7  
8  
9

10 **Figure 5.** Glutathione reductase activity in rainbow trout fed Ax (astaxanthin), Cx (canthaxanthin),  
11 and Ax + Cx diets for 8 weeks. Liver (L) or kidney (K) of each group was analyzed individually ( $n$   
12 = 10). The results are reported as mean and standard deviation. Statistical significant differences ( $p$   
13 <0.05) are indicated by the following symbols: between the same group of both experimental times,  
14 asterisk (\*), and between Ax, Cx and Ax + Cx samples at the same endpoint, different letters (a, b,  
15 c). T1= 4 weeks; T2 = 8 weeks.  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25

26 **Figure 6.** Glutathione S-transferase activity in rainbow trout fed Ax (astaxanthin), Cx  
27 (canthaxanthin), and Ax + Cx diets for 8 weeks. Liver (L) or kidney (K) of each group was  
28 analyzed individually ( $n = 10$ ). The results are reported as mean and standard deviation. Statistical  
29 significant differences ( $p <0.05$ ) are indicated by the following symbols: between the same group of  
30 both experimental times, asterisk (\*), and between Ax, Cx and Ax + Cx samples at the same  
31 endpoint, different letters (a, b, c). T1= 4 weeks; T2 = 8 weeks.  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41

42 **Figure 7.** Total glutathione concentration in rainbow trout fed Ax (astaxanthin), Cx  
43 (canthaxanthin), and Ax + Cx diets for 8 weeks. Liver (L) or kidney (K) of each group was  
44 analyzed individually ( $n = 10$ ). The results are reported as mean and standard deviation. Statistical  
45 significant differences ( $p <0.05$ ) are indicated by the following symbols: between the same group of  
46 both experimental times, asterisk (\*), and between Ax, Cx and Ax + Cx samples at the same  
47 endpoint, different letters (a, b, c). T1= 4 weeks; T2 = 8 weeks.  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




Figure 1

245x122mm (300 x 300 DPI)






Figure 2

245x120mm (300 x 300 DPI)




Figure 3

248x120mm (300 x 300 DPI)




Figure 4

245x120mm (300 x 300 DPI)




Figure 5

247x120mm (300 x 300 DPI)




Figure 6

250x120mm (300 x 300 DPI)




Figure 7

250x120mm (300 x 300 DPI)

Table 1- Length and weight of treated rainbow trout fed astaxanthin (Ax) and/or canthaxanthin (Cx) diets

	Length	Weight
C T1	31.05 ± 0.83 <sup>a*</sup>	306.5 ± 26.57 <sup>a</sup>
Ax T1	31.7 ± 1.00 <sup>a</sup>	329.5 ± 32.1 <sup>a</sup>
Cx T1	31.25 ± 1.11 <sup>a*</sup>	314.5 ± 22.66 <sup>a*</sup>
Ax + Cx T1	31.5 ± 0.88 <sup>a</sup>	315 ± 23.57 <sup>a</sup>
C T2	32.65 ± 1.44 <sup>a*</sup>	337.5 ± 48.95 <sup>a</sup>
Ax T2	33.1 ± 1.24 <sup>a</sup>	356 ± 45.39 <sup>a</sup>
Cx T2	33.7 ± 1.25 <sup>a*</sup>	367 ± 42.24 <sup>a*</sup>
Ax + Cx T2	33.2 ± 0.82 <sup>a</sup>	358.5 ± 32.83 <sup>a</sup>

The results are reported as mean and standard deviation. Statistical significant differences ( $p < 0.05$ ) are indicated by the following symbols: between the same group at both experimental times, asterisk (\*), and between Ax, Cx and Ax + Cx samples at the same endpoint, different letters (a, b, c). T1= 4 weeks; T2 = 8 weeks.