

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Highly defatted insect meal in Siberian sturgeon juveniles feeds

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1723444> since 2020-01-16T15:09:40Z

Publisher:

EAAP scientific committee

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Highly defatted insect meal in Siberian sturgeon juveniles feeds

Gasco L.^{1,4}, Schiavone A.², Renna M.¹, Lussiana C.¹, Dabbou S.², Meneguz M.¹, Malfatto V.¹, Prearo M.³, Bonaldo A.⁴, Zoccarato I.¹, Gai F.⁵

¹Department of Agricultural, Forest and Food Sciences. University of Turin. Italy

²Department of Veterinary Sciences. University of Turin. Italy

³Veterinary Medical Research Institute for Piemonte, Liguria and Valle D'Aosta. Italy

⁴Department of Veterinary Medical Sciences. University of Bologna. Italy

⁵Institute of Science of Food Production, National Research Council. Italy

Corresponding author: laura.gasco@unito.it

Recent investigations highlighted that insect protein meals can be a more sustainable alternative to conventional protein used so far in aquaculture. *Hermetia illucens* (HI) is a good candidate due to its valuable nutritional properties. The aim of this research was to evaluate the effects of fishmeal (FM) substitution by a highly defatted HI larvae meal in sturgeon juveniles feeds.

Four diets were formulated: a control (70% of FM - CF), two diets where FM was replaced by 25 (HI25) and 50% (HI50) of HI and a vegetable protein based diet without HI (CV).

352 *Acipenser baerii* were distributed in 16 fiberglass tanks. Each diet was assigned to 4 groups of 22 fish and feed was distributed to apparent satiation.

At the end of the trial (118 days) fish growth performances were calculated, and whole body (WBC) proximate and fatty acid (FA) composition were analyzed.

Data were statistically analyzed by one way ANOVA. Significance level was set at $P < 0.05$.

Results indicate that the inclusion of HI affected fish performances and WBC. Generally, up to 25% of FM substitution, fish performance was comparable to those of fish fed CF or CV while the 50% substitution induced a worsening of performance parameters and the same trend was observed for WBC. Lauric acid and total saturated FA contents were higher in fish fed HI when compared to CF and CV groups. Monounsaturated, polyunsaturated, and n3 and n6 FA contents showed differences among groups, with lowest values for CF; however, no differences were found in the n3/n6 FA ratio in WB of CF and HI25 groups.