

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

HPV8 Field Cancerization in a Transgenic Mouse Model Is due to Lrig1+ Keratinocyte Stem Cell Expansion

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1660828> since 2018-02-27T10:55:49Z

Published version:

DOI:10.1016/j.jid.2017.04.039

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

HPV8 Field Cancerization in a Transgenic Mouse Model is due to Lrig1+ Keratinocyte Stem Cell Expansion

Simone Lanfredini^{1*}, Carlotta Olivero^{2*}, Cinzia Borgogna², Federica Calati², Kathryn Powell¹, Kelli-Jo Davies¹, Marco De Andrea^{2,3}, Sarah Harries¹, Hiu Kwan Carolyn Tang¹, Herbert Pfister⁴, Marisa Gariglio², Girish K Patel¹.

1. European Cancer Stem Cell Research Institute, School of Biosciences, Cardiff University, Maindy Road, Cardiff, UK

2. Virology Unit, Department of Translational Medicine, Novara Medical School, Novara, Italy

3. Viral Pathogenesis Unit, Department of Public Health and Pediatric Sciences, Turin Medical School, Turin, Italy

4. Institute of Virology, University of Cologne, Cologne, Germany

* The first two authors contributed equally to this work.

** Marisa Gariglio and Girish K Patel are joint senior authors.

Reprint request and correspondence to: Dr Girish K Patel MD, European Cancer Stem Cell Research Institute, School of Biosciences, Cardiff University, Maindy Road, Cardiff, CF24 4HQ, UK. E-mail: patelgk@cardiff.ac.uk

Short title: HPV8 induces keratinocyte stem cell expansion

Key words: Human Papillomavirus, Field Cancerization, Squamous cell carcinoma, Keratinocyte stem cells

Abbreviations: human papillomavirus (HPV), Epidermodysplasia Verruciformis (EV), cutaneous squamous cell carcinoma (SCC), wild type (WT), hair follicle (HF), interfollicular epidermis (IFE), keratinocyte stem cell (KSC), actinic keratosis (AK).

Author contributions: GKP and MG designed research; SL, KP, CO, FC, KD, CB, MDA, SH and HKCT performed research; HP contributed new reagents; SL, CO, CB, MG and GKP analyzed data; and SL, CO, MG, and GKP wrote the manuscript.

Word Count: 2868 (excluding references and figure legends)

References: 42

Figures: 6 with legends less than 125 words.

ABSTRACT

β -HPV cause near ubiquitous latent skin infection within long-lived hair follicle keratinocyte stem cells (HF KSC). In patients with epidermodysplasia verruciformis (EV), β -HPV viral replication is associated with skin keratosis and cutaneous squamous cell carcinoma (SCC). To determine the role of HF KSC in β -HPV induced skin carcinogenesis, we utilized a transgenic mouse model in which the keratin 14 promoter drives expression of the entire HPV8 early region (HPV8tg). HPV8tg mice developed thicker skin in comparison to wild type littermates consistent with a hyperproliferative epidermis. HF keratinocyte proliferation was evident within the Lrig1+ KSC population (69 vs 55%, $p < 0.001$, $n = 6$), and not in the CD34+, LGR5+ and LGR6+ KSC populations. This was associated with a 2.8-fold expansion in Lrig1+ keratinocytes and 3.8 fold increased colony forming efficiency. Consistent with this, we observed nuclear p63 expression throughout this population and the HF infundibulum and adjoining IFE, associated with a switch from p63 TA isoforms to Δ Np63 isoforms in HPV8tg skin. EV keratosis and in some cases actinic keratoses demonstrated similar histology associated with β -HPV virus reactivation and nuclear p63 expression within the HF infundibulum and perifollicular epidermis. These findings would suggest that β -HPV field cancerization arises from the HF junctional zone and predispose to SCC.

INTRODUCTION

More than 200 human papillomavirus (HPV) types have been DNA sequenced and thus classified into five genera (*Alpha*-, *Beta*-, *Gamma*-, *Mu*- and *Nu*-papillomavirus) (Bernard *et al.*, 2010; <http://pave.niaid.nih.gov/#home>). High-risk α -HPV types have been established to be causative for cancer, notably in the ano-genital tract (Bosch *et al.*, 2013). In contrast β -HPV types are evolutionarily distinct as they do not integrate into the host genome and cause ubiquitous latent skin infection (Quint *et al.*, 2015). The likely reservoir for β -HPV latent infection is postulated to reside within long-lived hair follicle keratinocyte stem cells (HF-KSC), since plucked hair consistently demonstrates β -HPV DNA and KSC characteristics are enhanced by HPV 5 and 8 early region genes (Boxman *et al.*, 1997; Bouwes Bavinck *et al.*, 2008; Hufbauer *et al.*, 2013). However, the precise HF-KSC populations involved in β -HPV latent infection remain to be defined (Kranjec and Doorbar, 2016). Markers of β -HPV infection are uniformly observed in Epidermodysplasia Verruciformis (EV) keratosis and cutaneous squamous cell carcinoma (SCC), which represents the prototypic model of β -HPV induced skin carcinogenesis (Borgogna *et al.*, 2012, Borgogna *et al.*, 2014a). EV is a rare genodermatosis that has been included in the list of primary immunodeficiency characterized by defects in innate immunity (Notarangelo *et al.*, 2004). Inactivating bi-allelic mutations of either EVER1/TMC6 or EVER2/TMC8 have been described in several families suffering from EV and account for about half of described EV cases (Ramos *et al.*, 2002). Atypical cases of EV have been also reported and they usually display T-cell defects (Azzimonti *et al.*, 2005, Borgogna *et al.*, 2014a, Landini *et al.*, 2014). It has been postulated that β -HPV reactivation is also associated with skin carcinogenesis in organ transplant recipients (Quint *et al.*, 2015; Borgogna *et al.*, 2014b; Howley *et al.*, 2015). To determine the role of HF-KSC in β -HPV induced skin carcinogenesis, we utilized a transgenic mouse model in which the entire HPV8 early region genes are expressed under a keratin 14 promoter (HPV8tg)

(Schaper *et al.*, 2005; De Andrea *et al.*, 2010). HPV8tg mice develop multiple persistent papillomas within 8 weeks, of which 6% go on to develop spontaneous SCC with metastatic potential.

In this study, we provide a model of β -HPV-induced skin carcinogenesis that is based on the aberrant expansion of Lrig1+ KSC in the upper hair follicle that spill out into the surrounding epidermis. Similarly expansion of the junctional zone HF KSC population was identified by p63 labelling in human EV keratosis and actinic keratosis.

RESULTS

The HPV8tg mice skin displays HF proliferative epidermal hyperplasia

After birth, HPV8tg mice develop thicker skin in comparison to wild type (WT) littermates (Figure 1). Adult HPV8tg skin thickness of the ear was 0.6 ± 0.1 vs 0.4 ± 0.1 mm ($p < 0.05$, $n=9$), there was no difference in weight and tail width. More keratinocyte layers were evident in the HF infundibulum and adjoining interfollicular epidermis (IFE) in HPV8tg, 4.2 ± 0.47 vs 2.0 ± 0.0 and 3.8 ± 0.49 vs 1.5 ± 0.43 respectively ($p < 0.01$, $n=5$), but stratum corneum thickness measured on histological sections was not different. Consistent with a hyperproliferative epidermis, keratinocyte proliferation assessed by Ki67 positive cells per basal keratinocyte was markedly increased within the HF (41 ± 10.9 vs 23 ± 11.8 , $n=7$, $p=0.01$) and to a lesser extent the IFE (0.46 ± 0.18 vs 0.31 ± 0.11 , $n=15$, $p=0.01$). The expression of HPV8 early region genes in this transgenic mouse model has been previously described (Schaper *et al.*, 2005; De Andrea *et al.*, 2010). While this HPV8tg mouse model yield spontaneous SCC formation, other similar β -HPV transgenic models driven by K14 promoter do not develop SCC spontaneously (Viarisio *et al.* 2011). Consistent with this, levels of E6 and E7 expression in HPV8tg mouse skin were similar to that observed in HeLa cells with natural HPV18 infection (Supplementary Figure 1). Together these findings

suggested that HPV8 early region genes induce a proliferative epidermal hyperplasia, notably in the HF. HPV8 induced keratinocyte proliferation was greatest in the HF and immediately adjoining IFE, as determined by Ki67 expression, even though keratin 14 promoter driven HPV8 early region genes were uniformly expressed.

The Lrig1 KSC population is expanded in HPV8tg mice

Within the HF, the mean area of the infundibulum was markedly increased in HPV8tg compared to WT mice (Figure 2a), while there was no difference in HF length (Figure 2b).

To determine which hair follicle keratinocyte population become expanded in HPV8tg mice compared to wild type littermates, we labelled skin sections in whole mount analysis with a set of stem cell markers. Consistent with the observed HF infundibulum expansion, keratinocyte proliferation was evident within the Lrig1+ KSC population (69 vs 55%, $p < 0.001$, $n=6$), and not in the CD34+ (1 vs 1%), LGR5+ (1 vs 3%) and LGR6+ (29 vs 40%) KSC populations ($n=7$, Figures 2c-e Figures 3a&b). Flow cytometric analysis of dissociated skin confirmed a 2.8-fold increase in Lrig1+ keratinocytes in the HPV8tg mice, 7.4% \pm 2.2% vs 2.7% \pm 0.8%, $n=6$, $P < 0.05$, but no difference in CD34+ KSC numbers (0.81% \pm 0.24% vs 0.73% \pm 0.37%) (Figure 2c). Flow sorted Lrig1+ and CD34+ keratinocyte sub-populations had similar levels of K14 promoter driven early region genes mRNA expression (Figure 4a), despite the observed difference in proliferation. To exclude any difference in the EVER 1 and 2 gene expression levels in HPV8tg mice versus control, qRT-PCR analysis was performed with the RNA extracted from Lrig1+ sorted cells and found comparable levels as shown in Figure 4b. Flow sorted Lrig1+ keratinocytes from HPV8tg mice also demonstrated a 3.8 fold increased colony forming efficiency (Figures 3c&d), hence Lrig1+ cells retain KSC function. There was no significant difference in colony forming efficiency from flow sorted Lrig1 negative keratinocytes from HPV8tg versus WT mice (Supplementary Figure 2). In

keeping with Lrig1+ expansion and proliferation, we observed nuclear p63 expression throughout this population and the emanating keratinocytes of HF infundibulum and adjoining IFE (Figure 3e). RT-PCR and western blotting analysis confirmed the switch from p63 TA isoforms to Δ Np63 isoforms in HPV8tg skin (Figures 5a-c), consistent with earlier reports indicating HPV8 early proteins induce p63 expression (Meyers *et al.*, 2013). Thus Lrig1+ KSC proliferation through induction of Δ Np63 in HPV8tg skin resulted in KSC expansion into the overlying infundibulum and adjoining IFE.

β -HPV associated expansion of the HF junctional zone KSC population in human skin field cancerization

The dilated HF infundibulum with increased keratinocyte layers and the crowded perifollicular epidermis with hyperkeratosis were consistently observed in the HPV8tg mice, and resembled Freudenthal's funnel, the pathognomonic histological finding in actinic keratosis (n=28 mice, Supplementary Figure 3). EV keratosis uniformly demonstrated similar histology associated with β -HPV virus reactivation and nuclear p63 expression within the HF infundibulum and perifollicular epidermis (Figure 5d, 6 patients with 44 lesions, Supplementary Table 1). Likewise, in some cases of actinic keratosis from non-EV patients, we were able to detect nuclear p63 expression within the infundibulum and perifollicular epidermis in areas where β -HPV virus reactivation was well evident, as detected by expression of the viral marker E4 (Figure 5d, n=2 of 25). Hence, our findings suggest that similar to HPV8 mice, EV and some patients with AK demonstrate expansion of a junctional zone HF KSC population identified by p63 labelling. These findings would suggest that β -HPV field cancerization arises from the HF infundibulum and predispose to SCC. Indeed, we observed p63 positive cells in HPV8tg and human SCC (Figure 5e, n=5); the latter were from immunocompetent patient from sun-exposed sites.

DISCUSSION

The concept of “field cancerization” as introduced by Slaughter *et al.* was initially used to describe an area of upper aerodigestive tract in situ SCC, within which develop multiple invasive SCC foci (Slaughter *et al.*, 1953). In the skin, similar pre-neoplastic field cancerization is observed and is characterized by the presence of multiple AK’s (Stockfleth *et al.*, 2011). Although ultraviolet (UV) light induced DNA damage is the prime cause of AK, other factors, including β -HPV infection can also contribute to development of this skin disorder, especially in immunocompromised patients (Taguchi *et al.*, 1994; Weissenborn *et al.*, 2005; Banerjee *et al.*, 2008). Patients with EV and related primary T cell immunodeficiency syndrome, are predisposed to develop skin field cancerization with similar keratoses that are associated with re-activation of latent β -HPV infection (Azzimonti *et al.*, 2005; Landini *et al.*, 2014). In all cases of skin field cancerization, UV light is the key driver for transformation to SCC, as tumors typically arise on sun-exposed sites and contain p53 UV signature mutations (Jonason *et al.*, 1996). Intriguingly, we have previously observed β -HPV reactivation at the clinically unaffected skin and in situ carcinoma at the periphery of SCC lesions from EV and organ transplant recipient, as well as within EV SCC (Borgogna *et al.*, 2014b). In some cases, 2 of 25 cases studied of non-EV AK, β -HPV reactivation was observed. The HPV8tg mouse model with constitutive epithelial HPV8 expression develops keratosis like skin changes, with 6% of lesions progressing to spontaneous SCC (Schaper *et al.*, 2005). The rate of conversion to SCC is greater in HPV8tg than in EV patients and may reflect the higher expression of early region genes. As in human EV, UV radiation rapidly leads to SCC formation in HPV8tg and other HPV mouse models (Marcuzzi *et al.*, 2009; Viarisio *et al.*, 2011).

KSC reside within each compartment of the skin: the IFE, HF, sebaceous and sweat glands. Different mouse KSC pools are distributed along the HF, defined by the expression of

cell surface proteins that facilitate isolation and thus characterization. To date KSC have been identified within four HF regions: upper (Lrig1) and lower (LGR6) junctional zone, the bulge (CD34 and K15) and bulb (LGR5) (Solanas & Benitah, 2013; Kretzschmar & Watt, 2014) (Figure 6). In this study, we have identified the HF Lrig1+ KSC population as the putative target in HPV8tg mice skin carcinogenesis. Lrig1 expression defines a distinct multipotent stem cell population in mammalian epidermis, which resides in the HF junctional zone in mice (Jensen *et al.*, 2009). Although Lrig1 expression defines an IFE KSC population in humans, it is not expressed in the reciprocal HF junctional zone (Supplementary Figure 4) and so prevented further characterization of the β -HPV target in humans (Jensen and Watt, 2006). Reporter mice studies have shown that junctional zone Lrig1 cells give rise to keratinocytes in the infundibulum and adjoining IFE, akin to the expansion of this population in our HPV8tg mice, human EV keratosis and AK.

p63 is a p53 protein family member expressed primarily in epithelia as six distinct isoforms, due to alternative transcription and C terminus splicing (Yang *et al.*, 1998). Three p63 isoforms contain an N-terminal transcriptional activation (TA isoforms) sequence, while the other three do not (Δ N isoforms). All of the p63 isoforms, TA and Δ N, are transcriptionally active and the Δ N isoforms repress TA isoforms as dominant-negative molecules (Koster. 2010). Multiple lines of evidence support the role of p63 in KSC maintenance (Melino *et al.*, 2015): i) p63 null mice demonstrate a terminally differentiated epidermis with no proliferative basal layer containing stem cells (Yang *et al.*, 1999); ii) epidermal p63 knockdown induces differentiation, which is induced by TAp63 (Truong *et al.*, 2006); iii) Δ Np63 expression maintains the proliferative basal layer (Truong *et al.*, 2006); and iv) p63 nuclear accumulation is prominent in KSC and holoclones (Pellegrini *et al.*, 2001). UV directly and via p53 mutation results in p63 downregulation and loss of transcriptional activity (Liefer *et al.*, 2000). Recently it has been proposed that UV induced concomitant

activation of oncogenic Ras and TGF- β pathways can restore p63 activity in p53 mutant cells to promote tumor progression (Vasilaki et al., 2016). In keeping with this, UV induced AK and cutaneous SCC frequently demonstrate nuclear p63 (Abbas et al., 2011).

The frequency of nuclear p63 positivity is greater in HPV associated SCC of the oral and anogenital regions, promoted by HPV E6 degradation of p53 (Melino, 2011).

β -HPV early region genes E2, E6 and E7 can induce KSC proliferation through inhibition of Notch signaling and subsequent induction of p63 (Hufbauer *et al.*, 2013; Meyers *et al.*, 2013, Pfefferle *et al.*, 2008) providing a putative mechanism for this β -HPV induced field cancerization. Hence, we propose that β -HPV early region genes initiate proliferation of Lrig1+ KSC causing their expansion into the overlying HF infundibulum and overlying epidermis. β -HPV driven KSC proliferation results in EV keratosis and occasionally non-EV AK, which are predisposed to transformation into SCC whereupon the β -HPV episome and so gene expression is lost (Figure 6).

MATERIAL AND METHODS

Transgenic Mouse Model

HPV8tg and wild type litter mates were housed and managed under conditions approved by the Italian Animal Care Committee. Age and sex matched mice ear thickness and tail width were measured using Vernier caliper.

Whole mount skin preparation

Tail and back skin was cut into 0.5 cm² pieces and dissociated using 2.5 U/ml Dispase (Roche, UK) over night at 4°C. The epidermis was gently removed and fixed in 10% neutral buffered formalin for 2 hours at room temperature, tissue was labelled and mounted as previously described (Braun K *et al.*, 2003).

Immunofluorescence labelling

Immunofluorescence on whole mount, frozen and paraffin embedded section was performed using standard techniques as previously described (Borgogna *et al.* 2012). For the list of antibodies used see Supplementary Table 2.

Single cell suspension for flow cytometry and colony forming efficiency

Tail and back skin was cut into 0.5 cm² pieces and dissociated using 2.5 U/ml Dispase (Roche, UK) over night at 4°C. The epidermis was gently removed and further dissociated with TrypLE™ Express Enzyme (ThermoFisher, UK), and the supernatant passed through a 70 µm cell strainer (BD bioscience, UK). Enzymes were inactivated with DMEM with 10% FBS and keratinocytes cell suspension were re-suspended as required.

Flow cytometry, cell sorting and colony forming efficiency assay

Samples were analysed and flow sorted using BD LSRFORTESSA and BD FACSAria™ Fusion (BD Biosciences, USA). The data were analyzed using Flowjo software (Tree Star Inc). Keratinocytes were flow sorted for CD34-/Lrig1+ population and 3000 cells per well were seeded in 6-well plate and cultured for 15 days. Rheinwald and Green Media was

changed every 3 days. The colonies were stained with crystal violet 0.05%, scanned with GelCount™ (Oxford Optronix), and analyzed using ImageJ software (NIH, USA).

RT-PCR

Mouse skin was homogenized in Trizol (ThermoFisher Scientific, UK). RNA isolation from sorted cells and homogenized tissues was performed with RNeasy kit and cDNA was synthesized using QuantiTect Reverse Transcription Kit (Qiagen, UK). Semi-quantitative reverse transcription-PCR (RT-PCR) reactions were carried out using GoTaq® G2 Green Master Mix (Promega, UK) and specific primers sequence are listed in Supplementary Table 3. Real-time quantitative reverse transcription analysis (qRT-PCR) was performed on QuantStudio™ 7 Flex Real-Time PCR System (ThermoFisher Scientific UK). For the determination of E6 and E7 gene expression levels, SyGreen (PCRBIO SYSTEM, UK) was used. Cycling condition were previously described in Schaper et al., 2005, De Andrea et al., 2010. Total mouse-specific β -Actin was used as the housekeeping gene.

EVER1, EVER2, TAp63 and Δ Np63 transcription levels were analyzed with Taqman probes at standard conditions. Total mouse-specific GAPDH was used as the housekeeping gene. Primers details are described in Supplementary Table 3 and 4.

Western blotting

100 mg of shaved back skin was homogenized in 1 ml of RIPA buffer containing 1% Triton X-100, 1% Sodium Deoxycholate, 0.1% SDS, 1 mM EDTA, 160 mM NaCl, 20 mM Tris-HCl (pH 7.4) and 25 μ l/ml Protease Inhibitor Cocktail (Sigma, UK). Protein concentrations were analyzed using Pierce BCA Protein Assay (ThermoFisher, UK). Thirty μ g of protein lysate was loaded on 8% SDS-PAGE, and transferred to polyvinylidene difluoride membranes (Millipore, UK). For the list of antibodies used see Supplementary Table 2.

Human tissue samples

EV patients' keratosis formalin-fixed and paraffin embedded (FFPE) tissue sections were analyzed according to the protocol approved by the "Maggiore Hospital" Research Ethics Committee, Italy. Non-EV actinic keratosis (14-NW-1272) and squamous cell carcinoma (09-WSE-02-1) tissues were obtained after UK NHS R&D, Local Research Ethics Committee approval and informed written consent.

Statistical analysis

Paired t-tests were used to compare HPV8 and wild type litter mates, using GraphPad software (Prism).

CONFLICT OF INTEREST

The authors state no conflict of interest.

ACKNOWLEDGEMENTS

This work was supported by grants from the Italian Ministry for University and Research - MIUR (PRIN 2012 to CB), Compagnia di San Paolo (CSP2014 to CB), Associazione Italiana per la Ricerca sul Cancro - AIRC (grant IG 2012 and 2016 to MG) and Cancer Research UK grant (2014). Carlotta Olivero fellowship was funded by Compagnia di San Paolo.

We would like to thank Dr John Doorbar, University of Cambridge, for providing the anti-E4 antibodies.

REFERENCES

- Abbas O, Richards JE, Yaar R, Mahalingam M. Stem cell markers (cytokeratin 15, cytokeratin 19 and p63) in in situ and invasive cutaneous epithelial lesions. *Mod Pathol* 2011; 24(1):90-7
- Azzimonti B, Mondini M, De Andrea M, Gioia D, Dianzani U, Mesturini R, et al. CD8+ T-cell lymphocytopenia and lack of EVER mutations in a patient with clinically and virologically typical epidermodysplasia verruciformis. *Arch Dermatol* 2005; 141(10):1323-5.
- Banerjee M, Sarma N, Biswas R, Roy J, Mukherjee A, Giri AK. DNA repair deficiency leads to susceptibility to develop arsenic-induced premalignant skin lesions. *Int J Cancer* 2008 15; 123(2):283-7
- Bernard HU, Burk RD, Chen Z, van Doorslaer K, zur Hausen H, de Villiers EM. Classification of papillomaviruses (PVs) based on 189 PV types and proposal of taxonomic amendments. *Virology* 2010; 401:70–79.
- Borgogna C, Zavattaro E, De Andrea M, Griffin HM, Dell'Oste V, Azzimonti B, et al. Characterization of beta papillomavirus E4 expression in tumours from Epidermodysplasia Verruciformis patients and in experimental models. *Virology* 2012; 423:195–204
- Borgogna C, Landini MM, Lanfredini S, Doorbar J, Bouwes Bavinck JN, Quint KD, et al. Characterization of skin lesions induced by skin-tropic α - and β -papillomaviruses in a patient with epidermodysplasia verruciformis. *Br J Dermatol* 2014a;171:1550-4
- Borgogna C, Lanfredini S, Peretti A, De Andrea M, Zavattaro E, Colombo E, et al. Improved detection reveals active β -papillomavirus infection in skin lesions from kidney transplant recipients. *Mod Pathol* 2014b;27(8):1101-15
- Bosch FX, Broker TR, Forman D, Moscicki AB, Gillison ML, Doorbar J, et al. Comprehensive control of human papillomavirus infections and related diseases. *Vaccine* 2013;31 (suppl 7):H1–31

Bouwes Bavinck JN, Plasmeijer EI, Feltkamp MC. Beta-papillomavirus infection and skin cancer. *J Invest Dermatol* 2008;128:1355-8

Boxman IL, Berkhout RJ, Mulder LH, Wolkers MC, Bouwes Bavinck JN, Vermeer BJ et al. Detection of human papillomavirus DNA in plucked hairs from renal transplant recipients and healthy volunteers. *J Invest Dermatol* 1997;108:712–5

Braun KM, Niemann C, Jensen UB, Sundberg JP, Silva-Vargas V, Watt FM. Manipulation of stem cell proliferation and lineage commitment: visualisation of label-retaining cells in wholemounts of mouse epidermis. *Development*. 2003;130(21):5241-55.

De Andrea M, Rittà M, Landini MM, Borgogna C, Mondini M, Kern F et al. Keratinocyte-specific Stat3 heterozygosity impairs development of skin tumors in human papillomavirus 8 transgenic mice. *Cancer Res* 2010;70:7938-48

Howley PM, Pfister HJ. Beta genus papillomaviruses and skin cancer. *Virology* 2015 May;479-480:290-6.

Hufbauer M, Biddle A, Borgogna C, Gariglio M, Doorbar J, Storey A et al. Expression of betapapillomavirus oncogenes increases the number of keratinocytes with stem cell-like properties. *J Virol* 2013;87:12158–65

Jensen KB, Collins CA, Nascimento E, Tan DW, Frye M, Itami S et al. Lrig1 expression defines a distinct multipotent stem cell population in mammalian epidermis. *Cell Stem Cell* 2009;4:427–39

Jensen KB, Watt FM. Single-cell expression profiling of human epidermal stem and transit-amplifying cells: Lrig1 is a regulator of stem cell quiescence. *Proc Natl Acad Sci USA* 2006;103:11958–63

Jonason AS, Kunala S, Price GJ, Restifo RJ, Spinelli HM, Persing JA, Leffell DJ, Tarone RE, Brash DE. Frequent clones of p53-mutated keratinocytes in normal human skin. *Proc Natl Acad Sci U S A*. 1996 Nov 26;93(24):14025-9

Koster MI. p63 in skin development and ectodermal dysplasias. *J Invest Dermatol* 2010;130(10):2352-8.

Kranjec C, Doorbar J. Human papillomavirus infection and induction of neoplasia: a matter of fitness. *Curr Opin Virol* 2016;20:1-8

Kretzschmar K, Watt FM. Markers of epidermal stem cell subpopulations in adult mammalian skin. *Cold Spring Harb Perspect Med* 2014;4(10). pii: a013631

Landini MM, Borgogna C, Peretti A, Colombo E, Zavattaro E, Boldorini R et al. α - and β -papillomavirus infection in a young patient with an unclassified primary T-cell immunodeficiency and multiple mucosal and cutaneous lesions. *J Am Acad Dermatol* 2014;71(1):108-15

Liefer KM, Koster MI, Wang XJ, Yang A, McKeon F, Roop DR. Down-regulation of p63 is required for epidermal UV-B-induced apoptosis. *Cancer Res* 2000;1;60(15):4016-20

Marcuzzi GP, Hufbauer M, Kasper HU, Weissenborn SJ, Smola S, Pfister H. Spontaneous tumour development in human papillomavirus type 8 E6 transgenic mice and rapid induction by UV-light exposure and wounding. *J Gen Virol* 2009;90(Pt 12):2855-64

Melino G. p63 is a suppressor of tumorigenesis and metastasis interacting with mutant p53. *Cell Death Differ* 2011;18(9):1487-99

Melino G, Memmi EM, Pelicci PG, Bernassola F. Maintaining epithelial stemness with p63. *Sci Signal* 2015;8(387):re9

Meyers JM, Spangle JM, Munger K. The human papillomavirus type 8 E6 protein interferes with NOTCH activation during keratinocyte differentiation. *J Virol* 2013;87:4762-7

Notarangelo L, Casanova JL, Fischer A, Puck J, Rosen F, Seger R, et al. International Union of Immunological Societies Primary Immunodeficiency diseases classification committee.. Primary immunodeficiency diseases: an update. *J Allergy Clin Immunol*. 2004 Sep;114(3):677-87.

Pellegrini G, Dellambra E, Golisano O, Martinelli E, Fantozzi I, Bondanza S et al. p63 identifies keratinocyte stem cells. *Proc Natl Acad Sci U S A* 2001;98(6):3156-61

Pfefferle R, Marcuzzi GP, Akgul B, Kasper HU, Schulze F, Haase I et al. The human papillomavirus type 8 E2 protein induces skin tumors in transgenic mice. *J Invest Dermatol* 2008;128:2310-5

Quint KD, Genders RE, de Koning MNC, Borgogna C, Gariglio M, Bouwes Bavinck JN, et al. Human Beta-papillomavirus infection and keratinocyte carcinomas. *J Pathol* 2015; 235(2):342–54

Ramoz N, Rueda LA, Bouadjar B, Montoya LS, Orth G, Favre M. Mutations in two adjacent novel genes are associated with epidermodysplasia verruciformis. *Nat Genet.* 2002 Dec;32(4):579-81.

Schaper ID, Marcuzzi GP, Weissenborn SJ, Kasper HU, Dries V, Smyth N et al. Development of skin tumors in mice transgenic for early genes of human papillomavirus type 8. *Cancer Res* 2005;65:1394–1400

Slaughter DP, Southwick HW, Smejkal W. ‘Field cancerisation’ in oral stratified squamous epithelium. *Cancer* 1953;6:963–968

Solanas G, Benitah SA. Regenerating the skin: a task for the heterogeneous stem cell pool and surrounding niche. *Nat Rev Mol Cell Biol* 2013;14(11):737-48

Stockfleth E, Ortonne J-P, Alomar A. Actinic keratosis and field cancerisation. *Eur J Dermatol* 2011;21:3-12

Taguchi M, Watanabe S, Yashima K, Murakami Y, Sekiya T, Ikeda S. Aberrations of the tumor suppressor p53 gene and p53 protein in solar keratosis in human skin. *J Invest Dermatol.* 1994;103 (4): 500–3

Truong AB, Kretz M, Ridky TW, Kimmel R, Khavari PA. p63 regulates proliferation and differentiation of developmentally mature keratinocytes. *Genes Dev* 2006;20(22):3185-97

Vasilaki E, Morikawa M, Koinuma D, Mizutani A, Hirano Y, Ehata S, et al. Ras and TGF- β signaling enhance cancer progression by promoting the Δ Np63 transcriptional program. *Sci Signal* 2016; 9:ra84.

Viarisio D, Mueller-Decker K, Kloz U, Aengeneyndt B, Kopp-Schneider A, Gröne HJet al. E6 and E7 from beta HPV38 cooperate with ultraviolet light in the development of actinic keratosis-like lesions and squamous cell carcinoma in mice. *PLoS Pathog* 2011;7(7):e1002125

Weissenborn SJ, Nindl I, Purdie K, Harwood C, Proby C, Breuer J, et al. Human papillomavirus-DNA loads in actinic keratoses exceed those in non-melanoma skin cancers. *J Invest Dermatol* 2005;125 (1): 93–7

Yang A, Kaghad M, Wang Y, Gillett E, Fleming MD, Dötsch V, et al. p63, a p53 homolog at 3q27-29, encodes multiple products with transactivating, death-inducing, and dominant-negative activities. *Mol Cell* 1998;2: 305–316

Yang A, Schweitzer R, Sun D, Kaghad M, Walker N, Bronson RT et al. p63 is essential for regenerative proliferation in limb, craniofacial and epithelial development. *Nature* 1999;398(6729):714-8

FIGURE LEGENDS

Figure 1: Phenotypic and histological characterization of HPV8tg mice.

a, Schematic representation of the HPV8 transgenes, showing the human cytokeratin-14 gene promoter upstream of the open reading frames of the HPV8 early region genes. **b**, Hematoxylin and eosin staining of paraffin embedded skin sections from WT (left panel) and HPV8tg (right panel) mouse skin. **c**, Toluidine blue staining of mouse whole mount skin including the hair follicle and overlying epidermis from WT (left panel) and HPV8tg (right panel) from the tail. **d**, Ear thickness (upper graph) and tail width (bottom graph) measured using calipers on age matched WT and HPV8tg littermates (n=8), with mean \pm SD (*,p<0.05; unpaired t test). All the images were processed using ImageJ software (NIH, USA). All scale bars =100 μ m.

Figure 2: HPV8 transgenes induce hair follicle changes in HPV8tg mice.

a, b, Adult mice whole mount skin were photographed and analyzed for the area of HF regions and length, WT and HPV8tg were compared, with mean \pm SD, using an unpaired t-test (n=20, ** p<0.01). **c**, FACs analysis WT and HPV8tg mice skin keratinocyte isolates (n=6), labelled with Lrig1-FITC and CD34-647 antibodies, with DAPI to select live cells. **d**, The number of Lrig1 positive cells determined by FACs (n=6). (**, p<0.01; unpaired t test), with mean \pm SD **e**, Whole mount immunofluorescence of adult WT and HPV8tg tail skin for Ki67 (red) and HF-KSC markers (green). All the images were processed using ImageJ software (NIH, USA). All scale bars = 100 μ m.

Figure 3: Lrig1 keratinocyte stem cell proliferation in HPV8tg

a, Whole mount adult WT and HPV8tg (n=6) skin sections labelled with Ki67 (red), Lrig1 (green) antibodies and DAPI (blue). **b**, Number of proliferating cells within KSC populations (Lrig1, LGR6, CD34, and LGR5) was enumerated, with mean \pm SD, in WT and HPV8tg tissue sections (n=7); ** p<0.01. **c**, Keratinocyte colony forming assays from flow sorted Lrig1 cells from WT and HPV8tg (n=6) skin dissociates, with mean \pm SD and **d**, A representative image. **e**, p63 labelled frozen sections from adult WT and HPV8tg mice (n=3), the broken line indicates the basal layer. Immuno-labelled tissue sections were visualized and photographed by fluorescent microscope with x20 magnification then processed using ImageJ software (NIH, USA). All scale bars are 100 μ m.

Figure 4: HPV8 transgenes are uniformly expressed in the epidermis of HPV8tg mice.

a, qRT-PCR of early region genes of WT and HPV8tg mouse skin isolated KSC populations, with mean \pm SD. **b**, qRT-PCR of murine homologues of EVER1 (TMC6) and EVER2 (TMC8) genes of WT and HPV8tg mouse skin isolated Lrig1+ KSC populations, with mean \pm SD.

Figure 5: β -HPV induced keratinocyte stem cell expansion results in keratosis that are predisposed to SCC

a&b, RT-PCR with mean \pm SD (**, p<0.01), and **c**, Western blot gels of p63 isoforms from RNA and protein isolates respectively from WT, HPV8tg skin, papilloma (pap) and SCC (n=6). **d**, Images of tissue sections of HPV8tg mouse skin, human EV keratosis, and AK labelled with p63 (green), β -HPV E4 (red) specific antibodies, and DAPI (blue). **e**, Images of tissue sections of HPV8tg mouse and human SCCs labelled with K14 (red), p63 (green) specific antibodies and DAPI (blue). Immuno-labelled tissue sections were visualized and

photographed by fluorescent microscope with x20 magnification then processed using ImageJ software (NIH, USA). All scale bars are 100 μ m.

Figure 6: β -HPV Field Cancerization Model

The mouse HF (left image) is composed by at least 4 different KSC populations. In the WT HF, the LGR5 is expressed in the hair bulb, CD34+ and K15+ KSC are located in the bulge, and the Lrig1 and LGR6 KSC are located in the upper and lower junctional zone respectively. In HPV8tg, the Lrig1 KSC population expand beyond the junctional zone niche and no longer express Lrig1 but they maintain KSC function. Similarly, in human β -HPV reactivation (right image), the expanded population occupies the hair follicle infundibulum and adjoining interfollicular epidermis, as shown by the yellow arrow. These changes culminate in the histological phenotype known as Freudenthal's funnel, the pathognomonic finding in actinic keratosis and skin field cancerization.