

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Role and function of exosomal miR-9-3p in hepatocellular carcinoma

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1650095> since 2018-01-08T12:54:06Z

Published version:

DOI:10.23736/S0026-4806.17.05384-8

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This is an author version of the contribution published on:

Questa è la versione dell'autore dell'opera:

[Minerva Medica, 2017, 10.23736/S0026-4806.17.05384-8.]

ovvero [Caviglia GP, Fagoonee S, 2017, pagg.1-3]

The definitive version is available at:

La versione definitiva è disponibile alla URL:

[<https://www.minervamedica.it/it/riviste/minerva-medica/index.php>]

Editorial

Role and function of exosomal miR-9-3p in hepatocellular carcinoma

Gian Paolo CAVIGLIA ^{1,*}, Sharmila FAGOONEE ²

¹ Department of Medical Sciences, University of Turin, Turin, Italy

² Institute for Biostructures and Bioimages CNR c/o Molecular Biotechnology Center, University of Turin, Turin, Italy

*Corresponding author: Gian Paolo Caviglia, Department of Medical Sciences, University of Turin, Turin 10100, Italy. E-mail: caviglia.giampi@libero.it

Conflict of interest: None

Key words: Biomarker - Hepatocellular carcinoma - microRNA

MicroRNAs (miRNAs) are small (~20-22 nucleotides) non-coding RNAs able to regulate gene expression through repression or degradation of mRNAs.¹ MiRNAs are the biggest class of genetic regulators and they act at a post-transcriptional level with a mechanism based on sequence complementarities between miRNA and target messenger RNA (mRNA) (Figure 1).² MiRNAs are important regulator of the principal cell functions including growth, proliferation, differentiation and apoptosis, being involved in the oncogenic processes acting as oncogenes, tumor suppressor genes or both.³ In addition, there is a growing interest to explore circulating miRNAs as potential biomarkers for early hepatocellular carcinoma (HCC) detection.⁴

Sun *et al.* observed an increased expression of miR-9 (*i.e.* miR-9-5p) in HCC tissue compared to normal liver and adjacent non-cancerous tissues; higher miR-9 expression was associated with a poor prognosis of HCC.⁵ Furthermore, it has been reported that miR-9 could promote tumour cell migration and invasion in HCC cell lines.⁶

Recently, Tang *et al* investigated the biological function of exosomal miR-9-3p (the “minor” product of miR-9) with a focus on its involvement in the proliferation of HCC cells.⁷ Interestingly, in an experimental model authors observed that miR-9-3p, as miR-9, is overexpressed in HCC cell lines, but showing opposite biological effects. Indeed, miR-9-3p was able to inhibit HCC cells proliferation by directly suppressing heparin-binding growth factor-5 (*HBGF-5*) expression, a gene member of the fibroblast growth factor (FGF) family,⁸ and by inhibiting the extracellular signal-regulated protein kinase 1 and 2 (*ERK1/2*) expression, involved in the regulation mechanism of cell proliferation and apoptosis.⁹

In addition, Tang *et al* found that miR-9-3p levels were significantly higher in the serum exosomes of normal donors compared to patients with HCC. The accuracy for the discrimination between donors and HCC patients was 0.813 (area under the curve) indicating a good diagnostic performance for miR-9-3p.⁷ To date, several studies assessed the potential role of miRNAs as well as other classes of biomarkers for HCC detection and reported promising results.^{1, 10-13} However,

must be stressed that caution should be used when dealing with miRNAs and other epigenetic molecules,^{14, 15} since the complex miRNA-mRNA interaction may be affected by numerous other genetic and epigenetic features, consequently resulting in an alteration of miRNAs expression.¹⁶ In addition, the use of such molecules as cancer biomarkers is partly limited by lack of standardized method for their quantitation. Since there is still a need of reliable biomarkers for HCC detection, the use of traditional biomarkers assessed by novel highly sensitive and fully automated analytical methods, may be the most effective strategy to address this clinically relevant issue.

References

1. Petrini E, Caviglia GP, Abate ML, Fagoonee S, Smedile A, Pellicano R. MicroRNAs in HBV-related hepatocellular carcinoma: functions and potential clinical applications. *Panminerva Med* 2015;57:201-9.
2. Yu B, Liu X, Chang H. MicroRNA-143 inhibits colorectal cancer cell proliferation by targeting MMP7. *Minerva Med* 2017;108:13-9.
3. Takata A, Otsuka M, Yoshikawa T, Kishikawa T, Ohno M, Koike K. MicroRNAs and liver function. *Minerva Gastroenterol Dietol* 2013;59:187-203.
4. Abenavoli L, Boccuto L. New serum markers for detection of early hepatocellular carcinoma. *Panminerva Med* 2017. doi: 10.23736/S0031-0808.17.03356-0
5. Sun J, Fang K, Shen H, Qian Y. MicroRNA-9 is a ponderable index for the prognosis of human hepatocellular carcinoma. *Int J Clin Exp Med* 2015;8:17748-56.
6. Sun Z, Han Q, Zhou N, Wang S, Lu S, Bai C, *et al.* MicroRNA-9 enhances migration and invasion through KLF17 in hepatocellular carcinoma. *Mol Oncol* 2013;7:884-94.
7. Tang J, Li Y, Liu K, Zhu Q, Yang WH, Xiong LK, *et al.* Exosomal miR-9-3p suppresses HBGF-5 expression and is a functional biomarker in hepatocellular carcinoma. *Minerva Med* 2017; in press.
8. Kornmann M, Ishiwata T, Beger HG, Korc M. Fibroblast growth factor-5 stimulates mitogenic signaling and is overexpressed in human pancreatic cancer: evidence for autocrine and paracrine actions. *Oncogene* 1997;15:1417-24.
9. Liu X, Li G. MicroRNA-133b inhibits proliferation and invasion of ovarian cancer cells through Akt and Erk1/2 inactivation by targeting epidermal growth factor receptor. *Int J Clin Exp Pathol* 2015;8:10605-14.

10. Jia H, Yu H, Liu Q. Single nucleotide polymorphisms of MIR-149 gene rs2292832 contributes to the risk of hepatocellular carcinoma, but not overall cancer: a meta-analysis. *Minerva Med* 2016;107:259-69.
11. Caviglia GP, Abate ML, Gaia S, Petrini E, Bosco C, Olivero A, *et al.* Risk of hepatocellular carcinoma in HBV cirrhotic patients assessed by the combination of miR- 122, AFP and PIVKA-II. *Panminerva Med* 2017. doi: 10.23736/S0031-0808.17.03353-5.
12. Caviglia GP, Smedile A. Hepatitis B core-related antigen: a novel biomarker for chronic hepatitis B treatment. *Minerva Gastroenterol Dietol* 2017;63:169-71.
13. Caviglia GP, Abate ML, Petrini E, Gaia S, Rizzetto M, Smedile A. Highly sensitive alpha-fetoprotein, Lens culinaris agglutinin-reactive fraction of alpha-fetoprotein and des-gamma-carboxyprothrombin for hepatocellular carcinoma detection. *Hepatol Res* 2016;46:E130-5.
14. Song W, Wang K, Zhang RJ, Dai QX, Zou SB. Long noncoding RNA GAS5 can predict metastasis and poor prognosis: a meta-analysis. *Minerva Med* 2016;107:70-6.
15. Wu SW, Hao YP, Qui JH, Zhang DB, Yu CG, Li HW. High expression of long noncoding RNA CCAT2 indicates poor prognosis of gastric cancer and promotes cell proliferation and invasion. *Minerva Med* 2017. doi: 10.23736/S0026-4806.17.04703-6.
16. Lombardi A, Grimaldi A, Zappavigna S, Misso G, Caraglia M. Hepatocarcinoma: genetic and epigenetic features. *Minerva Gastroenterol Dietol* 2017. doi: 10.23736/S1121-421X.17.02408-4.

Figure 1.-Biogenesis and function of miRNAs.

MiRNAs are transcribed in the nucleus by RNA polymerase II as a stem-loop primary transcript (pri-miRNAs). Following a cleavage by the ribonucleoprotein complex Drosha/DGCR8, hairpin precursor-miRNAs (pre-miRNAs) are exported into the cytoplasm through exportin-5. Then, pre-miRNAs are further processed by another RNase III (Dicer) into an RNA-duplex consisting of the mature miRNA and the passenger miRNA strand. Subsequently, miRNA duplex is unwinded by helicase activity and the mature miRNA (arisen from the 5' arm) can be incorporated into the RNA-induced silencing complex (RISC). Usually, the “minor” passenger strand (arisen from the 3' arm) is not functional and cannot be incorporated into the RISC complex. The miRNA-RISC complex allow the interaction between the mature miRNA and the target mRNA. According to miRNA-mRNA complementarity, their interaction induces either mRNA translation repression or mRNA degradation.

Abbreviations: miRNA, microRNA; RISC, RNA-induced silencing complex.