

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Patterns of infections in older patients acutely admitted to medical wards: data from the REPOSI register

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1719068> since 2019-12-13T12:17:45Z

Published version:

DOI:10.1007/s11739-019-02178-1

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Patterns of infections in older patients acutely admitted to medical wards: data from the REPOSI register

Raffaella Rossio¹, Ilaria Ardoino², Carlotta Franchi², Alessandro Nobili², Pier Mannuccio Mannucci³, Flora Peyvandi^{4,5}, REPOSI Investigators

Keywords Older people · Infections · Hospitalizations · Comorbidities

Dear Editor,

In older adults infections are among the leading causes of emergency department visits, hospitalization, morbidity and mortality [1–3]. Infections also occur as adverse events during hospitalization, as highlighted by the large use of antibiotics in this setting, resulting in an increase of hospitalization length and mortality rate [4–6]. There is a paucity of studies, especially in European countries, that did offer a general pattern on all the types of infections occurring in acutely hospitalized older patients, being the literature mainly focused on single type of infections (i.e. pneumonia and urinary tract infections). To fill this gap of knowledge, we chose to observe and describe the prevalence and types of infections in a large cohort of older hospitalized patients in the frame of REPOSI (REGistro POLiterapie SIMI) register. REPOSI is a collaborative register that involves a large number of Italian internal medicine and geriatric wards. Briefly, patients aged 65 years or more acutely hospitalized during four index periods lasting one week in each season were enrolled and signed an informed consent. The attending physicians were required to compile for each patient a web-based Case Report Form, including the main socio-demographic data, specific diagnoses and their severity according to the Cumulative Illness Rating Scale (CIRS), functional status at hospital admission (measured by Barthel index [BI]) [7] and drug therapies during the whole hospitalization period. This study was approved by the Ethical Committees of all participating hospitals. More details on REPOSI were provided elsewhere [5, 6].

Patients recruited in the register in 2008, 2010, 2012, 2014 and 2016 were considered for the purpose of this study. Infections were considered when they were the reason for hospitalization, when reported in the CIRS at admission and also when they occurred during the hospital stay, whether reported as a relevant adverse event or as the indication for antimicrobial therapy. Infections were categorized using the International Classification of Diseases-Ninth Edition (ICD-

9) according to the physiological system of origin, on the basis of a classification provided in other studies (Supplementary Table 1) [8]. Data were summarized as prevalence (%), or means and standard deviations when pertinent. The 95% confidence intervals (CI) were provided as well. The analysis was performed using the SAS/STAT software Version 9.2 (SAS Institute Inc., Cary, NC, USA).

Among the 6047 patients enrolled in the REPOSI register, 2991 (49.5%, 95% CI 48.2–50.7%) were diagnosed with at least one infection, accounting for a total of 3554 infections. Overall 2522 patients (84.3%) had a single infection, 392 (13.1%) two and 77 (2.6%) three or more infections. Almost all the infections had a bacterial or suspected bacterial etiology, but 284 (8.0%) were viral, mainly hepatobiliary (N = 223) and lower-respiratory track (N = 41) infections. Socio-demographic and clinical characteristics of the patients with and without infections are reported in Table 1. A total of 1554 patients (52.0%) were females and their mean age was 80.0 years (SD = 7.5). Compared to patients without infections, almost the same prevalence of females was observed but they were slightly older, the mean age of patients without infections being 78.6 years (SD = 7.4) and they took more medications at hospital admission. While the Barthel index (BI) was not available in the 2008 REPOSI run, in the remaining years patients with infections were often functionally impaired, those with a severe

dependence in daily activities (BI < 49) having a prevalence of 71.5% compared to 28.4% in those without infections. Considering the different years of REPOSI, a substantial increase in the overall prevalence of patients with infections was observed, from 34.0% in the 2008 up to 52.1% in the 2010, with an almost stable prevalence during the following years (with a mean value of 54.5%) (P values < 0.0001 for the overall chi squared comparison and 0.24 for the comparison after excluding the 2008 REPOSI run).

Before admission 187 patients (6.5%) with infections

lived in nursing homes, those from nursing homes being 4.7% less among patients without infections. The greatest number of hospitalizations took place during the winter season, but a slightly higher prevalence of patients with infections was observed during summer (18.2% vs 14.7% in those without infections).

The most common type of infections were in the upper and lower respiratory tract (38.2%), of which 860 were pneumonia and bronchopneumonia, followed by those in the kidney and urinary tract (21.1%) and the hepatobiliary system (9.3%). Bloodstream infections accounted for the 5.5% of the infections (Table 2). An infection was the reason of hospitalization in 734 patients (24.5%, 95% CI 23.0–26.1%), about half of them being hospitalized for respiratory tract infections, urinary tract and gastrointestinal infections accounting for 35.0% of the remaining infections. Respiratory tract infections were also the most common infection occurring during hospital stay, followed by those of the urinary tract. When analysing the type of infection over time, there was an almost stable pattern, with an increase only for septicemia (Supplementary Fig. 1). When we analysed infection sites according to sex, urinary (25.5% vs 16.24%) and gastrointestinal (8.5% vs 7.2%) infections occurred more frequently in females, hepatobiliary infections in males (10.6% vs 8.0%), and respiratory infections with no marked differences for sex distribution (35.3% in females vs 41.4% in males). Respiratory tract (35.9% vs 43.4%) and urinary tract infections (20.1% vs 23.4%) were more prevalent among the oldest patients (≥ 85 years), while hepatobiliary infections (10.8% vs 5.9%) were more prevalent in those less than 85 years of age.

Among patients with infections 176 (5.9%) died during hospitalization.

All in all, approximately half of the older patients enrolled in the frame of the REPOSI register (49.5%) had at least one infection diagnosed at admission or during hospital stay, and over the 8-year study period their prevalence remained almost unchanged after an early increase. In Italy, Orlando et al. have previously emphasized as an emerging problem that an increasing number of older people with multiple chronic conditions is admitted to hospital with infections, but actual data on older people acutely hospitalized in medical wards were scanty [9]. In the frame of the different runs of the REPOSI register, respiratory tract infections were the most frequent type of infection being more than one third of the cases (38.2%), followed in prevalence by urinary tract infections throughout all the register runs. The same pattern was also observed by others [10]. Septicemia actually increased during the REPOSI runs from 11.3% in 2008 to 24.2% in 2016. This can be explained by an increasing attention for this diagnosis and the large use of artificial devices in older patients.

The patients most frequently hospitalized in internal medicine wards are often very old and characterized by multiple chronic conditions, the most frequent concomitant diseases being diabetes and chronic renal failure. Polypharmacy is often a consequence of multimorbidity and indeed at admission more than half of our patients took 5 or more drugs. Other studies identified an association between infections and drug intake and it is known that some drugs frequently prescribed to old people increase the risk of infections [11]. In spite of the fact that a relatively small number of our older patients had a severe or total dependence according to the Barthel index (17.5%), our data support the hypothesis that a compromised functional status and polypharmacy are associated with an increased prevalence of infections in the elderly [12]. In the previous studies, a poor

functional status was found to be an independent prognostic factor for mortality in older people with pneumonia and a risk factor for respiratory and urinary tract infections [13, 14]. Furthermore, another REPOSI study found that a low Barthel index increased the risk of death in old patients admitted for pneumonia [15]. Finally, there was a lower degree of physical ability in patients with pneumonia developing during hospitalization compared to those admitted with community acquired pneumonia, as previously reported in a REPOSI study [6]. On the whole, the results of the present study, that considered all types of infections occurring in older people, suggest that a low functional status may be associated with all types of infection, not only with those of the respiratory tract.

There are limitations in this merely descriptive study, such as the fact that the register does not collect details on etiology, outcomes and diagnostic methods for infections (e.g. microbiological cultures, X-ray imaging, etc.). On the other hand REPOSI is based upon data from more than 100 internal medicine and geriatric wards throughout Italy, thus making this study representative of the hospitalized older population of this country. Moreover, data collection started in 2008, so that changes over time were accounted for.

In conclusion infections are frequent events diagnosed in older hospitalized patients with a high number of comorbidities and on polypharmacy, respiratory tract infections being the most frequent followed by urinary tract infections.

Funding None.

Compliance with ethical standards

Conflict of interest FP has received honoraria for participating as a speaker at satellite symposia and educational meetings organized by Ablynx, Grifols, Novo Nordisk, Roche, Shire and Sobi. She has received consulting fees from Kedrion and she is member of the scientific advisory board of Ablynx. Other authors have no conflict of interest to declare.

Statements on human and animal rights REPOSI was accepted by the ethical committees of all participant hospitals.

Informed consent Each patient supplied informed consent.

References

1. Liang SY (2016) Sepsis and other infectious disease emergencies in the elderly. *Emerg Med Clin N Am* 34:501–522
2. Shi T, Denouel A, Tietjen AK, Lee JW, Falsey AR, Demont C, Nyawanda BO, Cai B, Fuentes R, Stoszek SK, Openshaw P, Campbell H, Nair H (2019) Global and regional burden of hospital admissions for pneumonia in older adults: a systematic review and meta-analysis. *J Infect Dis*. <https://doi.org/10.1093/infdis/jiz053>
3. Kardas P, Ratajczyk-Pakalska E (2003) Reasons for elderly patient hospitalization in departments of internal medicine in Lodz. *Aging Clin Exp Res* 15:25–31
4. Falcone M, Paul M, Yahav D, Orlando G, Tiseo G, Prendki V, Güerri-Fernández R, Gavazzi G, Mutters NT, Cookson B, Tinelli M (2019) Study Group for Infections in the elderly (ESGIE) anti-microbial consumption and impact of antimicrobial stewardship programmes in long-term care facilities. *Clin Microbiol Infect* 25:562–569
5. Ardoino I, Mannucci PM, Nobili A, Franchi C (2019) Antibiotic use and associated factors in a large sample of hospitalized older people. *J Glob Antimicrob Resist*. <https://doi.org/10.1016/j.jgar.2019.04.013>
6. Rossio R, Franchi C, Ardoino I, Djade CD, Tettamanti M, Pasina L, Salerno F, Marengoni A, Corrao S, Marcucci M, Peyvandi F, Biganzoli EM, Nobili A, Mannucci PM (2015) Adherence

to antibiotic treatment guidelines and outcomes in the hospitalized elderly with different types of pneumonia. *Eur J Intern Med* 26:330–337

7. Mahoney FI, Barthel DW (1965) Functional evaluation: the Barthel Index. *Md State Med J* 14:61–65

8. Christensen KL, Holman RC, Steiner CA, Sejvar JJ, Stoll BJ, Schonberger LB (2009) Infectious disease hospitalizations in the United States. *Clin Infect Dis* 49:1025–1035

9. Orlando G, Gubertini G, Negri C, Coen M, Ricci E, Galli M, Rizzardi G (2014) Trends in hospital admissions at a department for infectious diseases in Italy from 1995 to 2011 and implications for health policies. *Public Health* 14:980

10. Kucukardali Y, Oncul O, Kunter E, Turhan V, Solmazgul E, Terekeci HM, Sayan O, Oktenli C (2009) Community acquired infections in elderly population. *Cent. Eur J Med* 4:171–178

11. Maes ML, Fixen DR, Linnebur SA (2017) Adverse effects of proton-pump inhibitor use in older adults: a review of the evidence. *Ther Adv Drug Saf* 8:273–297

12. High K, Bradley S, Loeb M, Palmer R, Quagliarello V, Yoshikawa T (2005) A new paradigm for clinical investigation of infectious syndrome in older adults: assessing functional status as a risk factor and outcome measure. *J Am Geriatr Soc* 53:528–535

13. Osakwe ZT, Larson E, Shang J (2019) Urinary tract infection related hospitalization among older adults receiving home health care. *Am J Infect Control* 47:786–792

14. Mody L, Sun R, Bradley S (2006) Assessment of pneumonia in older adults: effect of functional status. *J Am Geriatr Soc* 54:1062–1067

15. Corrao S, Argano C, Natoli G, Nobili A, Corazza GR, Mannucci PM, Perticone F (2018) Disability, and not diabetes, is a strong predictor of mortality in oldest old patients hospitalized with pneumonia. *Eur J Intern Med* 54:53–59

Table 1 Socio-demographic and clinical characteristics of 2991 older patients with infections and 3056 without infections at hospital admission

Variables	Patients with infection N (%)	Patients without infections	Missing data
2008	453 (15.2%)	879 (28.8%)	
2010	719 (24.0%)	661 (21.6%)	
2012	703 (23.5%)	620 (20.3%)	
2014	674 (22.5%)	538 (17.6%)	
2016	442 (14.8%)	358 (11.7%)	
Women (n, %)	1554 (52.0%)	1572 (51.4%)	
Age (mean ± SD)	80 ± 7.5	78.6 ± 7.4	
Age group (n, %)			
Young old (≤ 75)	877 (29.3%)	1101 (36.0%)	
Old (76–84)	1231 (41.2%)	1246 (40.8%)	
Oldest old (≥ 85)	883 (29.5%)	709 (23.2%)	
Living arrangement (n, %)			219
Alone	660 (22.8%)	657 (22.4%)	
With family member/caregiver	2046 (70.7%)	2140 (72.9%)	
Nursing home	187 (6.5%)	139 (4.7%)	
Barthel index score (n, %)			1411 (1332 in 2008)
Total dependence (0–24)	354 (14.2%)	124 (5.8%)	
Severe dependence (25–49)	229 (9.2%)	108 (5.1%)	
Moderate dependence (50–74)	345 (13.8%)	241 (11.3%)	
Mild dependence (75–90)	465 (18.6%)	403 (18.9%)	
Negligible dependence (91–100)	1108 (44.3%)	1260 (59.0%)	
Number of drugs (n, %)			
0	81 (2.7%)	113 (3.7%)	
1–4	1048 (35.0%)	1252 (41.0%)	
≥ 5	1862 (62.3%)	1691 (55.3%)	
Number of concomitant comorbidities			
0	25 (0.8%)	73 (2.4%)	
1–2	229 (7.7%)	351 (11.5%)	
3–4	455 (15.2%)	674 (22.0%)	
≥ 5	2282 (76.3%)	1958 (64.1%)	
Season of hospitalization (n, %)			
Spring	800 (26.8%)	827 (27.1%)	
Summer	544 (18.2%)	450 (14.7%)	
Autumn	707 (23.6%)	825 (27%)	

Variables	Patients with infection N (%)	Patients without infections	Missing data
Winter	940 (31.4%)	954 (31.2%)	
Associated chronic diseases (n, %)			
Diabetes	844 (28.2%)	800 (26.2%)	
Heart failure	504 (16.8%)	519 (17.0%)	
Dementia	74 (2.5%)	488 (16.0%)	
Parkinson disease	105 (3.5%)	428 (14.0%)	
Chronic renal failure	567 (19.0%)	76 (2.5%)	
Cancer	538 (18.0%)	38 (1.2%)	
Patient status at discharge (n, %)			35
Home	2556 (85.8%)	2695 (88.9%)	
Dead during hospitalization	176 (5.9%)	86 (2.8%)	
Transferred to another medical ward	249 (8.4%)	250 (8.3%)	

Table 2 Prevalence of each infection typical in the REPOSI register from 2010 to 2016

Type of infection	Number (%)
Respiratory tract infections	1357 (38.2%)
Lower	1074
Upper	267
Not classified	16
Gastrointestinal tract infections	281 (7.9%)
Urinary tract infections	750 (21.1%)
Hepatobiliary infections	330 (9.3%)
Cardiac infections	32 (0.9%)
Septicemia	194 (5.5%)
Mycosis	178 (5.0%)
Others	432 (12.1%)