

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Letter: HBsAg kinetics-guided interferon therapy for chronic hepatitis D - authors' reply**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1640645> since 2017-05-31T17:25:45Z

*Published version:*

DOI:10.1111/apt.13892

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



# UNIVERSITÀ DEGLI STUDI DI TORINO

***This is an author version of the contribution published on:***

*Questa è la versione dell'autore dell'opera:*

*[Aliment Pharmacol Ther., 45(3), 2017, doi: 10.1111/apt.13892.]*

*ovvero [Niro GA, Smedile A, Andriulli A, Rizzetto M, 45, Wiley & Sons, 2017, pagg.481-482]*

***The definitive version is available at:***

*La versione definitiva è disponibile alla URL:*

*[<http://onlinelibrary.wiley.com.offcampus.dam.unito.it/doi/10.1111/apt.1>]*

**Letter: HBsAg kinetics-guided interferon therapy for chronic hepatitis D – authors' reply**

Niro GA<sup>1</sup>, Smedile A<sup>2</sup>, Andriulli A<sup>1</sup>, Rizzetto M<sup>2</sup>

<sup>1</sup> Gastroenterology Unit, IRCCS "Casa Sollievo Sofferenza" Hospital, San Giovanni Rotondo (FG), Italy.

<sup>2</sup> Division of Gastroenterology and Hepatology, Department of Medical Sciences, Città della Salute e della Scienza University Hospital, University of Turin, Turin, Italy.

Sirs,

We thank Dr Kao et al. for their interest in our work.[1, 2] Increased attention is now paid to quantitative HBsAg, to predict not only sero-clearance of HBV in chronic HBV infection, but also clearance of HDV in chronic HDV/HBV co-infection. HDV is highly infectious[3]: if a patient remains HBsAg positive after apparent successful elimination of HDV-RNA, very small amounts of the virus undetectable by current assays, can restore the infection and repeat liver damage.[4] On this basis, several authors have identified that a decreasing titre of serum HBsAg is a useful tool to predict and interpret the response to IFN, and to adapt the duration of IFN treatment with the ultimate aim to eradicate HDV through eradication of the HBsAg.[5-7]

The data of Kao et al. confirm this strategy. They report a HBV/HDV-positive patient who cleared both viruses after 35 months of a combination therapy with Peg-IFN plus entecavir, and a consolidation therapy after HBsAg loss.[7] Quantitative HBsAg was less than 300 IU/mL at baseline, and the progressive decline of HBsAg was followed by HBsAg loss at month 23, with a protective anti-HBs level 2 months later. Another HDV cirrhotic patient underwent treatment with interferon: quantitative HBsAg reduction of 0.205 at 6 months of therapy proved useful in predicting the HDV-RNA clearance

following therapy. In the last patient, HBsAg became negative at 3 months off therapy, and the treatment was stopped after 30 months due to intolerance.

The common denominator with our[1, 2, 7] and other studies[5, 6] is the decrease of circulating HBsAg at month 6 of therapy, the decline of which seems a useful criterion to decide whether to continue or withdraw IFN treatment. Prolonged treatments should be considered in patients who clear the HDV-RNA but remain HBsAg positive at declining titres during the first months. An unsolved question remains the need of consolidation therapy either after HBsAg clearance or seroconversion.

### **Acknowledgement**

The authors' declarations of personal and financial interests are unchanged from those in the original article.<sup>2</sup>

### **References**

- 1 Kao C-N, Su T-H, Kao J-H. Letter: HBsAg kinetics-guided interferon therapy for chronic hepatitis D. *Aliment Pharmacol Ther* 2017; 45: 480–1.
- 2 Niro GA, Smedile A, Fontana R, et al. HBsAg kinetics in chronic hepatitis D during interferon therapy: on-treatment prediction of response. *Aliment Pharmacol Ther* 2016; 44: 620–8.
- 3 Ponzetto A, Hoyer BH, Popper H, et al. Titration of the infectivity of hepatitis D virus in chimpanzees. *J Infect Dis* 1987; 155: 72–8.
- 4 Heidrich B, Yurdaydin C, Kabacam G, et al. Late HDV-RNA relapse after peginterferon alpha-based therapy of chronic hepatitis delta. *Hepatology* 2014; 60: 87–97.

5 Manesis EK, Schina M, Le Gal F, et al. Quantitative analysis of hepatitis D virus RNA and hepatitis B surface antigen serum levels in chronic delta hepatitis improves treatment monitoring.

*Antivir Ther* 2007; 12: 381–8.

6 Ouzan D, Pénaranda G, Joly H, Halfon P. Optimized HBsAg titer monitoring improves interferon therapy in patients with chronic hepatitis delta. *J Hepatol* 2013; 58: 1258–9.

7 Chen GY, Su TH, Kao JH. Successful treatment of chronic hepatitis B and D with pegylated-interferon plus entecavir. *J Formos Med Assoc* 2015; 114: 1140–1.