



AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Lagrange--Hermite interpolation on the real semiaxis

| This is the author's manuscript |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Original Citation: |
| |
| |
| |
| Availability: |
| This version is available http://hdl.handle.net/2318/1726698 since 2020-04-07T09:30:46Z |
| |
| |
| Published version: |
| DOI:10.1007/s10092-015-0147-y |
| Terms of use: |
| Open Access |
| Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law. |

(Article begins on next page)





This is the author's final version of the contribution published as:

Giuseppe Mastroianni, Incoronata Notarangelo and Pietro Pastore Lagrange– Hermite interpolation on the real semiaxis. *Calcolo*, 53 (2016), 235– 261. DOI: 10.1007/s10092-015-0147-y

The publisher's version is available at: [https://link.springer.com/article/10.1007/s10092-015-0147-y]

When citing, please refer to the published version.

Link to this full text: [http://hdl.handle.net/2318/1726698]

This full text was downloaded from iris-AperTO: https://iris.unito.it/

iris-AperTO

University of Turin's Institutional Research Information System and Open Access Institutional Repository

LAGRANGE–HERMITE INTERPOLATION ON THE REAL SEMIAXIS

G. MASTROIANNI, I. NOTARANGELO AND P. PASTORE

ABSTRACT. In order to approximate continuous functions on $[0, +\infty)$, we consider a Lagrange–Hermite polynomial, interpolating a finite section of the function at the zeros of some orthogonal polynomials and, with its first (r-1) derivatives, at the point 0. We give necessary and sufficient conditions on the weights for the uniform boundedness of the related operator. Moreover, we prove optimal estimates for the error of this process in the weighted L^p and uniform metric.

Keywords: Hermite–Lagrange interpolation; approximation by algebraic polynomials; orthogonal polynomials; generalized Laguerre weights; real semiaxis.

MCS classification (2000): 41A05, 41A10.

1. INTRODUCTION

In this paper we discuss the weighted polynomial interpolation of continuous functions on $[0, +\infty)$, which are (r-1)-times differentiable at 0 and can increase with order $\mathcal{O}\left(e^{x^{\beta}/2}\right), \beta > 1/2$, for $x \to \infty$.

The case $\beta = 1$ has been treated in [1, 5, 15, 20, 22, 24, 25, 26], where the Authors considered Lagrange polynomials based on Laguerre zeros (see also [8]). Here we choose as a tool the orthonormal system $\{p_m(w)\}_m$ in $(0, +\infty)$ related to a generalized Laguerre weight of the form $w(x) = x^{\alpha} e^{-x^{\beta}}$. From the numerical point of view, we observe that the weight w is nonclassical in general and, for the construction of the orthonormal system in the case $\beta \neq 1$, we can use the procedure introduced in [2] (see also [16]).

The presence of the derivatives of the function in 0 leads in a natural way to the construction of Lagrange–Hermite polynomial $\mathcal{L}_{m,r}(w, f)$ based at the zeros of the polynomial $p_m(w)$, 0 as a multiple node and another additional node. Applying the operator $\mathcal{L}_{m,r}(w)$ to a suitable finite section of the function f, we obtain a new interpolation process, that we will denote by $\mathcal{L}_{m,r}^*(w)$. This new operator is not a projector into the set of all polynomials of degree at most m + r, \mathbb{P}_{m+r} , but

The first author and the third authors were partially supported by University of Basilicata (local funds). The second author was supported by University of Basilicata (local funds) and by National Group of Computing Science GNCS–INDAM..

on a special subset $\mathcal{P}_{m+r}^* \subset \mathbb{P}_{m+r}$ that can replace the space \mathbb{P}_{m+r} , in the sense we are going to show. Thus the projector $\mathcal{L}_{m,r}^*(w)$ can be profitably used in quadrature rules and in the numerical treatment of Boundary Value Problems on $(0, \infty)$ (see, e.g., [6, 27]).

We estimate the error of the process in weighted L^p and uniform metric and compare it with the order of convergence of the best weighted polynomial approximation (that can be found in [21]). The error estimates are sharp for the considered classes of functions. All the results of this paper are new and cover the ones in the literature.

The paper is structured as follows. In Section 2 we recall some basic facts and give some preliminary results, in Section 3 we will state our main results and in Section 4 we will prove them.

2. Preliminary results

In the sequel \mathcal{C} will stand for a positive constant that can assume different values in each formula and we shall write $\mathcal{C} \neq \mathcal{C}(a, b, ...)$ when \mathcal{C} is independent of a, b, ... Furthermore $A \sim B$ will mean that if A and B are positive quantities depending on some parameters, then there exists a positive constant \mathcal{C} independent of these parameters such that $(A/B)^{\pm 1} \leq \mathcal{C}$. Finally, we denote by \mathbb{P}_m the set of all algebraic polynomials of degree at most m.

In order to introduce some interpolation processes we consider the weight

(2.1)
$$w(x) = x^{\alpha} e^{-x^{\beta}}, \quad x \in (0, +\infty),$$

with $\alpha > -1$, $\beta > \frac{1}{2}$, and the corresponding sequence of orthonormal polynomials $\{p_m(w)\}_m$, with positive leading coefficient γ_m . The zeros $x_k = x_{m,k}(w)$ of $p_m(w)$, $m \ge 1$, are located as follows

$$\left(\frac{\sqrt{a_m}}{m}\right)^2 < x_1 < x_2 < \ldots < x_m < a_m \left(1 - \frac{c}{m^{2/3}}\right),$$

where

$$a_m = a_m(\sqrt{w}) = \left[\frac{2^{2\beta-2}\Gamma(\beta)^2}{\Gamma(2\beta)}\right]^{1/\beta} (4m+2\alpha+1)^{1/\beta} \sim m^{1/\beta}$$

is the Mhaskar–Rakhmanov–Saff number related to the weight \sqrt{w} (see also [18, 19]). If $\beta = 1$ the nodes x_k are the Laguerre zeros. Nevertheless, in general w is a nonstandard weight and, for the computation of the zeros and the Christoffel numbers, we may use the Mathematica package "OrthogonalPolynomials" introduced in [2]. For any continuous function in $[0, +\infty)$, $f \in C^0([0, +\infty))$, and (r-1)-times differentiable at 0, briefly $f \in C_r^0$, $r \geq 1$, we define the Lagrange–Hermite polynomial $\mathcal{L}_{m,r}(w, f)$ as follows

$$\mathcal{L}_{m,r}(w,f,x_k) = f(x_k), \qquad k = 1, \dots, m+1,$$

with $x_{m+1} = a_m$, and

$$\mathcal{L}_{m,r}(w,f)^{(j)}(0) = f^{(j)}(0), \qquad j = 0, 1, \dots, r-1,$$

where $f^{(0)} \equiv f$. Here we used also an idea introduced by J. Szabados for Lagrange and Hermite–Fejér interpolation, adding the additional node a_m (see [28]).

The polynomial $\mathcal{L}_{m,r}(w, f)$ can be written as

$$\mathcal{L}_{m,r}(w, f, x) = \sum_{k=1}^{m+1} \frac{x^r}{x_k^r} \ell_k(w, x) f(x_k)$$

$$(2.2) + (a_m - x) p_m(w, x) \sum_{i=0}^{r-1} \frac{x^i}{i!} \left(\frac{f}{(a_m - \cdot) p_m(w)}\right)^{(i)}(0),$$

where

$$\ell_k(w,x) = \frac{p_m(w,x)}{(x-x_k)p'_m(w,x_k)} \frac{(a_m-x)}{(a_m-x_k)}, \qquad k = 1, 2, \dots, m,$$

and

$$\ell_{m+1}(w,x) = \frac{p_m(w,x)}{p_m(w,a_m)}$$

are the fundamental Lagrange polynomials. It is easily seen that

$$\mathcal{L}_{m,r}(w,P) = P, \qquad P \in \mathbb{P}_{m+r},$$

and $\mathcal{L}_{m,r}(w)$ is a projector from C_r^0 into \mathbb{P}_{m+r} .

We are now going to introduce another Lagrange–Hermite type operator $\mathcal{L}_{m,r}^*(w)$, modifying the operator $\mathcal{L}_{m,r}(w)$. To this end, for a fixed $\theta \in (0,1)$ and for a sufficiently large m we define an index $j = j(m,\theta)$ such that

(2.3)
$$x_j = \min_k \{x_k : x_k \ge \theta a_m\}.$$

Of course if m is small we let j = m. Hence we define the new Lagrange–Hermite type polynomial as follows

$$\mathcal{L}_{m,r}^{*}(w, f, x) = \sum_{k=1}^{j} \frac{x^{r}}{x_{k}^{r}} \ell_{k}(w, x) f(x_{k}) + (a_{m} - x) p_{m}(w, x) \sum_{i=0}^{r-1} \frac{x^{i}}{i!} \left(\frac{f}{(a_{m} - \cdot)p_{m}(w)}\right)^{(i)}(0)$$

By this definition,

$$\mathcal{L}_{m,r}^{*}(w, f, x_{k}) = f(x_{k}), \qquad k = 1, \dots, j,$$

 $\mathcal{L}_{m,r}^{*}(w, f, x_{k}) = 0, \qquad k = j + 1, \dots, m + 1,$

and

$$\mathcal{L}_{m,r}^{*}(w,f)^{(i)}(0) = f^{(i)}(0), \qquad i = 0, 1, \dots, r-1.$$

Moreover, we observe that $\mathcal{L}_{m,r}^*(w)$ does not preserve all the polynomials of degree at most m+r, for example $\mathcal{L}_{m,r}^*(w,1) \neq 1$.

Nevertheless, if we introduce the following set of polynomials

$$\mathcal{P}_{m,r}^* = \{ Q \in \mathbb{P}_{m+r} : Q(x_i) = 0, \, i > j \}$$

then it is easy to show that for any $f \in C_r^0$, $\mathcal{L}_{m,r}^*(w, f) \in \mathcal{P}_{m,r}^*$ and for any $Q \in \mathcal{P}_{m,r}^*$, $\mathcal{L}_{m,r}^*(w, Q) = Q$, therefore $\mathcal{L}_{m,r}^*(w, f)$ is a projector from C_r^0 into $\mathcal{P}_{m,r}^*$.

In order to show some approximation properties of the projectors defined above, we are going to define some function spaces.

2.1. Function spaces. Let us consider a weight of the form

(2.5)
$$u(x) = x^{\gamma} e^{-x^{\beta}/2}, \qquad x \in (0, +\infty),$$

with $\beta > \frac{1}{2}$ and the following weighted function spaces associated to u.

For $1 \leq p < \infty$ and $\gamma > -1/p$, by L_u^p we denote the set of all measurable functions f such that

$$||f||_{L^p_u} := ||fu||_p = \left(\int_0^{+\infty} |fu|^p(x) \,\mathrm{d}x\right)^{1/p} < \infty.$$

For $p = \infty$ and $\gamma \ge 0$, by a slight abuse of notation, we set

$$L_u^{\infty} := C_u = \left\{ f \in C^0(0, +\infty) : \lim_{\substack{x \to 0^+ \\ x \to +\infty}} f(x)u(x) = 0 \right\},$$

and we equip this space with the norm

$$||f||_{L^{\infty}_{u}} := ||fu||_{\infty} = \sup_{x \in (0, +\infty)} |f(x)u(x)|.$$

Note that the Weierstrass theorem implies the limit conditions in the definition of C_u .

Subspaces of L_u^p , $1 \le p \le \infty$, are the Sobolev spaces, given by

$$W_s^p(u) = \left\{ f \in L_u^p : \ f^{(s-1)} \in AC(0, +\infty), \ \|f^{(s)}\varphi^s u\|_p < \infty \right\}, \quad 1 \le s \in \mathbb{Z},$$

where $AC(0, +\infty)$ denotes the set of all functions which are absolutely continuous on every closed subset of $(0, +\infty)$ and $\varphi(x) = \sqrt{x}$. We equip these spaces with the norm

$$||f||_{W^p_s(u)} = ||fu||_p + ||f^{(s)}\varphi^s u||_p.$$

In order to define some further function spaces, we consider the s-th modulus of smoothness of $f \in L^p_u$, $1 \le p \le \infty$, $s \ge 1$,

$$\Omega_{\varphi}^{s}(f,t)_{u,p} = \sup_{0 < h \le t} \left\| \Delta_{h\varphi}^{s}(f) \, u \right\|_{L^{p}(I_{rh})},$$

where

$$\Delta_{h\varphi}^{s}f(x) = \sum_{i=0}^{s} (-1)^{i} {\binom{s}{i}} f\left(x + (s-i)h\varphi(x)\right)$$

is the forward finite difference of order s with variable step $h\varphi(x)$, $\varphi(x) = \sqrt{x}$ and $I_{sh} = [8s^2h^2, Ch^{-\frac{1}{(\beta-1/2)}}], h > 0.$

Let $f \in L^p_u$, $1 \le p \le \infty$. Then the following estimate

(2.6)
$$\Omega_{\varphi}^{s}(f,t)_{u,p} \leq \mathcal{C} \sup_{0 < h \leq t} h^{s} \|f^{(s)}\varphi^{s}u\|_{L^{p}(I_{sh})}$$

holds with $C \neq C(f, t)$, provided the norm on the right-hand side is finite (see [21]).

Using the previous moduli of smoothness we can define the Zygmund spaces as

$$Z_{\lambda}^{p}(u) = \left\{ f \in L_{u}^{p} : \sup_{t > 0} \frac{\Omega_{\varphi}^{s}(f, t)_{u, p}}{t^{\lambda}} < \infty, \ s > \lambda \right\}, \qquad \lambda \in \mathbb{R}, \ \lambda > 0,$$

with the norm

$$\|f\|_{Z^p_\lambda(u)}=\|fu\|_p+\sup_{t>0}\frac{\Omega^s_\varphi(f,t)_{u,p}}{t^\lambda}\,,\quad s>\lambda\,.$$

Let us denote by

$$E_m(f)_{u,p} = \inf_{P \in \mathbb{P}_m} \left\| (f - P) \, u \right\|_p$$

the error of best weighted polynomial approximation in L_u^p , $1 \le p \le \infty$. Estimates for the error of best weighted approximation have been proved in [21]. In particular it has been shown that

(2.7)
$$\lim_{m \to \infty} E_m(f)_{u,p} = 0 \qquad \forall f \in L^p_u$$

For our aims it is sufficient to recall the following weak Jackson inequality

(2.8)
$$E_m(f)_{u,p} \le \mathcal{C} \int_0^{\frac{\sqrt{a_m}}{m}} \frac{\Omega_{\varphi}^s(f,t)_{u,p}}{t} \, \mathrm{d}t$$

that holds for any function $f \in L^p_u$, $1 \le p \le \infty$.

In particular, if $f \in Z_{\lambda}^{p}$, by (2.8) we deduce

(2.9)
$$E_m(f)_{u,p} \le \mathcal{C}\left(\frac{\sqrt{a_m}}{m}\right)^{\lambda} \sup_{t>0} \frac{\Omega_{\varphi}^s(f,t)_{u,p}}{t^{\lambda}},$$

and by (2.6) we have

(2.10)
$$E_m(f)_{u,p} \le \mathcal{C}\left(\frac{\sqrt{a_m}}{m}\right)^s \|f^{(s)}\varphi^s u\|_p,$$

for any $f \in W_s^p$, $s \ge 1$ and $1 \le p \le \infty$.

We are now able to show an important property of the polynomials in $\mathcal{P}_{m,r}^*$. To this end we need some further notation. We say that P_M is a polynomial of quasi best approximation for $f \in L^p_u$, $1 \le p \le \infty$, if, for some $\mathcal{C} \ge 1$,

$$\left\| \left(f - P_M \right) u \right\|_p \le \mathcal{C} E_M(f)_{u,p}$$

Moreover, we denote by

$$E_{m,r}^{*}(f)_{u,p} = \inf_{Q \in \mathcal{P}_{m,r}^{*}} \| (f-Q)u \|_{p},$$

the error of best approximation by means of polynomials of $\mathcal{P}_{m,r}^*$. The relation between $E_m(f)_{u,p}$ and $E_m^*(f)_{u,p}$ is established by the following Lemma.

Lemma 1. Let w, u be the weights defined in (2.1) and (2.5) with $\alpha > -1$, $\beta > \frac{1}{2}$ and $u \in L^p$, $1 \leq p \leq \infty$. Let $f \in L^p_u$ and $P_M \in \mathbb{P}_M$ one of its polynomial of quasi best approximation, with

(2.11)
$$M = \left\lfloor \left(\frac{\theta}{\theta+1}\right)m\right\rfloor$$

for a fixed $\theta \in (0,1)$. Then we have (2.12)

$$E_{m,r}^{*}(f)_{u,p} \leq \|[f - \mathcal{L}_{m,r}^{*}(w, P_{M})]u\|_{p} \leq \mathcal{C}\{E_{M}(f)_{u,p} + e^{-cm}\|fu\|_{p}\},\$$

where C, c are independent of m, r, f. Moreover, for $s \ge 1$, we get (2.13)

$$\left(\frac{\sqrt{a_m}}{m}\right)^s \|\mathcal{L}_{m,r}^*(w, P_M)^{(s)}\varphi^s u\|_p \le \mathcal{C}\left\{e^{-cm}\|fu\|_p + \int_0^{\frac{\sqrt{a_m}}{m}} \frac{\Omega_{\varphi}^s(f, t)_{u,p}}{t} \,\mathrm{d}t\right\}.$$

Using Lemma 1 and (2.7) it follows that the order of convergence of $E_{m,r}^*(f)_{u,p}$ is the same as that of $E_{m,r}(f)_{u,p}$. Therefore, $\bigcup_m \mathcal{P}_{m,r}^*$ is dense in L_u^p . In the next Section we will show the behaviour of the operators $\mathcal{L}_{m,r}(w)$ and $\mathcal{L}_{m,r}^*(w)$ in different function spaces.

3. Main results

The following results hold true in weighted uniform norm. In order to state it we need a further definition. We say that a function is quasi increasing on I if there exists a constant C such that $f(x) \leq Cf(y)$ for $x < y, x, y \in I$.

Theorem 2. Let w and u be the previously defined weights, with $\gamma \geq 0$ and $\alpha > -1$. For every $f \in C_r^0 \cap C_u$, $r \geq 1$, we have (3.1)

$$\|\mathcal{L}_{m,r}(w,f)\,u\|_{\infty} \le \mathcal{C}\left\{ (\log m)\|fu\|_{\infty} + \left(\frac{\sqrt{a_m}}{m}\right)^{2\gamma} \sum_{i=0}^{r-1} \frac{|f^{(i)}(0)|}{i!} \left(\frac{\sqrt{a_m}}{m}\right)^{2i} \right\}$$

where $C \neq C(m, f)$, if and only if

(3.2)
$$\frac{\alpha}{2} + \frac{1}{4} \le \gamma + r \le \frac{\alpha}{2} + \frac{5}{4}$$

Moreover, under the assumptions (3.2), if $|f^{(i)}|$ is quasi increasing on $[0, \alpha_m]$, with $\alpha_m = a_m/m^2$ and i = 0, 1, ..., r-1, we get (3.3)

$$\left\| \left[f - \mathcal{L}_{m,r}\left(w,f\right) \right] u \right\|_{\infty} \le \mathcal{C}(\log m) E_{m+r}(f)_{u,\infty} + \mathcal{O}\left(\left(\frac{\sqrt{a_m}}{m}\right)^{r-1} \right)$$

where $\mathcal{C} \neq \mathcal{C}(m, f)$ and the constant in " \mathcal{O} " is independent of m.

Therefore, in weighted spaces of continuous functions, the behaviour of the operators $\{\mathcal{L}_{m,r}(w)\}_m$ is comparable with similar interpolation processes based on Jacobi zeros on bounded intervals (see, e.g., [15, 17]).

We emphasize that if r and γ are given, then, in order to approximate a function f, we can always choose some α such that the Lagrange– Hermite polynomial converges with the order of the best polynomial approximation times the extra factor log m. In fact, we can rewrite (3.2) as follows

$$2r + 2\gamma - \frac{5}{2} \le \alpha \le 2r + 2\gamma - \frac{1}{2}.$$

Moreover, since the Lebesgue constants related to the processes $\{\mathcal{L}_{m,r}^*(w)\}_m$ are bounded by the ones related to $\{\mathcal{L}_{m,r}(w)\}_m$, by Theorem 2 and Lemma 1, we deduce the following error estimate for the "truncated" process.

Corollary 3. Let $\theta \in (0,1)$, γ and α as in Theorem 2. Then, for every $f \in C_r^0$, $r \ge 1$, for m sufficiently large (say $m > m_0$) we have

$$\left\| \left[f - \mathcal{L}_{m,r}^*\left(w,f\right) \right] u \right\|_{\infty} \leq \mathcal{C}(\log m) E_M(f)_{u,\infty} + \mathcal{O}\left(\left(\frac{\sqrt{a_m}}{m} \right)^{r-1} \right) ,$$

where M given by (2.11), $\mathcal{C} \neq \mathcal{C}(m, f)$ and the constant in " \mathcal{O} " is independent of m.

In weighted L^p - spaces the behaviour of $\{\mathcal{L}_{m,r}(w)\}_m$ is not optimal, while for the new process $\{\mathcal{L}_{m,r}^*(w)\}_m$ we can state the following

Theorem 4. Let $u \in L^p$, $1 and <math>\theta \in (0, 1)$ be fixed. Then, for every function $f \in C_r^0$, $r \ge 1$, we have

(3.4)
$$\begin{aligned} \left\| \mathcal{L}_{m,r}^{*}(w,f) \, u \right\|_{p} &\sim \left(\sum_{k=1}^{j} \Delta x_{k} |fu|^{p}(x_{k}) \right)^{1/p} \\ &+ \left(\frac{\sqrt{a_{m}}}{m} \right)^{2(\gamma+1/p)} \sum_{i=0}^{r-1} \frac{|f^{(i)}(0)|}{i!} \left(\frac{\sqrt{a_{m}}}{m} \right)^{2i} \end{aligned}$$

if and only if

(3.5)
$$\frac{\alpha}{2} + \frac{1}{4} - \frac{1}{p} < \gamma + r < \frac{\alpha}{2} + \frac{5}{4} - \frac{1}{p}$$

where the constants in "~" depend on θ and are independent of m and f.

If the function $f \in L^p_u$ fulfills the additional assumption

(3.6)
$$\frac{\Omega^s_{\varphi}(f,t)_{u,p}}{t^{1+1/p}} \in L^1$$

for some $s \ge 1$, then f is continuous in $(0, +\infty)$ (see [21]). Moreover, using the same arguments as in [22], we obtain

$$\left(\sum_{k=1}^{j} \Delta x_k |fu|^p(x_k)\right)^{1/p} \le \mathcal{C}\left[\|fu\|_p + \left(\frac{\sqrt{a_m}}{m}\right)^{1/p} \int_0^{\frac{\sqrt{a_m}}{m}} \frac{\Omega_{\varphi}^s(f,t)_{u,p}}{t^{1+1/p}} \,\mathrm{d}t\right].$$

Hence, if $f \in L^p_u$ fulfills (3.6) and is (r-1)-times differentiable at 0, the bound (3.4) becomes

$$(3.\|\mathcal{L}_{m,r}^{*}(w,f) u\|_{p} \leq \mathcal{C} \left[\|fu\|_{p} + \left(\frac{\sqrt{a_{m}}}{m}\right)^{1/p} \int_{0}^{\frac{\sqrt{a_{m}}}{m}} \frac{\Omega_{\varphi}^{s}(f,t)_{u,p}}{t^{1+1/p}} dt + \left(\frac{\sqrt{a_{m}}}{m}\right)^{2(\gamma+1/p)} \sum_{i=0}^{r-1} \frac{|f^{(i)}(0)|}{i!} \left(\frac{\sqrt{a_{m}}}{m}\right)^{2i} \right],$$

where \mathcal{C} depends on $\theta \in (0, 1)$ (fixed) and is independent of m and f. Note that if $f \in \mathcal{P}_{m,r}^*$, (3.7) becomes an equivalence.

Theorem 5. Under the assumptions (3.5), let $f \in L^p_u \cap C^0_r$ satisfy (3.6) for some $s \ge r$ and $|f^{(i)}|$ be quasi increasing on $[0, \alpha_m]$, with $\alpha_m = a_m/m^2$ and $i = 0, 1, \ldots, r-1$. Then, for $m > m_0$, we have (3.8)

$$\|[f - \mathcal{L}_{m,r}^*(w,f)]u\|_p \le \mathcal{C}\left(\frac{\sqrt{a_m}}{m}\right)^{1/p} \int_0^{\frac{\sqrt{a_m}}{m}} \frac{\Omega_{\varphi}^s(f,t)_{u,p}}{t^{1+1/p}} \,\mathrm{d}t + \mathcal{O}\left(\left(\frac{\sqrt{a_m}}{m}\right)^{r-1}\right)$$

where C depends on θ and is independent of m and f, the constant in " \mathcal{O} " is independent of m.

In particular, under the assumptions of Theorem 5, if $f \in Z^p_{\lambda}(u)$, $1/p < \lambda \leq r - 1$, the error estimate (3.8) becomes

$$\left\| \left[f - \mathcal{L}_{m,r}^{*}\left(w,f\right) \right] u \right\|_{p} = \mathcal{O}\left(\left(\frac{\sqrt{a_{m}}}{m} \right)^{\lambda} \right),$$

where the constant in \mathcal{O} is independent of m and f, that is the order of the best approximation in $Z^p_{\lambda}(u)$ (see (2.9)).

Analogously, if $f \in W_s^p(u)$, $s \leq r-1$, from Theorem 5 we get

$$\|[f - \mathcal{L}_{m,r}^*(w, f)]u\|_p = \mathcal{O}\left(\left(\frac{\sqrt{a_m}}{m}\right)^s\right),$$

where the constant in \mathcal{O} is independent of m and f (see (2.10)). While, for $f \in W_r^p(u)$, the following corollary holds.

Corollary 6. Under the assumptions (3.5), let $f \in W_r^p(u) \cap C_r^0$, $1 , such that <math>|f^{(i)}|$ is quasi increasing on $[0, \alpha_m]$, $\alpha_m = a_m/m^2$, for $i = 0, 1, \ldots, r-1$ and m sufficiently large (say $m > m_0$). Then we have

(3.9)
$$\left\|\mathcal{L}_{m,r}^{*}\left(w,f\right)u\right\|_{p} \leq \mathcal{C}\left\{\|fu\|_{p} + \left(\frac{\sqrt{a_{m}}}{m}\right)^{r}\|f^{(r)}\varphi^{r}u\|_{p}\right\}$$

and

(3.10)
$$\left\| \left[f - \mathcal{L}_{m,r}^*\left(w,f\right) \right] u \right\|_p \le \mathcal{C} \left\{ \left(\frac{\sqrt{a_m}}{m} \right)^r \| f^{(r)} \varphi^r u \|_p + e^{-cm} \| f u \|_p \right\}$$

where C, c depend on θ and are independent of m and f.

Finally, the following remark is of some interest.

Remark 7. The behaviours of the two interpolation processes $\{\mathcal{L}_{m,r}(w)\}_m$ and $\{\mathcal{L}_{m,r}^*(w)\}_m$ are essentially equivalent in weighted uniform norm. Nevertheless, the "truncated" process $\{\mathcal{L}_{m,r}^*(w)\}_m$ has the advantage of dropping c m terms, c < 1. This turns out to be useful in the numerical treatment of functional equations, since the dimension of the matrices obtained from the discretization of operators are strongly reduced with an evident computational saving.

On the other hand, the two operators behave in a completely different way in weighted L^p -norm. In fact, the equivalence (3.4) of Theorem 4 is not true for the "nontruncated" operator $\{\mathcal{L}_{m,r}(w)\}_m$.

Moreover, all the constants in Corollary 3, Theorems 4 and 5 are independent of m and f, but depend on θ . For instance, the constant in the upper bound of (3.4) is $\mathcal{O}\left((1-\theta)^{-3/4}\right)$ for $\theta \to 1$ and so θ cannot assume value 1.

We refer to Section 4 for more details.

4. Proofs

First of all we recall some weighted polynomial inequalities with the weight u defined in (2.5) which will be used in the sequel (see [21]).

Let $1 \leq p \leq \infty$ and $a_m = a_m(u) \sim m^{1/\beta}$. For any $P_m \in \mathbb{P}_m$, the restricted range inequalities

(4.1)
$$||P_m u||_p \le \mathcal{C} ||P_m u||_{L^p(I_m)}, \qquad I_m = \left[c \, \frac{a_m}{m^2}, a_m\right],$$

and

(4.2)
$$||P_m u||_{L^p[a_m(1+\delta),+\infty)} \le C e^{-cm} ||P_m u||_p, \quad \delta > 0,$$

hold with $\mathcal{C} \neq \mathcal{C}(m, P_m)$ and $c \neq c(m, P_m)$. Moreover, we recall the Bernstein and Markov inequalities

(4.3)
$$\|P'_m \varphi u\|_p \le C \frac{m}{\sqrt{a_m}} \|P_m u\|_p, \quad \varphi(x) = \sqrt{x},$$

and

(4.4)
$$||P'_{m}u||_{p} \leq C \frac{m^{2}}{a_{m}} ||P_{m}u||_{p},$$

where $\mathcal{C} \neq \mathcal{C}(m, P_m)$ in both cases. Finally, for $1 \leq p < \infty$, we will need the following Nikolskii inequality

(4.5)
$$||P_m u||_{\infty} \leq \mathcal{C}\left(\frac{m^2}{a_m}\right)^{\frac{1}{p}} ||P_m u||_p, \quad \mathcal{C} \neq \mathcal{C}(m, P_m).$$

We also need some estimates for the polynomials of the orthonormal system $\{p_m(w)\}_{m\in\mathbb{N}}$, where w is the weight defined in (2.1). The generalized Laguerre weights w is a nonstandard weight and the estimates for the related orthogonal polynomials can be deduced by [7, 11, 12, 21, 22].

The estimate (which can be deduced from [9], see also [21, 19])

(4.6)
$$|p_m(w,x)| \left(x + \frac{a_m}{m^2}\right)^{\frac{\alpha}{2} + \frac{1}{4}} e^{-\frac{x^{\beta}}{2}} \sqrt[4]{|a_m - x|} + \frac{a_m}{m^{2/3}} \sim \frac{|x - x_d|}{\Delta x_{d\pm 1}}$$

holds with $x \in [0, +\infty)$ and x_d a zero closest to x. From (4.6) it follows that

(4.7)
$$|p_m(w,x)|\sqrt{w(x)}\sqrt{x(a_m-x)} \le \mathcal{C}, \qquad x \in I_m.$$

Moreover, the following proposition will be useful.

Proposition 8. Let w be the weight defined by (2.1) and $\{p_m(w)\}_m$ its related orthonormal system. Then we have

(4.8)
$$|p_m(w,0)| \sim \left(\frac{m}{\sqrt{a_m}}\right)^{\alpha+\frac{1}{2}} \frac{1}{\sqrt[4]{a_m}},$$

(4.9)
$$|p_m^{(k)}(w,0)| \le C \left(\frac{m}{\sqrt{a_m}}\right)^{2k} |p_m(w,0)|,$$

and

(4.10)
$$\left| \left(\frac{1}{p_m(w)} \right)^{(k)}(0) \right| \le \mathcal{C} \left(\frac{m}{\sqrt{a_m}} \right)^{2k} \frac{1}{|p_m(w,0)|}$$

where \mathcal{C} and the constant in " \sim " are independent of m.

Proof. Equivalence (4.8) easily follows from (4.6) for x = 0. In order to prove (4.9) we set

$$g(x) = \left(x + \frac{a_m}{m^2}\right)^{\frac{\alpha}{2} + \frac{1}{4}} e^{-\frac{x^{\beta}}{2}} \sqrt[4]{|a_m - x|} + \frac{a_m}{m^{2/3}}.$$

Then, using (4.4) with u replaced by g, by (4.6), we obtain

$$|p_m^{(k)}(w,x)| \le \mathcal{C}\left(\frac{m}{\sqrt{a_m}}\right)^{2k} \frac{\|p_m(w)g\|_{\infty}}{|g(x)|} \le \mathcal{C}\left(\frac{m}{\sqrt{a_m}}\right)^{2k} \frac{1}{|g(x)|}$$

Hence, for x = 0, by (4.8), we get

$$|p_m^{(k)}(w,0)| \le \mathcal{C}\left(\frac{m}{\sqrt{a_m}}\right)^{2k} \frac{1}{\left(\frac{\sqrt{a_m}}{m}\right)^{\alpha+\frac{1}{2}} \sqrt[4]{a_m}} \le \mathcal{C}\left(\frac{m}{\sqrt{a_m}}\right)^{2k} |p_m(w,0)|.$$

Finally, let us show (4.10) by induction. For k = 1, by (4.9) we have

$$\left| \left(\frac{1}{p_m(w)} \right)'(0) \right| = \frac{\left| p'_m(w,0) \right|}{\left| p_m(w,0) \right|^2} \le \mathcal{C} \left(\frac{m}{\sqrt{a_m}} \right)^2 \frac{1}{\left| p_m(w,0) \right|}$$

For k > 1, from the identity

$$\left(\frac{1}{p_m(w)}p_m(w)\right)^{(k)}(0) = \sum_{i=0}^k \binom{k}{i} \left(\frac{1}{p_m(w)}\right)^{(i)}(0)p_m^{(k-i)}(w,0) = 0$$

it follows that

$$-\left(\frac{1}{p_m(w)}\right)^{(k)}(0)p_m(w,0) = \sum_{i=0}^{k-1} \binom{k}{i} \left(\frac{1}{p_m(w)}\right)^{(i)}(0)p_m^{(k-i)}(w,0)$$

Then, using the induction hypothesis and (4.9), we obtain

$$|p_m(w,0)| \left| \left(\frac{1}{p_m(w)} \right)^{(k)}(0) \right| = \left| \sum_{i=0}^{k-1} \binom{k}{i} \left(\frac{1}{p_m(w)} \right)^{(i)}(0) p_m^{(k-i)}(w,0) \right|$$
$$\leq \mathcal{C} \left(\frac{m}{\sqrt{a_m}} \right)^{2k},$$

which completes the proof.

Furthermore, the following relation will be useful (see [10, 21])

(4.11)
$$\frac{1}{|p'_m(w, x_k)|\sqrt{w(x_k)}} \sim \Delta x_k \sqrt[4]{|a_m - x_k|} + \frac{a_m}{m^{2/3}}.$$

By (4.6) and (4.11) if $x \in \left[\left(\frac{\sqrt{a_m}}{m}\right)^2, a_m\right]$ we have (4.12)

$$u(x)\frac{|\ell_k(w,x)|}{u(x_k)} \le \mathcal{C}\left(\frac{x}{x_k}\right)^{\gamma-\frac{\alpha}{2}-\frac{1}{4}} \left(\frac{a_m-x_k}{a_m-x}\right)^{1/4} \frac{\Delta x_k}{|x-x_k|}, \quad (x \ne x_k).$$

Moreover, if x_d is a node closest to x, we get (see [4])

(4.13)
$$u(x)\frac{|\ell_d(w,x)|}{u(x_d)} \sim 1$$

Proof of Lemma 1. We are going to prove (2.12) only for $p < \infty$, since the case $p = \infty$ is simpler.

Let $P_M \in \mathbb{P}_M$, with $M = \lfloor \frac{\theta m}{\theta + 1} \rfloor$, be a polynomial of quasi best approximation for $f \in L^p_u$. So we can write

$$E_{m,r}^*(f)_{u,p} \le \left\| \left[f - \mathcal{L}_{m,r}^*(w, P_M) \right] u \right\|_p.$$

Since

(4.14)
$$\mathcal{L}_{m,r}^{*}(w, P_{M}) = P_{M} - \sum_{k=j+1}^{m} \frac{\ell_{k}(w, x)x^{r}}{x_{k}^{r}} P_{M}(x_{k})$$

then, letting $v^r(x) = x^r$, we get

$$E_{m,r}^{*}(f)_{u,p} \leq \mathcal{C}E_{M}(f)_{u,p} + \left\| \sum_{k=j+1}^{m} \frac{\ell_{k}(w)v^{r}u}{v^{r}(x_{k})u(x_{k})} (P_{M}u)(x_{k}) \right\|_{p}.$$

By (4.1) the second summand on the right-hand side is dominated by

$$\mathcal{C}a_m^{1/p} \|P_M u\|_{L^{\infty}[\theta a_m, +\infty)} \sup_{x \in I_m} \sum_{k=j+1}^m \frac{|\ell_k(w, x)| v^r(x) u(x)}{v^r(x_k) u(x_k)},$$

and, using (4.12) and (4.13), we have

(4.15)
$$\sup_{x \in I_m} \sum_{k=j+1}^m \frac{|\ell_k(w,x)| v^r(x) u(x)}{v^r(x_k) u(x_k)} \le \mathcal{C}m^{\tau},$$

for some $\tau > 0$. Moreover, by (4.2) and (4.5), we obtain

$$\left\|\sum_{k=j+1}^{m} \frac{\ell_k(w)v^r u}{v^r(x_k)u(x_k)} (P_M u)(x_k)\right\|_p \leq Cm^{\tau} a_m^{1/p} \|P_M u\|_{L^{\infty}[\theta a_m, +\infty)}$$

$$\leq Cm^{\tau} a_m^{1/p} e^{-cm} \|P_M u\|_{\infty}$$

$$\leq Cm^{\tau+\frac{2}{p}} e^{-cm} \|P_M u\|_p$$

$$\leq Ce^{-cm} \|fu\|_p.$$

Let us now prove (2.13). By (4.14) we have

$$\left(\frac{\sqrt{a_m}}{m}\right)^r \|\mathcal{L}_{m,r}^*(w, P_M)^{(r)} \varphi^r u\|_p$$

$$\leq \left(\frac{\sqrt{a_m}}{m}\right)^r \|P_M^{(r)} \varphi^r u\|_p + \left(\frac{\sqrt{a_m}}{m}\right)^r \left\| \left(\sum_{k=j+1}^m \frac{\ell_k(w)v^r}{v^r(x_k)} P_M(x_k)\right)^{(r)} \varphi^r u \right\|_p$$

For the first term on the right-hand side we use the following estimate (see [21])

$$\left(\frac{\sqrt{a_m}}{m}\right)^r \|P_M^{(r)}\varphi^r u\|_p \le \mathcal{C} \int_0^{\sqrt{a_m}/m} \frac{\Omega_{\varphi}^r (f,t)_{u,p}}{t} \, \mathrm{d}t \, .$$

While, for the second term on the right-hand side, using k times the Bernstein inequality (4.3) and (4.16), we obtain

$$\left(\frac{\sqrt{a_m}}{m}\right)^r \left\| \left(\sum_{k=j+1}^m \frac{\ell_k(w)v^r}{v^r(x_k)} P_M(x_k) \right)^{(r)} \varphi^r u \right\|_p \leq \mathcal{C} \left\| \sum_{k=j+1}^m \frac{\ell_k(w)v^r}{v^r(x_k)} P_M(x_k) u \right\|_p \leq \mathcal{C} e^{-cm} \|fu\|_p.$$

Hence we get (2.13).

We recall the following estimate, concerning the distance Δx_k between two consecutive zeros (see [10, 21]):

(4.17)
$$\Delta x_k = x_{k+1} - x_k \sim \frac{a_m}{m} \sqrt{x_k} \frac{1}{\sqrt{a_m - x_k + a_m m^{-2/3}}},$$

where the constants in " \sim " are independent of m and k. We remark that from (4.17), with j given by (2.3), it follows that

$$\Delta x_k \sim \frac{\sqrt{a_m}}{m} \sqrt{x_k} \qquad k = 1, 2, \dots j$$

Now, letting

(4.18)
$$\sigma_m(f) = \sum_{i=0}^{r-1} \frac{f^{(i)}(0)}{i!} \left(\frac{\sqrt{a_m}}{m}\right)^{2i}$$

and

(4.19)
$$A(x) = (a_m - x)p_m(w, x) \sum_{i=0}^{r-1} \frac{x^i}{i!} \left(\frac{f}{(a_m - \cdot)p_m(w)}\right)^{(i)}(0),$$

we can state the following proposition:

Proposition 9. With the notation (4.18) and (4.19), we have

(4.20)
$$||Au||_{\infty} \leq \mathcal{C}\left(\frac{\sqrt{a_m}}{m}\right)^{2\gamma} \sigma_m(f), \qquad \mathcal{C} \neq \mathcal{C}(m, f),$$

if and only if

(4.21)
$$\frac{\alpha}{2} + \frac{1}{4} \le \gamma + r \le \frac{\alpha}{2} + \frac{5}{4}.$$

Moreover, for $p \ge 1$, we get

(4.22)
$$||Au||_p \le \mathcal{C}\left(\frac{\sqrt{a_m}}{m}\right)^{2(\gamma+1/p)} \sigma_m(f), \qquad \mathcal{C} \neq \mathcal{C}(m, f),$$

if and only if

(4.23)
$$\frac{\alpha}{2} + \frac{1}{4} - \frac{1}{p} \le \gamma + r \le \frac{\alpha}{2} + \frac{5}{4} - \frac{1}{p}$$

Proof. We first prove (4.20). Recalling (4.19), we may write

(4.24)
$$|A(x)u(x)| \le \sum_{i=0}^{r-1} b_i \frac{|D_i|}{i!},$$

where

$$b_i = \|(a_m - \cdot)p_m(w, \cdot)uv^i\|_{\infty}, \qquad v^i(x) = x^i$$

and

$$|D_i| = \left| \left(\frac{f}{(a_m - \cdot)p_m(w)} \right)^{(i)}(0) \right|$$
$$= \left| \sum_{\eta=0}^i {i \choose \eta} f^{(\eta)}(0) \left(\frac{1}{(a_m - \cdot)p_m(w, \cdot)} \right)^{(i-\eta)}(0) \right|.$$

By (4.10) and (4.11) we get

$$\left| \left(\frac{1}{(a_m - \cdot)p_m(w, \cdot)} \right)^{(i-\eta)}(0) \right| = \left| \sum_{j=0}^{i-\eta} {i-\eta \choose j} \frac{1}{(a_m)^{j+1}} \left(\frac{1}{p_m(w, \cdot)} \right)^{(i-\eta-j)}(0) \right|$$
$$\leq \frac{\mathcal{C}}{|p_m(w, 0)|} \sum_{j=0}^{i-\eta} {i-\eta \choose j} \frac{1}{(a_m)^{j+1}} \left(\frac{m}{\sqrt{a_m}} \right)^{2(i-\eta-j)}$$
$$= \frac{\mathcal{C}}{a_m |p_m(w, 0)|} \left(\frac{1}{a_m} + \frac{m^2}{a_m} \right)^{i-\eta}$$
$$\leq \mathcal{C} \left(\frac{m}{\sqrt{a_m}} \right)^{2i-2\eta} \frac{1}{a_m |p_m(w, 0)|}.$$

Then we have

(4.25)
$$|D_i| \le \frac{\mathcal{C}}{a_m |p_m(w,0)|} \sum_{\eta=0}^i \binom{i}{\eta} |f^{(\eta)}(0)| \left(\frac{m}{\sqrt{a_m}}\right)^{2i-2\eta}.$$

On the other hand, by (4.1), (3.2) and (4.6) , we get

$$b_{i} \leq \mathcal{C} \max_{x \in I_{m}} \left| \sqrt{w(x)} p_{m}(w, x) (a_{m} - x) x^{i + \gamma - \frac{\alpha}{2}} \right|$$

$$\leq \mathcal{C} \max_{x \in I_{m}} \left| (a_{m} - x)^{\frac{3}{4}} x^{i + \gamma - \frac{\alpha}{2} - \frac{1}{4}} \right|$$

$$(4.26) \leq \mathcal{C} (a_{m})^{\frac{3}{4}} \left(\frac{\sqrt{a_{m}}}{m} \right)^{2i + 2\gamma - \alpha - \frac{1}{2}} \sim a_{m} \left(\frac{\sqrt{a_{m}}}{m} \right)^{2i + 2\gamma} |p_{m}(w, 0)|.$$

Combining (4.25) and (4.26) in (4.24), we obtain

$$\begin{aligned} |A(x)u(x)| &\leq \mathcal{C}\left(\frac{\sqrt{a_m}}{m}\right)^{2\gamma} \sum_{i=0}^{r-1} \frac{1}{i!} \sum_{\eta=0}^{i} \binom{i}{\eta} |f^{(\eta)}(0)| \left(\frac{\sqrt{a_m}}{m}\right)^{2\eta} \\ &= \mathcal{C}\left(\frac{\sqrt{a_m}}{m}\right)^{2\gamma} \sum_{i=0}^{r-1} \sum_{\eta=0}^{r-1} (i-\eta)_+^0 |f^{(\eta)}(0)| \left(\frac{\sqrt{a_m}}{m}\right)^{2\eta} \binom{i}{\eta} \frac{1}{i!} \\ &= \mathcal{C}\left(\frac{\sqrt{a_m}}{m}\right)^{2\gamma} \sum_{\eta=0}^{r-1} \left(\frac{\sqrt{a_m}}{m}\right)^{2\eta} |f^{(\eta)}(0)| \sum_{i=0}^{r-1} (i-\eta)_+^0 \binom{i}{\eta} \frac{1}{i!} \\ &\sim \left(\frac{\sqrt{a_m}}{m}\right)^{2\gamma} \sum_{\nu=0}^{r-1} \frac{|f^{(\nu)}(0)|}{\nu!} \left(\frac{\sqrt{a_m}}{m}\right)^{2\nu}, \end{aligned}$$

since $\sum_{i=\eta}^{r-1} {i \choose \eta} \frac{1}{i!} \sim 1.$

Let us now show (4.22). Recalling (4.25), we have that

$$\begin{aligned} \|(a_m - \cdot)p_m(w)uv^{i}\|_p &\leq \mathcal{C}\|(a_m - \cdot)p_m(w)uv^{i}\|_{L^p(I_m)} \\ &= \mathcal{C}\|\sqrt{w}p_m(w)(a_m - \cdot)v^{i+\gamma-\frac{\alpha}{2}}\|_{L^p(I_m)} \\ &\leq \mathcal{C}\|(a_m - \cdot)^{\frac{3}{4}}v^{i+\gamma-\frac{\alpha}{2}-\frac{1}{4}}\|_{L^p(I_m)} \\ &\leq \mathcal{C}(a_m)^{\frac{3}{4}}\left(\frac{\sqrt{a_m}}{m}\right)^{2i+2\gamma-\alpha-\frac{1}{2}+\frac{2}{p}} \\ &= \mathcal{C}\left(\frac{\sqrt{a_m}}{m}\right)^{2i+2\gamma+\frac{2}{p}}a_m|p_m(w,0)| \end{aligned}$$

where we have applied (4.1), (4.6), (4.8) and considered the assumption (4.23). Then, we deduce (4.22).

The necessity of the conditions (4.21) and (4.23) can be proved using standard arguments (see [22]). We omit the details.

Proof of Theorem 2. Let us first prove that conditions (3.2) imply inequality (3.1). Recalling (2.2) and Proposition 9, it remains to prove that

$$\left|\sum_{k=1}^{m+1} u(x) \frac{x^r}{x_k^r} \frac{\ell_k(w, x)}{u(x_k)} f(x_k) u(x_k)\right| \le \mathcal{C}(\log m) \max_{1 \le i \le m+1} |fu|(x_i).$$

Let $x \in I_m = [c_{\overline{m^2}}^{a_m}, a_m]$ and x_d be a zero closest to x. By using (4.7), (4.11) and (4.13), we get

$$\begin{split} \sum_{k=1}^{m+1} u(x) \frac{x^r}{x_k^r} \frac{|\ell_k(w,x)|}{u(x_k)} f(x_k) u(x_k) \\ &\leq \mathcal{C} \left[|p_m(w,x)| \sqrt{w(x)} \sqrt{x(a_m-x)} \sum_{k \neq d, m+1} \left(\frac{x}{x_k} \right)^{r+\gamma - \frac{\alpha}{2} - \frac{1}{4}} \left(\frac{a_m - x}{a_m - x_k} \right)^{\frac{3}{4}} \frac{\Delta x_k}{|x - x_k|} \\ &+ 1 + \left| \frac{p_m(w,x)}{p_m(w,a_m)} \right| \frac{x^r}{a_m^r} \frac{u(x)}{u(a_m)} \right] \max_{1 \leq i \leq m+1} |fu|(x_i) \,. \end{split}$$

By (4.6) the last summand on the right-hand side is bounded by $\left(\frac{x}{a_m}\right)^{r+\gamma-\frac{\alpha}{2}-\frac{1}{4}}$ and so the whole right-hand side is dominated by

$$\mathcal{C}\left(\sum_{k\neq d,m+1} \left(\frac{x}{x_k}\right)^{r+\gamma-\frac{\alpha}{2}-\frac{1}{4}} \left(\frac{a_m-x}{a_m-x_k}\right)^{\frac{3}{4}} \frac{\Delta x_k}{|x-x_k|} + 1 + \left(\frac{x}{a_m}\right)^{r+\gamma-\frac{\alpha}{2}-\frac{1}{4}}\right) \max_{1\leq i\leq m+1} |fu|(x_i).$$

Therefore, if the parameters r, γ and α fulfill the assumptions (3.2), the sum on the right-hand side is bounded by $C(\log m)$ and the last term is bounded by 1.

Let us now prove that inequality (3.1) implies conditions (3.2). To this aim, for any $f \in C_u$, consider the piecewise linear function f_1 defined as follows

$$\begin{cases} f_1^{(i)}(0) = 0, & i = 0, 1, \dots, r - 1, \\ f_1(x_k) = -|f(x_k)| \operatorname{sgn}\{p'_m(w, x_k)\} \neq 0, & \text{for } 1 \le x_k \le 2, \\ f_1(x_k) = 0, & \text{otherwise.} \end{cases}$$

By (3.1) we have (4.28)

$$\mathcal{C}(\log m) \max_{1 \le x_i \le 2} |f_1 u|(x_i) \ge \|\mathcal{L}_{m,r}(w, f_1) u\|_{\infty} \ge \|\mathcal{L}_{m,r}(w, f_1) u\|_{L^{\infty}[0,1]},$$

where, for $x \in [0, 1]$ and $x_k \in [1, 2]$,

$$\mathcal{L}_{m,r}(w, f_1, x)u(x) = -\sum_{1 \le x_k \le 2} \frac{x^r}{x_k^r} \frac{p_m(w, x)|f_1u|(x_k)}{|p'_m(w, x_k)|(x - x_k)} \left(\frac{a_m - x}{a_m - x_k}\right) \frac{u(x)}{u(x_k)}$$
$$= \sum_{1 \le x_k \le 2} \frac{x^r}{x_k^r} \frac{p_m(w, x)|f_1u|(x_k)}{|p'_m(w, x_k)||x - x_k|} \left(\frac{a_m - x}{a_m - x_k}\right) \frac{u(x)}{u(x_k)},$$

since $x - x_k < 0$. Now, if $x \in [0, 1]$ and $x_k \in [1, 2]$, then $|x - x_k| \le 2$ and $\frac{a_m-x}{a_m-x_k} \sim 1$. Hence, setting

$$\widetilde{A}(x) = p_m(w, x)\sqrt{w(x)\sqrt{x(a_m - x)}}$$

and using (4.11), we get

(4.29)

$$\|\mathcal{L}_{m,r}(w,f_1)u\|_{L^{\infty}[0,1]} \ge \mathcal{C}\|\widetilde{A}v^{r+\gamma-\frac{\alpha}{2}-\frac{1}{4}}\|_{L^{\infty}[0,1]} \sum_{1\le x_k\le 2} \frac{\Delta x_k |fu|(x_k)}{x_k^{r+\gamma-\frac{\alpha}{2}-\frac{1}{4}}}.$$

Combining (4.28) and (4.29), we obtain

$$\|\widetilde{A}v^{r+\gamma-\frac{\alpha}{2}-\frac{1}{4}}\|_{L^{\infty}[0,1]} \sum_{1 \le x_k \le 2} \frac{\Delta x_k |fu|(x_k)}{x_k^{r+\gamma-\frac{\alpha}{2}-\frac{1}{4}}} \le \mathcal{C}(\log m) \max_{1 \le x_i \le 2} |fu|(x_i)$$

with $\mathcal{C} \neq \mathcal{C}(m, f)$, whence letting $b_i = |fu|(x_i)$ and $||\underline{b}||_{\infty}^* = \max_{1 \le x_i \le 2} b_i$, we have

$$\|\widetilde{A}v^{r+\gamma-\frac{\alpha}{2}-\frac{1}{4}}\|_{L^{\infty}[0,1]}\sum_{1\leq x_{k}\leq 2}\frac{\Delta x_{k}}{x_{k}^{r+\gamma-\frac{\alpha}{2}-\frac{1}{4}}}\frac{b_{k}}{\|\underline{b}\|_{\infty}^{*}}\leq \mathcal{C}(\log m), \quad \mathcal{C}\neq \mathcal{C}(m,f)$$

and then

$$\|\widetilde{A}v^{r+\gamma-\frac{\alpha}{2}-\frac{1}{4}}\|_{L^{\infty}[0,1]} \sup_{\|\underline{d}\|_{\infty}^{*}=1} \sum_{1 \le x_{k} \le 2} \frac{\Delta x_{k}}{x_{k}^{r+\gamma-\frac{\alpha}{2}-\frac{1}{4}}} d_{k} \le \mathcal{C}(\log m), \quad \mathcal{C} \neq \mathcal{C}(m,f) \le C$$

Hence we get

$$\|\widetilde{A}v^{r+\gamma-\frac{\alpha}{2}-\frac{1}{4}}\|_{L^{\infty}[0,1]}\sum_{1\leq x_k\leq 2}\frac{\Delta x_k}{x_k^{r+\gamma-\frac{\alpha}{2}-\frac{1}{4}}}\leq \mathcal{C}(\log m).$$

Obviously the sum on the left-hand side is ~ 1 . Moreover, by (4.6), we have

$$\|\widetilde{A}v^{r+\gamma-\frac{\alpha}{2}-\frac{1}{4}}\|_{L^{\infty}[0,1]} \ge \left|\widetilde{A}\left(\frac{a_m}{m^2}\right)v^{r+\gamma-\frac{\alpha}{2}-\frac{1}{4}}\left(\frac{a_m}{m^2}\right)\right| \sim \left(\frac{a_m}{m^2}\right)^{r+\gamma-\frac{\alpha}{2}-\frac{1}{4}}.$$

So, from

$$\left(\frac{a_m}{m^2}\right)^{r+\gamma-\frac{\alpha}{2}-\frac{1}{4}} \le \mathcal{C}(\log m)$$

it follows that $r + \gamma \ge \frac{\alpha}{2} + \frac{1}{4}$. In order to prove the necessity of the other inequality in (3.2), we introduce the piecewise linear function f_2 defined as

(4.30)
$$\begin{cases} f_2^{(i)}(0) = 0, & i = 0, 1, \dots, r-1, \\ f_2(x_k) = |f(x_k)| \operatorname{sgn}\{p'_m(w, x_k)\}, & \text{for } 0 < x_k \le 1, \\ f_2(x_k) = 0, & \text{otherwise.} \end{cases}$$

Using similar arguments we obtain

$$\|\widetilde{A}v^{r+\gamma-\frac{\alpha}{2}-\frac{1}{4}}\|_{L^{\infty}[1,2]} \sum_{0 < x_{k} \le 1} \frac{\Delta x_{k}}{x_{k}^{r+\gamma-\frac{\alpha}{2}-\frac{1}{4}}} \le \mathcal{C}(\log m) \,.$$

Now, the sum on the left-hand side is $\sim \int_{x_1}^1 x^{-r-\gamma+\frac{\alpha}{2}+\frac{1}{4}} dx$, while the norm is bounded. So, from

$$\int_{x_1}^1 x^{-r-\gamma+\frac{\alpha}{2}+\frac{1}{4}} \, \mathrm{d}x \le \mathcal{C} \sum_{0 < x_k \le 1} \frac{\Delta x_k}{x_k^{r+\gamma-\frac{\alpha}{2}-\frac{1}{4}}} \le \mathcal{C}(\log m)$$

we deduce $r + \gamma \leq \frac{\alpha}{2} + \frac{5}{4}$ and the first part of the theorem follows. Let us now prove (3.3). Since $|f^{(i)}|$ is quasi increasing on $[0, \alpha_m]$, with $i \in \{0, 1, \ldots, r-1\}$ and $\alpha_m = a_m/m^2$, we have

$$\alpha_m^{i+\gamma} \left| f^{(i)}(0) \right| \le \mathcal{C} \alpha_m^{i+\gamma} \left| f^{(i)} \left(\frac{\alpha_m}{2} \right) \right| \le \mathcal{C} \alpha_m^{i+\gamma} \left\| f^{(i)} \right\|_{L^{\infty}[\frac{\alpha_m}{2}, \alpha_m]}$$

Recalling a formula in [3, p. 15], it follows that

$$\begin{aligned} \alpha_m^{i+\gamma} \left| f^{(i)}(0) \right| &\leq \mathcal{C}\alpha_m^{\gamma} \left\| f \right\|_{L^{\infty}\left[\frac{\alpha_m}{2},\alpha_m\right]} + \mathcal{C}\alpha_m^{r-1+\gamma} \left\| f^{(r-1)} \right\|_{L^{\infty}\left[\frac{\alpha_m}{2},\alpha_m\right]} \\ &\leq \mathcal{C} \left\| f u \right\|_{L^{\infty}\left[\frac{\alpha_m}{2},\alpha_m\right]} + \mathcal{C} \left(\frac{\sqrt{a_m}}{m}\right)^{r-1} \left\| f^{(r-1)} \varphi^{r-1} u \right\|_{L^{\infty}\left[\frac{\alpha_m}{2},\alpha_m\right]} \end{aligned}$$

Hence (3.1) becomes

$$\|\mathcal{L}_{m,r}(w,f)u\|_{\infty} \leq \mathcal{C}(\log m)\|fu\|_{\infty} + \mathcal{O}\left(\left(\frac{\sqrt{a_m}}{m}\right)^{r-1}\right).$$

Letting $P_{m+r} \in \mathbb{P}_{m+r}$ be the polynomial of best approximation for $f \in C_u$, it follows that

$$\begin{aligned} \left\| \left[f - \mathcal{L}_{m,r}\left(w,f\right) \right] u \right\|_{\infty} &= \left\| \left[f - P_{m+r} - \mathcal{L}_{m,r}\left(w,f - P_{m+r}\right) \right] u \right\|_{\infty} \\ &\leq \mathcal{C}(\log m) E_{m+r}(f)_{u,\infty} + \mathcal{O}\left(\left(\frac{\sqrt{a_m}}{m} \right)^{r-1} \right), \end{aligned}$$

i.e. (3.3).

In order to prove Theorem 4 we recall some known results concerning the Hilbert transform H(f,t). The Hilbert transform related to the interval (0, a) is defined as follows

$$H(f,t) = \int_0^a \frac{f(x)}{x-t} \, \mathrm{d}x, \qquad t \in (0,a) \,, \quad a > 0 \,,$$

where the integral is understood in the Cauchy principal value sense. Letting $v^{\rho}(x) = x^{\rho}$ and 1 , the bound

(4.31)
$$\| (Hf)v^{\rho} \|_{L^{p}(0,1)} \leq \mathcal{C} \| fv^{\rho} \|_{L^{p}(0,1)}, \qquad \mathcal{C} \neq \mathcal{C}(f),$$

holds for any f such that $fv^{\rho} \in L^{p}(0, a)$ if and only if $-\frac{1}{p} < \rho < 1 - \frac{1}{p}$ (see [23]).

The following lemma, that can be found in [21] (see also [8]), will be also useful in the proof of Theorem 4.

Lemma 10. Let $0 < \theta < 1$, j be given by (2.3), $u \in L^p$, $1 \leq p < \infty$ and $\Delta x_k = x_{k+1} - x_k$. Then, for an arbitrary polynomial $P \in \mathbb{P}_{lm}$ (where l is a fixed integer), there exists $\overline{\theta} \in (\theta, 1)$ such that

(4.32)
$$\sum_{k=1}^{j} \Delta x_k |Pu|^p(x_k) \le \mathcal{C} \int_{x_1}^{\bar{\theta}a_m} |Pu|^p(x) \,\mathrm{d}x,$$

with $\mathcal{C} \neq \mathcal{C}(m, f)$.

We want to emphasize that the first Marcinkiewicz inequality (4.32) does not hold if the sum on the left-hand side is extended to all the zeros of $p_m(w)$, i.e. $k = 1, \ldots, m$ (see [13, 14]).

Proof of Theorem 4. Taking into account Proposition 9 it remains to estimate the L_u^p -norm of the first term in (2.4). Letting $v^r(x) = x^r$ and

$$L_m^*(w, f, x) = \sum_{k=1}^j \ell_k(w, x) f(x_k), \quad \ell_k(w, x) = \frac{p_m(w, x)}{(x - x_k)p'_m(w, x_k)} \frac{(a_m - x)}{(a_m - x_k)},$$

this term can be rewritten as $v^r L_m^*(w, \frac{f}{v^r})$. So, using (4.1), we have

where $I_m = [c a_m m^{-2}, a_m]$ and

$$\Gamma(g) = \int_{I_m} x^r \sum_{k=1}^j \frac{\ell_k(w, x)}{x_k^r} f(x_k) g(x) u(x) \, \mathrm{d}x$$

= $\sum_{k=1}^j \frac{f(x_k) u(x_k)}{p'_m(w, x_k) (a_m - x_k) u(x_k) x_k^r} \int_{I_m} \frac{(a_m - x) p_m(w, x) x^r g(x) u(x)}{x - x_k} \, \mathrm{d}x.$

By using (4.11), we have

$$\begin{aligned} |\Gamma(g)| &\leq \mathcal{C} \sum_{k=1}^{j} \frac{\Delta x_{k} |fu|(x_{k})}{(a_{m}-x_{k})^{\frac{3}{4}} x_{k}^{r+\gamma-\frac{\alpha}{2}-\frac{1}{4}}} \left| \int_{I_{m}} \frac{(a_{m}-x) p_{m}(w,x) x^{r} g(x) u(x)}{x-x_{k}} \, \mathrm{d}x \right| \\ &\leq \frac{\mathcal{C}}{a_{m}^{3/4}} \sum_{k=1}^{j} \frac{\Delta x_{k} |fu|(x_{k})}{x_{k}^{r+\gamma-\frac{\alpha}{2}-\frac{1}{4}}} \left| \int_{I_{m}} \frac{(a_{m}-x) p_{m}(w,x) x^{r} g(x) u(x)}{x-x_{k}} \, \mathrm{d}x \right|. \end{aligned}$$

Denoting by $G(x_k)$ the integral on the right-hand side and using the Hölder inequality, we get (4.34)

$$|\Gamma(g)| \le \frac{\mathcal{C}}{a_m^{3/4}} \left(\sum_{k=1}^j \Delta x_k |fu|^p(x_k) \right)^{1/p} \left(\sum_{k=1}^j \Delta x_k \left| \frac{G(x_k)}{x_k^{r+\gamma-\alpha/2-1/4}} \right|^q \right)^{1/q}.$$

For any polynomial $0 < Q \in \mathbb{P}_{lm}$, with l a fixed integer, we can write

$$G(x_k) = \int_{I_m} \frac{(a_m - x)p_m(w, x)Q(x) - (a_m - x_k)p_m(w, x_k)Q(x_k)}{x - x_k} \frac{x^r g(x)u(x)}{Q(x)} dx$$

and G(t) is a polynomial of degree m + lm. Thus, using Lemma 10, we have

$$B = \left(\sum_{k=1}^{j} \Delta x_{k} \left| \frac{G(x_{k})}{x_{k}^{r+\gamma-\alpha/2-1/4}} \right|^{q} \right)^{1/q} \le \left(\int_{I_{\bar{\theta}m}} \left| \frac{G(t)}{t^{r+\gamma-\alpha/2-1/4}} \right|^{q} \mathrm{d}t \right)^{1/q},$$

where $0 < \theta < \bar{\theta} < 1$ and $I_{\bar{\theta}m} = [c a_m m^{-2}, \bar{\theta} a_m]$. Then we obtain

$$B \leq \left(\int_{I_{\bar{\theta}m}} \left| t^{-r-\gamma+\alpha/2+1/4} \int_{I_m} \frac{(a_m - x)p_m(w, x)x^r(gu)(x)}{x - t} \, \mathrm{d}x \right|^q \, \mathrm{d}t \right)^{1/q} \\ + \left(\int_{I_{\bar{\theta}m}} \left| t^{-r-\gamma+\alpha/2+1/4}(a_m - t)p_m(w, t)Q(t) \int_{I_m} \frac{x^r(gu)(x)}{(x - t)Q(x)} \, \mathrm{d}x \right|^q \, \mathrm{d}t \right)^{1/q} \\ =: B_{\mathrm{f}} + B_{\mathrm{f}2}.$$

Let us estimate the term B_1 . By (4.7) we have

$$\left|\widetilde{A}(x)\right| := |p_m(w,x)|\sqrt{w(x)\sqrt{x(a_m-x)}} \le \mathcal{C}, \qquad x \in I_m.$$

Moreover, under the assumptions (3.5), $t^{-r-\gamma+\frac{\alpha}{2}+\frac{1}{4}} \in L^q$ and $t^{r+\gamma-\frac{\alpha}{2}-\frac{1}{4}} \in L^p$, we can apply (4.31), obtaining

$$B_{1} \leq \left(\int_{I_{m}} \left| t^{-r-\gamma+\frac{\alpha}{2}+\frac{1}{4}} \int_{I_{m}} \frac{(a_{m}-x)^{\frac{3}{4}} \widetilde{A}(x) x^{r+\gamma-\frac{\alpha}{2}-\frac{1}{4}} g(x)}{x-t} \, \mathrm{d}x \right|^{q} \, \mathrm{d}t \right)^{1/q} \\ \leq \mathcal{C} \left(\int_{I_{m}} \left| (a_{m}-x)^{\frac{3}{4}} \widetilde{A}(t) g(t) \right|^{q} \, \mathrm{d}t \right)^{1/q} \\ (4.36) \quad \mathcal{C}a_{m}^{\frac{3}{4}} \|g\|_{L^{q}(I_{m})} \, .$$

In order to estimate B_2 we can construct a polynomial $Q \in \mathbb{P}_{lm}$ such that $Q(x) \sim x^r u(x)$ for $x \in I_m$. So, using (4.7), we have

$$\left|t^{-r-\gamma+\alpha/2+1/4}(a_m-t)p_m(w,t)Q(t)\right| \le Ca_m^{3/4}.$$

Then, by (4.31), we get (4.37)

$$B_2 \le C a_m^{3/4} \left(\int_{I_m} \left| \int_{I_m} \frac{x^r(gu)(x)}{(x-t)Q(x)} \, \mathrm{d}x \right|^q \, \mathrm{d}t \right)^{1/q} \le C a_m^{3/4} \|g\|_{L^q(I_m)}.$$

So, taking into account (4.36), (4.37) and (4.35), it follows that

$$\left(\int_{I_{\bar{\theta}m}} \left|\frac{G(t)}{t^{r+\gamma-\alpha/2-1/4}}\right|^q \, \mathrm{d}t\right)^{1/q} \le \mathcal{C}a_m^{3/4} \|g\|_{L^q(I_m)}$$

and combining with (4.34) and (4.33) the first term in (2.4) is bounded by

$$\mathcal{C}\left(\sum_{k=1}^{j} \Delta x_k |fu|^p(x_k)\right)^{1/p}$$

Therefore, recalling (4.22), we have that (3.5) implies (3.4).

In order to prove that (3.4) implies (3.5) we use arguments similar to those in the proof of Theorem 4. So, we are going to show only the main steps. Considering the function f_1 as in (4.27) we can write

$$\|\mathcal{L}_{m,r}^{*}(w,f_{1})u\|_{L^{p}[c\,a_{m}m^{-2},1]} \leq \|\mathcal{L}_{m,r}^{*}(w,f_{1})u\|_{p} \leq \mathcal{C}\left(\sum_{1\leq x_{k}\leq 2}\Delta x_{k}|f_{1}u|^{p}(x_{k})\right)^{1/p}$$

where $[c a_m m^{-2}, 1] = [0, 1] \cap I_m$. Recalling the expression of $\mathcal{L}^*_{m,r}(w, f_1, x)$ for $x \in [c a_m m^{-2}, 1]$, with

$$\left|\widetilde{A}(x)\right| = \left|p_m(w,x)\right| \sqrt{w(x)\sqrt{x(a_m-x)}},$$

we obtain (4.38)

$$\|\widetilde{A}v^{r+\gamma-\frac{\alpha}{2}-\frac{1}{4}}\|_{L^{p}[a_{m}m^{-2},1]} \sum_{1 \le x_{k} \le 2} \frac{\Delta x_{k}|f_{1}u|(x_{k})}{x_{k}^{r+\gamma-\frac{\alpha}{2}-\frac{1}{4}}} \le \mathcal{C}\left(\sum_{1 \le x_{k} \le 2} \Delta x_{k}|f_{1}u|^{p}(x_{k})\right)^{1/p}.$$

Now, to simplify the notation, we set

$$a_k = \frac{(\Delta x_k)^{1/q}}{x_k^{r+\gamma-\frac{\alpha}{2}-\frac{1}{4}}}, \ b_k = (\Delta x_k)^{1/p} |f_1 u|(x_k), \ \|\underline{b}\,\|_p^* = \left(\sum_{1 \le x_k \le 2} b_k^p\right)^{1/p}.$$

Hence (4.38) becomes

$$\|\widetilde{A}v^{r+\gamma-\frac{\alpha}{2}-\frac{1}{4}}\|_{L^{p}[c\,a_{m}m^{-2},1]}\sum_{1\leq x_{k}\leq 2}a_{k}\frac{b_{k}}{\|\underline{b}\|_{p}^{*}}\leq \mathcal{C}, \quad \mathcal{C}\neq \mathcal{C}(m,f).$$

Taking the supremum on \underline{b} , we get

$$\|\widetilde{A}v^{r+\gamma-\frac{\alpha}{2}-\frac{1}{4}}\|_{L^{p}[c\,a_{m}m^{-2},1]}\left(\sum_{1\leq x_{k}\leq 2}|a_{k}|^{q}\right)^{1/q}\leq \mathcal{C},$$

which implies $r + \gamma - \frac{\alpha}{2} - \frac{1}{4} > -\frac{1}{p}$, taking into account that, by (4.7), $\left| \widetilde{A}(x) \right| \leq \mathcal{C}, x \in [c \, a_m m^{-2}, 1].$

Analogously, considering the function f_2 defined in (4.30), we get

$$\|\widetilde{A}v^{r+\gamma-\frac{\alpha}{2}-\frac{1}{4}}\|_{L^{p}[1,2]}\left(\sum_{0 < x_{k} \leq 1} |a_{k}|^{q}\right)^{1/q} \leq \mathcal{C},$$

i.e.

$$\|\widetilde{A}v^{r+\gamma-\frac{\alpha}{2}-\frac{1}{4}}\|_{L^p[1,2]} \left(\sum_{0< x_k\leq 1} \frac{\Delta x_k}{x_k^{(r+\gamma-\alpha/2-1/4)q}}\right)^{1/q} \leq \mathcal{C},$$

whence $-r - \gamma + \frac{\alpha}{2} + \frac{1}{4} > -\frac{1}{q}$ that is $r + \gamma - \frac{\alpha}{2} - \frac{1}{4} < 1 - \frac{1}{p}$, which completes the proof.

Proof of Theorem 5. Since $|f^{(i)}|$ is quasi increasing on $[0, \alpha_m]$, with $i \in \{0, 1, \ldots, r-1\}$ and $\alpha_m = a_m/m^2$, using the Hölder inequality, we have

$$\begin{aligned} \alpha_m^{i+\gamma+1/p} \left| f^{(i)}(0) \right| &\leq \mathcal{C} \alpha_m^{i+\gamma+1/p} \left| f^{(i)} \left(\frac{\alpha_m}{2} \right) \right| &\leq \mathcal{C} \alpha_m^{i+\gamma+1/p-1} \int_{\frac{\alpha_m}{2}}^{\alpha_m} \left| f^{(i)}(x) \right| \mathrm{d}x \\ &\leq \mathcal{C} \alpha_m^{i+\gamma} \left\| f^{(i)} \right\|_{L^p[\frac{\alpha_m}{2},\alpha_m]} . \end{aligned}$$

Recalling a formula in [3, p. 15], it follows that

$$\begin{aligned} \alpha_{m}^{i+\gamma+1/p} \left| f^{(i)}(0) \right| &\leq \mathcal{C}\alpha_{m}^{\gamma} \left\| f \right\|_{L^{p}\left[\frac{\alpha_{m}}{2},\alpha_{m}\right]} + \mathcal{C}\alpha_{m}^{r-1+\gamma} \left\| f^{(r-1)} \right\|_{L^{p}\left[\frac{\alpha_{m}}{2},\alpha_{m}\right]} \\ &\leq \mathcal{C} \left\| f u \right\|_{L^{p}\left[\frac{\alpha_{m}}{2},\alpha_{m}\right]} + \mathcal{C}\left(\frac{\sqrt{a_{m}}}{m}\right)^{r-1} \left\| f^{(r-1)}\varphi^{r-1}u \right\|_{L^{p}\left[\frac{\alpha_{m}}{2},\alpha_{m}\right]} \end{aligned}$$

So inequality (3.7) becomes

$$\begin{aligned} \left\| \mathcal{L}_{m,r}^{*}\left(w,f\right) u \right\|_{p} &\leq \mathcal{C} \left[\|fu\|_{p} + \left(\frac{\sqrt{a_{m}}}{m}\right)^{1/p} \int_{0}^{\frac{\sqrt{a_{m}}}{m}} \frac{\Omega_{\varphi}^{s}(f,t)_{u,p}}{t^{1+1/p}} \,\mathrm{d}t \\ (4.39) &+ \left(\frac{\sqrt{a_{m}}}{m}\right)^{r-1} \left\| f^{(r-1)} \varphi^{r-1} u \right\|_{L^{p}[0,\alpha_{m}]} \right]. \end{aligned}$$

Let $P_M \in \mathbb{P}_M$, with $M = \lfloor \frac{\theta m}{\theta + 1} \rfloor$, be a polynomial of quasi best approximation for $f \in L^p_u$, and set $Q = \mathcal{L}^*_{m,r}(w, P_M)$. Hence, by (4.39), we get

$$\begin{split} \|[f - \mathcal{L}_{m,r}^{*}(w, f)]u\|_{p} &\leq \mathcal{C} \left[\|(f - Q)u\|_{p} + \left(\frac{\sqrt{a_{m}}}{m}\right)^{1/p} \int_{0}^{\frac{\sqrt{a_{m}}}{m}} \frac{\Omega_{\varphi}^{s}(f - Q, t)_{u,p}}{t^{1+1/p}} \, \mathrm{d}t \right. \\ &+ \left(\frac{\sqrt{a_{m}}}{m}\right)^{r-1} \left\| (f - Q)^{(r-1)} \varphi^{r-1}u \right\|_{L^{p}[0,\alpha_{m}]} \right] \end{split}$$

Now, to estimate the first summand at the right-hand side we can use (2.12) and (2.8). For the second summand we note that

$$\Omega_{\varphi}^{s}(f-Q,t)_{u,p} \leq \Omega_{\varphi}^{s}(f,t)_{u,p} + t^{s} \left\| Q^{(s)}\varphi^{s}u \right\|_{p}$$

So, by (2.13), taking into account that

$$\int_0^{\frac{\sqrt{a_m}}{m}} \frac{\Omega_{\varphi}^s(f,t)_{u,p}}{t} \, \mathrm{d}t \le \left(\frac{\sqrt{a_m}}{m}\right)^{1/p} \int_0^{\frac{\sqrt{a_m}}{m}} \frac{\Omega_{\varphi}^s(f,t)_{u,p}}{t^{1+1/p}} \, \mathrm{d}t \,,$$

ity (3.8) follows.

inequality (3.8) follows.

Proof of Corollary 6. Proceeding as in the proof of Theorem 5, since $|f^{(i)}|$ is quasi increasing on $[0, \alpha_m]$, with $i \in \{0, 1, \dots, r-1\}$ and $\alpha_m =$ a_m/m^2 , and $f \in W^p_r(u)$, we get

$$\begin{aligned} \alpha_m^{i+\gamma+1/p} \left| f^{(i)}(0) \right| &\leq \mathcal{C} \alpha_m^{\gamma} \left\| f \right\|_{L^p\left[\frac{\alpha_m}{2},\alpha_m\right]} + \mathcal{C} \alpha_m^{r+\gamma} \left\| f^{(r)} \right\|_{L^p\left[\frac{\alpha_m}{2},\alpha_m\right]} \\ (4.40) &\leq \mathcal{C} \left\| f u \right\|_{L^p\left[\frac{\alpha_m}{2},\alpha_m\right]} + \mathcal{C} \left(\frac{\sqrt{a_m}}{m} \right)^r \left\| f^{(r)} \varphi^r u \right\|_{L^p\left[\frac{\alpha_m}{2},\alpha_m\right]} . \end{aligned}$$

Hence, by (3.7), (2.6) and (4.40), we obtain (3.9).

Now, in order to prove (3.10), let $Q = \mathcal{L}_{m,r}^*(w, P_M)$, where $P_M \in \mathbb{P}_M$, with $M = \lfloor \frac{\theta m}{\theta + 1} \rfloor$, is a polynomial of quasi best approximation for $f \in L^p_u$. By (3.9) we have

$$\begin{split} \left\| \left[f - \mathcal{L}_{m,r}^*\left(w,f\right) \right] u \right\|_p &\leq \mathcal{C} \left\| \left(f - Q \right) u \right\|_p + \left\| \mathcal{L}_{m,r}^*\left(w,f-Q\right) u \right\|_p \\ &\leq \mathcal{C} \left\| \left(f - Q \right) u \right\|_p + \left(\frac{\sqrt{a_m}}{m} \right)^r \left\| f^{(r)} \varphi^r u \right\|_p + \left(\frac{\sqrt{a_m}}{m} \right)^r \left\| Q^{(r)} \varphi^r u \right\|_p \,. \end{split}$$

Using Lemma 1, (2.10) and (2.6), we obtain (3.10).

Acknowledgements. The authors would like to thank the referee for carefully reading the manuscript and for his/her suggestions which contributed in improving the presentation of the paper.

References

- K. Balázs, Lagrange and Hermite interpolation processes on the positive real line, J. Approx. Theory 50, no. 1, 18–24, (1987)
- [2] A. S. Cvetkovic and G. V. Milovanovic, The Mathematica package "OrthogonalPolynomials", Facta Univ. Ser. Math. Inform. No. 19, 17-36, (2004)
- [3] Z. Ditzian, V. Totik, *Moduli of smoothness*, Springer Series in Computational Mathematics 9, Springer Verlag, New York (1987)
- [4] P. Erdős and P. Turán, On interpolation. III. Interpolatory theory of polynomials, Ann. of Math. (2) 41, 510–553, (1940)
- [5] G. Freud, Convergence of the Lagrange interpolation on the infinite interval, Mat. Lapok 18, 289–292, (1967)
- [6] C. Frammartino, A Nyström method for solving a boundary value problem on [0,∞), in: Approximation and Computation (W. Gautchi, G. Mastroianni and T. Rassias, Eds.), 311– 325. Springer Optim. Appl., 42, Springer, New York (2011)
- [7] T. Kasuga and R. Sakai, Orthonormal polynomials with generalized Freud-type weights, J. Approx. Theory 121, 13–53, (2003)
- [8] C. Laurita and G. Mastroianni, L^p-Convergence of Lagrange interpolation on the semiaxis, Acta Math. Hungarica 120, no. 3, 249–273, (2008)
- [9] A. L. Levin and D. S. Lubinsky, Christoffel functions, orthogonal polynomials, and Nevai's conjecture for Freud weights, *Constr. Approx.* 8, no. 4, 463–535, (1992)
- [10] A. L. Levin and D. S. Lubinsky, Orthogonal polynomials for exponential weights, CSM Books in Mathematics/Ouvrages de Mathématiques de la SMC, 4. Springer-Verlag, New York (2001)
- [11] A.L. Levin and D.S. Lubinsky, Orthogonal polynomials for weights $x^{2\rho}e^{-2Q(x)}$ on [0, d), J. Approx. Theory **134**, no. 2, 199–256, (2005)
- [12] A.L. Levin and D.S. Lubinsky, Orthogonal polynomials for weights $x^{2\rho}e^{-2Q(x)}$ on [0, d), II, J. Approx. Theory **139**, no. 1, 107–143, (2006)
- [13] D. S. Lubinsky and G. Mastroianni, Mean convergence of extended Lagrange interpolation with Freud weights, Acta Math. Hungar. 84, no. 1–2, 47–63, (1999)
- [14] D. S. Lubinsky and G. Mastroianni, Converse quadrature sum inequalities for Freud weights. II, Acta Math. Hungar. 96, no. 1–2, 147–168, (2002)
- [15] G. Mastroianni and G. V. Milovanović, Interpolation processes. Basic theory and applications, Springer Monographs in Mathematics. Springer-Verlag, Berlin (2008)
- [16] G. Mastroianni and G. V. Milovanovic, Some numerical methods for second kind Fredholm integral equation on the real semiaxis, *IMA Journal of Numerical Analysis* 29, 1046–1066, (2009)
- [17] G. Mastroianni, G. V. Milovanović and I. Notarangelo, On an interpolation process of Lagrange–Hermite type, *Publication de l'Institut Mathématique*, Nouvelle série, tome 91 (105), 163–175, (2012)
- [18] G. Mastroianni and I. Notarangelo, A Lagrange-type projector on the real line, *Math. Comp.* 79, no. 269, 327–352, (2010)

- [19] G. Mastroianni and I. Notarangelo, Some Fourier-type operators for functions on unbounded intervals, Acta Math. Hungar. 127, no. 4, 347–375, (2010)
- [20] G. Mastroianni and D. Occorsio, Lagrange Interpolation at Laguerre zeros in some weighted uniform spaces, Acta Math. Hungar. 91, no. 1–2, 27–52, (2001)
- [21] G. Mastroianni and J. Szabados, Polynomial approximation on the real semiaxis with generalized Laguerre weights, *Studia Univ. "Babes-Bolyai"* Studia Europa, Vol. LII, Number 4, 89–103, (2007)
- [22] G. Mastroianni and P. Vértesi, Fourier sums and Lagrange interpolation on (0, +∞) and (-∞, +∞), in: Frontiers in Interpolation and Approximation, Dedicated to the memory of A. Sharma, (N.K. Govil, H.N. Mhaskar, R.N. Mohapatra, Z. Nashed and J. Szabados, eds.), pp. 307–344. Taylor & Francis Books, Boca Raton, Florida, (2006)
- [23] B. Muckenhoupt and L. Wheeden, Two weight function norm inequalities for the Hardy-Littlewood maximal function and the Hilbert transform, *Studia Math.* 55, no. 3, 279-294, (1976)
- [24] P. Nevai, On Lagrange interpolation based on Laguerre roots, [Hungarian] Mat. Lapok 22, 149–164, (1971)
- [25] P. Nevai, On the convergence of Lagrange interpolation based on Laguerre roots, [Russian] Publ. Math. 20, 235–239, (1973)
- [26] P. Nevai, Lagrange interpolation at zeros of orthogonal polynomials, in: Approximation Theory (G. G. Lorentz, Ed.), pp. 163–201. Academic Press, New York, (1976)
- [27] J. Shen, L.-L. Wang, Some recent advances on spectral methods for unbounded domains, Commun. Comput. Phys. 5, no. 2-4, 195-241 (2009)
- [28] J. Szabados, Weighted Lagrange and Hermite–Fejér interpolation on the real line, J. Inequal. Appl. 1, no. 2, 99–123, (1997)

DEPARTMENT OF MATHEMATICS, COMPUTER SCIENCES AND ECONOMICS, UNIVERSITY OF BASIL-ICATA, VIALE DELL'ATENEO LUCANO 10, 85100 POTENZA, ITALY