

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Willingness to Provide a Hair Sample for Drug Testing among Electronic Dance Music Party Attendees

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1730748> since 2020-02-25T11:45:42Z

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Willingness to Provide a Hair Sample for Drug Testing among Electronic Dance Music Party Attendees

Journal:	<i>Substance Abuse</i>
Manuscript ID	WSUB-2017-2994.R1
Manuscript Type:	Original Research
Keywords:	data collection methods, drug screening tests, epidemiology

SCHOLARONE™
Manuscripts

Review Only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

****CLEAN VERSION****

Running head: DRUG TESTING AND HAIR SAMPLE RESPONSE RATES

Willingness to Provide a Hair Sample for Drug Testing among
Electronic Dance Music Party Attendees

For Peer Review Only

Abstract

Background: Non-disclosure of drug use on surveys is common and many drug users unknowingly ingest adulterant or replacement drugs, which leads to underreporting of use of these drugs. Biological testing can complement survey research, and hair-testing is an appealing method as many drugs are detectable for months post-use. We examined willingness to donate a hair sample to be tested among those surveyed in a population at high risk for consuming adulterated drugs—electronic dance music (EDM) party attendees.

Methods: We surveyed 933 adults entering EDM parties in New York City in 2017. Hair donation response rates and reasons for refusal were examined from this cross-sectional study.

Results: A third (n=312; 33.4%) provided a hair sample. Lack of interest (21.0%), lack of time (19.8%), not wanting a lock of hair cut (17.7%), and disinterest in having hair cut in public (13.8%) were the main reported reasons for refusal. 4.7% refused because they could not receive results. Past-year drug users were more likely to fear identification than non-users (p<.001). Asian participants were at lower odds of providing a hair sample (aOR=0.53, 95% CI=0.32-0.87), and those reporting past-year use of LSD (aOR=1.62, 95% CI=1.11-2.35), opioids (nonmedical; aOR=1.93, 95% CI=1.25-2.99), and/or methamphetamine (aOR=3.43, 95% CI=1.36-8.62) were at higher odds of providing a sample than non-users of these drugs.

Conclusions: Only a third provided a hair sample and we found individual-level differences regarding willingness to provide a sample. Factors contributing to refusal should be considered to increase response rates and generalizability of results.

Key words: data collection methods; drug screening tests; epidemiology

Introduction

Accurate data on prevalence of drug use is important to inform prevention and harm reduction efforts. Surveys are the most common source of data in drug epidemiology studies, but not all survey respondents provide accurate information about drug use. Dishonest reporting about drug use has always been a concern among researchers.^{1,2} However, a newer concern is inaccurate reporting in which an individual may have thought he or she used a particular drug, but in fact used a different drug, or a drug containing adulterant drugs. It is important to investigate what drugs a user believes he or she used, but it is also important for epidemiologists to know what drugs were actually used. Biological testing is common in intervention research, but most surveys do not include biological measures—especially measures capable of detecting new psychoactive substances (NPS) and fentanyl analogs, which continue to emerge.⁵⁻⁸ In this paper, we examine willingness of individuals to provide a hair sample to be analyzed for NPS during a survey of a population at high risk for unintentionally using adulterated drugs—electronic dance music (EDM) party attendees.^{9,10}

Testing of drug products (e.g., pills, powders) has been the most common method used to test for drug adulteration. While seized drugs are typically tested for their contents,^{7,8} drug content information is frequently sought by users of certain drugs to inform harm reduction.^{8,11-15} Self-testing kits for drugs such as ecstasy have been popular for decades,¹⁶ and drug checking has increased in popularity with some harm reduction organizations conducting on-site drug-testing outside of EDM festivals.¹⁷⁻¹⁹ Other services throughout Europe allow users to drop off or send in drugs to be tested, and these organizations provide drug content results to users.^{8,11-14} Testing illegal drugs, however, can lead to legal risk for both users and testers, so biological

1
2
3 specimen testing is another option to determine drug adulteration. While this method can only
4
5 test for drugs already used, such testing can serve as an objective measure of drug use in
6
7 studies and test results can be used to inform prevalence estimates of use.²⁰ Providing users
8
9 with their results can also inform prevention and harm reduction regarding future use.
10
11

12
13 Individuals in the EDM scene are not only at high risk for use of drugs such as
14
15 ecstasy/Molly and various other drugs,²¹⁻²⁶ but they are also at high risk for using drugs
16
17 adulterated with NPS.^{9,10,17,19,27} Few studies, however, have added biological testing to surveys
18
19 conducted at or outside of nightlife parties. Multiple studies focusing on nightclub scenes have
20
21 tested attendees' saliva for various common drugs via oral assay,^{23,28-30} and Mohr et al²⁷ tested
22
23 saliva, urine, and blood of individuals entering EDM festivals for NPS and other drugs. However,
24
25 test results were not compared to self-reported use in these studies, so it is unknown whether
26
27 drugs were used intentionally or unintentionally. Palamar et al^{9,10} added hair testing to two
28
29 epidemiology surveys focusing on EDM attendees to determine the extent of unintentional use
30
31 of NPS among ecstasy users, but like many studies, only a convenience sample of survey
32
33 respondents were tested. While these studies incorporating biological assays are informative,
34
35 most results are based on convenience sampling, which limits generalizability of findings.
36
37 Systematic methods of obtaining biological specimens in addition to larger epidemiology
38
39 surveys are needed.
40
41
42
43
44
45

46
47 Hair testing is beneficial because unlike urine, saliva, or blood, in which drugs are only
48
49 detectable for days or hours post-use,³¹⁻³⁴ many drugs (e.g., NPS such as "bath salts") are
50
51 detectable in hair for months post-use.³⁵ Likewise, hair can be collected quickly, without pain,
52
53 in almost any environment (e.g., in the dark, cold, or rain). Thus, collection is typically not as
54
55
56
57
58
59
60

1
2
3 burdensome to participants as other biological testing methods. However, while hair-testing is
4
5 an appealing method to test for both known and unknown drug use, studies have not examined
6
7 acceptability of hair collection in at-risk scenes. Response rates and reasons for refusal have
8
9 been examined in population and household studies,³⁶⁻³⁹ but acquiring such data in at-risk
10
11 scenes is needed to inform future studies that incorporate this method as an objective
12
13 measure. We examined willingness of survey respondents in this high-risk scene to provide a
14
15 hair sample for analysis as part of an epidemiology survey. We also examined barriers to
16
17 participants providing samples by querying reasons for unwillingness to provide a sample, and
18
19 we examined characteristics of individuals most likely to provide a sample. These findings can
20
21 inform future studies that incorporate biological testing in a systematic manner.
22
23
24
25
26
27
28
29

30 **Methods**

31 *Procedure*

32
33
34
35 We utilized time-space sampling to recruit participants, which produces a probability
36
37 sample of visits to parties within the sampling frame.⁴⁰ Specifically, every week, a list of
38
39 upcoming EDM parties was created (based on EDM ticket websites). Parties at nightclubs and
40
41 festivals were randomly selected each week to survey potential participants. Individuals were
42
43 deemed eligible for this cross-sectional intercept survey study if they were 1) 18-40 years of
44
45 age, and 2) were about to enter the randomly selected party. Passersby—who were alone or in
46
47 groups—were approached by recruiters, and if confirmed eligible, were asked if they would like
48
49 to take a drug survey. Participants provided informed consent and self-administered the
50
51 surveys on tablets. A total of 933 participants completed the survey. The survey was conducted
52
53
54
55
56
57
58
59
60

1
2
3 from June through September of 2017 and the survey response rate (for those approached)
4
5 was 74%. This study was approved by the institutional review board of the first author's
6
7 institution.
8
9

10 11 12 13 *Sociodemographics and drug use*

14
15 The survey first asked about sociodemographics including age, gender, race/ethnicity,
16
17 educational attainment, and sexual orientation, and participants were asked to report level of
18
19 nightclub/festival/rave/party attendance in the past year. We also recorded whether the
20
21 participant was surveyed outside of a nightclub or festival. Participants were asked about past-
22
23 year use of various drugs via a validated rapid survey instrument developed by the lead author.
24
25 In this analysis we focus on some of the most prevalent drugs in the scene;²¹ specifically, we
26
27 focus on use of ecstasy/MDMA/Molly, LSD, powder cocaine, methamphetamine, and
28
29 nonmedical use of opioids.
30
31
32
33

34 35 *Willingness to provide a hair sample*

36
37 Near the end of the survey, participants were asked their willingness to provide a hair
38
39 sample. Specifically, the survey asked, "Would you be willing to donate a small hair sample to
40
41 us when you complete this survey to be tested for a variety of drugs including 'bath salts'? If so,
42
43 the recruiter will cut a small lock of hair from your head (or elsewhere). A lock is typically taken
44
45 from the lower back of the head from close to the scalp and it is very rarely visible." It was
46
47 further explained that we will use these results to inform prevention and harm reduction in the
48
49 EDM scene and that the recruiter can answer any questions they may have. Answer options for
50
51 willingness to provide a sample were "yes" and "no" and the response was piped to the last
52
53
54
55
56
57
58
59
60

1
2
3 page of the survey to inform the recruiter whether the participant was willing to provide a
4
5 sample. Those checking they were unwilling were provided with a checklist of 13 potential
6
7 reasons for unwillingness to provide a sample. One reason was “other reason” with an option
8
9 to type in a response. We created this checklist based on qualitative responses provided by
10
11 participants in a similar study the previous year.
12
13

14
15 Regardless of their survey response regarding willingness to provide hair, participants
16
17 were still asked verbally by the recruiter if they were willing to provide a sample. Participants
18
19 were either asked shortly before completing the survey or when they had completed the
20
21 survey. This was done for three reasons. First, recruiters were not always able to view the
22
23 participants’ piped responses regarding willingness because some participants closed the last
24
25 page of the survey containing the piped response. Second, sometimes recruiters asked
26
27 participants’ willingness shortly before finishing the survey in order to save time for the
28
29 participant and collect the specimen early. Third, recruiters asked participants their willingness,
30
31 verbally, regardless of their survey response, as many misunderstood what providing a hair
32
33 sample entailed or they required additional information from the recruiter. For example, some
34
35 participants asked recruiters to confirm anonymity, that we were not analyzing their DNA, or
36
37 that hair could be cut from nonvisible areas.
38
39
40
41
42
43
44
45
46

47 *Hair collection*

48
49 If the participant verbally agreed, the recruiter cut a small lock of hair from the
50
51 participant—as close to the scalp as possible using a scissor. Since willingness to provide a
52
53 sample was the main factor examined, unlike some previous epidemiology studies,^{20,36,37} short
54
55
56
57
58
59
60

1
2
3 hair (e.g., <1cm) did not preclude sample donation eligibility.³⁸ In some cases, male participants
4
5 agreed to have hair clipped from the beard, arm, chest, or leg. Some participants also provided
6
7 pubic hair. Hair was folded in a piece of tin foil and placed in a small envelope labeled with the
8
9 participant's study ID number. This ID was linked to the participant's survey responses.
10
11
12 Participants were compensated \$10 USD for completing the survey, but they were not provided
13
14 extra compensation for providing a hair sample. Samples were collected to record response
15
16 rates to inform a larger future study and participants did not receive results.
17
18
19
20
21
22

23 *Statistical analysis*

24
25 We first examined sample characteristics, hair response rates, and reasons for
26
27 unwillingness to provide a hair sample. We then determined whether there were differences in
28
29 reasons for unwillingness according to each demographic covariate and by whether use of any
30
31 of the five drugs included in this analysis (coded into an overall drug use variable) were used in
32
33 the past year. This was done using chi-square and Fisher's exact tests. To prevent inflation of
34
35 Type I Error which may result from repeated testing on 13 separate variables indicating
36
37 different reasons, we utilized a Bonferroni correction ($\alpha=.05/13=.004$).
38
39
40
41

42 We then examined the degree of overlap between those who checked off willingness to
43
44 provide a sample and those who actually provided a sample, and examined whether
45
46 sociodemographic and drug use covariates were related to whether or not a hair sample was
47
48 provided. This was examined using chi-square. Finally, we computed a multivariable logistic
49
50 regression model with hair donation (yes/no) as the outcome, and covariates that were
51
52 significant or approached significance ($p<.10$) in bivariable models were included as
53
54
55
56
57
58
59
60

1
2
3 independent variables. Each of the covariates included in the model is thus adjusted for all the
4
5 other covariates in that model. Data were analyzed using Stata 13 SE (StataCorp, 2013).
6
7
8
9

10 **Results**

11
12 Sample characteristics are shown in Table 1. The majority (56.3%) of the sample
13
14 identified as white, and nearly six out of ten (59.2%) reported having a college degree or higher.
15
16 Most (70.3%) were surveyed outside of a nightclub, and past-year drug use was prevalent.
17
18
19

20 A third (33.4%, n=312) provided a hair sample. However, while 83.3% of those providing
21
22 a sample (260 of 312) indicated willingness to provide a hair sample on the survey, 16.7%
23
24 (n=52) of participants providing a hair sample originally reported unwillingness on the survey.
25
26 Of those not providing a sample (n=621), 94.4% (n=586) provided a concordant response on the
27
28 survey regarding willingness; 5.6% (n=35) reported willingness to provide a sample and then did
29
30 not provide one.
31
32
33
34

35 Table 2 summarizes reasons for refusal among the two-thirds (n=638) of the sample
36
37 reporting unwillingness to provide a hair sample on the survey question. It also presents
38
39 comparisons according to gender and drug use. Lack of interest (21.0%), lack of time (19.8%),
40
41 not wanting a lock of hair cut (17.7%), and disinterest in having hair cut in public (13.8%) were
42
43 the main reported reasons for refusal. In addition, 8.0% refused because they thought we
44
45 would not detect any drugs, 7.4% feared identification, 6.6% thought their hair was too short,
46
47 4.7% refused because they could not receive results, and 2.8% refused because they thought
48
49 drugs would be detected. Three out of ten (30.7%) reported that they were unwilling for none
50
51 of the listed reasons. We also determined that males (10.1%) were more likely than females
52
53
54
55
56
57
58
59
60

1
2
3 (2.9%) to think their hair was too short ($\chi^2(1)=13.89$, $p<.001$), and past-year drug users (3.6%)
4
5 were less likely than non-past-year users (12.1%) to report unwillingness because they thought
6
7 we would not detect drugs ($\chi^2(1)=14.11$, $p<.001$). Past-year drug users (23.8%) were also less
8
9 likely than non-past-year users (37.2%) to check off no listed reasons ($\chi^2(1)=13.40$, $p<.001$).
10
11 Drug users (11.4%) were more likely to fear identification than non-users (3.6%; $\chi^2(1)=14.11$,
12
13 $p<.001$) and drug users (9.5%) were more likely to fear that testing may lead them to get into
14
15 trouble compared to non-users (0.9%; $\chi^2(1)=24.38$, $p<.001$). Comparisons according to all other
16
17 demographic covariates are presented in Supplemental Table 1 and no comparisons were
18
19 significant.
20
21
22
23
24

25 With regard to differences in hair donation according to sample characteristics, as
26
27 shown in Table 3, there was a difference according to sexual orientation with those identifying
28
29 as bisexual or other sexuality being more likely to provide a sample (16.4% vs. 9.6% not
30
31 providing a sample; $\chi^2(2)=10.01$, $p=.007$). Compared to non-past-year users, past-year users of
32
33 ecstasy (43.3% vs. 34.1%, $\chi^2(1)=7.41$, $p=.006$), LSD (34.0% vs. 21.3%, $\chi^2(1)=17.68$, $p<.001$),
34
35 powder cocaine (39.7% vs. 33.2%, $\chi^2(1)=3.92$, $p=.048$), opioids (nonmedical; 16.7% vs. 8.2%,
36
37 $\chi^2(1)=15.11$, $p<.001$), and/or of methamphetamine (4.8% vs. 1.3%, $\chi^2(1)=110.70$, $p=.001$) were
38
39 more likely to provide a sample. We then fit sexual orientation and use of ecstasy, LSD, cocaine,
40
41 opioids, and methamphetamine into the multivariable model, along with race/ethnicity
42
43 ($p=.086$) and venue type ($p=.060$), as these variables approached significance in bivariable
44
45 models. The model met all statistical assumptions and, as a group, predictors were significantly
46
47 associated with providing a hair sample ($\chi^2=53.63$, $p<.001$). Compared to white participants,
48
49 those identifying as Asian were at half the odds of providing a sample (aOR=0.53, $p=.012$) and
50
51
52
53
54
55
56
57
58
59
60

1
2
3 compared to those surveyed outside of a nightclub, those surveyed outside of a festival were at
4
5 higher odds of providing a sample (aOR=1.39, p=.036). Sexual orientation was no longer
6
7 significant; however, past-year use of three different drugs was associated with increased odds
8
9 of providing a sample. Specifically, past-year users of LSD (aOR=1.62, p=.011), opioids
10
11 (nonmedical; aOR=1.93, p=.003), and/or methamphetamine (aOR=3.43, p=.009) were at higher
12
13 odds for providing a hair sample.
14
15
16
17
18
19

20 Discussion

21
22
23 Only a third of individuals surveyed in a high-risk scene were willing to provide a hair
24
25 sample to be analyzed without being provided extra compensation. We believe results of this
26
27 study yield important information to inform future research.
28
29

30
31 Previous studies have found that black or Hispanic participants were less willing (or less
32
33 able) to provide a hair sample than white participants.^{36,37,39} We, however, found that Asian
34
35 Americans were less likely to provide a sample. Previous research suggests that Asian
36
37 Americans are more likely to participate in research if they trust or know the researcher,⁴¹ so
38
39 additional effort may be required to increase trust. Beliefs among individuals of various
40
41 ethnicities should also be considered.⁴²
42
43
44

45
46 Those surveyed outside of festivals were more likely to provide a sample than those
47
48 surveyed outside of nightclubs. Festivals tend to have over 10,000 attendees per day^{43,44} and
49
50 have a wider audience than most nightclubs. It is possible that this diversity led to increased
51
52 response rates. Festivals also take place during the day and attendees may be more willing to
53
54
55
56
57
58
59
60

1
2
3 participate in the daylight. They may also be in less of a hurry, compared to when attending
4
5 parties at night.
6
7

8 Past-year users of drugs such as LSD, opioids, and methamphetamine were more likely
9
10 to provide a hair sample and this corroborates an older study focusing on a general population
11
12 sample.³⁶ While these differences in response rates can in fact bias results, drug users being
13
14 more willing to provide a sample is an important finding because it is these individuals who are
15
16 more likely to use (adulterated) drugs. Such willingness among users suggests potential concern
17
18 about adulterants; therefore, interactions with users in a study setting may present an
19
20 opportunity to provide information (e.g., providing them their results in the future or providing
21
22 them results from past studies in real-time) to inform prevention or harm reduction.
23
24
25
26

27 Reasons for refusal to provide hair were queried to inform the design of future studies
28
29 and some reasons appear to be more addressable than others. For example, lack of overall
30
31 interest was reported by a fifth of our sample. Participant compensation is often provided as a
32
33 means to increase motivation or willingness to take a survey and additional compensation for
34
35 providing a biological specimen may increase willingness.³⁶ A fifth of the sample also indicated
36
37 they were unwilling to provide a sample due to lack of time. We believe verbally asking
38
39 participants their willingness to provide a hair sample earlier during the survey instead of at or
40
41 near the end of the survey may increase willingness to provide a sample during survey
42
43 administration. This would help ensure that the participant will not have to wait until after the
44
45 survey to provide a sample.
46
47
48
49
50

51 Regarding participants not wanting a researcher "messing with" their hair, similar to
52
53 past studies, sometimes this was due to general disinterest and other times it was due to
54
55
56
57
58
59
60

1
2
3 specific hairstyles.³⁹ For example, braids or dreadlocks may prevent donation.³⁷ Qualitatively,
4
5 we learned that some participants were unaware that hair is usually cut from a nonvisible area
6
7 of the head (typically loose hair above the nape). Some participants, primarily males, thought
8
9 their hair was too short to provide, and short hair has been a barrier in previous studies.^{37,39}
10
11 While a shaved or bald head can in fact limit hair collection, in many cases we were still able to
12
13 collect body hair or beard hair. Embarrassment or being uncomfortable having one's hair cut in
14
15 public were also concerns of some participants. These concerns are difficult to address in a
16
17 street-intercept survey, but these concerns may apply less in other studies that are conducted
18
19 in private.
20
21
22
23
24

25 Only 3% reported unwillingness because they thought we would detect drugs. This was
26
27 unexpected considering this was an anonymous drug survey conducted on the street. This may
28
29 be related to two other reasons for refusal—fear of identification and fear of getting into
30
31 trouble. Drug users in particular reported not providing hair for these reasons, likely due to fear
32
33 of legal sanctions, and perhaps additional information is needed to demonstrate that results
34
35 are anonymous. Five percent indicated unwillingness to provide a hair sample because we
36
37 were unable to provide them with their results. Drug checking is increasing in popularity and
38
39 users are provided with results of drug content. A recent study asking festival-attending drug
40
41 users about drug checking found that 94% would use such services, yet of these, almost two-
42
43 thirds would not submit drugs if they could not receive results.⁴⁵ Providing the option for users
44
45 to receive their results would likely increase willingness. While our study was not a drug-
46
47 checking study, it is possible that such biological testing can serve as a proxy for drug checking
48
49 in a retrospective manner to inform prevention and harm reduction.
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Regarding the efficacy of querying willingness to provide a hair sample via survey, many
4 participants who reported unwillingness on the survey changed their mind and provided hair
5 after receiving additional information. We believe providing the initial information and gauging
6 one's initial willingness may have a positive effect on response rates at the end of the survey
7 when verbally asked willingness to provide a sample.
8
9
10
11
12
13
14
15
16
17

18 **Limitations**

19
20 Results may not be generalizable to collection of biological specimens other than hair
21 and results may not be generalizable beyond the EDM party scene. We were unable to record
22 reasons why 5% expressed willingness to provide a sample and then did not provide one.
23 Likewise, we were unable to track what questions were asked or what misconceptions about
24 the study were most commonly discussed with recruiters. The reasons for refusal list was not
25 validated and participants were unable to rate or rank level of importance for each reason for
26 refusal.
27
28
29
30
31
32
33
34
35
36
37
38
39

40 **Conclusions**

41
42 Findings have implications for future research. Only a third of those surveyed provided a
43 hair sample and we found individual-level differences regarding willingness to provide a
44 sample, which can further limit results. So while hair analysis has utility in resolving issues with
45 self-report,^{9,10,20} generalizing findings to populations is limited due to selection bias. Similar
46 surveys incorporating hair collection have found similar biases although response rates tended
47 to be higher. For example, Fendrich et al³⁹ had a response rate of 57% during a household
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 survey; however, \$10 incentive was offered for hair submission in that study. We could not
4
5 offer extra incentive for hair donation in this study, which likely contributed to lower response
6
7 rates. While research is needed to determine adequate amounts for incentives, we recommend
8
9 incentives as a method to potentially increase response rates. Higher response rates in
10
11 household surveys are also expected as it can be difficult obtaining agreement to collect a hair
12
13 sample late at night on the street when a participant is rushing into a party. Race/ethnicity,
14
15 gender, and type of party also need to be considered, and tactics may need to be tested to
16
17 increase response rates among those less likely to respond. We have learned, for example, that
18
19 allowing submissions of body hair increases response rates among those with short head hair.
20
21
22
23
24

25 Much of the public is unfamiliar with hair testing and many individuals have negative
26
27 attitudes toward drug-testing in general as their experience is likely limited to testing in the
28
29 workplace or by law enforcement.⁴⁶ We believe that in this population it is necessary to acquire
30
31 trust that collection is for research purposes and to inform prevention and harm reduction.
32
33 Recruiters can also help response rates by “going the extra mile”³⁹ and trying harder to explain
34
35 the importance of providing a sample to participants. Regardless, the reader must keep in mind
36
37 that both surveying and collecting hair samples via street-intercept late at night can be difficult.
38
39
40
41

42 In sum, while hair collection is a non-invasive method of biological specimen collection,
43
44 it is in fact a challenging method when added to street-intercept surveys. Hair testing has high
45
46 utility in adding information to survey data, but future research needs to examine how to
47
48 increase feasibility and response rates in such at-risk scenes.
49
50
51
52
53
54
55
56
57
58
59
60

References

1. Darke S. Self-report among injecting drug users: a review. *Drug Alcohol Depend.* 1998;51:253-263.
2. Harrison L. The validity of self-reported drug use in survey research: an overview and critique of research methods. *NIDA Res Monogr.* 1997;167:17-36.
3. Pichini S, Solimini R, Berretta P, Pacifici R, Busardo FP. Acute intoxications and fatalities from illicit fentanyl and analogues: an update. *Ther Drug Monit.* 2018;40:38-51.
4. Presence of fentanyl in cocaine contributing to increase in drug overdose deaths [press release]. 2017.
5. European Monitoring Centre for Drugs and Drug Addiction. *European Drug Report 2017.* Lisbon, 2017.
6. U.S. Drug Enforcement Administration DCD. *National Forensic Laboratory Information System: Year 2016 Annual Report.* Springfield, VA; 2017.
7. Drug Enforcement Administration. *Emerging Threat Report: Mid-year 2017.* 2017.
8. Brunt TM, Nagy C, Bucheli A, et al. Drug testing in Europe: monitoring results of the Trans European Drug Information (TEDI) project. *Drug Test Anal.* 2016;9:188-198.
9. Palamar JJ, Salomone A, Vincenti M, Cleland CM. Detection of "bath salts" and other novel psychoactive substances in hair samples of ecstasy/MDMA/"Molly" users. *Drug Alcohol Depend.* 2016;161:200-205.
10. Palamar JJ, Salomone A, Gerace E, Di Corcia D, Vincenti M, Cleland CM. Hair testing to assess both known and unknown use of drugs amongst ecstasy users in the electronic dance music scene. *Int J Drug Policy.* 2017;48:91-98.
11. Wood DM, Stribley V, Dargan PI, Davies S, Holt DW, Ramsey J. Variability in the 3,4-methylenedioxymethamphetamine content of 'ecstasy' tablets in the UK. *Emerg Med J.* 2011;28:764-765.
12. Vogels N, Brunt TM, Rigter S, van Dijk P, Vervaeke H, Niesink RJ. Content of ecstasy in the Netherlands: 1993-2008. *Addiction.* 2009;104:2057-2066.
13. Caudevilla-Gálligo F, Ventura M, Indave Ruiz BI, Fornís I. Presence and composition of cathinone derivatives in drug samples taken from a drug test service in Spain (2010-2012). *Hum Psychopharmacol.* 2013;28:341-344.
14. Vidal Gine C, Ventura Vilamala M, Fornis Espinosa I, et al. Crystals and tablets in the Spanish ecstasy market 2000-2014: are they the same or different in terms of purity and adulteration? *Forensic Sci Int.* 2016;263:164-168.
15. Renfro CL. MDMA on the street: Analysis Anonymous. *J Psychoactive Drugs.* 1986;18:363-369.
16. Murray RA, Doering PL, Boothby LA, et al. Putting an ecstasy test kit to the test: harm reduction or harm induction? *Pharmacotherapy.* 2003;23:1238-1244.
17. Saleemi S, Pennybaker SJ, Wooldridge M, Johnson MW. Who is 'Molly'? MDMA adulterants by product name and the impact of harm-reduction services at raves. *J Psychopharmacol (Oxford, England).* 2017;31:1056-1060.
18. Gine CV, Vilamala MV, Measham F, et al. The utility of drug checking services as monitoring tools and more: a response to Pirona et al. *Int J Drug Policy.* 2017;45:46-47.

19. Martins D, Barratt MJ, Pires CV, et al. The detection and prevention of unintentional consumption of DOx and 25x-NBOMe at Portugal's Boom Festival. *Hum Psychopharmacol*. 2017;32.
20. Fendrich M, Johnson TP, Wislar JS, Hubbell A, Spiehler V. The utility of drug testing in epidemiological research: results from a general population survey. *Addiction*. 2004;99:197-208.
21. Palamar JJ, Griffin-Tomas M, Ompad DC. Illicit drug use among rave attendees in a nationally representative sample of US high school seniors. *Drug Alcohol Depend*. 2015;152:24-31.
22. Palamar JJ, Acosta P, Ompad DC, Cleland CM. Self-reported ecstasy/MDMA/"molly" use in a sample of nightclub and dance festival attendees in New York City. *Subst Use Misuse*. 2017;52:82-91.
23. Miller BA, Byrnes HF, Branner AC, Voas R, Johnson MB. Assessment of club patrons' alcohol and drug use: the use of biological markers. *Am J Prev Med*. 2013;45:637-643.
24. Surratt HL, Kurtz SP, Buttram M, Levi-Minzi MA, Pagano ME, Cicero TJ. Heroin use onset among nonmedical prescription opioid users in the club scene. *Drug Alcohol Depend*. 2017;179:131-138.
25. Kelly BC, Parsons JT, Wells BE. Prevalence and predictors of club drug use among club-going young adults in New York City. *J Urban Health*. 2006;83:884-895.
26. Yamamoto T, Kawsar A, Ramsey J, Dargan PI, Wood DM. Monitoring trends in recreational drug use from the analysis of the contents of amnesty bins in gay dance clubs. *QJM*. 2013;106:1111-1117.
27. Mohr ALA, Friscia M, Yeakel JK, Logan BK. Use of synthetic stimulants and hallucinogens in a cohort of electronic dance music festival attendees. *Forensic Sci Int*. 2018;282:168-178.
28. Miller P, Curtis A, Jenkinson R, Droste N, Bowe SJ, Pennay A. Drug use in Australian nightlife settings: estimation of prevalence and validity of self-report. *Addiction*. 2015;110:1803-1810.
29. Miller BA, Furr-Holden D, Johnson MB, Holder H, Voas R, Keagy C. Biological markers of drug use in the club setting. *J Stud Alcohol Drugs*. 2009;70:261-268.
30. Johnson MB, Voas RA, Miller BA, Holder HD. Predicting drug use at electronic music dance events: self-reports and biological measurement. *Eval Rev*. 2009;33:211-225.
31. Jufer R, Walsh SL, Cone EJ, Sampson-Cone A. Effect of repeated cocaine administration on detection times in oral fluid and urine. *J Anal Toxicol*. 2006;30:458-462.
32. Smith-Kielland A, Skuterud B, Mørland J. Urinary excretion of 11-nor-9-carboxy-delta9-tetrahydrocannabinol and cannabinoids in frequent and infrequent drug users. *J Anal Toxicol*. 1999;23:323-332.
33. Vindenes V, Yttredal B, Oiestad EL, et al. Oral fluid is a viable alternative for monitoring drug abuse: detection of drugs in oral fluid by liquid chromatography-tandem mass spectrometry and comparison to the results from urine samples from patients treated with Methadone or Buprenorphine. *J Anal Toxicol*. 2011;35:32-39.
34. Caplan YH, Goldberger BA. Alternative specimens for workplace drug Testing. *J Anal Toxicol*. 2001;25:396-399.

- 1
- 2
- 3
- 4 35. Kintz P, Salomone A, Vincenti M. *Hair Analysis in Clinical and Forensic Toxicology*. San
- 5 Diego, CA: Academic Press; 2015.
- 6 36. Fendrich M, Johnson TP, Wislar JS, Hubbell A. Drug test feasibility in a general
- 7 population household survey. *Drug Alcohol Depend*. 2004;73:237-250.
- 8 37. Ford JL, Boch SJ, McCarthy DO. Feasibility of hair collection for cortisol measurement in
- 9 population research on adolescent health. *Nurs Res*. 2016;65:249-255.
- 10 38. Colon HM, Robles RR, Sahai H. The validity of drug use responses in a household survey
- 11 in Puerto Rico: comparison of survey responses of cocaine and heroin use with hair
- 12 tests. *Int J Epidemiol*. 2001;30:1042-1049.
- 13 39. Fendrich M, Johnson T, Wislar JS, Sudman S. The feasibility of hair testing in a household
- 14 survey of drug abuse. In: Mieczkowski T, ed. *Drug testing technology: Assessment of*
- 15 *field applications*. Boca Raton, FL: CRC Press; 1999.
- 16 40. MacKellar DA, Gallagher KM, Finlayson T, Sanchez T, Lansky A, Sullivan PS. Surveillance
- 17 of HIV risk and prevention behaviors of men who have sex with men--a national
- 18 application of venue-based, time-space sampling. *Public Health Rep*. 2007;122:39-47.
- 19 41. George S, Duran N, Norris K. A systematic review of barriers and facilitators to minority
- 20 research participation among African Americans, Latinos, Asian Americans, and Pacific
- 21 Islanders. *Am J Public Health*. 2014;104:e16-31.
- 22 42. Coetzee B, Kagee A, Tomlinson M, Warnich L, Ikediobi O. Reactions, beliefs and concerns
- 23 associated with providing hair specimens for medical research among a South African
- 24 sample: a qualitative approach. *Future Virol*. 2012;7:1135-1142.
- 25 43. Ridpath A, Driver CR, Nolan ML, et al. Illnesses and deaths among persons attending an
- 26 electronic dance-music festival - New York City, 2013. *MMWR Morb Mortal Wkly Rep*.
- 27 2014;63:1195-1198.
- 28 44. Lund A, Turris SA. Mass-gathering medicine: risks and patient presentations at a 2-day
- 29 electronic dance music event. *Prehosp Disaster Med*. 2015;30:271-278.
- 30 45. Barratt MJ, Bruno R, Ezard N, Ritter A. Pill testing or drug checking in Australia:
- 31 acceptability of service design features. *Drug Alcohol Rev*. 2018;37:226-236.
- 32 46. Fendrich M, Kim JYS. The experience and acceptability of drug testing: poll trends. *J Drug*
- 33 *Issues*. 2002;32:81-95.
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60

Table 1. Sample characteristics (*N*=933).

	N	%
Age		
18-24	424	45.4
25-40	509	54.6
Sex		
Male	480	51.5
Female	453	48.5
Race/Ethnicity		
White	525	56.3
Black	72	7.7
Hispanic	169	18.1
Asian	108	11.6
Other/Mixed	59	6.3
Education		
High School or Less	120	12.9
Some College	261	28.0
College Degree	419	44.9
Graduate School	133	14.3
Sexual Orientation		
Heterosexual	756	81.0
Gay/Lesbian	66	7.1
Bisexual/Other Sexuality	111	11.9
Level of EDM Party Attendance		
Never or a Couple of Times per Year	195	20.9
Every Couple of Months	245	26.3
Monthly	176	18.9
Biweekly	166	17.8
Weekly or More Often	151	16.2
Type of Venue Where Surveyed		
Nightclub	656	70.3
Festival	277	29.7
Past-Year Drug Use		
Ecstasy/MDMA/Molly	347	37.2
LSD	238	25.5
Powder Cocaine	330	35.4
Opioids (nonmedical)	103	11.0
Methamphetamine	23	2.5

Table 2. Reasons for being unwilling to provide a hair sample (N=638).

	Full Sample		Sex		χ^2	Past-Year Drug Use		
	N	%	Male % (n)	Female % (n)		No % (n)	Yes % (n)	χ^2
I'm simply not interested	134	21.0	17.9 (58)	24.2 (76)	3.82	17.8 (59)	24.4 (75)	4.19
I don't have the time	126	19.8	21.0 (68)	18.5 (58)	0.64	15.7 (52)	24.1 (74)	7.08
I don't want anyone messing with my hair	113	17.7	14.8 (48)	20.7 (65)	3.79	17.2 (57)	18.2 (56)	0.11
I would not want to have a piece of my hair cut in public	88	13.8	10.5 (34)	17.2 (54)	6.03	12.7 (42)	15.0 (46)	0.71
I know you won't find any drugs in it	51	8.0	6.5 (21)	9.6 (30)	2.05	12.1 (40)	3.6 (11)	15.65*
I fear I will be identified	47	7.4	8.0 (26)	6.7 (21)	0.42	3.6 (12)	11.4 (35)	14.11*
It may be embarrassing	45	7.1	9.0 (29)	5.1 (16)	3.61	5.1 (17)	9.1 (28)	3.86
My hair is probably too short	42	6.6	10.2 (33)	2.9 (9)	13.89*	7.0 (23)	6.2 (19)	0.15
I fear I may somehow get in trouble	32	5.0	5.6 (18)	4.5 (14)	0.40	0.9 (3) [†]	9.5 (29)	24.38*
Because you're not able to give me the results	30	4.7	4.6 (15)	4.8 (15)	0.01	2.4 (8)	7.2 (22)	8.02
I know you will find drugs in it	18	2.8	2.8 (9)	2.9 (9)	0.00	1.5 (5)	4.2 (13)	4.31
Other reason	12	1.9	2.5 (8) [†]	1.3 (4)	1.23	1.2 (4) [†]	2.6 (8)	1.72
None of the above	196	30.7	34.6 (112)	26.8 (84)	4.58	37.2 (123)	23.8 (73)	13.40*

Note. Only those who checked that they were unwilling to provide a hair sample were asked for reasons for unwillingness. Percentages according to gender and drug use are absolute percentages. [†] Indicates that Fisher's Exact Test p-value was used instead of chi-square when there were fewer than 5 participants in a cell. The majority (79.3%) of participants checked off one reason for unwillingness, 10.5% checked off two reasons and 10.2% reported three or more reasons. The mean number of reasons reported was 1.46 (SD = 1.23). "Past-year" drug use as defined as reporting past-year use of ecstasy, powder cocaine, LSD, opioids (nonmedical), or methamphetamine. A Bonferroni statistical correction was applied to reduce Type I Error due to multiple testing (alpha = .05 / 13 outcomes = .004). *p < .004

Table 3. Sociodemographic and drug use characteristics according to whether a hair sample was provided.

	Raw Proportions		Multivariable Model		
	Did Not Provide a Sample, % (n)	Provided a Sample, % (n)	χ^2	aOR	95% CI
Age			0.15		
18-24 (reference)	45.9 (285)	44.6 (139)			
25-40	54.1 (336)	55.4 (173)			
Sex			0.24		
Male (reference)	52.0 (323)	50.3 (157)			
Female	48.0 (298)	49.7 (155)			
Race/Ethnicity			8.17		
White (reference)	54.1 (336)	60.6 (189)		1.00	
Black	7.4 (46)	8.3 (26)		1.20	(0.71, 2.04)
Hispanic	18.4 (114)	17.6 (55)		0.85	(0.58, 1.25)
Asian	13.5 (84)	7.7 (24)		0.53*	(0.32, 0.87)
Other/Mixed	6.6 (41)	5.8 (18)		0.76	(0.42, 1.38)
Education			2.11		
High School or Less (reference)	12.2 (76)	14.1 (44)			
Some College	27.5 (171)	28.9 (90)			
College Degree	44.9 (279)	44.9 (140)			
Graduate School	15.3 (95)	12.2 (38)			
Sexual Orientation			10.01**		
Heterosexual (reference)	82.5 (512)	78.2 (244)		1.00	
Gay/Lesbian	7.9 (49)	5.5 (17)		0.54	(0.29, 1.01)
Bisexual/Other Sexuality	9.7 (60)	16.4 (51)		1.47	(0.97, 2.25)
Level of EDM Party Attendance			5.29		
Never /Couple of Times per Year (reference)	22.2 (138)	18.3 (57)			
Every Couple of Months	26.6 (165)	25.6 (80)			
Monthly	17.9 (111)	20.8 (65)			
Biweekly	18.5 (115)	16.4 (51)			
Weekly or More Often	14.8 (92)	18.9 (59)			
Type of Venue Where Surveyed			3.53		
Nightclub (reference)	72.3 (449)	66.4 (207)		1.00	
Festival	27.7 (172)	33.6 (105)		1.39*	(1.02, 1.89)
Past-Year Drug Use					
Ecstasy/MDMA/Molly			7.41**		
No (reference)	65.9 (409)	56.7 (177)		1.00	
Yes	34.1 (212)	43.3 (135)		1.12	(0.79, 1.59)
LSD			17.68***		
No (reference)	78.7 (489)	66.0 (206)		1.00	
Yes	21.3 (132)	34.0 (106)		1.62*	(1.11, 2.35)
Powder Cocaine			3.92*		
No (reference)	66.8 (415)	60.3 (188)		1.00	

Yes	33.2 (206)	39.7 (124)	0.88	(0.62, 1.27)
Opioids (nonmedical)			15.11***	
No (reference)	91.8 (570)	83.3 (260)	1.00	
Yes	8.2 (51)	16.7 (52)	1.93**	(1.25, 2.99)
Methamphetamine			10.70***	
No (reference)	98.7 (613)	95.2 (297)	1.00	
Yes	1.3 (8)	4.8 (15)	3.43**	(1.36, 8.62)

Note. aOR = adjusted odds ratio (controlling for all covariates); CI = confidence interval. Asterisks next to covariate name indicate significance in bivariable model. Only bivariable findings that were significant or approached significance ($p < .10$) were included in the multivariable model.

* $p < .05$, ** $p < .01$, *** $p < .001$

Supplemental Table 1. Reasons for being unwilling to provide a hair sample (N=638).

	Age			Race/Ethnicity					χ^2
	18-24 % (n)	25-40 % (n)	χ^2	White % (n)	Black % (n)	Hispanic % (n)	Asian % (n)	Other % (n)	
I'm simply not interested	22.0 (65)	20.2 (69)	0.30	20.3 (71)	6.7 (3)	26.7 (31)	18.9 (17)	31.6 (12)	10.76
I don't have the time	18.9 (56)	20.5 (70)	0.24	19.2 (67)	22.2 (10)	17.2 (20)	21.1 (19)	26.3 (10)	1.84
I don't want anyone messing with my hair	18.6 (55)	17.0 (58)	0.29	18.3 (64)	24.4 (11)	15.5 (18)	12.2 (11)	23.7 (9)	4.67
I would not want to have a piece of my hair cut in public	12.2 (36)	15.2 (52)	1.24	13.8 (48)	6.7 (3)	16.4 (19)	15.6 (14)	10.5 (4)	3.15
I know you won't find any drugs in it	7.8 (23)	8.2 (28)	0.04	8.6 (30)	13.3 (6)	5.2 (6)	7.8 (7)	5.3 (2)	3.56
I fear I will be identified	5.4 (16)	9.1 (31)	3.11	7.7 (27)	4.4 (2)	5.2 (6)	7.8 (7)	13.2 (5)	3.34
It may be embarrassing	8.8 (26)	5.6 (19)	2.52	8.0 (28)	4.4 (2)	5.2 (6)	7.8 (7)	5.3 (2)	1.85
My hair is probably too short	5.7 (17)	7.3 (25)	0.63	6.3 (22)	13.3 (6)	6.9 (8)	4.4 (4)	5.3 (2)	4.17
I fear I may somehow get in trouble	5.1 (15)	5.0 (17)	0.00	5.7 (20)	2.2 (1)	3.5 (4)	4.4 (2)	7.9 (3)	2.43
Because you're not able to give me the results	5.4 (16)	4.1 (14)	0.61	3.2 (11)	4.4 (2)	10.3 (12)	4.4 (4)	2.6 (1)	10.50
I know you will find drugs in it	3.0 (9)	2.6 (18)	0.10	3.7 (13)	4.4 (2)	2.6 (3)	0.0 (0)	0.0 (0)	5.21
Other reason	2.4 (7)	1.5 (5)	0.68	1.2 (4)	6.7 (3)	2.6 (3)	2.2 (2)	0.0 (0)	7.66
None of the above	34.1 (101)	27.8 (95)	3.00	30.4 (106)	26.7 (12)	31.0 (36)	33.3 (30)	31.6 (12)	0.67

	Education				χ^2	Sexual Orientation			χ^2
	High School or Less % (n)	Some College % (n)	College Degree % (n)	Graduate School % (n)		Hetero- sexual % (n)	Gay/lesbian % (n)	Bisexual/ Other % (n)	
I'm simply not interested	23.3 (17)	23.2 (42)	22.0 (63)	12.2 (12)	5.47	20.8 (109)	15.7 (8)	27.4 (17)	2.43
I don't have the time	20.6 (15)	18.8 (34)	21.00 (60)	17.4 (17)	0.77	19.6 (103)	23.5 (12)	17.7 (11)	0.62
I don't want anyone messing with my hair	12.3 (9)	23.8 (43)	14.3 (41)	20.4 (20)	8.72	16.6 (87)	27.5 (14)	19.4 (12)	3.90
I would not want to have a piece of my hair cut in public	9.6 (7)	12.7 (23)	16.1 (46)	12.2 (12)	2.72	13.3 (70)	15.7 (8)	16.1 (10)	0.53
I know you won't find any drugs in it	1.4 (1)	9.4 (17)	7.3 (21)	12.2 (12)	7.41	8.0 (42)	5.9 (3)	9.7 (6)	0.55
I fear I will be identified	5.5 (4)	6.1 (11)	7.7 (22)	10.2 (10)	2.02	6.5 (34)	13.7 (7)	9.7 (6)	4.12
It may be embarrassing	11.0 (8)	8.3 (15)	5.9 (17)	5.1 (5)	3.22	6.7 (35)	9.8 (5)	8.1 (5)	0.81
My hair is probably too short	4.1 (3)	6.6 (12)	5.9 (17)	10.2 (10)	3.01	5.9 (31)	11.8 (6)	8.1 (5)	2.84
I fear I may somehow get in trouble	5.5 (4)	3.9 (7)	5.2 (15)	6.1 (6)	0.82	4.6 (24)	7.8 (4)	6.5 (4)	1.34
Because you're not able to give me the results	5.5 (4)	7.2 (13)	3.5 (10)	3.1 (3)	4.10	4.0 (21)	7.8 (4)	8.1 (5)	3.26

I know you will find drugs in it	1.4 (1)	3.9 (7)	2.5 (7)	3.1 (3)	1.45	1.9 (10)	7.8 (4)	6.5 (4)	9.28
Other reason	1.4 (1)	2.2 (4)	2.1 (6)	1.0 (1)	0.69	1.5 (8)	2.0 (1)	4.9 (3)	3.40
None of the above	41.1 (30)	32.6 (59)	27.6 (79)	28.6 (28)	5.49	32.6 (171)	25.5 (13)	19.4 (12)	5.26
	Type of Venue Where Surveyed			Level of EDM Party Attendance					
	Nightclub % (n)	Festival % (n)	χ^2	Never or rarely % (n)	Every few months % (n)	Monthly % (n)	Biweekly % (n)	Weekly or more % (n)	χ^2
I'm simply not interested	20.7 (96)	21.8 (38)	0.10	20.4 (28)	17.1 (29)	27.5 (30)	24.2 (29)	17.7 (18)	5.83
I don't have the time	18.8 (87)	22.4 (39)	1.07	17.5 (24)	19.4 (33)	25.7 (28)	17.5 (21)	19.6 (20)	3.25
I don't want anyone messing with my hair	16.4 (76)	21.3 (37)	2.07	13.1 (18)	19.4 (33)	17.4 (19)	16.7 (20)	22.6 (23)	4.04
I would not want to have a piece of my hair cut in public	13.8 (64)	13.8 (24)	0.00	13.9 (19)	12.4 (21)	7.3 (8)	18.3 (22)	17.7 (18)	7.47
I know you won't find any drugs in it	8.0 (37)	8.1 (14)	0.00	10.2 (14)	6.5 (11)	9.2 (10)	7.5 (9)	6.9 (7)	1.88
I fear I will be identified	9.1 (42)	2.9 (5)	7.08	2.9 (4)	7.1 (12)	8.3 (9)	11.7 (14)	7.8 (8)	7.41
It may be embarrassing	7.3 (34)	6.3 (11)	0.20	5.8 (8)	6.5 (11)	6.4 (7)	10.8 (13)	5.9 (6)	3.29
My hair is probably too short	6.5 (30)	6.9 (12)	0.04	8.8 (12)	7.1 (12)	6.4 (7)	5.8 (7)	3.9 (4)	2.41
I fear I may somehow get in trouble	6.0 (28)	2.3 (4)	3.71	2.2 (3)	4.7 (8)	7.3 (8)	5.0 (6)	6.9 (7)	4.30
Because you're not able to give me the results	4.3 (20)	5.8 (10)	0.58	4.4 (6)	5.3 (9)	3.7 (4)	5.0 (6)	4.9 (5)	0.46
I know you will find drugs in it	3.5 (16)	1.2 (2)	2.44	1.5 (2)	1.8 (3)	4.6 (5)	3.3 (4)	3.9 (4)	3.42
Other reason	2.2 (10)	1.2 (2)	0.70	2.9 (4)	0.6 (1)	0.9 (1)	3.4 (4)	2.0 (2)	4.27
None of the above	32.3 (150)	26.4 (46)	2.06	32.9 (45)	32.9 (56)	26.6 (29)	30.0 (36)	29.4 (30)	1.66

Note. Only those who checked that they were unwilling to provide a hair sample were asked for reasons for unwillingness. A Bonferroni statistical correction was applied to reduce Type I Error due to multiple testing ($\alpha = .05 / 13$ outcomes = .004). * $p < .004$

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Running head: DRUG TESTING AND HAIR SAMPLE RESPONSE RATES

For Peer Review Only

Willingness to Provide a Hair Sample for Drug Testing among
Electronic Dance Music Party Attendees

Abstract

Background: [Non-disclosure of drug use on surveys is common and](#) many drug users unknowingly ingest adulterant or replacement drugs, which leads to underreporting of use of these drugs. Biological testing can complement survey research, and hair-testing is an [appealing](#) method as many drugs are detectable for months post-use. We examined willingness to donate a hair sample to be tested among those surveyed in a population at high risk for consuming adulterated drugs—electronic dance music (EDM) party attendees.

Methods: We surveyed 933 adults entering EDM parties in New York City in 2017. Hair donation response rates and reasons for refusal were examined [from this cross-sectional study](#).

Results: A third (n=312; 33.4%) provided a hair sample. Lack of interest (21.0%), lack of time (19.8%), not wanting a lock of hair cut (17.7%), and disinterest in having hair cut in public (13.8%) were the main reported reasons for refusal. 4.7% refused because they could not receive results. Past-year drug users were more likely to fear identification than non-users (p<.001). Asian participants were at lower odds of providing a hair sample ([aOR=0.53, 95% CI=0.32-0.87](#)), and those reporting past-year use of LSD ([aOR=1.62, 95% CI=1.11-2.35](#)), opioids (nonmedical; [aOR=1.93, 95% CI=1.25-2.99](#)), and/or methamphetamine ([aOR=3.43, 95% CI=1.36-8.62](#)) were at higher odds of providing a sample than non-users of these drugs.

Conclusions: [Only a third provided a hair sample and we found individual-level differences regarding willingness to provide a sample. Factors contributing to refusal should be considered to increase response rates and generalizability of results.](#)

Key words: [data collection](#) methods; [drug screening tests](#); [epidemiology](#)

Introduction

Accurate data on prevalence of drug use is important to inform prevention and harm reduction efforts. Surveys are the most common source of data in drug epidemiology studies, but not all survey respondents provide accurate information about drug use. Dishonest reporting about drug use has always been a concern among researchers.^{1,2} However, a newer concern is inaccurate reporting in which an individual may have thought he or she used a particular drug, but in fact used a different drug, or a drug containing adulterant drugs. It is important to investigate what drugs a user believes he or she used, but it is also important for epidemiologists to know what drugs were actually used. Biological testing is common in intervention research, but most surveys do not include biological measures—especially measures capable of detecting new psychoactive substances (NPS) and fentanyl analogs, which continue to emerge.⁵⁻⁸ In this paper, we examine willingness of individuals to provide a hair sample to be analyzed for NPS during a survey of a population at high risk for unintentionally using adulterated drugs—electronic dance music (EDM) party attendees.^{9,10}

Testing of drug products (e.g., pills, powders) has been the most common method used to test for drug adulteration. While seized drugs are typically tested for their contents,^{7,8} drug content information is frequently sought by users of certain drugs to inform harm reduction.^{8,11-}

¹⁵ Self-testing kits for drugs such as ecstasy have been popular for decades,¹⁶ and drug checking has increased in popularity with some harm reduction organizations conducting on-site drug-testing outside of EDM festivals.¹⁷⁻¹⁹ Other services throughout Europe allow users to drop off or send in drugs to be tested, and ~~all of~~ these organizations provide drug content results to users.^{8,11-14} Testing illegal drugs, however, can lead to legal risk for both users and testers, so

1
2
3 biological specimen testing is another option to determine drug adulteration. While this
4
5 method can only test for drugs already used, such testing can serve as an objective measure of
6
7 drug use in studies and test results can be used to inform prevalence estimates of use.²⁰
8
9
10 Providing users with their results can also inform prevention and harm reduction regarding
11
12 future use.
13

14
15 Individuals in the EDM scene are not only at high risk for use of drugs such as
16
17 ecstasy/Molly and various other drugs,²¹⁻²⁶ but they are also at high risk for using drugs
18
19 adulterated with NPS.^{9,10,17,19,27} Few studies, however, have added biological testing to surveys
20
21 conducted at or outside of nightlife parties. Multiple studies focusing on nightclub scenes have
22
23 tested attendees' saliva for various common drugs via oral assay,^{23,28-30} and Mohr et al²⁷ tested
24
25 saliva, urine, and blood of individuals entering EDM festivals for NPS and other drugs. However,
26
27 test results were not compared to self-reported use in these studies, so it is unknown whether
28
29 drugs were used intentionally or unintentionally. Palamar et al^{9,10} added hair testing to two
30
31 epidemiology surveys focusing on EDM attendees to determine the extent of unintentional use
32
33 of NPS among ecstasy users, but like many studies, only a convenience sample of survey
34
35 respondents were tested. While these studies incorporating biological assays are informative,
36
37 most results are based on convenience sampling, which limits generalizability of findings. **More**
38
39 Systematic methods of obtaining biological specimens in addition to larger epidemiology
40
41 surveys are needed.
42
43
44
45
46
47
48

49
50 Hair testing is beneficial because unlike urine, saliva, or blood, in which drugs are only
51
52 detectable for days or hours post-use,³¹⁻³⁴ many drugs (e.g., NPS such as "bath salts") are
53
54 detectable in hair for months post-use.³⁵ Likewise, hair can be collected quickly, without pain,
55
56
57
58
59
60

1
2
3 in almost any environment (e.g., in the dark, cold, or rain). Thus, collection is typically not as
4 burdensome to participants as other biological testing methods. However, while hair-testing is
5 an appealing method to test for both known and unknown drug use, studies have not examined
6 acceptability of hair collection in at-risk scenes. Response rates and reasons for refusal have
7 been examined in population and household studies,³⁶⁻³⁹ but acquiring such data in at-risk
8 scenes is needed to inform future studies that incorporate this method as an objective
9 measure. We examined willingness of survey respondents in this high-risk scene to provide a
10 hair sample for analysis as part of an epidemiology survey. We also examined barriers to
11 participants providing samples by querying reasons for unwillingness to provide a sample, and
12 we examined characteristics of individuals most likely to provide a sample. These findings can
13 inform future studies that incorporate biological testing in a systematic manner.
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31

32 **Methods**

33 *Procedure*

34
35
36
37 We utilized time-space sampling to recruit participants, which produces a probability
38 sample of visits to parties within the sampling frame.⁴⁰ Specifically, every week, a list of
39 upcoming EDM parties was created (based on EDM ticket websites). Parties at nightclubs and
40 festivals were randomly selected each week to survey potential participants. Individuals were
41 deemed eligible for this cross-sectional intercept survey study if they were 1) 18-40 years of
42 age, and 2) were about to enter the randomly selected party. Passersby—who were alone or in
43 groups—were approached by recruiters, and if confirmed eligible, were asked if they would like
44 to take a drug survey. Participants provided informed consent and self-administered the
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 surveys on tablets. A total of 933 participants completed the survey. The survey was conducted
4
5 from June through September of 2017 and the survey response rate (for those approached)
6
7 was 74%. This study was approved by the institutional review board of the first author's
8
9 institution.
10
11
12
13
14

15 *Sociodemographics and drug use*

16
17 The survey first asked about sociodemographics including age, gender, race/ethnicity,
18
19 educational attainment, and sexual orientation, and participants were asked to report level of
20
21 nightclub/festival/rave/party attendance in the past year. We also recorded whether the
22
23 participant was surveyed outside of a nightclub or festival. Participants were asked about past-
24
25 year use of various drugs via a validated rapid survey instrument developed by the lead author.
26
27
28
29

30 In this analysis we focus on some of the most prevalent drugs in the scene;²¹ specifically, we
31
32 focus on use of ecstasy/MDMA/Molly, LSD, powder cocaine, methamphetamine, and
33
34 nonmedical use of opioids.
35
36

37 *Willingness to provide a hair sample*

38
39 Near the end of the survey, participants were asked their willingness to provide a hair
40
41 sample. Specifically, the survey asked, "Would you be willing to donate a small hair sample to
42
43 us when you complete this survey to be tested for a variety of drugs including 'bath salts'? If so,
44
45 the recruiter will cut a small lock of hair from your head (or elsewhere). A lock is typically taken
46
47 from the lower back of the head from close to the scalp and it is very rarely visible." It was
48
49 further explained that we will use these results to inform prevention and harm reduction in the
50
51 EDM scene and that the recruiter can answer any questions they may have. Answer options for
52
53
54
55
56
57
58
59
60

1
2
3 willingness to provide a sample were “yes” and “no” and the response was piped to the last
4
5 page of the survey to inform the recruiter whether the participant was willing to provide a
6
7 sample. Those checking they were unwilling were provided with a checklist of 13 potential
8
9 reasons for unwillingness to provide a sample. One reason was “other reason” with an option
10
11 to type in a response. We created this checklist based on qualitative responses provided by
12
13 participants in a similar study the previous year.
14
15
16
17

18 Regardless of their survey response regarding willingness to provide hair, participants
19
20 were still asked verbally by the recruiter if they were willing to provide a ~~hair~~ sample.
21
22 Participants were either asked shortly before completing the survey or when they had
23
24 completed the survey. This was done for three reasons. First, recruiters were not always able to
25
26 view the participants’ piped responses regarding willingness because some participants closed
27
28 the last page of the survey containing the piped response. Second, sometimes recruiters asked
29
30 participants’ willingness shortly before finishing the survey in order to save time for the
31
32 participant and collect the specimen early. Third, recruiters asked participants their willingness,
33
34 verbally, regardless of their survey response, as many misunderstood what providing a hair
35
36 sample entailed or they required additional information from the recruiter. For example, some
37
38 participants asked recruiters to confirm anonymity, that we were not analyzing ~~or saving~~ their
39
40 DNA, or that hair could be cut from nonvisible areas.
41
42
43
44
45
46
47
48
49

50 *Hair collection*

51
52 If the participant verbally agreed, the recruiter cut a small lock of hair from the
53
54 participant—as close to the scalp as possible using a ~~clean~~ scissor. Since willingness to provide a
55
56
57
58
59
60

1
2
3 sample was the main factor examined, unlike some previous epidemiology studies,^{20,36,37} short
4 hair (e.g., <1cm) did not preclude sample donation eligibility.³⁸ In some cases, male participants
5
6 agreed to have hair clipped from the beard, arm, chest, or leg. Some participants also provided
7
8 pubic hair. Hair was folded in a piece of tin foil and placed in a small envelope labeled with the
9
10 participant's study ID number. This ID was linked to the participant's survey responses.
11
12

13
14
15 Participants were compensated \$10 USD for completing the survey, but they were not provided
16
17
18 extra compensation for providing a hair sample. Samples were collected to record response
19
20 rates to inform a larger future study and participants did not receive results.
21
22

23 24 25 *Statistical analysis*

26
27 We first examined sample characteristics, hair response rates, and reasons for
28
29 unwillingness to provide a hair sample. We then determined whether there were differences in
30
31 reasons for unwillingness according to each demographic covariate and by whether use of any
32
33 of the five drugs included in this analysis (coded into an overall drug use variable) were used in
34
35 the past year. This was done using chi-square and Fisher's exact tests. To prevent inflation of
36
37 Type I Error which may result from repeated testing on 13 separate variables indicating
38
39 different reasons, we utilized a Bonferroni correction ($\alpha=.05/13=.004$).
40
41
42
43

44
45 We then examined the degree of overlap between those who checked off willingness to
46
47 provide a sample and those who actually provided a sample, and examined whether
48
49 sociodemographic and drug use covariates were related to whether or not a hair sample was
50
51 provided. This was examined using chi-square. Finally, we computed a multivariable binary
52
53 logistic regression model with hair donation (yes/no) as the outcome, and covariates that were
54
55
56
57
58
59
60

1
2
3 significant or approached significance ($p < .10$) in bivariable models were included as
4
5 independent variables. Each of the covariates included in the model is thus adjusted for all the
6
7 other covariates in that model. Data were analyzed using Stata 13 SE (StataCorp, 2013).
8
9
10

11 12 13 **Results**

14
15 Sample characteristics are shown in Table 1. The majority (56.3%) of the sample
16
17 identified as white, and nearly six out of ten (59.2%) reported having a college degree or higher.
18
19 Most (70.3%) were surveyed outside of a nightclub, and past-year drug use was prevalent.
20
21

22
23 A third (33.4%, $n=312$) provided a hair sample. However, while 83.3% of those providing
24
25 a sample (260 of 312) indicated willingness to provide a hair sample on the survey, 16.7%
26
27 ($n=52$) of participants providing a hair sample originally reported unwillingness on the survey.
28
29 Of those not providing a sample ($n=621$), 94.4% ($n=586$) provided a concordant response on the
30
31 survey regarding willingness; 5.6% ($n=35$) reported willingness to provide a sample and then did
32
33 not provide one.
34
35

36
37 Table 2 summarizes reasons for refusal among the two-thirds ($n=638$) of the sample
38
39 reporting unwillingness to provide a hair sample on the survey question. It also presents
40
41 comparisons according to gender and drug use. Lack of interest (21.0%), lack of time (19.8%),
42
43 not wanting a lock of hair cut (17.7%), and disinterest in having hair cut in public (13.8%) were
44
45 the main reported reasons for refusal. In addition, 8.0% refused because they thought we
46
47 would not detect any drugs, 7.4% feared identification, 6.6% thought their hair was too short,
48
49 4.7% refused because they could not receive results, and 2.8% refused because they thought
50
51 drugs would be detected. Three out of ten (30.7%) reported that they were unwilling for none
52
53
54
55
56
57
58
59
60

of the listed reasons. We also determined that males (10.1%) were more likely than females (2.9%) to think their hair was too short ($\chi^2(1)=13.89$, $p<.001$), and past-year drug users (3.6%) were less likely than non-past-year users (12.1%) to report unwillingness because they thought we would not detect drugs ($\chi^2(1)=14.11$, $p<.001$); ~~and they~~ Past-year drug users (23.8%) were also less likely than non-past-year users (37.2%) to check off no listed reasons ($\chi^2(1)=13.40$, $p<.001$). Drug users (11.4%) were more likely to fear identification than non-users (3.6%; $\chi^2(1)=14.11$, $p<.001$) and drug users (9.5%) were more likely to fear that testing may lead them to get into trouble compared to non-users (0.9%; $\chi^2(1)=24.38$, $p<.001$). Comparisons according to all other demographic covariates are presented in Supplemental Table 1 and no comparisons were significant.

With regard to differences in hair donation according to sample characteristics, as shown in Table 3, there was a difference according to sexual orientation with those identifying as bisexual or other sexuality being more likely to provide a sample (16.4% vs. 9.6% not providing a sample; $\chi^2(2)=10.01$, $p=.007$). Compared to non-past-year users, past-year users of ecstasy (43.3% vs. 34.1%, $\chi^2(1)=7.41$, $p=.006$), LSD (34.0% vs. 21.3%, $\chi^2(1)=17.68$, $p<.001$), powder cocaine (39.7% vs. 33.2%, $\chi^2(1)=3.92$, $p=.048$), opioids (nonmedical; 16.7% vs. 8.2%, $\chi^2(1)=15.11$, $p<.001$), and/or of methamphetamine (4.8% vs. 1.3%, $\chi^2(1)=110.70$, $p=.001$) were more likely to provide a sample. We then fit sexual orientation and use of ecstasy, LSD, cocaine, opioids, and methamphetamine into the multivariable model, along with race/ethnicity ($p=.086$) and venue type ($p=.060$), as these variables approached significance in bivariable models. The model met all statistical assumptions and, as a group, predictors were significantly associated with providing a hair sample ($\chi^2=53.63$, $p<.001$). Compared to white participants,

1
2
3 those identifying as Asian were at half the odds of providing a sample (aOR=0.53, p=.012) and
4
5 compared to those surveyed outside of a nightclub, those surveyed outside of a festival were at
6
7 higher odds of providing a sample (aOR=1.39, p=.036). Sexual orientation was no longer
8
9 significant; however, past-year use of three different drugs was associated with increased odds
10
11 of providing a sample. Specifically, past-year users of LSD (aOR=1.62, p=.011), opioids
12
13 (nonmedical; aOR=1.93, p=.003), and /or methamphetamine (aOR=3.43, p=.009) were at higher
14
15 odds for providing a hair sample.
16
17
18
19
20
21
22

23 Discussion

24
25 Only a third of individuals surveyed in a high-risk scene were willing to provide a hair
26
27 sample to be analyzed without being provided extra compensation. We believe results of this
28
29 study yield important information to inform future research.
30
31

32
33 Previous studies have found that black or Hispanic participants were less willing (or less
34
35 able) to provide a hair sample than white participants.^{36,37,39} We, however, found that Asian
36
37 Americans were less likely to provide a sample. Previous research suggests that Asian
38
39 Americans are more likely to participate in research if they trust or know the researcher,⁴¹ so
40
41 additional effort may be required to increase trust. Beliefs among individuals of various
42
43 ethnicities should also be considered.⁴²
44
45
46

47
48 Those surveyed outside of festivals were more likely to provide a sample than those
49
50 surveyed outside of nightclubs. Festivals tend to have over 10,000 attendees per day^{43,44} and
51
52 have a wider audience than most nightclubs. It is possible that this diversity led to increased
53
54 response rates. Festivals also take place during the day and attendees may be more willing to
55
56
57
58
59
60

1
2
3 participate in the daylight. They may also be in less of a hurry, compared to when attending
4
5 parties at night.
6

7
8 Past-year users of drugs such as LSD, opioids, and methamphetamine were ~~also~~ more
9
10 likely to provide a hair sample and this corroborates an older study focusing on a general
11
12 population sample.³⁶ While these differences in response rates can in fact bias results, drug
13
14 users being more willing to provide a sample is an important finding because it is these
15
16 individuals who are more likely to use (adulterated) drugs. Such willingness among users
17
18 suggests potential concern about adulterants; therefore, interactions with users in a study
19
20 setting may present an opportunity to provide information (e.g., providing them their results in
21
22 the future or providing them results from past studies in real-time) to inform prevention or
23
24 harm reduction.
25
26
27
28
29

30 Reasons for refusal to provide hair were queried to inform the design of future studies
31
32 and some reasons appear to be more addressable than others. For example, lack of overall
33
34 interest was reported by a fifth of our sample. Participant compensation is often provided as a
35
36 means to increase motivation or willingness to take a survey and additional compensation for
37
38 providing a biological specimen may increase willingness.³⁶ A fifth of the sample also indicated
39
40 they were unwilling to provide a sample due to lack of time. We believe verbally asking
41
42 participants their willingness to provide a hair sample earlier during the survey instead of at or
43
44 near the end of the survey may increase willingness to provide a sample during survey
45
46 administration. This would help ensure that the participant will not have to wait until after the
47
48 survey to provide a sample.
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Regarding participants not wanting a researcher "messaging with" their hair, similar to
4
5 past studies, sometimes this was due to general disinterest and other times it was due to
6
7 specific hairstyles.³⁹ For example, braids or dreadlocks may prevent donation.³⁷ Qualitatively,
8
9 we learned that some participants were unaware that hair is usually cut from a nonvisible area
10
11 of the head (typically loose hair above the nape). Some participants, primarily males, thought
12
13 their hair was too short to provide, and short hair has been a barrier in previous studies.^{37,39}
14
15 While a shaved or bald head can in fact limit hair collection, in many cases we were still able to
16
17 collect body hair or beard hair. Embarrassment or being uncomfortable having one's hair cut in
18
19 public were also concerns of some participants. These concerns are difficult to address in a
20
21 street-intercept survey, but these concerns may apply less in other studies that are conducted
22
23 in private.
24
25
26
27
28
29

30 Only 3% reported unwillingness because they thought we would detect drugs. This was
31
32 unexpected considering this was an anonymous drug survey conducted on the street. This may
33
34 be related to two other reasons for refusal—fear of identification and fear of getting into
35
36 trouble. Drug users in particular reported not providing hair for these reasons, likely due to fear
37
38 of legal sanctions, and perhaps additional information is needed to demonstrate that results
39
40 are anonymous. Five percent indicated unwillingness to provide a hair sample because we
41
42 were unable to provide them with their results. Drug checking is increasing in popularity and
43
44 users are provided with results of drug content. A recent study asking festival-attending drug
45
46 users about drug checking found that 94% would use such services, yet of these, almost two-
47
48 thirds would not submit drugs if they could not receive results.⁴⁵ Providing the option for users
49
50 to receive their results would likely increase willingness. While our study was not a drug-
51
52
53
54
55
56
57
58
59
60

1
2
3 checking study, it is possible that such biological testing can serve as a proxy for drug checking
4
5
6 in a retrospective manner to inform prevention and harm reduction.

7
8 Regarding the efficacy of querying willingness to provide a hair sample via survey, many
9
10 participants who reported unwillingness on the survey changed their mind and provided hair
11
12 after receiving additional information. We believe providing the initial information and gauging
13
14 one's initial willingness may have a positive effect on response rates at the end of the survey
15
16 when verbally asked willingness to provide a sample.
17
18
19
20
21
22

23 **Limitations**

24
25 Results may not be generalizable to collection of biological specimens other than hair
26
27 and results may not be generalizable beyond the EDM party scene. We were unable to record
28
29 reasons why 5% expressed willingness to provide a sample and then did not provide one.
30
31

32 Likewise, we were unable to track what questions were asked or what misconceptions about
33
34 the study were most commonly discussed with recruiters. The reasons for refusal list was not
35
36 validated and participants were unable to rate or rank level of importance for each reason for
37
38 refusal.
39
40
41
42
43
44

45 **Conclusions**

46
47 Findings have implications for future research. Only a third of those surveyed provided a
48
49 hair sample and we found individual-level differences regarding willingness to provide a
50
51 sample, which can further limit results. So while hair analysis has utility in resolving issues with
52
53 self-report,^{9,10,20} generalizing findings to populations is limited due to selection bias. Similar
54
55
56
57
58
59
60

1
2
3 surveys incorporating hair collection have found similar biases although response rates tended
4 to be higher. For example, Fendrich et al³⁹ had a response rate of 57% during a household
5 survey; however, \$10 incentive was offered for hair submission in that study. We could not
6 offer extra incentive for hair donation in this study, which likely contributed to lower response
7 rates. While research is needed to determine adequate amounts for incentives, we recommend
8 incentives as a method to potentially increase response rates. Higher response rates in
9 household surveys are also expected as it can be difficult obtaining agreement to collect a hair
10 sample late at night on the street when a participant is rushing into a party. Race/ethnicity,
11 gender, and type of party also need to be considered, and tactics may need to be tested to
12 increase response rates among those less likely to respond. We have learned, for example, that
13 allowing submissions of body hair increases response rates among those with short head hair.

14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30 Much of the public is unfamiliar with hair testing and many individuals have negative
31 attitudes toward drug-testing in general as their experience is likely limited to testing in the
32 workplace or by law enforcement.⁴⁶ We believe that in this population it is necessary to acquire
33 trust that collection is for research purposes and to inform prevention and harm reduction.
34
35
36
37
38
39
40 Recruiters can also help response rates by “going the extra mile”³⁹ and trying harder to explain
41 the importance of providing a sample to participants. Regardless, the reader must keep in mind
42 that both surveying and collecting hair samples via street-intercept late at night can be difficult.

43
44
45
46
47 In sum, while hair collection is a non-invasive method of biological specimen collection,
48 it is in fact a challenging method when added to street-intercept surveys. Hair testing has high
49 utility in adding information to survey data, but future research needs to examine how to
50 increase feasibility and response rates in such at-risk scenes.
51
52
53
54
55
56
57
58
59
60

References

1. Darke S. Self-report among injecting drug users: a review. *Drug Alcohol Depend.* 1998;51:253-263.
2. Harrison L. The validity of self-reported drug use in survey research: an overview and critique of research methods. *NIDA Res Monogr.* 1997;167:17-36.
3. Pichini S, Solimini R, Berretta P, Pacifici R, Busardo FP. Acute intoxications and fatalities from illicit fentanyl and analogues: an update. *Ther Drug Monit.* 2018;40:38-51.
4. Presence of fentanyl in cocaine contributing to increase in drug overdose deaths [press release]. 2017.
5. European Monitoring Centre for Drugs and Drug Addiction. *European Drug Report 2017.* Lisbon, 2017.
6. U.S. Drug Enforcement Administration DCD. *National Forensic Laboratory Information System: Year 2016 Annual Report.* Springfield, VA; 2017.
7. Drug Enforcement Administration. *Emerging Threat Report: Mid-year 2017.* 2017.
8. Brunt TM, Nagy C, Bucheli A, et al. Drug testing in Europe: monitoring results of the Trans European Drug Information (TEDI) project. *Drug Test Anal.* 2016;9:188-198.
9. Palamar JJ, Salomone A, Vincenti M, Cleland CM. Detection of "bath salts" and other novel psychoactive substances in hair samples of ecstasy/MDMA/"Molly" users. *Drug Alcohol Depend.* 2016;161:200-205.
10. Palamar JJ, Salomone A, Gerace E, Di Corcia D, Vincenti M, Cleland CM. Hair testing to assess both known and unknown use of drugs amongst ecstasy users in the electronic dance music scene. *Int J Drug Policy.* 2017;48:91-98.
11. Wood DM, Stribley V, Dargan PI, Davies S, Holt DW, Ramsey J. Variability in the 3,4-methylenedioxymethamphetamine content of 'ecstasy' tablets in the UK. *Emerg Med J.* 2011;28:764-765.
12. Vogels N, Brunt TM, Rigter S, van Dijk P, Vervaeke H, Niesink RJ. Content of ecstasy in the Netherlands: 1993-2008. *Addiction.* 2009;104:2057-2066.
13. Caudevilla-Gálligo F, Ventura M, Indave Ruiz BI, Fornís I. Presence and composition of cathinone derivatives in drug samples taken from a drug test service in Spain (2010-2012). *Hum Psychopharmacol.* 2013;28:341-344.
14. Vidal Gine C, Ventura Vilamala M, Fornis Espinosa I, et al. Crystals and tablets in the Spanish ecstasy market 2000-2014: are they the same or different in terms of purity and adulteration? *Forensic Sci Int.* 2016;263:164-168.
15. Renfro CL. MDMA on the street: Analysis Anonymous. *J Psychoactive Drugs.* 1986;18:363-369.
16. Murray RA, Doering PL, Boothby LA, et al. Putting an ecstasy test kit to the test: harm reduction or harm induction? *Pharmacotherapy.* 2003;23:1238-1244.
17. Saleemi S, Pennybaker SJ, Wooldridge M, Johnson MW. Who is 'Molly'? MDMA adulterants by product name and the impact of harm-reduction services at raves. *J Psychopharmacol (Oxford, England).* 2017;31:1056-1060.
18. Gine CV, Vilamala MV, Measham F, et al. The utility of drug checking services as monitoring tools and more: a response to Pirona et al. *Int J Drug Policy.* 2017;45:46-47.

19. Martins D, Barratt MJ, Pires CV, et al. The detection and prevention of unintentional consumption of DOx and 25x-NBOMe at Portugal's Boom Festival. *Hum Psychopharmacol*. 2017;32.
20. Fendrich M, Johnson TP, Wislar JS, Hubbell A, Spiehler V. The utility of drug testing in epidemiological research: results from a general population survey. *Addiction*. 2004;99:197-208.
21. Palamar JJ, Griffin-Tomas M, Ompad DC. Illicit drug use among rave attendees in a nationally representative sample of US high school seniors. *Drug Alcohol Depend*. 2015;152:24-31.
22. Palamar JJ, Acosta P, Ompad DC, Cleland CM. Self-reported ecstasy/MDMA/"molly" use in a sample of nightclub and dance festival attendees in New York City. *Subst Use Misuse*. 2017;52:82-91.
23. Miller BA, Byrnes HF, Branner AC, Voas R, Johnson MB. Assessment of club patrons' alcohol and drug use: the use of biological markers. *Am J Prev Med*. 2013;45:637-643.
24. Surratt HL, Kurtz SP, Buttram M, Levi-Minzi MA, Pagano ME, Cicero TJ. Heroin use onset among nonmedical prescription opioid users in the club scene. *Drug Alcohol Depend*. 2017;179:131-138.
25. Kelly BC, Parsons JT, Wells BE. Prevalence and predictors of club drug use among club-going young adults in New York City. *J Urban Health*. 2006;83:884-895.
26. Yamamoto T, Kawsar A, Ramsey J, Dargan PI, Wood DM. Monitoring trends in recreational drug use from the analysis of the contents of amnesty bins in gay dance clubs. *QJM*. 2013;106:1111-1117.
27. Mohr ALA, Friscia M, Yeakel JK, Logan BK. Use of synthetic stimulants and hallucinogens in a cohort of electronic dance music festival attendees. *Forensic Sci Int*. 2018;282:168-178.
28. Miller P, Curtis A, Jenkinson R, Droste N, Bowe SJ, Pennay A. Drug use in Australian nightlife settings: estimation of prevalence and validity of self-report. *Addiction*. 2015;110:1803-1810.
29. Miller BA, Furr-Holden D, Johnson MB, Holder H, Voas R, Keagy C. Biological markers of drug use in the club setting. *J Stud Alcohol Drugs*. 2009;70:261-268.
30. Johnson MB, Voas RA, Miller BA, Holder HD. Predicting drug use at electronic music dance events: self-reports and biological measurement. *Eval Rev*. 2009;33:211-225.
31. Jufer R, Walsh SL, Cone EJ, Sampson-Cone A. Effect of repeated cocaine administration on detection times in oral fluid and urine. *J Anal Toxicol*. 2006;30:458-462.
32. Smith-Kielland A, Skuterud B, Mørland J. Urinary excretion of 11-nor-9-carboxy-delta9-tetrahydrocannabinol and cannabinoids in frequent and infrequent drug users. *J Anal Toxicol*. 1999;23:323-332.
33. Vindenes V, Yttredal B, Oiestad EL, et al. Oral fluid is a viable alternative for monitoring drug abuse: detection of drugs in oral fluid by liquid chromatography-tandem mass spectrometry and comparison to the results from urine samples from patients treated with Methadone or Buprenorphine. *J Anal Toxicol*. 2011;35:32-39.
34. Caplan YH, Goldberger BA. Alternative specimens for workplace drug Testing. *J Anal Toxicol*. 2001;25:396-399.

- 1
- 2
- 3
- 4 35. Kintz P, Salomone A, Vincenti M. *Hair Analysis in Clinical and Forensic Toxicology*. San
- 5 Diego, CA: Academic Press; 2015.
- 6 36. Fendrich M, Johnson TP, Wislar JS, Hubbell A. Drug test feasibility in a general
- 7 population household survey. *Drug Alcohol Depend*. 2004;73:237-250.
- 8 37. Ford JL, Boch SJ, McCarthy DO. Feasibility of hair collection for cortisol measurement in
- 9 population research on adolescent health. *Nurs Res*. 2016;65:249-255.
- 10 38. Colon HM, Robles RR, Sahai H. The validity of drug use responses in a household survey
- 11 in Puerto Rico: comparison of survey responses of cocaine and heroin use with hair
- 12 tests. *Int J Epidemiol*. 2001;30:1042-1049.
- 13 39. Fendrich M, Johnson T, Wislar JS, Sudman S. The feasibility of hair testing in a household
- 14 survey of drug abuse. In: Mieczkowski T, ed. *Drug testing technology: Assessment of*
- 15 *field applications*. Boca Raton, FL: CRC Press; 1999.
- 16 40. MacKellar DA, Gallagher KM, Finlayson T, Sanchez T, Lansky A, Sullivan PS. Surveillance
- 17 of HIV risk and prevention behaviors of men who have sex with men--a national
- 18 application of venue-based, time-space sampling. *Public Health Rep*. 2007;122:39-47.
- 19 41. George S, Duran N, Norris K. A systematic review of barriers and facilitators to minority
- 20 research participation among African Americans, Latinos, Asian Americans, and Pacific
- 21 Islanders. *Am J Public Health*. 2014;104:e16-31.
- 22 42. Coetzee B, Kagee A, Tomlinson M, Warnich L, Ikediobi O. Reactions, beliefs and concerns
- 23 associated with providing hair specimens for medical research among a South African
- 24 sample: a qualitative approach. *Future Virol*. 2012;7:1135-1142.
- 25 43. Ridpath A, Driver CR, Nolan ML, et al. Illnesses and deaths among persons attending an
- 26 electronic dance-music festival - New York City, 2013. *MMWR Morb Mortal Wkly Rep*.
- 27 2014;63:1195-1198.
- 28 44. Lund A, Turris SA. Mass-gathering medicine: risks and patient presentations at a 2-day
- 29 electronic dance music event. *Prehosp Disaster Med*. 2015;30:271-278.
- 30 45. Barratt MJ, Bruno R, Ezard N, Ritter A. Pill testing or drug checking in Australia:
- 31 acceptability of service design features. *Drug Alcohol Rev*. 2018;37:226-236.
- 32 46. Fendrich M, Kim JYS. The experience and acceptability of drug testing: poll trends. *J Drug*
- 33 *Issues*. 2002;32:81-95.
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60

Table 1. Sample characteristics (*N*=933).

	N	%
Age		
18-24	424	45.4
25-40	509	54.6
Sex		
Male	480	51.5
Female	453	48.5
Race/Ethnicity		
White	525	56.3
Black	72	7.7
Hispanic	169	18.1
Asian	108	11.6
Other/Mixed	59	6.3
Education		
High School or Less	120	12.9
Some College	261	28.0
College Degree	419	44.9
Graduate School	133	14.3
Sexual Orientation		
Heterosexual	756	81.0
Gay/Lesbian	66	7.1
Bisexual/Other Sexuality	111	11.9
Level of EDM Party Attendance		
Never or a Couple of Times per Year	195	20.9
Every Couple of Months	245	26.3
Monthly	176	18.9
Biweekly	166	17.8
Weekly or More Often	151	16.2
Type of Venue Where Surveyed		
Nightclub	656	70.3
Festival	277	29.7
Past-Year Drug Use		
Ecstasy/MDMA/Molly	347	37.2
LSD	238	25.5
Powder Cocaine	330	35.4
Opioids (nonmedical)	103	11.0
Methamphetamine	23	2.5

Table 2. Reasons for being unwilling to provide a hair sample (N=638).

	Full Sample		Sex		χ^2	Past-Year Drug Use		
	N	%	Male % (n)	Female % (n)		No % (n)	Yes % (n)	χ^2
I'm simply not interested	134	21.0	17.9 (58)	24.2 (76)	3.82	17.8 (59)	24.4 (75)	4.19
I don't have the time	126	19.8	21.0 (68)	18.5 (58)	0.64	15.7 (52)	24.1 (74)	7.08
I don't want anyone messing with my hair	113	17.7	14.8 (48)	20.7 (65)	3.79	17.2 (57)	18.2 (56)	0.11
I would not want to have a piece of my hair cut in public	88	13.8	10.5 (34)	17.2 (54)	6.03	12.7 (42)	15.0 (46)	0.71
I know you won't find any drugs in it	51	8.0	6.5 (21)	9.6 (30)	2.05	12.1 (40)	3.6 (11)	15.65*
I fear I will be identified	47	7.4	8.0 (26)	6.7 (21)	0.42	3.6 (12)	11.4 (35)	14.11*
It may be embarrassing	45	7.1	9.0 (29)	5.1 (16)	3.61	5.1 (17)	9.1 (28)	3.86
My hair is probably too short	42	6.6	10.2 (33)	2.9 (9)	13.89*	7.0 (23)	6.2 (19)	0.15
I fear I may somehow get in trouble	32	5.0	5.6 (18)	4.5 (14)	0.40	0.9 (3) [†]	9.5 (29)	24.38*
Because you're not able to give me the results	30	4.7	4.6 (15)	4.8 (15)	0.01	2.4 (8)	7.2 (22)	8.02
I know you will find drugs in it	18	2.8	2.8 (9)	2.9 (9)	0.00	1.5 (5)	4.2 (13)	4.31
Other reason	12	1.9	2.5 (8) [†]	1.3 (4)	1.23	1.2 (4) [†]	2.6 (8)	1.72
None of the above	196	30.7	34.6 (112)	26.8 (84)	4.58	37.2 (123)	23.8 (73)	13.40*

Note. Only those who checked that they were unwilling to provide a hair sample were asked for reasons for unwillingness. Percentages according to gender and drug use are absolute percentages. [†] Indicates that Fisher's Exact Test p-value was used instead of chi-square when there were fewer than 5 participants in a cell. The majority (79.3%) of participants checked off one reason for unwillingness, 10.5% checked off two reasons and 10.2% reported three or more reasons. The mean number of reasons reported was 1.46 (SD = 1.23). "Past-year" drug use as defined as reporting past-year use of ecstasy, powder cocaine, LSD, opioids (nonmedical), or methamphetamine. A Bonferroni statistical correction was applied to reduce Type I Error due to multiple testing (alpha = .05 / 13 outcomes = .004). *p < .004

Table 3. Sociodemographic and drug use characteristics according to whether a hair sample was provided.

	Raw Proportions		Multivariable Model		
	Did Not Provide a Sample, % (<i>n</i>)	Provided a Sample, % (<i>n</i>)	χ^2	aOR	95% CI
Age			<u>0.15</u>		
18-24 (<u>reference</u>)	45.9 (285)	44.6 (139)			
25-40	54.1 (336)	55.4 (173)			
Sex			<u>0.24</u>		
Male (<u>reference</u>)	52.0 (323)	50.3 (157)			
Female	48.0 (298)	49.7 (155)			
Race/Ethnicity			<u>8.17</u>		
White (<u>reference</u>)	54.1 (336)	60.6 (189)		1.00	
Black	7.4 (46)	8.3 (26)		1.20	(0.71, 2.04)
Hispanic	18.4 (114)	17.6 (55)		0.85	(0.58, 1.25)
Asian	13.5 (84)	7.7 (24)		0.53*	(0.32, 0.87)
Other/Mixed	6.6 (41)	5.8 (18)		0.76	(0.42, 1.38)
Education			<u>2.11</u>		
High School or Less (<u>reference</u>)	12.2 (76)	14.1 (44)			
Some College	27.5 (171)	28.9 (90)			
College Degree	44.9 (279)	44.9 (140)			
Graduate School	15.3 (95)	12.2 (38)			
Sexual Orientation**			<u>10.01**</u>		
Heterosexual (<u>reference</u>)	82.5 (512)	78.2 (244)		1.00	
Gay/Lesbian	7.9 (49)	5.5 (17)		0.54	(0.29, 1.01)
Bisexual/Other Sexuality	9.7 (60)	16.4 (51)		1.47	(0.97, 2.25)
Level of EDM Party Attendance			<u>5.29</u>		
Never /Couple of Times per Year (<u>reference</u>)	22.2 (138)	18.3 (57)			
Every Couple of Months	26.6 (165)	25.6 (80)			
Monthly	17.9 (111)	20.8 (65)			
Biweekly	18.5 (115)	16.4 (51)			
Weekly or More Often	14.8 (92)	18.9 (59)			
Type of Venue Where Surveyed			<u>3.53</u>		
Nightclub (<u>reference</u>)	72.3 (449)	66.4 (207)		1.00	
Festival	27.7 (172)	33.6 (105)		1.39*	(1.02, 1.89)
Past-Year Drug Use					
Ecstasy/MDMA/Molly**			<u>7.41**</u>		
<u>No</u> (<u>reference</u>)	65.9 (409)	56.7 (177)		1.00	
<u>Yes</u>	34.1 (212)	43.3 (135)		1.12	(0.79, 1.59)
LSD***			<u>17.68***</u>		
<u>No</u> (<u>reference</u>)	78.7 (489)	66.0 (206)		1.00	
<u>Yes</u>	21.3 (132)	34.0 (106)		1.62*	(1.11, 2.35)
Powder Cocaine*			<u>3.92*</u>		
<u>No</u> (<u>reference</u>)	66.8 (415)	60.3 (188)		1.00	

<u>Yes</u>	33.2 (206)	39.7 (124)	0.88	(0.62, 1.27)
Opioids (nonmedical) ^{***}			<u>15.11^{***}</u>	
<u>No (reference)</u>	<u>91.8 (570)</u>	<u>83.3 (260)</u>	<u>1.00</u>	
<u>Yes</u>	<u>8.2 (51)</u>	<u>16.7 (52)</u>	1.93 ^{**}	(1.25, 2.99)
Methamphetamine ^{**}			<u>10.70^{***}</u>	
<u>No (reference)</u>	<u>98.7 (613)</u>	<u>95.2 (297)</u>	<u>1.00</u>	
<u>Yes</u>	<u>1.3 (8)</u>	<u>4.8 (15)</u>	3.43 ^{**}	(1.36, 8.62)

Note. aOR = adjusted odds ratio (controlling for all covariates); CI = confidence interval. Asterisks next to covariate name indicate significance in bivariable model. Only bivariable findings that were significant or approached significance ($p < .10$) were included in the multivariable model.

* $p < .05$, ** $p < .01$, *** $p < .001$