

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Proper gravel management may counteract population decline of the Collared Sand Martin  
Riparia riparia**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1728400> since 2020-02-18T23:34:36Z

*Published version:*

DOI:10.30456/AVO.2019204

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



# UNIVERSITÀ DEGLI STUDI DI TORINO

**This is an author version of the contribution published on:**

**Questa è la versione dell'autore dell'opera:**

[Avocetta 43: 139-147 (2019), DOI: <https://doi.org/10.30456/AVO.2019204>]

**The definitive version is available at:**

**La versione definitiva è disponibile alla URL:**

<http://www.avocetta.org/articles/vol-43-2-ems-a-new-alliance-between-a-riparian-species-and-gravel-diggers-might-counteract-population-declines/>

**Proper gravel management may counteract population decline of the Collared Sand**

**Martin** *Riparia riparia*

Giulia Masoero<sup>1</sup>, Giovanni Boano<sup>2</sup>, Alberto Tamietti<sup>3</sup>, Enrico Caprio<sup>4</sup>

<sup>1</sup> Section of Ecology, Department of Biology, FI-20014 University of Turku, Finland

<sup>2</sup> Museo Civico di Storia Naturale di Carmagnola, Via San Francesco di Sales 188, Carmagnola (TO), Italy

<sup>3</sup> Aree Protette del Po Torinese, Corso Trieste 98, Moncalieri (TO), Italy

<sup>4</sup> Università degli Studi di Torino, Dipartimento di Scienze della Vita e Biologia dei Sistemi, Via Accademia Albertina 13, 10123 Torino, Italy

Corresponding author: Giulia Masoero [giulia.masoero@gmail.com](mailto:giulia.masoero@gmail.com)

ORCID:

Giulia Masoero 0000-0003-4429-7726

Giovanni Boano 0000-0002-2670-4351

Enrico Caprio 0000-0002-5997-5959

Short title: Sand martins, rivers and quarries

**ABSTRACT**

Riparian habitats have gone through major structural changes, and related bird populations had to suffer the consequences or adapt to the newfound conditions. Here, we present the results of the analysis of on the river and quarry evolution, in relation to a long-term monitoring (1970-2016) of a Sand Martins population nesting along the River

Po in northern Italy. During the course of the study, the population changed breeding site, preferring more anthropogenic sites in the surrounding quarries to the natural river banks. The alteration of the river dynamic and linearization of the course, alongside the development of the sand quarries, may have caused this change. We conclude with a consideration on the ways to support the survival of populations of riparian birds. Functional nesting habitat along the river should be better preserved, and potential nesting areas in the surrounding quarries should be protected with long term restoration projects. Sand quarries are not free of threats, and cooperation among stakeholders has proven to be of the utmost importance to ensure the success of the population breeding in the area.

**Keywords:** nest site selection; river dynamic; *Riparia riparia*; quarry restoration.

## INTRODUCTION

Habitat loss, due to destruction and alteration, is one of the most important anthropogenic impacts on animal populations (Newton 1998, Baillie et al. 2004, Curry Lindahl 1972). Among the most affected habitats, riparian ecosystems have been deeply altered with radical changes to their courses, flow and dynamics. The regulation of the flow and course forces rivers to become more and more linear, losing their meanders and changing the erosion of river banks (Brookes 1988). Changes in the riparian system can affect many aquatic animals, and have proven to affect the structure of breeding bird community (Figarski & Kajtoch 2015, Girvetz 2010).

The Collared Sand Martin (*Riparia riparia*; hereafter “Sand Martin”) is riparian species particularly specialised in nesting habitat (Moffatt et al. 2005). This colonial

species usually excavates nest holes in sandy banks, normally in the vicinity of water bodies, such as the eroding banks of meandering rivers and streams, but also in artificial sand deposits (Cramp 1988). The most recent IUCN assessment of the global trend categorized Sand Martin populations as Decreasing (BirdLife International 2016). According to the most recent population assessment published by BirdLife International in 2015, Sand Martin populations are decreasing in many European countries, including Italy (Brichetti & Fracasso 2007, Campedelli et al. 2012). This trend appears even more concerning since the species has also experienced large scale population declines since the early 1960s (Cowley 1979, Jones 1987a).

Climate change has often been considered as the major driver behind Sand Martin population changes and this trans-Saharan migrant has been well studied in terms of the influence that climatic factors in the African wintering quarters have on its survival (Cowley 1979, Svensson 1986, Persson 1987a, Szép 1995, Norman & Peach 2013). In more recent years, there has not been any strong drought in the Sahel region (Evan et al. 2014, Sanogo et al. 2015) that might have driven the global decrease in Sand Martin population sizes. Habitat change might therefore be the cause behind more recent declines. However, despite it being recognised as a cause of population decline (Garrison 1999, del Hoyo et al. 2004), few papers have analysed habitat alteration on the breeding grounds. In California, where the species population is declining mainly due to habitat alteration, Moffatt and colleagues (2005) studied the importance of riparian systems and the effects of different restoration strategies on the species. Sand martins were reported to breed in sandy quarry complexes all over Europe (Alves 1997, Cowley & Siriwardena 2005, Heneberg 2007, Jones 1987b, Krištofík et al. 1994, Kitowski et al. 2015, Morgan 1979, Norman & Peach 2013, Persson 1987b, 1987a). Some of the eastern populations of

Sand Martin, however, were still recently found breeding in natural sites in Poland (Figarski & Kajtoch 2015) and, in large numbers, along the banks of the Tisza River in Hungary (Szép 1995, Szép et al. 2003). Despite this important change in the selection of breeding sites in some areas, to our knowledge there are no studies that have analysed the causes behind this phenomenon.

In this paper, we have analysed survey data, collected over 45 years, of a Sand Martin population in north-western Italy. The monitored population did not present signs of decline during the study, and its fluctuations in numbers appeared to have no correlation with climate (Masoero et al. 2016). We have also studied the river dynamic of the past 68 years and the evolution of the quarry complexes, from 1960 until the present, in the nesting area of the Sand Martins. Our aim was to understand the relationship between Sand Martins and their nesting habitat and how the modifications of the river and of the quarries could affect them.

## **METHODS**

### **Study area and the quarries**

The study area was included about 25 km of the River Po South of the city of Turin, Italy (from 44.971 N 7.693 E to 44.832 N 7.636 E) and all the 15 surrounding quarries, nine active during the study and six already restored to a more natural state (Fig. 1). The area is located in the River Po Torinese Regional Park. The River Po and its tributaries transported and deposited materials during the Middle Pleistocene and the Holocene creating the river channel deposits that currently constitute this area of the Po plain. The sediments at the top layer are 25-100 m thick and are exploited by quarries complexes (Castiglioni et al. 1999). The raw materials, such as sand and gravel, are excavated from

the ground to be used as construction aggregates. After the extraction from the ground, sand and gravel are separated in a range of sizes and stored into stockpiles, creating heaps and banks in which the Sand Martins can build their nests (Fig. 2). The quarries are usually worked in progressive phases, in order to minimise the exposed areas. This method also ensures timely restoration of the already excavated sites while the work is still ongoing. The Park oversees the work of the quarries under its jurisdiction and controls the restoration of the excavated banks and surfaces.

We georeferenced seven different maps of the River Po (from Casalgrasso 44.830 N 7.634 E to Moncalieri 45.008 N 7.679 E) from the years 1922, 1948, 1955, 1963, 1978, 1991 and 2008. To assess river dynamism, we calculated the sinuosity index, i.e. the measures of the degree of sinuosity of meandering rivers calculated as the ratio of the length of the midline of the channel and the air-line distance (Friend & Sinha 1993, Alabyan & Chalov 1998). Values of the sinuosity index close to or equal to 1 correspond to a linear river, whereas higher values correspond to a more winding river.

Data concerning the changes in the quarries were obtained from the databases on extraction activities of the Ministry of Industry and of the Piedmont Region. From 1960 to 1985, the Italian Ministry of Industry collected the data on the amount of extracted construction aggregates for the whole region of Piedmont. After 1982, the Piedmont Region started collecting the data regarding the quarries, obtaining therefore more precise data and we could retrieve data concerning the situation in the 15 quarries located in the study area. We decided to use the amount of extracted material as an index of the quarry activity, and therefore of the amount of potential available nesting habitat because of two main reasons. First, the studied quarries are authorised to extract materials also going below the level of the aquifer, i.e. sand and gravel can still be extracted with the surface

of the active quarry remaining stable. Second, sand martins often colonised the sand heaps of extracted materials and seldom the excavated cliffs.

### **Census of breeding colonies and data analysis**

From 1970 up to 2016 the riverbanks and the sand quarries were surveyed each year to identify breeding sites and their location. The census of the colonies with an exhaustive count of all active nests started in 2000, whereas in earlier years (i.e, since 1970), the monitoring of the river was not conducted quantitatively, due to the difficulty to reach every area of the river. From the end of April to July all the possible breeding sites were checked during multiple occasions, both along the river Po and in active and abandoned quarries, and all active nests were counted. Almost every year there was a change in the nesting location chosen by the population, which sometimes split among two or more quarries if the nesting cliffs were not wide enough.

We used generalized additive models (GAMs; Hastie & Tibshirani 1990) to test for the effects of year on the number of nests in riverbanks assuming a Poisson data distribution. GAMs were performed using R 3.3.1 (R Core Team 2017), with the package mgcv (Wood 2011). The default cubic smoothing spline method with four degrees of freedom was used to smooth the year component. We also evaluated Collared Sand Martin population trends between 2002 and 2016 (years with a complete census of the population) using the software TRIM (Trend & Indices for Monitoring data, TRIM 3.54; Pannekoek & Van Strien 2006). TRIM estimates annual indices and evaluates trends in these indices implementing log-linear models, an approach commonly employed in temporal series analysis. We also tested for a linear relationship between the number of



161 nests in riverbanks and quarries and the amount of gravel and sand extracted from gravel  
162 pits using a generalized linear model, specifying a Poisson data distribution.

## 164 **RESULTS**

### 165 **River dynamic and quarry evolution**

166 The River Po course changed markedly in the study area from 1922 until 1978 (Fig.  
167 3), after which it remained largely unchanged. The sinuosity index of the River Po  
168 decreased from a value of 2.19 in 1922, to a minimum of 1.59 calculated in 2008 (Fig.  
169 4), showing that the river had become increasingly linear. Two large floods happened in  
170 2000 and 2016, and created a few suitable nesting areas that were not colonised by Sand  
171 Martins. The new nesting cliffs were nevertheless very limited compared to the previous  
172 floods, the bank defences did not allow more substantial changes.

173 From the 1960-1985 data from the Italian Minister of Industry it emerged that until  
174 1966 the production of construction materials in Piedmont region was low, and always  
175 below 3 Mm<sup>3</sup>. In the subsequent 10 years it grew steeply, reaching a maximum of 14  
176 Mm<sup>3</sup> in 1976. It started then to decrease, but always maintaining values above the 6 Mm<sup>3</sup>  
177 (Fig. 5a). After 1982, more precise data from the Piedmont Region are available and it  
178 was possible to look in detail at the situation in the 15 quarries present in the study area  
179 (Fig. 5b). During the years from 1982 to 2007, the amount of extracted construction  
180 aggregates presented a fluctuating trend, ranging from a minimum of 110000 m<sup>3</sup> in 1995  
181 and a maximum of 1.71 Mm<sup>3</sup> in 1983. After 2007, the production almost constantly  
182 declined.

### 184 **Census of breeding colonies**

The size of colonies along the riverbanks varied from a maximum of 244 active nests recorded in 1973 to a minimum of 30 nests in 1983 (Fig. 4). No active nests were recorded after 2002. A Generalized Additive Model of the number of nests in riverbanks against year revealed a significant non-linear negative relationship ( $\chi^2_{7.058} = 276.5$ ,  $p < 0.001$ , deviance explained = 79%). We cannot assert that the population breeding along the river was smaller than the actual population, due to incomplete surveys carried on in the last century, but we can assert that after 2002 all the colonies are breeding in the quarries. Colonies in sand quarries were detected the first time in 1983 with 50 active nests and were found always in active quarries. The number of active nests varied from a minimum of 20 nests in 1993 to a maximum of 934 nests in 2007 (Fig. 4). A Generalized Additive Model of the number of nests in sand quarries against year revealed a significant non-linear positive relationship ( $\chi^2_{8.108} = 3960$ ,  $p < 0.001$ , deviance explained = 93%). Using TRIM, the population trend from 2002-2016 is uncertain (multiplicative overall slope model: 1.0067, SE:  $\pm 0.0349$ ). The number of nests counted in sand quarries were positively related with the amount of construction aggregates extracted in the study area (with the standardised variable, intercept:  $6.02 \pm 0.01$ , beta:  $0.40 \pm 0.01$ ;  $F_{1,19}$ : 6.07,  $p = 0.023$ ).

## DISCUSSION

In this paper we analysed breeding observations of Sand Martins along a section of the river Po and the surrounding quarries. At the same time, we looked into detailed data regarding the river course and the development of the quarries. Overall, we think this provides a reasonable explanation to the change in breeding sites and to the lack of decrease in size previously shown by the studied population (Masoero et al. 2016).

## **River and quarries**

By looking at map of the River Po courses along the years and at the values of the sinuosity index we can notice a few things. First, a decrease in the river dynamism in the study area, i.e. the river changed its course markedly until 1978 and after that, the course remained largely unchanged, even after two flooding events in the last 15 years. Second, the low values of the index in the later years show a linearization of the river when compared to the earlier values. The fixed course of the river and the lower sinuosity index indicates a decrease in river dynamism, and as a consequence, a decrease in river functionality (Fehér et al. 2012, Yu et al. 2015). The history behind these changes, both regarding the river and the surrounding quarries, is quite complex and has to be told alongside. Between the 1960s and the 1980s, Italy experienced an economic boom. As the surrounding plain became more anthropized, towns grew. An increasing number of flood protections were built around the surrounding cities, houses, crops and poplar plantations. Without any space left to change its course, the river assumed a more fixed and linear path. In the meanwhile, the amount of construction aggregates excavated was also following the market needs. Up until 1960 the extraction took place mainly inside the river with limited volumes. In the early 1960s, the demand for aggregates slowly started to grow and the quantity of sand and gravel extracted from the riverbed was no longer sufficient and the quarry companies started exploiting the sediments present along the river course and the first lakes created by gravel excavations appeared in the alluvial plain. After 1966, the economic boom lead to a strong increase in the volumes of excavated materials. In the 1970s, the excavation of the riverbed started to decrease, due to more severe environmental laws, until it came to a stop. From the 1980s the extraction

of construction aggregates continued along the river in quarries outside of the riverbed. The existing quarries expanded both in area and depth and the amount of extracted construction aggregates presented a fluctuating trend, following the marked needs. Around 2007, to the economic crisis hit the construction market. Some quarries were closed and the active ones started extract less sand and gravel. Inactive quarries are currently under a restoration process coordinated by the River Po Torinese Regional Park. A small increase in the extracted material in 2016 might be the sign of a mild economic recovery.

#### **Sand martins and breeding habitat**

The evolution of the river habitat and of its quarries had a deep impact on Sand Martins. The population investigated in our study, as other studied populations in many European countries (Cowley & Siriwardena 2005, Norman & Peach 2013), has shown a change in the selection of breeding sites, although the general area remained unchanged. It was not possible to obtain overall population trends for the study area because the census on the riverbanks were not always performed with the same methodology. Despite we reckon this may be a limit for previous years, but we still believe that we were able to identify (qualitatively) the magnitude of the reduction of the nests in the riverbanks, and correctly censused the nests since 2000. In the early years, the population was exclusively breeding in the banks of the River Po. It then began to progressively colonize the adjacent sand quarry complexes, and in 2003 the change had become permanent as the traditional riverbank sites were definitively abandoned. This could be explained by considering the decrease in river dynamism (Moffatt et al. 2005). Lower dynamism means less bank erosion, so new sand cliffs are not created and the old ones, remaining

undisturbed, start to become covered in vegetation. At the same time, the river became more linear, determining a change in the river regime, and as a consequence, faster and more destructive floods that could have had an impact on the population survival and breeding success. In the meanwhile, the need for construction aggregates grew because of the economic boom, and new sand and gravel pits began to spread along the course of the river. Sand quarries offered clear sand cliffs, free of vegetation, apparently the perfect substitute to the traditional breeding sites. Nevertheless, nesting in an active sand quarry presents some potential threats. If the workers are careless or not aware of the presence of a colony, they could destroy it. In Italy, active nests are protected by the law (L. 157/92), but people are often unaware of the law or of the presence of nests. Quarries can therefore turn into an ecological trap, i.e. Sand Martins can start nesting in a good sand cliff and the nests can be destroyed. Another problem can follow from the quarry management which can change following the market needs. In the quarry environment, a good management, alongside the protection and the monitoring of the colonies, is crucial, because such quarries act as surrogates for riverbanks not only for the Sand Martin, but also for other species such as the European Bee-eater (*Merops apiaster*) and the Common Kingfisher (*Alcedo atthis*). With simple initiatives (i.e. leaving undisturbed the sand heaps with active nests during May, June and July) that have very low impact on quarry management, it is possible to combine economic interests with nature conservation. All of the above has been possible in the study area thanks to a positive interaction between the people in charge of wildlife protection and those working in the quarries. The River Po Torinese Regional Park is actively protecting the colonies having these sand quarries under its jurisdiction, and it is also monitoring them. In 2000, the Park staff have also started to raise awareness among the gravel diggers about the presence and the importance

of Sand Martins, and thus, the nests in the gravel pits have been left undisturbed allowing Sand Martins to successfully fledge.

After the quarry exploitation, it is important that the restoration plan allows the Sand Martins to nest for many years to come. Quarry restoration is indeed necessary because of the extensive land disturbances and negative impacts in terms of both safety and environment (Milgrom 2008, Zuquette et al. 2002). Reclamation techniques often try to address this issues by creating a safe environment with more natural slopes and by creating of appropriate surfaces for the establishment of vegetation (Down & Stocks 1978, Haywood 1979, Gunn & Bailey 1993, Legwaila et al. 2015, Muzzi & Mongardi 2016). Nowadays, the fact that colonies were never found in inactive quarries suggests that the sites are not restored in a way that provides suitable nesting habitat. In fact, the sand cliffs and heaps of active quarries used as breeding sites are no longer available after the restoration process.

### **Concluding remarks**

Diversity and abundance of riparian birds is usually increased by a renaturalisation of the river channels often favoured by floods (Kajtoch and Figarski 2013). In most industrialised countries, rivers have been showing signs of decrease in dynamism and functionality, which has led to loss in suitable nesting habitat for the Sand Martins (Moffatt et al. 2005). Not finding suitable habitat along the river, the studied population of Sand Martins probably started nesting in the adjacent sand quarries, which have increased their activity around the same years (Figs. 4 and 5). After 2007, fewer and fewer building materials were needed by the building industry, and many of the quarries present in the area began to close and being reclaimed. The population already lost the traditional

nesting sites along the river: what will happen when all the currently active quarries will be closed and restored?

A thorough analysis of the available literature lead to the conclusion that there are two possible alternatives: to leave sand cliffs close to the water available to the Sand Martins to build their nests, or to build artificial breeding cliffs (Gulickx et al. 2007, Bachmann et al. 2008). The first solution can last a few years, but the cliff can collapse or the vegetation can soon invade it. In any case, it has to be managed throughout the years. The second seems a more stable solution, but comes with another type of problem: ectoparasites. Parasites that infest nest holes include different species, including the tick *Ixodes lividus* (Szép & Møller 1999). Newly formed nest sites do not present high levels of infestation by ectoparasites, but as colonies get older the prevalence can approach 100% (e.g. Szép and Møller 1999). In the traditional breeding sites and in the quarries the colony is usually used for only a year since it is destroyed after the nesting season by floods or by quarry activities, and the parasite load does not approach such high levels. Artificial nests are used from one year to another and can therefore develop very high concentrations of ectoparasites. In a colony of nests artificially excavated in the limestone in the UK, all the nest holes were washed out using a water hose (Gulickx et al. 2007), but the effectiveness of this measure remain to be proven.

A better management of rivers and of the riverbanks in particular should be promoted to help the Sand Martins in their breeding grounds, and it would be beneficial also for many other species of birds (Jankowiak and Ławicki 2014; Figarski and Kajtoch 2015). In areas where the Sand Martins are present and already nesting in quarries, restoration projects should consider the creation of artificial colonies (as in Bachmann et al. 2008). Moreover, active quarries hosting Sand Martin colonies should provide cliffs that should

remain untouched during the whole breeding season, and create new ones every year. This management could help preserving a large number of nests over a long period of time as happened in our study.

### Acknowledgements

We thank all the people that helped collecting the data and the River Po Torinese Regional Park. We are grateful to all the Park staff (in particular E. Castello, F. Nobili and S. Zaghi), and to S. Blasutta, I. Ellena, G. Ramello, S. Tozzi and G. Vaschetti. We acknowledge L. Mancuso and M. Moccagatta for their help in recovering the Piemonte Region data from the database. Thanks also to D.E. Chamberlain, his comments and corrections on an early version of the manuscript were greatly appreciated.

### References

- Alabyan A.M. & Chalov R.S., 1998. Types of river channel patterns and their natural controls. *Earth Surf. Process. Landf.* 23: 467–474.
- Alves M.A.S., 1997. Effects of ectoparasites on the Sand Martin *Riparia riparia* nestlings. *Ibis*. 139: 494–496.
- Bachmann S., Haller B., Lötscher R., Rehsteiner U., Spaar R. & Voge C., 2008. Guide de promotion de l’hirondelle de rivage en Suisse. Conseils pratiques pour la gestion des colonies dans les carrières et la construction de parois de nidification. Fondation Paysage et Gravier, Uttigen, Association Suisse de l’industrie des Graviers et du Béton, Berne, Association Suisse pour la Protection des Oiseaux ASPO/BirdLife Suisse, Cudrefin, Station ornithologique suisse, Sempach.



- 353 Baillie J., Hilton-Taylor C. & Stuart S.N., 2004. 2004 IUCN red list of threatened  
 354 species: a global species assessment Gland, Switzerland: IUCN, pp. 191.
- 355 BirdLife International, 2016. *Riparia riparia*. The IUCN Red List of  
 356 Threatened Species 2016: e.T103815961A87438023.  
 357 [http://dx.doi.org/10.2305/IUCN.UK.2016-](http://dx.doi.org/10.2305/IUCN.UK.2016-3.RLTS.T103815961A87438023.en)  
 358 [3.RLTS.T103815961A87438023.en](http://dx.doi.org/10.2305/IUCN.UK.2016-3.RLTS.T103815961A87438023.en). Downloaded on 05 November 2019.
- 359 BirdLife International, 2017. European birds of conservation concern: populations,  
 360 trends and national responsibilities. BirdLife International, Cambridge.
- 361 Brichetti P. & Fracasso G., 2007. Ornitologia italiana. Vol. 4 Apodidae –  
 362 Prunellidae. Alberto Perdisa Editore. Bologna.
- 363 Brookes A., 1988. Channelized Rivers: Perspectives for Environmental  
 364 Management. Wiley, Chichester.
- 365 Campedelli T., Buvoli L., Bonazzi P., Calabrese L., Calvi G., Celada C., Cutini S.,  
 366 De Carli E., Fornasari L., Fulco E., La Gioia G., Londi G., Rossi P., Silva L. & Tellini  
 367 Florenzano G., 2012. Andamenti di popolazione delle specie comuni nidificanti in  
 368 Italia: 2000-2011. Avocetta 36: 121–143.
- 369 Castiglioni G.B., Biancotti A., Bondesan M., Cortemiglia G.C., Elmi C., Favero V.,  
 370 Gasperi G., Marchetti G., Orombelli G., Pellegrini G.B. & Tellini C., 1999.  
 371 Geomorphological map of the Po Plain, Italy, at a scale of 1:250 000. Earth Surf.  
 372 Process. Landf. 24: 115–1120.
- 373 Cowley E., 1979. Sand Martin population trends in Britain, 1965–1978. Bird Study  
 374 26: 113–116.
- 375 Cowley E. & Siriwardena G.M., 2005. Long-term variation in survival rates of Sand  
 376 Martins *Riparia riparia*: dependance on breeding and wintering ground weather, age

- and sex, and their population consequences. *Bird Study* 52: 237–251.
- Cramp S., 1988. *Handbook of the Birds of Europe, the Middle East and North Africa: the birds of the Western Palearctic. Volume V. Tyrant Flycatchers to Thrushes.* Oxford University Press, Oxford.
- Curry-Lindahl K., 1972. *Conservation for survival. An ecological strategy.* William Morrow, London.
- Down C.G. & Stocks J., 1978. *Environmental impact of mining.* Applied Science Publishers, London.
- Evan A.T., Flamant C., Lavaysse C., Kocha C. & Saci A., 2014. Water Vapor–Forced Greenhouse Warming over the Sahara Desert and the Recent Recovery from the Sahelian Drought. *Journal of Climate*, 28, 108–123.
- Fehér J., Gáspár J., Veres K.S., Kiss A., Globevnik L., Peterlin M., Kirn T., Stein U., Prins T., Spiteri C., Laukkonen E., Heiskanen A.-S., Austner K., Semerádova S. & Künitzer A., 2012. Hydromorphological alterations and pressures in European rivers, lakes, transitional and coastal waters: thematic assessment for EEA Water 2012 report. VITUKI, NIVA, IWRS, Ecologic, Ecologic Institute, Stichting Deltares, SYKE, CENIA, Prague.
- Figarski T. & Kajtoch Ł., 2015. Alterations of riverine ecosystems adversely affect bird assemblages. *Hydrobiologia* 744: 287–296.
- Friend P.F. & Sinha R., 1993. Braiding and meandering parameters. *Geol. Soc. London Spec. Publ.* 75: 105–111.
- Garrison B.A., 1999. Bank Swallow (*Riparia riparia*). In A. Poole, F. Gill (Eds.), *The birds of North America*, 414, Inc., Philadelphia.
- Girvetz E.H., 2010. Removing erosion control projects increases bank swallow

- 401 (*Riparia riparia*) population viability modeled along the Sacramento River, California,  
 402 USA. Biol. Conserv. 143: 828–838.
- 403 Gulickx M.M.C., Beecroft R. & Green A., 2007. Creation of artificial Sand Martin  
 404 *Riparia riparia* burrows at Kingfishers Bridge. Conservation Evidence. 4: 51–53.
- 405 Gunn J. & Bailey D., 1993. Limestone quarrying and quarry reclamation in Britain.  
 406 Environ. Geol. 21: 167–172.
- 407 Hastie T. & Tibshirani R., 1990. Generalized Additive Models. Chapman and Hall,  
 408 New York.
- 409 Haywood S.M., 1979. Mineral landscapes: The next ten years? Environ. Geochem.  
 410 Health 1: 25–30.
- 411 Heneberg P., 2007. Sand martin (*Riparia riparia*) in the Czech Republic at the turn  
 412 of the millenium. Linzer biol. Beitr. 39: 293–312.
- 413 del Hoyo J., Elliott A. & Christie D.A., 2004. Handbook of the birds of the world,  
 414 Volume 9. Cotingas to pipits and wagtails. Lynx Editions, Barcelona.
- 415 Jankowiak Ł & Ławicki Ł, 2014. Marginal habitats as important refugia for riparian  
 416 birds during flood years. Bird Study. 61: 125–129.
- 417 Jones G., 1987a. Selection against large size in the Sand Martin. Ibis. 129: 274–280.
- 418 Jones G., 1987b. Selection against large size in the Sand Martin *Riparia riparia*  
 419 during a dramatic population crash. Ibis. 129: 274–280.
- 420 Kajtoch Ł & Figarski T, 2013. Short-term restoration of riverine bird assemblages  
 421 after a severe flood. Bird Study. 60: 327–334.
- 422 Kitowski I., Komosa A. & Chodorowski J., 2015. Absorbed radiation dose from  
 423 radon to the Sand Martin *Riparia riparia* during breeding at the sand mines in Eastern  
 424 Poland. Int. J. Environ. Res. 9: 1097–1106.

- 425 Krištofík J., Šustek Z. & Gajdoš P., 1994. Arthropods in nests of the Sand Martin  
 426 (*Riparia riparia* Linnaeus, 1758) in south Slovakia. Biol. Bratislava 49: 683–690.
- 427 Legwaila I.A., Lange E. & Cripps J., 2015. Quarry Reclamation in England: a  
 428 Review of techniques. Jasmr 4: 55–79.
- 429 Masoero G., Tamietti A., Boano G. & Caprio E., 2016. Apparent constant adult  
 430 survival of a Sand Martin *Riparia riparia* population in relation to climatic variables.  
 431 Ardea 104: 253–262.
- 432 Milgrom T., 2008. Environmental aspects of rehabilitating abandoned quarries: Israel  
 433 as a case study. Landsc. Urban Plan. 87: 172–179.
- 434 Moffatt K.C., Crone E.E., Holl K.D., Schlorff R.W. & Garrison B.A., 2005.  
 435 Importance of hydrologic and landscape heterogeneity for restoring bank swallow  
 436 (*Riparia riparia*) colonies along the Sacramento River, California. Restor. Ecol. 13:  
 437 391–402.
- 438 Morgan R.A., 1979. Sand Martin nest record cards. Bird Study 26: 129–132.
- 439 Muzzi E. & Mongardi G., 2016. Comparison of revegetation techniques on mineral  
 440 clay soil: analysis of quantitative response of vegetation cover. Ital. J. Agron. 11: 164–  
 441 170.
- 442 Newton I., 1998. Population Limitation in Birds. Academic Press, London.
- 443 Norman D. & Peach W.J., 2013. Density-dependent survival and recruitment in a  
 444 long-distance Palaearctic migrant, the Sand Martin *Riparia riparia*. Ibis. 155: 284–296.
- 445 Pannekoek J. & Van Strien, A., 2006. TRIM version 3.54 (Trends & Indices for  
 446 Monitoring data). Voorburg: Statistics Netherlands.
- 447 Persson C., 1987a. Population processes in south-west Scanian Sand martins  
 448 (*Riparia riparia*). J. Zool. 1: 671–691.

- 449 Persson C., 1987b. Age structure, sex ratios and survival rates in a south Swedish  
450 Sand martin (*Riparia riparia*) population, 1964 to 1984. J. Zool. 1: 639–670.
- 451 R Core Team, 2017. R: A language and environment for statistical computing. R  
452 Foundation for Statistical Computing. Available at: <https://www.r-project.org/>
- 453 Sanogo S., Fink A.H., Omotosho J.A., Ba A., Redl R. & Ermert V., 2015. Spatio-  
454 temporal characteristics of the recent rainfall recovery in West Africa. Int. J. Climatol.  
455 35: 4589–4605.
- 456 Svensson S., 1986. Number of pairs, timing of egg-laying and clutch size in a  
457 subalpine Sand Martin *Riparia riparia* colony, 1968-1985. Ornis Scand. 17: 221–229.
- 458 Szép T., 1995. Survival rates of Hungarian Sand Martins and their relationship with  
459 Sahel rainfall. J. Appl. Stat. 22: 891–904.
- 460 Szép T. & Møller A.P., 1999. Cost of parasitism and host immune defence in the  
461 Sand Martin *Riparia riparia*: A role for parent-offspring conflict? Oecologia 119: 9–15.
- 462 Szép T., Szabó D.Z. & Vallner J., 2003. Integrated population monitoring of Sand  
463 Martin *Riparia riparia* - an opportunity to monitor the effects of environmental  
464 disasters along the river Tisza. Ornis Hungarica 12: 169–182.
- 465 Wood S.N., 2011. Fast stable restricted maximum likelihood and marginal likelihood  
466 estimation of semiparametric generalized linear models. J. R. Stat. Soc. Ser. B 73: 3–36.
- 467 Yu G.-A., Disse M., Tong L. & Yu Y., 2015. Evolution of channel networks and  
468 morphology of a dryland River under human impacts – A case from Tarim River in  
469 Northwest China. In E-proceedings of the 36th IAHR World Congress, 28 June - 3 July  
470 p 11. The Hague, the Netherlands.
- 471 Zuquette L.V., Colares O.J. & Pejon O.J., 2002. Environmental degradation related  
472 to human activities, Fortaleza metropolitan region, State of Ceará, Brazil. Bull. Eng.

473 Geol. Environ. 61: 241–251.

474

475

476

## Figure captions

Fig. 1

Location of the study area, showing the River Po, and the active (circles) and abandoned (squares) quarries.

Fig. 2

Breeding sites of Sand Martins in the quarries.

Fig. 3

Changes course of the River Po from 1922 to 2008. Size of lines is for presentational purposes only (i.e. it is not related to channel width).

Fig. 4

Number of nests per year from 1970 to 2016 counted during the censuses of the breeding colonies in riverbanks (light grey) and in quarries (dark grey) in relation to the amount of sand and gravel extracted from quarries from 1980 and 2016 (dashed) and to the trend of the sinuosity index of the River Po course from 1922 to 2008 (thick solid line). The sinuosity index was calculated for the River Po courses in the years 1922, 1948, 1955, 1963, 1978, 1991 and 2008. The solid lines (light grey for riverbanks and dark for quarries) are smooths ( $\text{edf} = 7.06$  and  $8.11$  respectively) fitted from a GAM specifying a Poisson data distribution, and broken lines are the 95% confidence interval around the smoothed trend. Note that the census of the colonies with an exhaustive count of all active nests started in 2000.

501 Fig. 5 (a) Millions of  $\text{m}^3$  (Mm<sup>3</sup>) of extracted construction aggregates in Piedmont from  
502 1960 to 1985 (source: Regione Piemonte - Documento di programmazione delle attività  
503 estrattive I Stralcio – Relazione -  
504 [http://www.regione.piemonte.it/industria/cave/dpae\\_1.htm](http://www.regione.piemonte.it/industria/cave/dpae_1.htm) accessed on 27 October  
505 2016).

506 (b) Thousands of  $\text{m}^3$  of extracted construction aggregates in the 15 quarries of the  
507 study area from 1982 to 2016 (data from the Piemonte Region's database).