

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Optimization and validation of a method based on QuEChERS extraction and liquid chromatographic-tandem mass spectrometric analysis for the determination of perfluoroalkyl acids in strawberry and olive fruits, as model crops with different matrix characteristics

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1740724> since 2020-06-09T10:18:07Z

Published version:

DOI:10.1016/j.chroma.2020.461038

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 **Optimization and validation of a method based on QuEChERS extraction and liquid**
2 **chromatographic-tandem mass spectrometric analysis for the determination of perfluoroalkyl**
3 **acids in strawberry and olive fruits, as model crops with different matrix characteristics**

4

5 **Cristina Vanessa Agata Scordo^a, Leonardo Checchini^a, Lapo Renai^a, Serena Orlandini^a, Maria**
6 **Concetta Bruzzoniti^b, Donatella Fibbi^c, Laila Mandi^d, Naaila Ouazzani^d, Massimo Del Bubba^a**

7

8 ^a Department of Chemistry “Ugo Schiff”, University of Florence, Via della Lastruccia 3 – 50019
9 Sesto Fiorentino, Florence, Italy

10 ^b Department of Chemistry, University of Turin, Via Pietro Giuria 5 – 10125 Turin, Italy

11 ^c GIDA S.p.A., Via di Baciacavallo 36, 59100 Prato, Italy

12 ^d National Center for Research and Studies on Water and Energy (CNEREE), Cadi Ayyad University,
13 Marrakech, Morocco

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35 **Abstract**

36 A QuEChERS method was optimized and validated for the LC-MS/MS analysis of perfluoro-n-
37 pentanoic acid (PFPeA), perfluoro-1-butanesulfonic acid (PFBuS), perfluoro-n-hexanoic acid
38 (PFHxA), perfluoro-n-heptanoic acid (PFHpA), perfluoro-1-hexanesulfonic acid (PFHxS),
39 perfluoro-n-octanoic acid (PFOA), perfluoro-n-nonanoic acid (PFNA), perfluoro-1-octanesulfonic
40 acid (PFOS) and perfluoro-n-decanoic acid (PFDA) in freeze-dried strawberry and olive, as model
41 fruits characterized by very different chemical compositions. The method was evaluated for apparent
42 recovery, intra-day and inter-day precision, matrix effect and recovery. The method optimized for
43 strawberry provided for most compounds absolute values of matrix effect ($|ME\%| \leq 11\%$), except for
44 PFHxA, which showed a signal suppression of 22%. The extraction efficiency was tested at the spike
45 levels 500-5000 pg/g d.w. for PFPeA, PFBuS, and PFHxA, and 100-1000 pg/g d.w. for the other
46 target analytes, evidencing as a whole recoveries in the range of 65-89%. For olive fruits, due to their
47 high fat content, an ultrasound-assisted extraction was necessary to obtain an efficient sample
48 disgregation so as to increase the extraction yield and its precision. Moreover, a d-SPE clean-up with
49 GCB allowed to achieve $|ME\%| \leq 8\%$ (except for PFBuS, which showed a signal enhancement of
50 19%) and recoveries calculated at the aforementioned spike levels were in the range 75-97%. The
51 two methods provided very good linearity ($R^2 \geq 0.9984$) from 10000 pg/g down to compound specific
52 quantification limits, which were included in the ranges of 2.9-393 pg/g and 2.6-127 pg/g for
53 strawberry and olive fruit, respectively. The methods were applied to the analysis of PFAAs in
54 strawberry and olive fruits commercially available in two Italian supermarkets, as well as obtained
55 under irrigation with various treated wastewaters (TWWs), evidencing in both cases a higher PFAAs
56 occurrence in olives than in strawberry. However, PFAAs concentrations determined in the
57 investigated fruit matrixes were quite low, being their sum 1.9 ng/g d.w. in the worst case (i.e. olive
58 fruits grown under irrigation with TWWs).

59

60

61

62

63

64 **Keywords:** LC-MS/MS; d-SPE; perfluorinated compounds; fruit safety; organic micropollutants;
65 irrigation by wastewater.

66 **1 Introduction**

67 Food safety is a very important and topical issue, since foodstuffs can be contaminated through
68 different ways, alongside production and distribution stages [1]. This contamination does not only
69 concern chemical substances intentionally used within the food production chain, such as pesticides
70 in agriculture. In fact, foodstuffs can come into contact with various environmental micropollutants
71 before human consumption [2].

72 Among organic micropollutants, those most recently identified in environmental matrices and/or
73 recognized as environmentally hazardous, are obviously of major concern in the scientific community
74 and have been included in the so-called group of "emerging contaminants" (ECs). ECs comprise a
75 large group of chemicals, such as pharmaceuticals, personal care products, and also perfluoroalkyl
76 acids (PFAAs) [3, 4], the latter widely used since 1960 for the industrial production of various
77 consumer products [5-8]. The large industrial use of PFAAs derives from their peculiar
78 physicochemical properties. In fact, PFAAs are synthetic chemicals in which carbon-hydrogen bonds
79 are replaced by carbon-fluorine bonds, which is known as one of the strongest linkage in organic
80 chemistry and entails extremely stable at high temperatures (typically up to 150°C) [9]. Moreover,
81 they are neither flammable nor degradable under strong acidic or basic conditions, resistant to
82 oxidizing agents and photolysis and not subjected to biodegradation processes [10, 11]. As a
83 consequence of their large use, as well as their chemical and biological recalcitrance, PFAAs are
84 characterized by an incomplete removal in wastewater treatment plants (WWTPs) and a wide
85 occurrence in different water ecosystems have been reported [12-14].

86 Results of toxicological studies on animals have indicated that two of the most common PFCs,
87 perfluorooctanesulfonate (PFOS) and perfluorooctanoic acid (PFOA), may affect fatty acid
88 metabolism and the reproductive system as well as induce adverse effects in liver and other tissues
89 [15, 16]. Furthermore, human biomonitoring data evidenced that PFAAs bind to proteins and, for
90 PFOA and PFOS, has been estimated a bio-elimination time from the human body of 3.8 and 5.4
91 years, respectively [17-21]. These characteristics have led the European Food Safety Authority
92 (EFSA) to establish maximum tolerable weekly intakes (TWIs) of 13 ng/kg body weight for PFOS
93 and 6 ng/kg body weight for PFOA [22]. Moreover, PFOS has been added to the list of persistent
94 organic pollutants (POPs) under the Stockholm Convention on POPs [23].

95 It should also be remarked that physicochemical properties (e.g. lipophilicity and acidity) of these
96 compounds greatly vary depending on the chain length and the acidic group present in the molecule.
97 Based on these considerations, PFAAs represent an interesting group of model micropollutants to be
98 investigated for their possible transfer to crops. Moreover, PFAAs have been found at low

99 concentrations in crops [24, 25], and therefore very sensitive methods are needed for their
100 determination, thus highlighting the challenging character of this issue in analytical chemistry.

101 Few studies regarding crop contamination by PFAAs are reported in literature [24-30], some of them
102 using the QuEChERS approach for the extraction of target analytes [25, 29]. In these works, however,
103 the entire analytical procedure (i.e. extraction and clean-up) is applied without evaluating the
104 influence of the two individual steps on matrix effect (ME). In this way, it is not clear whether the
105 clean-up procedure leads to an effective advantage in terms of reduction of the matrix effect (ME),
106 the investigation of which, on the other hand, is overlooked in most cases [24, 25, 28-30]. It should
107 also be noted that the only methodological work concerning PFAAs determination in food, focuses
108 on the analysis of a wide range of products (e.g. fruit, vegetables and cheese), proposing for all the
109 matrices investigated the same complex clean-up procedures [27]. However, such a procedure,
110 necessary for fatty foods, might be superfluous in the case of simple matrices with low or null fat
111 content.

112 Among crops that could be adopted as models for investigating the contamination of fruits, strawberry
113 (*Fragaria x ananassa*) and olive (*Olea europaea* L.) are surely very interesting, due to their
114 remarkably different matrix characteristics. In fact, strawberry is characterized by a very high water
115 content, which accounts for 85-90% irrespective of the cultivar considered [31]. Conversely, olive is
116 one of the most lipophilic fruit, being it rich in fats, with a water content approximately included
117 between 50% and 70%, depending on the cultivar considered [32]. Hence, olive fruit is very
118 interesting to be investigated for the transfer of hydrophobic compounds, such as longer chain
119 perfluorocarboxylic acids. In this regard, according to literature, the cultivars showing among the
120 highest fruit fat content and corresponding low water percentage were “Frantoio” (<50%), “Picual”
121 and “Koroneiki” (about 50%) [32].

122 Based on the considerations reported above, the aim of this study was to develop a rapid, sensitive
123 and green method for the determination of nine PFAAs in strawberry and olive fruits using the
124 QuEChERS extraction and the dispersive solid-phase extraction (d-SPE) as clean-up approach,
125 followed by liquid chromatography coupled to tandem mass spectrometry (LC-MS/MS). The
126 optimized method was applied to the evaluation of PFAA occurrence in: (i) strawberry and olive
127 fruits commercially available in Italian supermarkets, and (ii) in strawberry (cultivar “Camarosa”)
128 and olive fruits (cultivar “Frantoio”) obtained under irrigation with various treated wastewaters
129 (TWWs). In fact, even though the wastewater reuse for irrigation is an efficient tool of reducing water
130 shortage and it is widely adopted in various Mediterranean countries [33], the presence of PFAAs in
131 TWWs [12, 34, 35] could negatively affect human health, owing to their possible transfer from
132 recycled water to crops.

133 2 Materials and methods

134 2.1 Chemicals and materials

135 Perfluoro-n-pentanoic acid (PFPeA, CAS 2706-90-3), perfluoro-1-butanesulfonic acid (PFBuS, CAS
136 375-73-5), perfluoro-n-hexanoic acid (PFHxA, CAS 307-24-4), perfluoro-n-heptanoic acid (PFHpA,
137 CAS 375-85-9), perfluoro-1-hexanesulfonic acid (PFHxS, CAS 355-46-4), perfluoro-n-octanoic acid
138 (PFOA, CAS 335-67-1), perfluoro-n-nonanoic acid (PFNA, CAS 375-95-1), perfluoro-1-
139 octanesulfonic acid (PFOS, CAS 1763-23-1) and perfluoro-n-decanoic acid (PFDA, CAS 335-76-2)
140 were purchased from Wellington Laboratories Inc. (Guelph, ON, Canada) as methanolic solutions,
141 each one at the concentration of 50 µg/mL. The ¹³C mass-labelled PFAA mixture containing the
142 aforementioned compounds, each one at 2 µg/mL in methanol, was purchased from Wellington
143 Laboratories Inc.

144 Sodium chloride (purity ≥ 99.0%) and anhydrous magnesium sulphate (purity ≥ 99.5%) used for
145 QuEChERS extraction were obtained from Sigma-Aldrich (St. Louis, MO, USA). Water, methanol
146 and acetonitrile (LC-MS grade) used for LC analyses and/or QuEChERS extraction, were obtained
147 from Carlo Erba (Milan, Italy). Ammonium acetate and formic acid (MS grade, purity ≥ 99.0%)
148 respectively used for the preparation of the LC eluents and the acidification of the extracts, were
149 purchased from Sigma-Aldrich.

150 Endcapped octadecylsilica (C18), primary secondary amine (PSA) and graphitized carbon black
151 (GCB) used for d-SPE clean-up of the QuEChERS extracts were purchased from Agilent
152 Technologies (Santa Clara, CA, USA). Minisart[®] regenerated cellulose syringe filters, used for the
153 filtration at 0.2 µm of the QuEChERS extracts, were supplied by Sartorius (Gottinga, Germany).

154 2.2 Samples

155 The strawberry (cultivar “Camarosa”) and olive (cultivar “Frantoio”) fruit samples used for the
156 optimization and validation of the analytical method were collected within an experimentation
157 conducted at the scientific campus of the University of Florence (Italy) and aimed at testing the
158 possible transfer of chemical contamination in fruits irrigated with four different types of TWWs and
159 tap water (TW) as control. More in detail, in this study, the effluents from the following WWTPs
160 located in the area of Prato (Italy), were used: (i) “Baciacavallo” activated sludge plant (TWW₁); (ii)
161 “Macrolotto 1” refining system of the Baciacavallo effluent (TWW₂); (iii) “Macrolotto 2” refining
162 system of the Baciacavallo effluent (TWW₃) and (iv) “Calice” activated sludge plant (TWW₄). Full
163 details of the treatment stages implemented in the four aforementioned WWTPs are reported in the
164 paragraph S.1 of the *Supplementary material* section. All the TWWs contained PFAAs at
165 concentrations ranging from a few to tens of ng/L.

166 Strawberry and olive fruits were harvested in July and November 2018, respectively. After collection,
167 the fruits were transported to the laboratory, immersed in liquid nitrogen, freeze-dried and finally
168 stored at -20 °C until they were analysed. Moreover, to test the applicability of the developed method,
169 commercially available strawberry and olive fruits were purchased in two supermarkets.

170 2.3 LC-MS/MS analysis

171 Separation of target analytes was performed on a Shimadzu (Kyoto, Japan) chromatographic system
172 consisting of two LC-20AD pumps (maximum allowed backpressure 600 bar), a SIL-30AC
173 autoinjector (maximum allowed backpressure 1000 bar) equipped with a 20 µL loop, a CTO/20AC
174 thermostatted column compartment and a CBM-20A module controller. A delay column ZORBAX
175 SB-C18 (Agilent Technologies, 4.6 x 75 mm, particle size 3.5 µm) was installed between the mixer
176 and the sample injector, in order to minimize the PFAA blank contributions due to the LC system
177 (i.e. solvents and/or tubing).

178 Chromatographic separation was obtained using a KINETEX XB-C18 column (100 x 3 mm, particle
179 size 2.6 µm) equipped with a guard column of the same phase (Phenomenex, Torrance, CA, USA),
180 thermostatted at 50°C. Eluent A: aqueous solution of 2 mM ammonium acetate. Eluent B: methanolic
181 solution of 2 mM ammonium acetate. A flow rate of 0.35 mL/min was adopted.
182 The chromatographic gradient was the following: 10% B for 2 min, from 10% to 90% in 9.20 min,
183 from 90% to 100% in 0.01 min, 100% for 6.79 min, from 100% to 10% in 2.50 min and final isocratic
184 10% for 5.5 min to allow system re-equilibration. Total analysis time (including system re-
185 equilibration) was 26 min and maximum backpressure 180 bar. The injection volume was set to 10
186 µL. In order to minimize MS source contamination, from 0 min to 8 min and after the elution of the
187 last peak until the start the system re-equilibration, the LC eluate was diverted to waste by means of
188 a Vici (Schenk, Switzerland) two-position six-port valve model HT (maximum allowed
189 backpressure 500 bar), installed upstream the mass spectrometer.

190 The Shimadzu LC system was coupled to a 5500 QTrap mass spectrometer (Sciex, Framingham, MA,
191 USA), by a Turbo V® interface equipped with an electrospray ionization (ESI) probe. MS/MS
192 analysis was carried out using a time-scheduled Multiple Reaction Monitoring (MRM) mode in
193 negative ionization. For each investigated compound, the most intense transition was used for
194 quantification and the second most intense, when present, for confirming the identification. All the
195 optimized MRM parameters are reported in **Table S1** of the *Supplementary material*.

196 Source dependent parameters were optimized in flow injection analysis at optimal LC flow and
197 mobile phase composition, and were as follows: Curtain Gas (CUR) 50, Collision Gas (CAD)
198 medium (8), Temperature (TEM) 650°C, Gas 1 (GS1) 50, Gas 2 (GS 2) 50, and IonSpray Voltage
199 (IS) -4500 V. Chromatograms were acquired by the 1.6.2 version of software Analyst (Sciex). Criteria

200 proposed by the Commission Decision 2002/657/CE were adopted for identity confirmation [36].
201 Peak attribution and quantitative determination were performed using MultiQuant software version
202 3.0.2 (ABSciex).

203 2.4 QuEChERS extraction and clean-up procedure

204 Polypropylene (PP) and polyethylene (PE) plastic ware was used for all the analytical steps. Under
205 the optimized experimental conditions, the following extraction and clean-up procedures were
206 adopted for strawberry and olive fruits, respectively.

207 2.4.1 *Strawberry*

208 One gram of freeze-dried fruit, which appears as a homogeneous powder, was weighed into a 50 mL
209 centrifuge tube and 5 mL of LC-MS water is added. The mixture was hand shaken for 15 s and vortex-
210 mixed for 1 min. Then, 10 mL of acetonitrile were added and the mixture was further hand shaken
211 for 15 s and vortex-mixed for 1 min. Afterwards, 2 g of NaCl and 2 g of MgSO₄ were added, and the
212 obtained mixture underwent to additional hand shaking (15 s) and vortex mixing (1 min). The tube
213 was centrifuged at 9000×g and T=4°C for 5 min. The acetonitrile supernatant is collected and finally
214 filtered through a 0.2 μm filter in a 15 mL graduated tube. Hence, the extraction procedure lasted
215 about 10 min. The filtered extract was acidified with formic acid (0.1% v/v) before LC-MS/MS
216 analysis.

217 2.4.2 *Olive*

218 One gram of freeze-dried fruit, which appears as an inhomogeneous oily solid, was weighed into a
219 50 mL centrifuge tube and 5 mL of LC-MS water is added. The mixture was hand shaken for 15 s
220 and vortex-mixed for 1 min. Then, 10 mL of acetonitrile were added and the mixture was further hand
221 shaken for 15 s, and sonicated for 90 sec (pulsed mode, 10 sec on and 5 sec off, power 750 W), using
222 a VC750 sonication probe (Sonics & Materials, Newtown, CT, USA). Afterwards, 2 g of NaCl and 2
223 g of MgSO₄ were added, and the obtained mixture underwent to additional hand shaking (15 s) and
224 vortex mixing (1 min). The tube was centrifuged at 9000×g and T=4°C for 5 min.

225 The acetonitrile supernatant was treated with 400 mg of GCB and 150 mg of MgSO₄ per mL of
226 extract. The resulting mixture was hand-shaken for 15 s, vortex mixed for 1 min, centrifuged at
227 9000×g and T=4°C for 5 min. and the recovered solvent filtered at 0.2 μm. The extract is finally made
228 up to the original volume underwent to the clean-up, using the acetonitrile of the procedural blank,
229 which is in our case the solvent that does not contain the matrix, brought through the entire analytical
230 procedure in the same manner as a test sample [37]. As a whole, the overall procedure lasted about
231 20 min. Before LC-MS/MS analysis, the filtered extract, was acidified to pH=2.50±0.05 with formic
232 acid (0.1% v/v)

233 2.5 Method performance evaluation and validation

234 In this work, the different method optimization steps were evaluated and validated by calculating the
235 performance parameters following reported. In this regard, taking into account that the QuEChERS
236 extracts, both as such and cleaned-up, contain salts which may affect the ionization of analytes,
237 performance evaluations of the various analytical optimization steps were performed, when
238 appropriate, in comparison with standard solutions prepared in the corresponding matrix-free
239 procedural blank.

240 2.5.1 Linearity of the methods

241 The analysis of strawberry and olive fruits is performed using two methods, which differ for the
242 presence of the d-SPE clean-up in the case of olives. For both matrixes, the method linearity was
243 assessed by spiking in triplicate 1 g d.w. aliquots with ¹³C labelled PFAA standards at the following
244 concentrations: 10, 50, 100, 500, 1000, 5000, and 10000 pg/g d.w. The spiked samples underwent to
245 the overall analytical protocol and finally analysed by LC-MS/MS after addition of formic acid 0.1%
246 (v/v).

247 2.5.2 Apparent recovery

248 The apparent recovery percentage (AR%) [38] of the methods were assessed, following the IUPAC
249 indications [39], by spiking in pentaplicate 1 g d.w. aliquots with ¹³C labelled PFAA standards at 500
250 (level 1) and 5000 (level 2) pg/g d.w. for PFPeA, PFBU_S, and PFHxA, and at 100 (level 1) and 1000
251 (level 2) pg/g d.w. for the other PFAAs (C_{spiked}). PFAAs concentrations in the spiked samples were
252 then quantified (C_{found}) using a standard calibration curve prepared in the matrix-free procedural blank
253 and AR% calculated according to the following equation.

$$254 \quad AR\% = \frac{C_{\text{found}}}{C_{\text{spiked}}} \cdot 100$$

255 Based on the procedure described above, apparent recovery takes into account the combined effect
256 of both matrix effect and reduced recovery during partition stages (e.g. between water and acetonitrile
257 or acetonitrile and d-SPE sorbent) due to the presence of matrix components [40].

258 2.5.3 Instrumental precision

259 Intra-day and inter-day instrumental precision were assessed by analysing from run-to-run over 1 and
260 5 days the final extracts deriving from the study of the apparent recovery in strawberry and olive
261 fruits, at level 1 concentrations.

262 2.5.4 Matrix effect

263 Matrix effect percentage (ME%) was defined as:

$$264 \quad ME\% = \left(\frac{S_{\text{matrix}}}{S_{\text{solvent}}} \cdot 100 \right) - 100$$

265 where S_{matrix} and S_{solvent} are the slopes of calibration lines in matrix and in procedural blank,
266 respectively [12]. Accordingly, ME% values higher or lower than zero indicate the presence of signal
267 enhancement or suppression in comparison with the instrumental response observed in procedural
268 blank. In fact, matrix components co-eluting with target analytes may alter their ionization, thus
269 affecting the sensitivity and accuracy of the method when real samples are analysed [40].

270 2.5.5 Recovery

271 The recovery percentage (R%) was defined as [41]:

$$272 \quad R\% = AR\% - ME\%$$

273 Standard deviation of R% ($\sigma_{R\%}$) was calculated according to error propagation rules

$$274 \quad \sigma_{R\%} = \sqrt{\sigma_{AR\%}^2 + \sigma_{ME\%}^2 + 2\sigma_{AR\%ME\%}}$$

275 where: $\sigma_{AR\%}$, $\sigma_{ME\%}$ are the experimental standard deviations of AR% and ME% and $\sigma_{AR\%ME\%}$ is the
276 covariance of the two variables.

277 2.5.6 Method detection and quantification limits

278 For each target PFAA, instrumental detection limit (IDL) and quantification limit (IQL) were
279 evaluated by replicated (n=5) analysis of procedural blanks, according to the following equation [42]:

$$280 \quad IQL \text{ (IDL)} = \frac{k \cdot \sigma_b}{S}$$

281 where: k is the critical value given by $2t_{(1-\alpha)}$ with n-1=4 degree of freedom and $\alpha=0.05$ (k= 4.264 and
282 k=10, for IDLs and IQLs, respectively); σ_b is the standard deviation of the procedural blank and S is
283 the slope of the calibration curve in procedural blank. The method detection (MDL) and quantification
284 (MQL) limits in fruit samples were obtained by multiplying IDL and IQL by the value of AR% and
285 dividing by 100.

286 2.5.7 Method accuracy and uncertainty

287 To the best of our knowledge, no reference material is available for the analysis of PFAAs in
288 strawberries and olives. Consequently, in accordance with the specifications of Eurachem [43], the
289 evaluation of the accuracy and uncertainty of the two methods was performed adopting for both
290 matrices the spiking procedure. TW samples, which showed PFAAs concentrations below MDLs or
291 MQLs, were chosen for performing the evaluation of method accuracy. Fruit aliquots of 1g d.w. were
292 spiked in pentaplicate with ^{13}C labelled and unlabelled PFAA standards at 500 pg/g d.w. for PFPeA,
293 PFBuS, and PFHxA, and at 100 pg/g d.w. for the other PFAAs. The spiked samples were then
294 extracted following the procedures described in paragraph 2.4 and analysed for labelled and
295 unlabelled spiked PFAAs using standard calibration curves prepared in the matrix-free procedural
296 blank. The concentrations determined in each spiked sample for labelled compounds allowed

297 calculating AR%, as described in paragraph 2.5.2. The concentrations of unlabelled PFAAs
298 determined through the standard calibration curve ($C_{\text{unlabelled}}$) were corrected for AR% using the
299 equation reported below, thus obtaining the actual concentration in each spiked samples (C_{actual}).

$$300 \quad C_{\text{actual}} = \frac{C_{\text{unlabelled}}}{\text{AR}\%} \cdot 100$$

301 Accuracy (Ac%) of the methods was finally calculated as the mean percentage ratio between C_{actual}
302 and C_{spiked} , determined in each spiked sample according to the following equation.

$$303 \quad \text{Ac}\% = \frac{C_{\text{actual}}}{C_{\text{spiked}}} \cdot 100$$

304 2.5.8 Analysis of real samples

305 Quantification of target analytes in real samples was performed by the isotope dilution method by
306 spiking 500 pg/g of ^{13}C labelled standards of each target analyte in 1 g d.w. aliquots of fruit samples
307 and keeping them in the dark at 4°C overnight before extraction, in order to calculate the AR% of the
308 whole analytical procedure. Quantification was performed by means of external calibration lines,
309 prepared in the corresponding procedural blank, taking into account the AR% of each analyte.

310 2.6 Statistical analysis

311 The plot of data, as well as of calibration lines and the evaluation of linearity parameters were
312 performed using Microsoft® Excel 2016 (Redmond, WA, USA). Analysis of variance and contrast
313 tests for comparisons of means were performed by the Games-Howell non-parametric test, using the
314 Minitab software packages version 17.0.1 (Minitab Inc., State College, PA, USA).

315 3 Results and discussion

316 PFAAs herein investigated are reported in **Table 1**, which provides compound abbreviation,
317 molecular structures, molecular masses, pKa, and values of the logarithm of the octanol-water
318 partition constants ($\log K_{\text{OW}}$). Target analytes included both sulfonic and carboxylic derivatives,
319 covering a wide range of polarity ($\log K_{\text{OW}}$ at pH=7 included from -1.81 to 4.15).

320 3.1 Optimization of the chromatographic conditions

321 Chromatographic analysis of PFAAs is usually affected by contaminations derived from eluents and
322 instrument parts made of fluorinated polymers (e.g. eluent lines, degassing apparatus, low pressure
323 mixing chamber). This problem is commonly faced by using the so-called “delay columns”,
324 containing a stationary phase able to strongly interact with PFAAs and installed between the eluent
325 mixer and the sample injector [12], thus allowing PFAAs released by the instrument and/or present
326 in the eluent to be delayed compared to those in the sample. As a drawback of this option, the
327 backpressure of the chromatographic system is more or less incremented, depending both on the
328 characteristics of the “delay” and of the analytical columns.

329 In this study, to reduce problems of contamination, LC pumps without any internal component made
330 of fluorinated polymers were used, all the eluent pipelines were in stainless steel and no on-line
331 degasser was employed. Moreover, an analytical pellicular column was used allowing to obtain
332 chromatographic performances comparable to those previously achieved using fully porous columns
333 with sub-2 μm particle size, but with much lower backpressure values, even though with an analysis
334 time about 6 min. longer [12]. Hence, it is possible to use high retentive trap columns of various
335 diameters and lengths packed with fully porous particles of relatively small size (e.g. 3.5 μm),
336 maintaining backpressures values below 200 bar, which translates in a lesser needing of maintenance
337 of LC system.

338 The LC elution gradient adopted in this study was properly optimized based on the conditions
339 elsewhere reported for the analysis of PFAAs in water samples [12]. Furthermore, the
340 chromatographic shape of the early-eluted PFAAs (i.e. PFBus and especially PFPeA) was strongly
341 influenced by the acidity of the injected samples. Accordingly, formic acid was added to the extracts
342 at 0.1% (v/v) to obtain narrower and quite symmetric peaks of PFPeA and PFBus (see **Figure 1**).
343 **Figure 2** illustrates a reconstructed LC-MS/MS MRM chromatogram of the quantifier and qualifier
344 transitions of target PFAAs obtained by injecting 10 μL of a standard solution (500 ng/L each) in
345 acetonitrile acidified with 0.1% (v/v) formic acid.

346 3.2 Optimization of the QuEChERS procedure

347 The QuEChERS approach generally involves two steps: (i) a water/acetonitrile salting-out
348 liquid/liquid partition of target compounds extracted from the solid matrix and (ii) a d-SPE for the
349 clean-up of the acetonitrile extract. Sonication can be also applied to enhance extraction efficiency
350 [44]. Within the general QuEChERS approach, these analytical steps must be properly optimized
351 considering the characteristics of both target analytes and matrix [45].

352 The QuEChERS method has been originally developed for the analysis of polar pesticides in fresh
353 fruit and vegetables [46, 47], which have a high water content. When dried samples are analysed,
354 their rehydration before QuEChERS extraction is therefore recommended for increasing the recovery
355 of target analytes [44]. The anhydrification of the samples and its subsequent rehydration with known
356 volumes of water should be preferred to the analysis of fresh samples. In fact, this approach allows
357 performing the analysis under standardized conditions from the point of view of the amount of water
358 present in the extraction mixture. It should be also noted that the quantity of water vary considerably
359 from fruit to fruit and within the same fruit, even from variety to variety [32]. In this latter regard, it
360 should be emphasized that for commodities with less than 80% of water content, the addition of an
361 extra-amount of water is important in order to weaken interactions of analytes with matrix and to
362 ensure their adequate partitioning [48]. The amount of rehydration water herein added to the freeze-

363 dried samples – i.e. 5 mL of water to 1 g d.w. of fruit – was chosen in order to restore the original
364 water percentage of strawberry (about 90%), obtaining at the same time a water excess in olive fruits
365 compared to its original content (about 50%).

366 3.2.1 Matrix-free partition tests

367 The development of the QuEChERS method involved preliminarily the evaluation of the water-to-
368 acetonitrile partition of target analytes. In particular, using 5 mL-aliquots of a standard solution of
369 PFAAs (500 ng/L each) in LC-MS water and 2 g of NaCl and MgSO₄, the extent of partition was
370 evaluated using 2.5, 5 and 10 mL of acetonitrile. The tests were carried out as fully described in the
371 *Supplementary material*. The results of these experiments highlighted that the variation of acetonitrile
372 volume did not significantly affect the recovery values (**Figure S1** of the *Supplementary material*).
373 Moreover, in order to evaluate the effect of water phase pH on the partition, 5 mL-aliquots of a PFAA
374 standard solution in LC-MS water as such (pH=6.57±0.05), as well as acidified with 0.1% formic
375 acid (v/v) (pH=2.50±0.05) were extracted with 10 mL of acetonitrile, following the aforementioned
376 procedure. The acidification gave rise to a slight recovery increase for the more polar
377 perfluorocarboxylates (e.g. PFPeA and PFHxA), which was however not statistically significant (data
378 not shown).

379 3.2.2 Strawberry fruits

380 The sample/acetonitrile ratio is a crucial parameter affecting the recovery of target analytes and an
381 excess of solvent compared to the amount of dried sample underwent to analysis is generally
382 suggested for improving the extraction efficiency [44]. Hence, the QuEChERS extraction procedure
383 described in the paragraph 2.4.1 was evaluated for AR% and ME%, using the aforementioned
384 different volumes of acetonitrile (i.e. sample/water/acetonitrile 1/5/2.5, 1/5/5 and 1/5/10 w/v/v),
385 keeping constant to 2 g each the amount of NaCl and MgSO₄ as salting-out agents. These tests were
386 performed by spiking 1 g-aliquots of fruits with 100 µL of a ¹³C-labelled PFAA working solution in
387 methanol containing each target analyte at 50 ng/mL and keeping them in the dark at 4°C overnight
388 before analysis.

389 **Figure 3** illustrates the results obtained for AR% and ME% in strawberry. The mean values of AR%
390 (**Fig. 3A**) showed an increasing trend with the increase of the acetonitrile volume. In particular, only
391 the use of 10 mL provided in almost all cases statistically higher recoveries compared to the lower
392 solvent volumes. These recoveries were in the ranges of 48-77% (2.5 mL), 57-78% (5 mL) and 67-
393 93% (10 mL) that are in most cases lower than those observed in the matrix-free partition tests. This
394 finding evidenced that the matrix exerted a significant influence on (i) the water-to-acetonitrile
395 partition of target analytes and/or (ii) their electrospray mass ionization, the latter being identified as
396 ME%. Accordingly, ME% was evaluated obtaining the results reported in **Fig. 3B**. In almost all cases,

397 a signal suppression due to the matrix was highlighted, being the only significant exception PFDA.
398 However, the extent of matrix suppression was compound-dependent and for PFBuS, PFOA, PFNA
399 PFOS and PFDA, $|ME| < 20\%$ were observed irrespective of the volume of acetonitrile used. In this
400 regard, it should be noted that signal enhancement/suppressions of 20% are commonly considered to
401 have a negligible influence on the performances of analytical methods [49-51]. PFPeA and PFHpA
402 showed $ME \leq 20\%$ with the use of 5-10 mL of acetonitrile. Conversely, for PFHxA and PFHxS, only
403 the use of 10 mL of acetonitrile allowed to decrease the signal suppression to values $\leq 20\%$, whereas
404 solvent volumes of 5 mL, and above all 2.5 mL, entailed suppressive ME% as high as 30-45%.
405 Accordingly, for strawberry, the sample/water/acetonitrile ratio of 1/5/10 *w/v/v* permits the direct
406 PFAA LC-MS/MS analysis of the QuEChERS extract without any clean-up process, thus increasing
407 the analytical throughput of the procedure (total analysis time of about 36 min.).

408 3.2.3 *Olive fruits*

409 As regards the analysis of olive fruits, the freeze-dried matrix appears as an inhomogeneous oily solid
410 that persists as such also after the rehydration process and the vortex extraction with acetonitrile.
411 Hence, this classic QuEChERS extraction was compared with an ultrasound-assisted procedure
412 (USAE-QuEChERS) consisting of the sonication of the sample for 90 sec (pulsed mode, 10 sec on
413 and 5 sec off, power 750 W) by the use of a sonication probe. Both QuEChERS and USAE-
414 QuEChERS gave rise to an oily and strongly pigmented extract (see **Figure S2** in the *Supplementary*
415 *material* section), even using 10 mL of acetonitrile, thus evidencing the need of a clean-up step before
416 LC-MS/MS analysis. This finding is in agreement with elsewhere reported recommendation for the
417 analysis of fruits characterized by high contents of fats, waxes and pigments, such as olives [48].
418 Preliminary matrix-free tests were performed in order to evaluate how PSA, C18 and GCB interact
419 with target analytes in the absence of matrix. These phases were selected as the ones commonly
420 employed in the clean-up step of QuEChERS protocols for the analysis of various food matrices [44],
421 including olive fruits [48]. Moreover, the use of PSA/C18 and PSA/GCB 1:1 (*w/w*) mixtures is
422 reported in official methods for pesticides residues analysis in foods [52]. In these procedures,
423 variable ratios between the volume of extract and the mass of sorbent have been used, depending on
424 the matrix and target analytes considered. In this regard, olive fruits can be considered a very complex
425 matrix, since the clean-up of olive oil extracts for PFAA analysis needed amounts of d-SPE sorbents
426 as high as 200 mg per mL of extract, even after a gel permeation pre-purification step [53].
427 The matrix-free tests were herein conducted using 10 mL-aliquots of a standard solution of PFAAs
428 (500 ng/L each) in acetonitrile (derived from the USAE-QuEChERS procedural blank), employing
429 150 mg of MgSO₄ and 300 mg of sorbent per mL of solvent. **Figure S3** of the *Supplementary material*
430 section illustrates the results of these tests. For most analytes, the use of PSA gave rise to recoveries

431 well below 50% (range 14-47%, median 37%). This finding is in accordance with the nature of anion
432 exchanger of PSA, and the specific indication of its use in d-SPE clean-up of QuEChERS extracts for
433 the removal of organic acids [52]. Conversely, C18 and GCB showed a very similar behaviour with
434 recoveries included in the ranges of 89-106% and 88-102%, respectively. Hence, both phases were
435 tested for the evaluation of ME%.

436 The optimization of the d-SPE clean-up was carried out by evaluating the extent of ME% in both
437 QuEChERS and USAE-QuEChERS approaches after the clean-up and then assessing the AR% of
438 the overall procedure. More in detail, the evaluation of ME% was performed by extracting 1 g of
439 freeze-dried matrix with 10 mL of acetonitrile and 2 g each of NaCl and MgSO₄, and treating the
440 resulting extract with 150 mg of MgSO₄, together with 200, 300 and 400 mg of C18 or GCB per mL
441 of extract. The cleaned-up extract was spiked with ¹³C-labelled PFAAs at 0, 50, 100, 500 and 1000
442 ng/L, comparing the slopes of these calibration lines in matrix with those in the procedural blank
443 prepared by vortex mixing and centrifuging LC-MS acetonitrile together with the d-SPE sorbent.

444 **Figure S4** and **Figure S5** of the *Supplementary material* illustrate the results of the d-SPE clean-up
445 using C18 (**Figs. S4A and S5A**) and GCB (**Figs. S4B and S5B**), on QuEChERS and USAE-
446 QuEChERS extracts, respectively. For both extraction approaches, the use of C18 showed |ME%| in
447 most cases higher than 20%, irrespective of the amount of sorbent used. Conversely, for GCB, a
448 decreasing trend with increasing the amount of the added sorbent was observed. This finding is
449 consistent with the fact that GCB is known to adsorb compounds which exhibit strong hydrophobic
450 interactions, such as pigments, which cannot be efficiently removed by C18 [53] and are probably
451 responsible for the residual matrix effect observed in **Fig. S4A and Fig. S5A**. In particular, the use
452 of 400 mg GCB allowed for achieving a strong reduction of the matrix effect, with |ME%| well below
453 20% for most analytes and in the worst cases of about +20% (i.e. for PFBuS with both procedures),
454 being them as a whole considered negligible. However, generally higher standard deviations were
455 observed for the QuEChERS method, compared to the USAE approach.

456 Based on the results of ME, the AR% was then evaluated for both QuEChERS and USAE-
457 QuEChERS procedures, using 400 mg of GCB per mL of extract for the d-SPE clean-up (see **Figure**
458 **4**). The USAE-QuEChERS procedure provided extraction efficiencies slightly higher than the
459 QuEChERS one and characterized by a much higher precision. These results were probably due to
460 the sample disgregation achieved by using the ultrasound probe, which allow for obtaining a
461 homogenous fine suspension of the sample in the extraction mixture.

462 Therefore, LC-MS/MS determination of PFAAs in olive fruits was performed by using the USAE-
463 QuEChERS procedure, with a total analysis time of about 50 min.

464 3.3 Method validation

465 After method optimization, a validation study was performed for both investigated matrixes to assess:
466 (i) linearity of the methods optimized for strawberry and olive fruits over proper spike ranges,
467 depending on the compound investigated; (ii) AR% at two spike levels; (iii) instrumental intra-day
468 and inter-day precision at two spike levels, expressed as relative standard deviation of peak area
469 (RSD%); (iv) ME% over proper linear calibration ranges, depending on the compound investigated;
470 (v) R% at two spike levels; (vi) MDLs and MQLs; (vii) Ac% at spike level 1. Note that different
471 spike levels were chosen depending on matrix and compound considered (**Table 2**). More in detail,
472 in strawberry, level 1 and level 2, were respectively chosen as 500 and 5000 pg/g d.w. for PFPeA,
473 PFBuS, and PFHxA, whereas for the other analytes they were 100 pg/g d.w. and 1000 pg/g d.w. In
474 olive, AR% of PFHxS, PFNA and PFDA was evaluated at 100 pg/g d.w. and 1000 pg/g d.w., whilst
475 for the other analytes the levels were 500 and 5000 pg/g d.w.

476 The method showed very high linearity, with determination coefficients in all cases ≥ 0.9984 . Intra-
477 day precision at level 1, as measured by RSD%, was in the ranges of 2.9-7.1 and 1.8-7.2 for strawberry
478 and olive fruits, respectively. At the level 2 the precision was slightly higher (RSD% of 2.1-6.1 for
479 strawberry and 1.3-5.0 for olive), in agreement with its higher concentration (one order of magnitude).
480 The inter-day RSD% were higher than the corresponding ones determined in the same day, remaining
481 however in most cases below 9%. The evaluation of ME% confirmed the absence of significant signal
482 suppression/enhancement in strawberry extracts, even without applying any clean-up process. It
483 should also be noted that the d-SPE treatment adopted for olive extract was able to reduce for almost
484 all analytes the |ME%| to values below 10%, with the only exception of PFBuS which showed a
485 mean amplification effect of 19%. Hence, for both fruits, the matrix matched calibration is not
486 necessary. The extraction procedures developed for the two matrices allowed not only to obtain small
487 alterations of the ESI ionization capacity due to the matrix (|ME%| < 20%), but also to efficiently
488 recover PFAAs from both fruits. The R% values, in fact, were generally high, being in any cases
489 >70%, with the only exceptions of PFHpA (65%) in the strawberry at the lowest fortification level
490 (100 pg/g d.w.). The higher recovery observed for olive fruit (75-97%) compared to strawberry (65-
491 89%) is probably due to the adoption of the USAE-QuEChERS. This procedure, in fact, led to the
492 sample disgregation, thus allowing to obtain a better recovery efficiency, which on average was
493 however limited to 10%. The proposed methods were also characterized by a very high sensitivity,
494 being MQLs in most cases included in between ppt and tens of ppt, with very few exceptions that
495 however showed values well below ppb levels. The evaluation of the overall Ac% of the methods
496 highlighted values included in the ranges of 99-109 and 98-114 for strawberry and olive, respectively.
497 The values higher than 100% found for most analytes in both matrices correspond to native
498 concentrations in TW samples in between ppt and tens of ppt, that are lower than MDLs or MQLs.

499 3.4 Comparison with previously published methods

500 The main characteristics of the method herein proposed can be compared with those reported in the
501 elsewhere published papers regarding method development for assessing fruit contamination by
502 PFAAs. However, as far as we know, no data have been published on PFAA determination in olive
503 or other “fatty fruits”, whilst very few papers are present in literature concerning PFAAs analysis in
504 fruits with high water content [25, 27-29] (**Table 3**). One of these researches focused on PFAAs
505 determination in strawberry [28], employing a method previously developed for vegetables [54]. In
506 this study a laborious extraction procedure was proposed, using a high percentage of dichloromethane
507 in the extraction mixture, thus providing a non-green analytical approach and making necessary
508 dryness evaporation in order to achieve compatibility with the complex clean-up process adopted and
509 LC-MS/MS instrumental analysis. As a whole, the analytical protocol lasted for about 8 hours and
510 provided quite low apparent recoveries (on average 45%), which could be ascribed to the complexity
511 of the method, while it is not possible to estimate the contribution of the matrix effect, since it has not
512 been investigated. Moreover, the sensitivity resulted fairly low, being MQLs in the range of 27-2910
513 pg/g w.w.

514 Much lower MQLs (1-4 pg/g w.w.) were reported by D'Hollander and co-workers [24] in a study
515 regarding strawberry and other water-rich fruits (e.g. orange, apple and melons). However, the
516 method envisaged a very high extraction time (about 16 h) and no information was reported regarding
517 method performances in terms of recovery and matrix effect.

518 Genualdi et al. [29] proposed a classical QuEChERS protocol for the analysis of PFAAs in cranberry
519 (*Vaccinium macrocarpon* Ait.), based on acidic water-to-acetonitrile partition of target analytes
520 followed by dispersive solid-phase extraction (d-SPE) with a sorbent mixture of PSA and GCB and
521 LC-MS/MS determination. The entire analytical procedure was quite rapid, since it lasted about 40
522 min. However, the method showed very high MDLs (240-2300 pg/g w.w. for the same PFAAs herein
523 analysed, corresponding to extrapolated MQLs in the range 792-7590 pg/g w.w., see **Table 3**), despite
524 the very high intrinsic sensitivity of the mass detector used and the high AR% obtained. In this latter
525 regard, it should be noted that the high recovery values reported have been obtained notwithstanding
526 the use of PSA, which was herein found not suitable for the d-SPE step due to its strong sorption
527 properties towards target analytes.

528 A more complex QuEChERS procedure was optimized by Sznajder-Katarzynska and co-workers [25]
529 for the determination of PFAAs in apple, on the basis of a method previously developed for the
530 analysis of PFOA and PFOS in honey [55]. Major differences compared to the classical QuEChERS
531 approach were the use of an ultrasound assisted partition of target compounds and a d-SPE clean-up

532 step with styrene-divinylbenzene. The optimized method achieved a sensitivity in the range 6-27 pg/g
533 w.w. with a total analysis time of about 70 min.

534 An analytical method aimed at determining PFAAs in very different kinds of food, including some
535 fruits, was developed by Ballesteros et al. [27], whose method was characterized by high apparent
536 recoveries, moderate matrix effect and MQLs included in the range 3.5-60 pg/g w.w. In order to make
537 the procedure suitable for a wide range of matrices, including fatty food, a complex clean-up strategy
538 was required, with a consequent increase of the total analysis time (about 2 hours). However, results
539 herein obtained showed that such a complex clean-up procedure is unnecessary in the case of
540 relatively simple matrices with low or null fat content such as strawberry, and also for fatty fruits
541 such as olives it seems oversized.

542 Based on the discussion reported above, the method herein proposed represented a general
543 improvement in terms of simplicity and total analysis time (36 min), as well as sensitivity in
544 comparison with previously published methods focusing on the determination of PFAAs in fruits.

545 3.5 Method application to real samples

546 The optimized methods have been applied to the analysis of strawberry and olive fruits purchased in
547 two Italian supermarkets, and obtained under controlled irrigation conditions with TW and different
548 TWWs (see paragraph 2.2). PFAAs concentrations found in these samples are shown in **Table 4**, as
549 mean and standard deviation of three independent determinations of each fruit sample. As an
550 example, **Figure 5** shows the overlapped quantifier and qualifier transitions of PFOA and PFOS in
551 marketed strawberry and olive fruits, compared to spiked procedural blanks.

552 As regards strawberry, in both commercial samples PFPeA and PFBuS were not detected. Moreover,
553 sample labelled as “Market 2” showed a higher number of detected analytes and generally higher
554 concentrations than “Market 1”. PFHxS was by far the most abundant compound determined in
555 commercially available strawberry, being it present at concentrations of 148 and 790 pg/g d.w., in
556 “Market 1” and “Market 2”, respectively. Regarding PFOA and PFOS, for which TWIs have been
557 provided by EFSA (6 and 13 ng/kg body weight), they were detected only in “Market 2” sample, at
558 27 and 90 pg/g d.w. These concentrations are low from the food intake point of view even assuming
559 a very high daily dose of strawberry, such as 500 g of fresh fruit, since they correspond to daily
560 ingestions approximately 30-40 times lower than the TWI for PFOS and PFOA, respectively. Only
561 one paper reported data regarding PFAAs occurrence in marketed strawberries, collected in Norway,
562 Belgium and Czech Republic [24]. In that study, PFAAs were found to be present in only one sample,
563 which evidenced the occurrence of PFBuS, PFHxA, PFHxS, PFOA and PFOS (58-192 pg/g w.w.)
564 and a total PFAAs concentration of 611 pg/g w.w., corresponding to about 6110 pg/g d.w. that is a
565 value higher than the ones herein found in fruits from Italian markets. Interestingly, strawberries

566 cultivated using TWWs did not show PFAAs concentrations above the quantification limits
567 irrespective of the kind of wastewater used for irrigation, suggesting that the contamination observed
568 in marketed strawberries could derive from contamination sources other than irrigation water. In this
569 regard, it must be considered that the absence of contamination or its very low extent have been
570 observed in the presence of irrigation with TWWs containing PFAAs at very different concentration
571 levels, ranging from few to hundreds of ng/L.

572 A different scenario was observed for PFAAs concentrations in olives (see **Table 4**), compared to
573 strawberries, since higher values were found in olive samples, both in the case of marketed products
574 and the ones obtained under irrigation with TWWs. In particular, both commercial samples showed
575 higher concentrations of PFPeA and PFBuS, compared to the other PFAAs. Moreover, “Market 2”
576 sample contained all the investigated PFAAs at concentrations well above the quantification limits,
577 with PFOA and PFOS at 124-137 and 39-95 pg/g d.w. These values, corresponding to about 60-70
578 and 20-50 pg/g w.w., even though higher than those previously reported for strawberry, can be
579 considered low in relation to TWIs established by EFSA for PFOA and PFOS, considering that olives
580 are a food much less consumed, compared to strawberry. Olive samples irrigated with the same
581 TWWs applied to strawberries showed a higher occurrence of PFAAs. More in detail, PFPeA and
582 PFHxA were the analytes determined at the highest concentrations, whereas PFOA and PFOS were
583 never quantified. The higher accumulation of PFAAs in olive fruits than in strawberry could be
584 related to their very different composition in terms of water and fat content.

585 **4 Conclusions**

586 The method optimized and validated in this research represents an improvement in terms of simplicity
587 and/or analysis time and/or sensitivity, compared to previously published analytical procedures
588 focusing on the determination of PFAAs in fruits. Moreover, the method herein proposed for PFAAs
589 analysis in olives represents the first validated analytical approach for this fruit matrix.

590 Since the analytical approach proposed has been validated on two types of fruit, which are very
591 different for their content of water and fatty substances, it is reasonable to assume that it can be
592 successfully applied to a wide range of fruit products.

593 The application of the proposed method to the analysis of real samples evidenced that fruits
594 commercially available at supermarkets contained concentrations generally higher than those grown
595 under controlled watering conditions, using tap water but also TWWs for their irrigation, thus
596 suggesting that commercial crops can be exposed to a wide range of PFAAs contamination sources,
597 in addition to irrigation water. In any case, the quantities of PFOA and PFOS that can be ingested
598 through the analysed fruits, even taking into account much higher daily doses than those calculated

599 on the basis of surveys carried out by major international organizations, are lower than the TWIs
600 established by EFSA. However, considering that the intake of these compounds derive from the
601 ingestions of various foodstuffs, all of them possibly contaminated by PFOA and PFOS [22], the
602 concentrations herein determined could be considered noteworthy.

603 **Acknowledgements**

604 This research was supported by the grant 13-069 (IRRIGATIO project) under the ERANET MED
605 2014 call, which is gratefully acknowledged. Ministero dell'Istruzione, dell'Università e della Ricerca
606 (MIUR) is also acknowledged. Authors also wish to thank Susan Mary Cadby for her linguistic
607 revision of the paper.

608 **References**

- 609 [1] P.A. Luning, F. Devlieghere, Safety in the agri-food chain, Wageningen Academic Pub, 2006.
610 [2] L.A. Thompson, W.S. Darwish, Environmental Chemical Contaminants in Food: Review of a
611 Global Problem, *J Toxicol*, 2019 (2019) 2345283-2345283.
612 [3] M. Gavrilescu, K. Demnerová, J. Aamand, S. Agathos, F. Fava, Emerging pollutants in the
613 environment: present and future challenges in biomonitoring, ecological risks and bioremediation,
614 *New biotechnology*, 32 (2015) 147-156.
615 [4] Y. Luo, W. Guo, H.H. Ngo, L.D. Nghiem, F.I. Hai, J. Zhang, S. Liang, X.C. Wang, A review on
616 the occurrence of micropollutants in the aquatic environment and their fate and removal during
617 wastewater treatment, *Science of the total environment*, 473 (2014) 619-641.
618 [5] F. Ye, Y. Zushi, S. Masunaga, Survey of perfluoroalkyl acids (PFAAs) and their precursors
619 present in Japanese consumer products, *Chemosphere*, 127 (2015) 262-268.
620 [6] Z. Wang, I.T. Cousins, M. Scheringer, K. Hungerbuehler, Hazard assessment of fluorinated
621 alternatives to long-chain perfluoroalkyl acids (PFAAs) and their precursors: status quo, ongoing
622 challenges and possible solutions, *Environment international*, 75 (2015) 172-179.
623 [7] K. Prevedouros, I.T. Cousins, R.C. Buck, S.H. Korzeniowski, Sources, fate and transport of
624 perfluorocarboxylates, *Environmental science & technology*, 40 (2006) 32-44.
625 [8] R.C. Buck, J. Franklin, U. Berger, J.M. Conder, I.T. Cousins, P. De Voogt, A.A. Jensen, K.
626 Kannan, S.A. Mabury, S.P. van Leeuwen, Perfluoroalkyl and polyfluoroalkyl substances in the
627 environment: terminology, classification, and origins, *Integrated environmental assessment and
628 management*, 7 (2011) 513-541.
629 [9] C. Lau, J.L. Butenhoff, J.M. Rogers, The developmental toxicity of perfluoroalkyl acids and their
630 derivatives, *Toxicology and applied pharmacology*, 198 (2004) 231-241.
631 [10] R.C. Buck, P.M. Murphy, M. Pabon, Chemistry, properties, and uses of commercial fluorinated
632 surfactants, in: *Polyfluorinated chemicals and transformation products*, Springer, 2012, pp. 1-24.
633 [11] A.B. Lindstrom, M.J. Strynar, E.L. Libelo, *Polyfluorinated compounds: past, present, and future*,
634 in, ACS Publications, 2011.
635 [12] L. Ciofi, L. Renai, D. Rossini, C. Ancillotti, A. Falai, D. Fibbi, M.C. Bruzzoniti, J.J. Santana-
636 Rodriguez, S. Orlandini, M. Del Bubba, Applicability of the direct injection liquid chromatographic
637 tandem mass spectrometric analytical approach to the sub-ng L⁻¹ determination of perfluoro-alkyl
638 acids in waste, surface, ground and drinking water samples, *Talanta*, 176 (2018) 412-421.
639 [13] G.B. Post, P.D. Cohn, K.R. Cooper, Perfluorooctanoic acid (PFOA), an emerging drinking water
640 contaminant: a critical review of recent literature, *Environmental research*, 116 (2012) 93-117.
641 [14] J. Elmoznino, P. Vlahos, M. Whitney, Occurrence and partitioning behavior of perfluoroalkyl
642 acids in wastewater effluent discharging into the Long Island Sound, *Environmental pollution*, 243
643 (2018) 453-461.

644 [15] G.L. Kennedy, J.L. Butenhoff, G.W. Olsen, J.C. O'Connor, A.M. Seacat, R.G. Perkins, L.B.
645 Biegel, S.R. Murphy, D.G. Farrar, The toxicology of perfluorooctanoate, *Critical reviews in*
646 *toxicology*, 34 (2004) 351-384.

647 [16] A.M. Seacat, P.J. Thomford, K.J. Hansen, G.W. Olsen, M.T. Case, J.L. Butenhoff, Subchronic
648 toxicity studies on perfluorooctanesulfonate potassium salt in cynomolgus monkeys, *Toxicological*
649 *Sciences*, 68 (2002) 249-264.

650 [17] J. Li, F. Guo, Y. Wang, J. Liu, Z. Cai, J. Zhang, Y. Zhao, Y. Wu, Development of extraction
651 methods for the analysis of perfluorinated compounds in human hair and nail by high performance
652 liquid chromatography tandem mass spectrometry, *Journal of Chromatography A*, 1219 (2012) 54-
653 60.

654 [18] P.D. Jones, W. Hu, W. De Coen, J.L. Newsted, J.P. Giesy, Binding of perfluorinated fatty acids
655 to serum proteins, *Environmental toxicology and chemistry*, 22 (2003) 2639-2649.

656 [19] J. Liu, J. Li, Y. Liu, H.M. Chan, Y. Zhao, Z. Cai, Y. Wu, Comparison on gestation and lactation
657 exposure of perfluorinated compounds for newborns, *Environment international*, 37 (2011) 1206-
658 1212.

659 [20] J.P. Giesy, K. Kannan, Global distribution of perfluorooctane sulfonate in wildlife,
660 *Environmental science & technology*, 35 (2001) 1339-1342.

661 [21] G.W. Olsen, J.M. Burris, D.J. Ehresman, J.W. Froehlich, A.M. Seacat, J.L. Butenhoff, L.R.
662 Zobel, Half-life of serum elimination of perfluorooctanesulfonate, perfluorohexanesulfonate, and
663 perfluorooctanoate in retired fluorochemical production workers, *Environmental health perspectives*,
664 115 (2007) 1298.

665 [22] EFSA Panel on Contaminants in the Food Chain, Risk to human health related to the presence
666 of perfluorooctane sulfonic acid and perfluorooctanoic acid in food, *EFSA Journal*, 16 (2018) e05194.

667 [23] Conference of the Parties of the Stockholm Convention, Listing of perfluorooctane sulfonic acid,
668 its salts and perfluorooctane sulfonyl fluoride, in: *Decision SC-4/17*, 2009.

669 [24] W. D'Hollander, D. Herzke, S. Huber, J. Hajslova, J. Pulkrabova, G. Brambilla, S.P. De Filippis,
670 L. Bervoets, P. de Voogt, Occurrence of perfluorinated alkylated substances in cereals, salt, sweets
671 and fruit items collected in four European countries, *Chemosphere*, 129 (2015) 179-185.

672 [25] K. Sznajder-Katarzyńska, M. Surma, E. Cieślík, W. Wiczowski, The perfluoroalkyl substances
673 (PFASs) contamination of fruits and vegetables, *Food Additives & Contaminants: Part A*, 35 (2018)
674 1776-1786.

675 [26] I. Aparicio, J. Martín, C. Abril, J.L. Santos, E. Alonso, Determination of household and industrial
676 chemicals, personal care products and hormones in leafy and root vegetables by liquid
677 chromatography-tandem mass spectrometry, *Journal of Chromatography A*, 1533 (2018) 49-56.

678 [27] A. Ballesteros-Gómez, S. Rubio, S. van Leeuwen, Tetrahydrofuran–water extraction, in-line
679 clean-up and selective liquid chromatography/tandem mass spectrometry for the quantitation of
680 perfluorinated compounds in food at the low picogram per gram level, *Journal of Chromatography*
681 *A*, 1217 (2010) 5913-5921.

682 [28] A.C. Blaine, C.D. Rich, E.M. Sedlacko, K.C. Hyland, C. Stushnoff, E.R. Dickenson, C.P.
683 Higgins, Perfluoroalkyl acid uptake in lettuce (*Lactuca sativa*) and strawberry (*Fragaria ananassa*)
684 irrigated with reclaimed water, *Environmental science & technology*, 48 (2014) 14361-14368.

685 [29] S. Genualdi, N. Jeong, L. Dejager, T. Begley, Investigation into perfluoroalkyl substances
686 (PFASs) in a cranberry bog: method development and sampling results, *Food Additives &*
687 *Contaminants: Part A*, 34 (2017) 2181-2189.

688 [30] S. Lasee, S. Subbiah, W.A. Thompson, A. Karnjanapiboonwong, J. Jordan, P. Payton, T.A.
689 Anderson, Plant Uptake of Per- and Polyfluoroalkyl Acids under a Maximum Bioavailability
690 Scenario, *Environmental Toxicology and Chemistry*, 38 (2019) 2497-2502.

691 [31] L.H. Mosquera, G. Moraga, N. Martínez-Navarrete, Critical water activity and critical water
692 content of freeze-dried strawberry powder as affected by maltodextrin and arabic gum, *Food Research*
693 *International*, 47 (2012) 201-206.

694 [32] K. Zeleke, R. Mailer, P. Eberbach, J. Wünsche, Oil content and fruit quality of nine olive (*Olea*
695 *europaea* L.) varieties affected by irrigation and harvest times, *New Zealand journal of crop and*
696 *horticultural science*, 40 (2012) 241-252.

697 [33] L. Rivoira, M. Castiglioni, A. Kettab, N. Ouazzani, E. Al-Karablieh, N. Boujelben, D. Fibbi, E.
698 Coppini, E. Giordani, M. Del Bubba, Impact of Effluents from Wastewater Treatments Reused for
699 Irrigation: Strawberry as Case Study, *Environmental Engineering & Management Journal (EEMJ)*,
700 18 (2019).

701 [34] O.S. Arvaniti, A.S. Stasinakis, Review on the occurrence, fate and removal of perfluorinated
702 compounds during wastewater treatment, *Science of the Total Environment*, 524 (2015) 81-92.

703 [35] C. Gallen, G. Eaglesham, D. Drage, T.H. Nguyen, J. Mueller, A mass estimate of perfluoroalkyl
704 substance (PFAS) release from Australian wastewater treatment plants, *Chemosphere*, 208 (2018)
705 975-983.

706 [36] T.C.O.T.E. COMMUNITIES, Commission Decision of 12 August 2002 implementing Council
707 Directive 96/23/EC concerning the performance of analytical methods and the interpretation of results
708 (2002/657/EC). (2002).

709 [37] J. Inczedy, T. Lengyel, A.M. Ure, A. Gelencsér, A. Hulanicki, *Compendium of analytical*
710 *nomenclature*, The Orange Book, 3rd Edn., (1998).

711 [38] D.T. Burns, K. Danzer, A. Townshend, Use of the term "recovery" and "apparent recovery" in
712 analytical procedures (IUPAC Recommendations 2002), *Pure and applied chemistry*, 74 (2002) 2201-
713 2205.

714 [39] M. Thompson, S.L. Ellison, A. Fajgelj, P. Willetts, R. Wood, Harmonized guidelines for the use
715 of recovery information in analytical measurement, *Pure and applied chemistry*, 71 (1999) 337-348.

716 [40] L. Ciofi, C. Ancillotti, U. Chiuminatto, D. Fibbi, B. Pasquini, M.C. Bruzzoniti, L. Rivoira, M.
717 Del Bubba, Fully automated on-line solid phase extraction coupled to liquid chromatography–tandem
718 mass spectrometry for the simultaneous analysis of alkylphenol polyethoxylates and their carboxylic
719 and phenolic metabolites in wastewater samples, *Analytical and bioanalytical chemistry*, 408 (2016)
720 3331-3347.

721 [41] L. Ciofi, C. Ancillotti, U. Chiuminatto, D. Fibbi, L. Checchini, S. Orlandini, M. Del Bubba,
722 Liquid chromatographic–tandem mass spectrometric method for the simultaneous determination of
723 alkylphenols polyethoxylates, alkylphenoxy carboxylates and alkylphenols in wastewater and
724 surface-water, *Journal of Chromatography A*, 1362 (2014) 75-88.

725 [42] T. Wenzl, J. Haedrich, A. Schaechtele, P. Robouch, J. Stroka, *Guidance Document on the*
726 *Estimation of LOD and LOQ for Measurements in the Field of Contaminants in Feed and Food*,
727 *Publications Office of the European Union: Luxemburg*, (2016).

728 [43] B. Magnusson, *The fitness for purpose of analytical methods: a laboratory guide to method*
729 *validation and related topics* (2014), in, *Eurachem*, 2014.

730 [44] M.C. Bruzzoniti, L. Checchini, R.M. De Carlo, S. Orlandini, L. Rivoira, M. Del Bubba,
731 QuEChERS sample preparation for the determination of pesticides and other organic residues in
732 environmental matrices: a critical review, *Analytical and Bioanalytical Chemistry*, 406 (2014) 4089-
733 4116.

734 [45] R.M. De Carlo, L. Rivoira, L. Ciofi, C. Ancillotti, L. Checchini, M. Del Bubba, M.C. Bruzzoniti,
735 Evaluation of different QuEChERS procedures for the recovery of selected drugs and herbicides from
736 soil using LC coupled with UV and pulsed amperometry for their detection, *Analytical and*
737 *bioanalytical chemistry*, 407 (2015) 1217-1229.

738 [46] M. Anastassiades, S.J. Lehotay, D. Štajnbaher, F.J. Schenck, Fast and easy multiresidue method
739 employing acetonitrile extraction/partitioning and “dispersive solid-phase extraction” for the
740 determination of pesticide residues in produce, *Journal of AOAC international*, 86 (2003) 412-431.

741 [47] S.J. Lehotay, A.d. Kok, M. Hiemstra, P.v. Bodegraven, Validation of a fast and easy method for
742 the determination of residues from 229 pesticides in fruits and vegetables using gas and liquid
743 chromatography and mass spectrometric detection, *Journal of AOAC International*, 88 (2005) 595-
744 614.

745 [48] T. Rejczak, T. Tuzimski, A review of recent developments and trends in the QuEChERS sample
746 preparation approach, *Open Chemistry*, 13 (2015).

747 [49] V.C. Fernandes, V.F. Domingues, N. Mateus, C. Delerue-Matos, Multiresidue pesticides
748 analysis in soils using modified QuEChERS with disposable pipette extraction and dispersive solid-
749 phase extraction, *Journal of separation science*, 36 (2013) 376-382.

750 [50] A.G. Frenich, R. Romero-González, M.L. Gómez-Pérez, J.L.M. Vidal, Multi-mycotoxin analysis
751 in eggs using a QuEChERS-based extraction procedure and ultra-high-pressure liquid
752 chromatography coupled to triple quadrupole mass spectrometry, *Journal of Chromatography A*,
753 1218 (2011) 4349-4356.

754 [51] M. Mei, D. Zhen-Xia, C. Yun, QuEChERS-ultra-performance liquid chromatography tandem
755 mass spectrometry for determination of five currently used herbicides, *Chinese Journal of Analytical*
756 *Chemistry*, 39 (2011) 1659-1664.

757 [52] European Committee for Standardization. EN 15662:2018, Foods of plant origin - Multimethod
758 for the determination of pesticide residues using GC- and LC-based analysis following acetonitrile
759 extraction/partitioning and clean-up by dispersive SPE - Modular QuEChERS-method. , in, Brussel
760 (Belgium), 2018.

761 [53] L. Yang, F. Jin, P. Zhang, Y. Zhang, J. Wang, H. Shao, M. Jin, S. Wang, L. Zheng, J. Wang,
762 Simultaneous determination of perfluorinated compounds in edible oil by gel-permeation
763 chromatography combined with dispersive solid-phase extraction and liquid chromatography–tandem
764 mass spectrometry, *Journal of agricultural and food chemistry*, 63 (2015) 8364-8371.










765 [54] A.C. Blaine, C.D. Rich, L.S. Hundal, C. Lau, M.A. Mills, K.M. Harris, C.P. Higgins, Uptake of
766 perfluoroalkyl acids into edible crops via land applied biosolids: field and greenhouse studies,
767 *Environmental science & technology*, 47 (2013) 14062-14069.

768 [55] M. Surma, W. Wiczkowski, E. Cieřlik, H. Zieliński, Method development for the determination
769 of PFOA and PFOS in honey based on the dispersive Solid Phase Extraction (d-SPE) with micro-
770 UHPLC–MS/MS system, *Microchemical Journal*, 121 (2015) 150-156.

771

772

Table 1 – Abbreviations, molecular structures, molecular masses, pK_a* and log K_{ow}* values (calculated at pH=7) of perfluoroalkyl acids (PFAAs) investigated in this study.

PFAAs	Abbreviation	Molecular Structure	Molecular Mass	pK _a	log K _{ow}
Perfluoro-n-pentanoic	PFPeA		264.05	0.40	0.64
Perfluoro-1-butananesulfonic	PFBuS		300.10	-3.57	-1.81
Perfluoro-n-hexanoic	PFHxA		314.05	0.42	1.24
Perfluoro-n-heptanoic	PFHpA		364.06	0.47	1.97
Perfluoro-1-hexanesulfonic	PFHxS		400.11	-3.34	-0.45
Perfluoro-n-octanoic	PFOA		414.07	0.50	2.69
Perfluoro-n-nonanoic	PFNA		464.08	0.52	3.42
Perfluoro-1-octanesulfonic	PFOS		500.13	-3.27	1.01
Perfluoro-n-decanoic	PFDA		514.08	0.52	4.15

* Calculated using Advanced Chemistry Development (ACD/Labs) Software V11.02 (© 1994-2019 ACD/Labs)

Table 2 – Linearity of the method expressed as determination coefficient (R^2) calculated over proper spike ranges^(a), apparent recovery (AR%), instrumental intra-day and inter-day precisions (expressed as relative standard deviation of peak area, RSD %), matrix effect (ME%), recovery (R%), method detection (MDLs) and quantification (MQLs) limits (pg/g d.w.), accuracy (Ac%). For AR%, ME%, R%, and Ac%, values in bracket represent the standard deviation. AR%, RSD% and R% were calculated at two spike levels, whereas Ac% at the spike level 1^(b). MDLs and MQLs were calculated based on AR% of the lowest spike level (level 1).

Compound	R^2	AR%		RSD% ^{intra-day}		RSD% ^{inter-day}		ME%	R%		MDLs	MQLs	Ac%
		level 1	level 2	level 1	level 2	level 1	level 2		level 1	level 2			
Strawberry													
PFPeA	0.9992	73 (8)	74 (6)	7.1	5.4	10.9	7.6	-5 (4)	78 (11)	79 (11)	109	393	109 (9)
PFBuS	0.9987	85 (7)	76 (6)	5.9	5.3	9.8	6.9	-4 (5)	89 (12)	80 (8)	29	104	107 (7)
PFHxA	0.9984	59 (5)	63 (7)	4.3	4.1	7.3	5.5	-22 (3)	81 (9)	85 (9)	40	146	106 (5)
PFHpA	0.9993	74 (4)	89 (8)	3.7	2.9	9.4	7.1	9 (5)	65 (9)	80 (13)	7.2	26	103 (7)
PFHxS	0.9988	66 (6)	71 (2)	5.4	2.1	9.7	8.2	-7 (2)	73 (8)	78 (4)	10	37	103 (7)
PFOA	0.9989	64 (3)	70 (3)	2.9	3.1	7.9	6.5	-11 (3)	75 (5)	81 (5)	8.6	31	107 (5)
PFNA	0.9993	64 (4)	72 (5)	4.1	3.6	8.6	7.0	-8 (4)	72 (7)	80 (7)	1.3	4.9	106 (6)
PFOS	0.9985	76 (8)	81 (8)	6.4	6.1	9.2	8.1	2 (6)	74 (11)	79 (14)	5.7	20	108 (9)
PFDA	0.9988	66 (4)	67 (4)	3.8	3.3	6.1	4.5	-4 (3)	71 (6)	72 (6)	0.8	2.9	99 (5)
Olive													
PFPeA	0.9997	93 (4)	88 (2)	3.0	2.2	6.8	6.1	-4 (2)	97 (6)	92 (4)	35	127	110 (6)
PFBuS	0.9991	105 (9)	102 (3)	1.8	1.3	5.5	4.3	19 (3)	86 (11)	83 (4)	24	85	107 (7)
PFHxA	0.9994	82 (3)	84 (3)	5.4	2.2	8.6	4.8	-4 (4)	86 (7)	88 (7)	26	94	114 (7)
PFHpA	0.9992	82 (5)	93 (4)	7.2	5.0	8.2	7.1	4 (3)	78 (5)	89 (6)	21	74	98 (9)
PFHxS	0.9997	83 (5)	91 (2)	5.1	4.2	8.9	7.3	-6 (4)	89 (7)	97 (5)	2.4	8.6	102 (8)
PFOA	0.9997	67 (7)	84 (3)	3.4	3.0	6.7	5.4	-8 (3)	75 (10)	92 (7)	22	78	106 (5)
PFNA	0.9997	86 (6)	89 (6)	3.1	1.8	4.8	3.6	-3(2)	89 (7)	92 (8)	0.7	2.6	102 (6)
PFOS	0.9994	87 (8)	92 (4)	5.3	1.3	9.6	3.5	3 (3)	84 (9)	89 (5)	5.8	21	99 (7)
PFDA	0.9997	87 (4)	87 (4)	4.1	3.7	7.3	6.4	4 (3)	83 (6)	83 (6)	2.7	9.8	105 (7)

^(a) R^2 values were calculated in the following ranges. Strawberry – (i) PFNA and PFDA 10-10000 pg/g d.w.; (ii) PFHpA, PFHxS, PFOA and PFOS 50-10000 pg/g d.w.; (iii) PFBuS, PFPeA and PFHxA 500-10000 pg/g d.w. Olive – (i) PFHxS, PFNA and PFDA 10-10000 pg/g d.w.; (ii) PFBuS, PFHxA, PFHpA, PFOA and PFOS 50-10000 pg/g d.w.; (iii) PFPeA 500-10000 pg/g d.w.

^(b) Spike levels for PFPeA, PFBuS, and PFHxA were 500 pg/g d.w. (level 1) and 5000 pg/g d.w. (level 2), whereas for the other analytes were 100 pg/g d.w. (level 1) and 1000 pg/g d.w. (level 2).

Table 3 – Main characteristics of the analytical method herein proposed for the analysis of target PFAAs in strawberry, in comparison with those provided by elsewhere published studies in various fruit matrixes, using different extraction/clean-up methods followed by LC-MS/MS instrumental determination. AR%=apparent recovery percentage. ME% = matrix effect percentage; MQL = method quantification limit (pg/g wet weight); n.i. = not investigated; n.r. = not reported.

Matrix	Method details	PFAAs	AR%	ME%	MQL	Reference
Strawberry	QuEChERS without clean-up. Total analysis time 36 min.	PFPeA	73	-5	39	This study
		PFBuS	85	-4	10	
		PFHxA	59	-22	15	
		PFHpA	74	9	2.6	
		PFHxS	66	-7	3.7	
		PFOA	64	-11	3.1	
		PFNA	64	-8	0.5	
		PFOS	76	2	2.0	
Strawberry	Ultrasonic assisted extraction with CH ₃ OH+1% NH ₄ OH/CH ₂ Cl ₂ 1/1 (v/v). Multiple clean-up processes. Total analysis time ≈ 8 h.	PFPeA			143 ^b	[28]
		PFBuS			71 ^b	
		PFHxA			1450 ^b	
		PFHpA	45 ^a	n.i.	2910 ^b	
		PFHxS			27 ^b	
		PFOA			737 ^b	
		PFNA			79 ^b	
		PFOS			29 ^b	
Strawberry	Ultrasonic treatment and shaking extraction with CH ₃ OH+10 mM KOH. Poly(styrene-co-pyrrolidone-co-divinylbenzene) functionalized with piperazine SPE clean-up. Total analysis time ≈ 17-16 h.	PFBuS			2	[24]
		PFHxA			2	
		PFHpA			4	
		PFHxS	n.i.	n.i.	4	
		PFOA			1	
		PFNA			1	
		PFOS			2	
		PFDA			4	
Cranberry Bog	QuEChERS with PSA/GCB 2/1 (w/w) d-SPE clean-up. Total analysis time ≈ 40 min.	PFPeA	106		2112 ^c	[29]
		PFBuS	104		7590 ^c	
		PFHxA	101		792 ^c	
		PFHpA	101	n.i.	3102 ^c	
		PFHxS	91		2607 ^c	
		PFOA	94		1386 ^c	
		PFOS	109		1782 ^c	
Apple	QuEChERS with poly(styrene-co-divinylbenzene) SPE clean-up. Total analysis time ≈ 70 min.	PFPeA	89		14	[25]
		PFBuS	89		27	
		PFHxA	91		24	
		PFHpA	97		20	
		PFHxS	85	n.r.	17	
		PFOA	89		12	
		PFNA	95		14	
		PFOS	91		6	
Apple and Orange	Ultrasonic assisted extraction with THF/water 75/25 (v/v). Multiple clean-up processes. Total analysis time ≈ 2 h.	PFPeA	n.r.		60	[27]
		PFBuS	n.r.		50	
		PFHxA	93-103		15	
		PFHpA	n.r.		15	
		PFHxS	81-120	≤10 ^d	25	
		PFOA	82-98		10	
		PFNA	82-103		5	
		PFOS	81-120		3.5	
PFDA	84-96		5			

^a Average recovery across all analytes. ^b Calculated from method ~~detection-quantification~~ limits reported in the original manuscript on a dry weight basis, dividing by 10 (i.e. the wet weight-to-dry weight ratio, considering a humidity percentage of 90%). ^c Calculated from method detection limits reported in the original manuscript, multiplying by 3.3. ^d Absolute value of matrix effect reported for all target analytes without specifications.

Table 4 – Mean concentrations (pg/g d.w.) and standard deviation (n=3, in bracket) of PFAAs in strawberry and olive fruits commercially available at supermarkets and grown under irrigation with tap water (TW) and treated wastewaters (TWWs).

Compound	Market 1	Market 2	TW	TWW ₁	TWW ₂	TWW ₃	TWW ₄
<u>Strawberry</u>							
PFPeA	<109 ^a	<109 ^a	<109 ^a	<109 ^a	<109 ^a	<109 ^a	<109 ^a
PFBuS	<29 ^a	<29 ^a	<29 ^a	<29 ^a	<29 ^a	<29 ^a	<29 ^a
PFHxA	<40 ^a	40 ^a -146 ^b	<40 ^a	40 ^a -146 ^b	<40 ^a	<40 ^a	40 ^a -146 ^b
PFHpA	<7.2 ^a	7.2 ^a -26 ^b	<7.2 ^a	<7.2 ^a	<7.2 ^a	<7.2 ^a	<7.2 ^a
PFHxS	148	790	<10 ^a	<10 ^a	<10 ^a	<10 ^a	<10 ^a
PFOA	<8.6 ^a	27	<8.6 ^a	8.6 ^a -31 ^b	8.6 ^a -31 ^b	8.6 ^a -31 ^b	8.6 ^a -31 ^b
PFNA	<1.3 ^a	47	<1.3 ^a	<1.3 ^a	<1.3 ^a	<1.3 ^a	<1.3 ^a
PFOS	<5.7 ^a	90	<5.7 ^a	<5.7 ^a	<5.7 ^a	<5.7 ^a	<5.7 ^a
PFDA	54	35	<0.8 ^a	<0.8 ^a	<0.8 ^a	<0.8 ^a	<0.8 ^a
Total PFAAs	202	989	n.d.	n.d.	n.d.	n.d.	n.d.
<u>Olive</u>							
PFPeA	227	682	35 ^a -127 ^b	1408	740	576	1019
PFBuS	403	185	24 ^a -85 ^b	<24 ^a	24 ^a -85 ^b	<24 ^a	114
PFHxA	<26 ^a	33	26 ^a -94 ^b	481	209	103	256
PFHpA	<21 ^a	107	<21 ^a	21 ^a -74 ^b	21 ^a -74 ^b	<21 ^a	21 ^a -74 ^b
PFHxS	<2.4 ^a	22	<2.4 ^a	2.4 ^a -8.6 ^b	2.4 ^a -8.6 ^b	<2.4 ^a	<2.4 ^a
PFOA	124	137	22 ^a -78 ^b	22 ^a -78 ^b	22 ^a -78 ^b	<22 ^a	22 ^a -78 ^b
PFNA	42	35	<0.7 ^a	<0.7 ^a	<0.7 ^a	<0.7 ^a	<0.7 ^a
PFOS	39	95	<5.8 ^a	5.8 ^a -21 ^b	5.8 ^a -21 ^b	<5.8 ^a	5.8 ^a -21 ^b
PFDA	2.7 ^a -9.8 ^b	15	<2.7 ^a	20	2.7 ^a -9.8 ^b	<2.7 ^a	12
Total PFAAs	835	1311	n.d.	1909	949	679	1401

(^a) MDL in real sample. (^b) MQL in real sample. n.d. = not determined.




Figure 1 – MRM chromatograms of (A) PFPeA and (B) PFBuS reference standard solutions (concentration 500 ng/L) prepared in the QuEChERS procedural blank as such (dashed line) and acidified with 0.1% (v/v) formic acid (solid line).




Figure 2 – Reconstructed MRM chromatogram of the quantifier (solid line) and qualifier (dashed line) transitions of (A) target PFAAs obtained by injecting 10 μL of a standard mixture in the procedural blank acidified with 0.1% formic acid, at a concentration level of 500 ng L^{-1} each. (1) PFPeA; (2) PFBuS; (3) PFHxA; (4) PFHpA; (5) PFHxS; (6) PFOA; (7) PFNA; (8) PFOS; (9) PFDA. Magnifications of peaks (7) and (8) are shown in boxes (B) and (C), respectively.




Figure 3 – Mean values ($n=3$) of apparent recoveries (A) and matrix effects (B) of the QuEChERS extraction performed on 1 g dry weight-aliquots of strawberry fruits spiked with ^{13}C -labelled PFAAs at 5 ng/g each, rehydrated with 5 mL of LC-MS water and extracted using 2.5, 5 and 10 mL of acetonitrile, and 2 g each of NaCl and MgSO₄. Error bars represent standard deviation. Within a same compound, different letters indicate statistically significant differences according to the Games-Howell non-parametric test ($P < 0.05$).




Figure 4 – Mean values (n=3) of apparent recoveries determined for the QuEChERS and USAE-QuEChERS extractions, coupled with d-SPE clean-up performed with 400 mg of GCB sorbent per mL of olive fruit extract. The extraction was performed on 1 g dry weight-aliquots of olive fruits spiked with ¹³C-labelled PFAAs at 5 ng/g, rehydrated with 5 mL of LC-MS water and treated with 10 mL of acetonitrile and 2 g each of NaCl and MgSO₄. Error bars represent standard deviation.




Figure 5 – Overlapped MRM quantifier (solid line) and qualifier (dashed line) transitions, retention time (t_R , min) and ion ratio of: (A) PFOA, 100 ng/L solution in strawberry procedural blank; (B) PFOA in strawberry “Market 2” sample; (C) PFOA, 100 ng/L solution in olive procedural blank; (D) PFOA in olive “Market 2” sample; (E) PFOS, 100 ng/L solution in strawberry procedural blank; (F) PFOS in strawberry “Market 2” sample; (G) PFOS, 100 ng/L solution in olive procedural blank; (H) PFOS in olive “Market 2” sample.

Supplementary material of the Manuscript:

“Optimization and validation of a method based on QuEChERS extraction and liquid chromatographic-tandem mass spectrometric analysis for the determination of perfluoroalkyl acids in strawberry and olive fruits, as model crops with different matrix characteristics”

by Cristina Vanessa Agata Scordo et al.

S.1 Treatment stages adopted for the production of treated wastewaters

The Baciacavallo wastewater treatment plant (WWTP) is the core of the centralized treatment system of the textile industrial district and domestic sewage of the Prato area (Italy). It is essentially constituted by equalization, primary sedimentation, biological oxidation, sedimentation, flocculation and a final refinement with ozone to remove colour and residual organic micropollutants (TWW₁). The TWW₁ is further treated by clariflocculation, sand filtration and activated carbon obtaining the TWW₂ (Macrolotto 1 refining system). In parallel, TWW₁ is treated by clariflocculation, sand filtration and finally mixed with the Bisenzio river water, obtaining the TWW₃ (Macrolotto 2 refining system). Calice WWTP is the second largest facility in the aforementioned area, devoted to the treatment of both domestic and industrial wastewater, together with leachate from landfills and sewages from septic tanks after a pre-treatment with a membrane biological reactor. The Calice WWTP essentially consists of equalization, primary sedimentation, denitrification, biological oxidation, sedimentation, clariflocculation, sand filtering and ozonation (TWW₄).

S.2 Optimization of mass parameters

The precursor and product ions, as well as compound-dependent parameters used for tandem mass spectrometric analysis of PFAAs, are reported in **Table S1**.

Table S1 – Optimized MRM parameters for the investigated analytes. Letters A and B in bracket after each compound abbreviation refer to the quantifier and qualifier transitions, respectively. Q1: precursor ion; Q3: product ion; DP: declustering potential; EP: entrance potential; CE: collision energy; CXP: collision exit potential.

PFAAs	Q1 (Da)	Q3 (Da)	DP (V)	EP (V)	CE (eV)	CXP (V)
PFBuS (A)	299	80	-120	-10	-66	-12
PFBuS (B)	299	99	-120	-10	-37	-13
PFPeA (A)	263	219	-53	-10	-12	-20
PFHxA (A)	313	269	-50	-10	-12	-23
PFHxA (B)	313	119	-50	-10	-25	-13
PFHxS (A)	399	80	-30	-10	-85	-10
PFHxS (B)	399	99	-30	-10	-42	-12
PFHpA (A)	363	319	-50	-10	-13	-28
PFHpA (B)	363	169	-50	-10	-25	-15
PFOA (A)	413	369	-55	-10	-13	-32
PFOA (B)	413	169	-55	-10	-24	-14
PFOS (A)	499	80	-120	-10	-105	-7
PFOS (B)	499	99	-120	-10	-95	-9
PFNA (A)	463	419	-60	-10	-15	-38
PFNA (B)	463	219	-60	-10	-25	-17
PFDA (A)	513	469	-67	-10	-15	-40
PFDA (B)	513	219	-67	-10	-25	-19
¹³ C3-PFBuS (A)	302	80	-110	-10	-70	-12
¹³ C3-PFBuS (B)	302	99	-110	-10	-35	-13
¹³ C5- PFPeA (A)	268	223	-40	-10	-15	-20
¹³ C5-PFHxA (A)	318	273	-45	-10	-13	-23
¹³ C5-PFHxA (B)	318	120	-45	-10	-30	-13
¹³ C3-PFHxS (A)	402	80	-110	-10	-85	-10
¹³ C3-PFHxS (B)	402	99	-110	-10	-45	-12
¹³ C4-PFHpA (A)	367	322	-60	-10	-15	-28
¹³ C4-PFHpA (B)	367	169	-60	-10	-20	-15
¹³ C8-PFOA (A)	421	376	-70	-10	-14	-32
¹³ C8-PFOA (B)	421	172	-70	-10	-28	-14
¹³ C8-PFOS (A)	507	80	-110	-10	-113	-7
¹³ C8-PFOS (B)	507	99	-110	-10	-85	-9
¹³ C9-PFNA (A)	472	427	-60	-10	-17	-38
¹³ C9-PFNA (B)	472	223	-60	-10	-28	-17
¹³ C6-PFDA (A)	519	474	-80	-10	-17	-40
¹³ C6-PFDA (B)	519	219	-80	-10	-28	-19

S.3 Optimization of the QuEChERS procedure

The development of the QuEChERS method involved preliminarily the evaluation of the water-to-acetonitrile partition of target analytes in the presence of NaCl and MgSO₄. In particular, using 5 mL-aliquots of a standard solution of PFAAs (500 ng/L each) in LC-MS water and 2 g of each salt, the extent of partition was evaluated using 2.5, 5 and 10 mL of acetonitrile. The tests were carried out as following described. Acetonitrile was added to the PFAA standard solution and the QuEChERS procedure mentioned above (paragraph 2.4 of the main text) was performed. The acetonitrile supernatant derived from the centrifugation is collected and made up to 10 mL with acetonitrile of the procedural blank in order to compensate the different suppressive ME due to the diverse relative proportions among water, salts and acetonitrile evaluated in the partition experiments. The extracts are then filtered at 0.2 µm and analysed by LC-MS/MS, comparing the chromatographic areas of the partition experiments with calibration lines based on PFAA standard solutions prepared with the corresponding procedural blanks. **Figure S1** illustrates the results of these experiments, which highlighted recovery percentages in the ranges of 88-104%, 83-99% and 84-108% for the acetonitrile volumes of 2.5 mL, 5 mL and 10 mL, respectively. In particular, it should be noted that high recovery values were achieved also for PFBuS and PFHxS, which showed negative values of the log K_{OW} and that, for almost all PFAAs, the variation of acetonitrile volume did not significantly affect the recovery values.




Figure S1 – Mean recovery values (n=3) of the water-to-acetonitrile partition experiments performed on PFAA standard solutions (500 ng/L) prepared in LC-MS water (pH=6.57±0.05), using 2.5, 5.0 and 10 mL of acetonitrile. Error bars represent standard deviation. Within a same compound, different letters indicate statistically significant differences according to the Games-Howell non-parametric test ($P < 0.05$).

Figure S2 highlights the detail of the oily and strongly pigmented acetonitrile extract obtained treating one gram of freeze-dried olive fruit (cultivar “Frantoio”) as specified in the paragraph 2.4.2 of the main text. Briefly, the freeze-dried fruit is rehydrated with 5 mL of LC-MS water and the mixture hand shaken for 15 s and vortex-mixed for 1 min. Then, 10 mL of acetonitrile are added and the mixture is further hand shaken for 15 s, and sonicated for 90 sec. Afterwards, 2 g of NaCl and 2 g of MgSO₄ are added, and the obtained mixture additionally hand shaken (15 s) and vortex mixed (1 min). The tube was centrifuged at 9000×g and T=4°C for 5 min, obtaining the aforementioned oily and pigmented extract.



Figure S2 – Detail of the oily and strongly pigmented QuEChERS extract obtained in this study for the extraction of the “Frantoio” cultivar olives.

The results of the matrix-free tests clean-up procedure employing PSA, C18 and GCB sorbents were illustrated in **Figure S3**.

Figure S4 and **Figure S5** illustrate the trends of the matrix effect (ME%), expressed as percentage of suppression (negative values) or enhancement (positive values) of the signal, determined for the QuEChERS and USAE- QuEChERS extracts of olive fruits.




Figure S3 – Mean recovery values (n=3) of the matrix-free d-SPE partition experiments performed on PFAA standard solutions (500 ng/L in LC-MS acetonitrile deriving from the QuEChERS extraction procedural blank), containing 150 mg/mL of MgSO₄, using 300 mg/mL of PSA, C18 and GCB. Error bars represent standard deviation. Within a same compound, different letters indicate statistically significant differences according to the Games-Howell non-parametric test ($P < 0.05$).




Figure S4 – Mean values (n=3) of matrix effects determined for the optimized QuEChERS extraction, coupled with d-SPE clean-up performed with 200, 300 and 400 mg of C18 (A) and GCB (B) sorbents per mL of olive fruit extract. Error bars represent standard deviation. Within a same compound, different letters indicate statistically significant differences according to the Games-Howell non-parametric test ($P < 0.05$).




Figure S5 – Mean values (n=3) of matrix effects determined for the optimized USAE-QuEChERS extraction, coupled with d-SPE clean-up performed with 200, 300 and 400 mg of C18 (A) and GCB (B) sorbents per mL of olive fruit extract. Error bars represent standard deviation. Within a same compound, different letters indicate statistically significant differences according to the Games-Howell non-parametric test ($P < 0.05$).

The QuEChERS protocols optimized and validated for the determination of PFAAs in freeze-dried strawberry and olive fruits are illustrated in **Figure S6**.




Figure S6 – Scheme of the QuEChERS protocols for the analysis of PFAAs in (A) strawberry and (B) olive fruits.

Table S2 – Characteristics of the calibration lines in the procedural blank.

Compound	Range (pg/g d.w.)	R ²	Slope	Slope error	Significance	Intercept	Intercept error	Significance
<u>Strawberry</u>								
PFPeA	500-10000	0.9992	43785	396	<0.05	-12892	2225	<0.05
PFBuS	500-10000	0.9987	42947	488	<0.05	3335	2742	0.25
PFHxA	500-10000	0.9984	81419	1043	<0.05	694	5858	0.90
PFHpA	50-10000	0.9993	103633	690	<0.05	2257	3169	0.48
PFHxS	50-10000	0.9988	61379	539	<0.05	1343	2474	0.60
PFOA	50-10000	0.9989	134297	1111	<0.05	911	5100	0.86
PFNA	10-10000	0.9993	104241	653	<0.05	917	2774	0.74
PFOS	50-10000	0.9985	54473	481	<0.05	-5833	2209	<0.05
PFDA	10-10000	0.9988	127352	1027	<0.05	581	4363	0.89
<u>Olive</u>								
PFPeA	500-10000	0.9997	33086	151	<0.05	1351	693	0.07
PFBuS	50-10000	0.9991	32628	250	<0.05	404	1151	0.73
PFHxA	50-10000	0.9994	84044	665	<0.05	-11339	3048	<0.05
PFHpA	50-10000	0.9992	101963	738	<0.05	-6669	3389	0.06
PFHxS	10-10000	0.9997	72345	298	<0.05	-5114	1267	<0.05
PFOA	50-10000	0.9997	102589	432	<0.05	-5354	1980	<0.05
PFNA	10-10000	0.9997	104497	442	<0.05	-2047	1879	0.28
PFOS	50-10000	0.9994	54980	462	<0.05	-3210	2119	0.15
PFDA	10-10000	0.9997	154130	660	<0.05	-11863	2804	<0.05