

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

A case of adrenal vein sampling in primary aldosteronism with homolateral suppression

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1742450> since 2020-06-29T12:53:51Z

Published version:

DOI:10.1210/js.2016-1105

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 **A CASE OF ADRENAL VEIN SAMPLING IN PRIMARY ALDOSTERONISM WITH**
2 **“HOMOLATERAL SUPPRESSION”**

3

4 Andrea Viola, Silvia Monticone, Denis Rossato, Karine Versace, Isabella Castellano, Jacopo
5 Burrello, Fabrizio Buffolo, Franco Veglio, Paolo Mulatero*.

6

7 Department of Medical Sciences, Division of Internal Medicine and Hypertension (A.V., S.M., J.B.,
8 F.B., F.V., P.M.), Service of Radiology (D.R., K.V.), and service of Pathology (I.C.), University of
9 Torino, Torino, 10126 Italy.

10

11 *Corresponding author:

12 Paolo Mulatero, Department of Medical Sciences, Division of Internal Medicine and Hypertension,
13 University of Torino, Via Genova 3, 10126, Torino, Italy.

14 e-mail: paolo.mulatero@unito.it

15 fax:-39-011-6336931

16 ph:-39-011-6336959

17

18 Word count: abstract 233; text 1472 excluding references, legends and figure captions.

19 Number of Tables: 2; Number of Figures: 2; Supplemental file: 1.

20 Running title: adrenal sampling in primary aldosteronism

21 Keywords: aldosterone producing adenoma; adrenal venous sampling; cosyntropin stimulation;
22 aldosterone, endocrine hypertension.

23

24 Source of founding: none.

25 Financial disclosure: none.

26

1 **ABSTRACT**

2 **Context:** Adrenal venous sampling (AVS) is regarded as the gold standard in primary
3 aldosteronism (PA) subtype diagnosis, although some authors have questioned its diagnostic
4 accuracy and highlighted its lack of standardization of procedure protocol and interpretation
5 criteria. In particular, the usefulness of cosyntropin stimulation and benefit of super-selective
6 adrenal vein catheterization have been hotly debated.

7 **Objective:** we report herein a case that highlights the potential pitfalls of super-selective sampling
8 and demonstrates a negligible effect of cosyntropin stimulation on aldosterone secretion in non-
9 adenomatous adrenal tissue when an aldosterone producing adenoma (APA) is present.

10 **Case Report:** a 38-year old man with PA and a single right macro-adenoma underwent AVS at our
11 Centre. The procedure was performed both under basal conditions and during cosyntropin
12 stimulation. Right adrenal vein angiography demonstrated two branches of the main adrenal vein
13 trunk, one draining the nodule and one draining the right adrenal gland. Hormonal assays confirmed
14 adrenal origin of left-sided and all right-sided samples, and were consistent with lateralization on
15 the right side with suppression of aldosterone secretion in the left adrenal gland and in the non-
16 adenomatous right adrenal tissue. Cosyntropin-stimulated AVS results were similar to those of the
17 unstimulated procedure.

18 **Conclusions:** cosyntropin stimulation does not significantly affect aldosterone secretion from non-
19 adenomatous adrenal tissue when an APA is present and can therefore be used during AVS for PA.
20 Super-selective AVS should be performed with caution and interpreted by expert clinicians.

21

1 **INTRODUCTION**

2 Diagnostic work-up in primary aldosteronism (PA) requires subtype differentiation that represents
3 the key for optimal therapeutic plan [1]. Adrenal venous sampling (AVS) is the recommended
4 procedure to distinguish unilateral from bilateral forms. Unfortunately AVS still lacks of a
5 standardized protocol, despite recent efforts to achieve consensus on this topic [2,3]. Different
6 Centres adopt different schedules in performing the sampling (e.g., bilateral simultaneous sampling
7 versus sequential cannulation of adrenal veins; central vein versus super-selective sampling from
8 secondary adrenal vein branches; use of cosyntropin stimulation) and several different cut-off
9 values of selectivity index and lateralization ratio have been proposed to define successful adrenal
10 vein cannulation and lateralized aldosterone overproduction [2,3].

11 The case we report herein allows discussion and provides useful answers about two debated issues
12 on performance and interpretation of AVS, namely the use of super-selective adrenal veins
13 cannulation and the benefit of cosyntropin stimulation.

14
15 **CASE REPORT**

16 We report the case of a 38-years-old man referred to our Hypertension Unit with a confirmed PA
17 diagnosis (Table 1). Contrast-enhanced abdominal computed-tomography scanning revealed a
18 normal left adrenal gland and a exophytic right adrenal nodule 22 mm in diameter (Figure 1). AVS
19 was performed under basal conditions and during continuous cosyntropin infusion (50 µg/h). On the
20 right side the radiologist placed the catheter at first just at the outlet of the main adrenal vein trunk
21 in the inferior vena cava (IVC), being able to stain by venography two secondary adrenal vein
22 branches: one, in the cranial position, seemed to drain the body and the limbs of the gland, whereas
23 a second branch, caudally directed, seemed to drain just the right nodule (Figure 2). Samples were
24 obtained initially placing the catheter tip at the outlet of the main adrenal vein trunk in the IVC
25 (Sample 1 – Table 2) and then by deeply inserting the catheter in both of the secondary branches

1 (Sample 2 from the upper branch draining the body and the limbs and Sample 3 from the caudal
2 branch draining the nodule – Table 2). Two of the three samples from the right side (the one
3 collected from the main right adrenal vein trunk and the one from the nodule-draining secondary
4 branch) showed a high cortisol-corrected aldosterone concentration, whereas the other (the one
5 collected from the whole gland-staining secondary branch) showed an aldosterone/cortisol ratio
6 (ACR) that was lower than that measured in IVC. The left-sided sample showed a low cortisol-
7 corrected aldosterone concentration as well, that was again inferior to that measured in IVC.
8 Interestingly, both in the unstimulated and in the cosyntropin-stimulated procedures, this condition
9 of “suppression” of aldosterone production could be observed. The lateralization index (LI, defined
10 as the ACR from dominant adrenal over the ACR from non-dominant adrenal) using sample 3 on
11 the right side was 7.2 under basal conditions and 13.8 during cosyntropin infusion; the ACR using
12 the sample 2 on the right side resulted inferior to the ACR from the IVC, indicating suppression of
13 the aldosterone production, both under basal and cosyntropin-stimulated conditions, similarly to the
14 left adrenal vein. The patient underwent right total laparoscopic adrenalectomy.

15 Immunohistochemical staining of the adrenal nodule and of the surrounding adrenal gland, using
16 specific antibodies for 11 β -hydroxylase and aldosterone synthase (from Prof. Celso Gomez-
17 Sanchez, University of Mississippi, Jackson, MS) [4] showed non-homogeneous aldosterone
18 synthase staining exclusively present in the adrenal nodule, and it was absent in the surrounding
19 adrenal cortex (Supplemental Figure S1A), indicating that the source of aldosterone excess was the
20 nodule and that aldosterone production in the surrounding adrenal zona glomerulosa was
21 suppressed. 11 β -hydroxylase staining was present both in the adrenal surrounding the nodule and,
22 less strongly, inside the nodule (Supplemental Figure S1B). After surgery, blood pressure and
23 potassium levels were normalized and the patient is now free from medication. The study was
24 approved by the local ethic committee and the patient gave his written consent.

1 A case with similar findings and “homolateral suppression” despite the presence of an aldosterone-
2 producing cell cluster outside the main nodule is provided in the supplemental file (supplemental
3 Table S1 and Figure S2).

4

5 **DISCUSSION**

6 In this case report, AVS results showed low cortisol-corrected aldosterone concentration not only
7 on the left side, contralateral to the adenoma, but also in one of two different sites of blood
8 sampling on the right side, both of adrenal origin. Both in the left-sided sample and in the right-
9 sided sample the ACR was less than that in IVC. This has been called “contralateral suppression”
10 when applied to the side contralateral to an adrenal adenoma [2,3]. Contralateral suppression has
11 been proposed as an additional criterion to detect lateralized aldosterone production [2,3] and some
12 authors consider it a necessary prerequisite before adrenalectomy [5,6]. These findings can be
13 interpreted considering that blood in the different samples from the right adrenal vein come from
14 different regions of the gland, among which only one was producing high amounts of aldosterone;
15 consequently, not only left-sided adrenal tissue aldosterone production, but also right non-
16 adenomatous tissue aldosterone production was suppressed.

17 We obtained right blood samples at the outlet of the main right adrenal vein trunk; we then inserted
18 the catheter at first in the gland-staining cranial secondary branch, and then in the nodule-staining
19 caudal secondary branch to perform a “super-selective” adrenal vein cannulation, retrieving
20 different samples of blood from different portions of the right-sided adrenal tissue. Selective adrenal
21 vein cannulation has been shown to be necessary when right adrenal vein and accessory hepatic
22 veins share the same point of entry into the IVC, to avoid excessive dilution of adrenal blood [3].
23 Super-selective adrenal venous sampling has been already described in literature and advocated as
24 accurate method to allow the localization of adrenal tissue involved in aldosterone hypersecretion
25 [7]. As shown in Figure 2, deep insertion and staining of the catheter just in the cranial vein branch

1 would have apparently demonstrated the whole right adrenal gland. An inexperienced radiologist
2 could have sampled blood from this site only, thereby missing the actual aldosterone-producing site,
3 resulting in a misleading AVS finding of bilateral suppression of aldosterone production. Instead,
4 when we sampled blood just from the outlet of the adrenal vein in the IVC, we were able to obtain a
5 high aldosterone concentration and consequently not miss the right aldosterone overproduction.
6 Therefore, super-selective AVS is a useful technique in some selected situations but should be
7 performed with the caution of not missing any adrenal vein branches. Furthermore, super-selective
8 AVS increases the occurrence of adrenal hemorrhage [7], a complication that is very rare in
9 standard AVS procedure [8].

10 Cosyntropin stimulation has a proposed favorable effect in maximizing the adrenal-to-peripheral
11 cortisol gradient, stimulating aldosterone secretion from the adenoma, increasing the LI and
12 reducing time-dependent fluctuations in hormone secretion. Recently, a multicentric study
13 demonstrated that cosyntropin administration during AVS does not affect significantly LI and gives
14 a similar diagnosis as with unstimulated AVS [9]. In the present clinical case we observed, both
15 under basal conditions and after cosyntropin stimulation, a consistent pattern of suppression not
16 only on the side contralateral to the adenoma, but also from a sampling site on the same side of the
17 node, draining the extra-nodal adrenal tissue. This observation reinforces suggests that cosyntropin
18 stimulation does not significantly affect aldosterone gradients between the two adrenal glands when
19 an adenoma is present and, therefore, it does not interfere with the final diagnosis [6].

20 Finally, the immunohistochemical analysis demonstrated that the APA was also cosecreting some
21 cortisol, still suppressible during overnight 1 mg dexamethasone suppression test and not sufficient
22 to inhibit both ACTH secretion and 11 β -hydroxylase expression outside the adenoma. Of particular
23 interest, the cosecretion of cortisol from the APA did not interfere with the diagnosis of unilateral
24 PA both under basal conditions and during cosyntropin infusion. APA cosecreting cortisol have
25 been described previously [10,11]. It has been suggested that this phenomenon is more frequent in

1 APA of greater size and therefore in these cases a dexamethasone suppression test before AVS
2 performance is warranted.

3 AVS, and particularly super-selective AVS, is a challenging technique not available in most centres.
4 In future, non-invasive procedures able to distinguish between unilateral and bilateral PA, such as
5 PET-CT scanning using CYP11B2-specific radiolabelled tracer (12), may substitute AVS in PA
6 subtype differentiation.

7

8 **CONCLUSION**

9 In conclusion, this case demonstrates some potential pitfalls and offers new insights into the use and
10 interpretation of AVS in the diagnostic PA work-up. This case shows the potential risks of super-
11 selective cannulation when it is not accurately performed and may therefore result in misleading
12 diagnosis. Furthermore, we observed that cosyntropin stimulation did not stimulate aldosterone
13 production from extra-nodal tissue homolateral to the adenoma; rather, cortisol-corrected
14 aldosterone concentration appeared consistent with suppression both in veins draining the extra-
15 nodal tissue homolateral to the adenoma and the contralateral adrenal gland. This observation
16 reinforces the indication that cosyntropin stimulation does not affect the diagnosis of unilateral PA
17 and therefore can be used during AVS.

1

2 **REFERENCES**

3 1. Funder JW, Carey RM, Mantero F, Murad MH, Reincke M, Shibata H, Stowasser M, Young WF
4 Jr. The Management of Primary Aldosteronism: Case Detection, Diagnosis, and Treatment: An
5 Endocrine Society Clinical Practice Guideline. *J Clin Endocrinol Metab* 2016; 101: 1889-916.

6 2. Monticone S, Viola A, Rossato D, Veglio F, Reincke M, Gomez-Sanchez C, Mulatero P. Adrenal
7 vein sampling in primary aldosteronism: towards a standardised protocol. *Lancet Diabetes
8 Endocrinol* 2015; 3: 296-303.

9 3. Rossi GP, Auchus RJ, Brown M, Lenders JW, Naruse M, Plouin PF, Satoh F, Young WF Jr. An
10 expert consensus statement on use of adrenal vein sampling for the subtyping of primary
11 aldosteronism. *Hypertension* 2014; 63: 151-60.

12 4. Monticone S, Castellano I, Versace K, Lucatello B, Veglio F, Gomez-Sanchez CE, Williams TA,
13 Mulatero P. Immunohistochemical, genetic and clinical characterization of sporadic aldosterone-
14 producing adenomas. *Mol Cell Endocrinol* 2015; 411: 146-54.

15 5. Wolley MJ, Gordon RD, Ahmed AH, Stowasser M. Does contralateral suppression at adrenal
16 venous sampling predict outcome following unilateral adrenalectomy for primary aldosteronism? A
17 retrospective study. *J Clin Endocrinol Metab* 2015; 100: 1477-84.

18 6. Monticone S, Satoh F, Viola A, Fischer E, Vonend O, Bernini G, Lucatello B, Quinkler M,
19 Ronconi V, Morimoto R, Kudo M, Degenhart C, Gao X, Carrara D, Willenberg HS, Rossato D,
20 Mengozzi G, Riestler A, Paci E, Iwakura Y, Burrello J, Maccario M, Giacchetti G, Veglio F, Ito S,
21 Reincke M, Mulatero P. Aldosterone suppression on contralateral adrenal during adrenal vein
22 sampling does not predict blood pressure response after adrenalectomy. *J Clin Endocrinol Metab*
23 2014; 99: 4158-66.7.

24 7. Satoh F, Morimoto R, Seiji K, Satani N, Ota H, Iwakura Y, Ono Y, Kudo M, Nezu M, Omata K,
25 Tezuka Y, Kawasaki Y, Ishidoya S, Arai Y, Takase K, Nakamura Y, McNamara K, Sasano H, Ito

- 1 S. Is there a role for segmental adrenal venous sampling and adrenal sparing surgery in patients
2 with primary aldosteronism? *Eur J Endocrinol* 2015; 173: 465-77.
- 3 8. Monticone S, Satoh F, Dietz AS, Goupil R, Lang K, Pizzolo F, Gordon RD, Morimoto R,
4 Reincke M, Stowasser M, Mulatero P. Clinical Management and Outcomes of Adrenal Hemorrhage
5 Following Adrenal Vein Sampling in Primary Aldosteronism. *Hypertension* 2016; 67: 146-52.
- 6 9. Monticone S, Satoh F, Giacchetti G, Viola A, Morimoto R, Kudo M, Iwakura Y, Ono Y, Turchi
7 F, Paci E, Veglio F, Boscaro M, Rainey W, Ito S, Mulatero P. Effect of adrenocorticotropic
8 hormone stimulation during adrenal vein sampling in primary aldosteronism. *Hypertension* 2012;
9 59: 840-6.
- 10 10. Fallo F, Bertello C, Tizzani D, Fassina A, Boulkroun S, Sonino N, Monticone S, Viola A,
11 Veglio F, Mulatero P. Concurrent primary aldosteronism and subclinical cortisol hypersecretion: a
12 prospective study. *J Hypertens* 2011; 29: 1773-7.
- 13 11. Hiraishi K, Yoshimoto T, Tsuchiya K, Minami I, Doi M, Izumiyama H, Sasano H, Hirata Y.
14 Clinicopathological features of primary aldosteronism associated with subclinical Cushing's
15 syndrome. *Endocr J* 2011; 58: 543-51.
- 16 12. Abe T, Naruse M, Young WF Jr, Kobashi N, Doi Y, Izawa A, Akama K, Okumura Y, Ikenaga
17 M, Kimura H, Saji H, Mukai K, Matsumoto H. A Novel CYP11B2-Specific Imaging Agent for
18 Detection of Unilateral Subtypes of Primary Aldosteronism. *J Clin Endocrinol Metab.*
19 2016;101:1008-15.

1

Clinical and biochemical parameters	At diagnosis	Normal values
SBP/DBP (mmHg)	175/115	≤140/90
Number of drugs	2	-
Serum Potassium (mmol/L)	2.6	3.5-5.5
Creatinine (mg/dL)	1.03	0.7-1.4
DRC (μU/mL]	3.8	7-76
Serum Aldosterone (ng/dL)	34.8	7-30
ACTH (pg/mL)	19.1	17-70
Cortisol at 8.00 a.m. (μg/dL)	28.4	5-25
Cortisol after DST (μg/dL)	0.1	≤1.8
Serum Aldosterone post-SLT (ng/dL)	49.7	≤5

2

3 **Table 1. Clinical and biochemical parameters of the patient.** SBP= Systolic Blood Pressure;
4 DBP= diastolic blood pressure; DRC= direct renin concentration; ACTH= adrenocorticotropic
5 hormone; DST= dexamethasone suppression test.

AVS hormonal values					
Unstimulated AVS	Routine Cortisol (µg/dl)	Aldosterone (ng/dl)	SI	ACR	IR
Infra-renal inferior vena cava	18.3	27.6	-	1.5	-
Left adrenal vein	112	52	6.1	0.5	0.3
Right adrenal vein – Sample 1	519	1277	28.4	2.5	1.6
Right adrenal vein – Sample 2	428	86	23.4	0.2	0.1
Right adrenal vein – Sample 3	625	2237	34.2	3.6	2.4
Cosyntropin-stimulated AVS	Routine Cortisol (µg/dl)	Aldosterone (ng/dl)	SI	ACR	IR
Infra-renal inferior vena cava	32.1	60	-	1.9	-
Left adrenal vein	312	139	9.7	0.4	0.2
Right adrenal vein – Sample 1	840	2927	26.2	3.5	1.8
Right adrenal vein – Sample 2	718	209	22.4	0.3	0.2
Right adrenal vein – Sample 3	1019	5581	31.7	5.5	2.9

2

3 **Table 2. Hormonal measurements from adrenal venous sampling.** AVS= Adrenal Venous

4 Sampling; SI= Selectivity Index; ACR= Aldosterone/Cortisol Ratio; IR= Ipsilateral Ratio.

1 **LEGENDS TO FIGURES**

2 **Legend to figure 1.** CT-scan images of the right adrenal gland. A: frontal view; arrow-head:
3 adrenal gland body; arrow: right adrenal nodule. B: sagittal view; arrow-head: adrenal gland body;
4 arrow: right adrenal nodule. CT: see text.

5 **Legend to figure 2.** Venography during AVS. A: cannulation of the right adrenal vein branch
6 draining the right adrenal gland; arrow-head indicates adrenal gland venography; a pale staining of
7 right adrenal vein branches surrounding right adrenal adenoma is also seen (due to partial passage
8 of contrast medium in the other adrenal vein branch). B: selective cannulation of the right adrenal
9 vein branch draining the right adrenal nodule; the arrow points at adrenal vein branches surrounding
10 the right adrenal nodule.