

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Taxonomic and functional homogenisation of macroinvertebrate communities in recently intermittent Alpine watercourses

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1755616> since 2023-01-27T10:42:07Z

Published version:

DOI:10.1111/fwb.13605

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 **Patterns in taxonomic and functional diversity highlight the “mediterraneization”**
2 **of macroinvertebrate communities in recently intermittent Alpine watercourses**

3 Elena Piano*^{1,2}, Alberto Doretto^{1,2}, Stefano Mammola³, Elisa Falasco^{1,4}, Stefano Fenoglio^{1,4}, Francesca
4 Bona^{1,4}

5 ¹ALPSTREAM – Alpine Stream Research Center, Ostana, I-12030, Italy

6 ²DISIT, University of Piemonte Orientale, Viale Teresa Michel 25, I-15121, Alessandria, Italy

7 ³Molecular Ecology Group (MEG), Water Research Institute (IRSA), National Research Council (CNR), Largo
8 Tonolli 50, I-28922, Pallanza, Italy

9 ⁴DBIOS, University of Turin, Via Accademia Albertina 13, I-10123, Turin, Italy

10 *corresponding author: elena.piano@unito.it

11
12 **Abstract**

- 13 1. Mountain streams are currently shifting from perennial to intermittent regime due to the combined
14 effect of climate change and local anthropogenic pressures. Studies on the effect of flow
15 intermittency on macroinvertebrates are as yet inconclusive due to the large heterogeneity in
16 geographical areas and spatial scales at which this relationship has been investigated. Given that
17 flow intermittency is a recently documented phenomenon in the Alps, only few studies have
18 investigated functional and taxonomical diversity of benthic invertebrate communities in recently
19 intermittent Alpine streams.
- 20 2. We used a hierarchical sampling design to investigate patterns in taxonomic and functional
21 diversity of benthic invertebrate communities in 13 recently intermittent Alpine streams in NW-
22 Italy. In each of these streams, we sampled in April 2017 benthic communities in two reaches with
23 different hydrological conditions: a control reach, with permanent flow, and an intermittent reach,
24 which recently experienced non-flow periods in summer.

- 25 3. We tested for the response of taxonomic richness at multiple spatial scales by partitioning total
26 diversity into the average richness of local communities and the richness due to variation
27 among local communities both within and among reaches. By partitioning total diversity (γ) into
28 its local (α) and turnover (β) components we showed a decrease in local and regional species
29 richness both within and among reaches, whereas variation among communities was significantly
30 lower in intermittent reaches only at the reach scale.
- 31 4. The analysis of the multidimensional trait space of macroinvertebrates in reaches with different
32 hydrological conditions further revealed a significant reduction of functional diversity, dispersion,
33 and evenness in intermittent reaches. There was trait overdispersion in intermittent reaches, as
34 these hosted both typical Alpine taxa and organisms adapted to flow intermittency. In particular,
35 we observed the replacement of taxa with aquatic respiration and preferring medium- to fast-
36 flowing oligotrophic waters by taxa adapted to lentic habitats and with larval dormancy phases.
- 37 5. These results provide strong support that recent flow intermittency determines drastic changes of
38 benthic invertebrate communities in Alpine streams which are facing a “mediterraneization”
39 process. Our work highlights the importance of considering multiple spatial scales and of
40 integrating taxonomic and functional diversity to thoroughly assess the impacts of flow
41 intermittency.

42 **Keywords:** flow intermittency, drying, diversity partitioning, nestedness, functional traits

43

44

45 INTRODUCTION

46 Stream ecosystems are particularly sensitive to climate change, as changes in precipitations,
47 temperature and evaporation direct translate in shifts in river flow regimes and their physical conditions
48 (Lake, 2003). The raising of water temperatures and the disruption of hydrologic cycles, further
49 exacerbated by anthropogenic activities (e.g. deforestation, water capture, streambed alterations; Wohl,
50 2006) can lead, in turn, to stochastic events (Scheffer et al., 2001), like floods and droughts. The observed
51 and predicted climate warming is also increasingly associated to changes in distribution and phenology of
52 biota (Heino et al., 2009; Whitehead et al., 2009).

53 Among flow-related disturbances, water scarcity, and consequent riverbed desiccation, is one of
54 the most pressing environmental issues related to climate change, with expected drastic consequences on
55 biodiversity (Lake, 2003). The necessity of understanding such complex dynamics have recently pushed
56 more and more researchers to study the effect of flow intermittency on the taxonomic richness of benthic
57 invertebrate communities. Yet contrasting results have so far been obtained, as either negative
58 relationships (Datry et al., 2014; Leigh & Datry, 2017; Datry et al. 2016a), no relationship (Bonada et al.,
59 2007; Bogan et al. 2015; Datry et al., 2016b; Leigh et al., 2016) or fluctuating relationships (Bogan et al.
60 2017) were observed. This heterogeneity of results suggests that the effect of flow intermittency may
61 strongly depend on the scale at which this trend is studied. Specifically, in streams experiencing flow
62 intermittency, we can observe the local extinction of species lacking traits conferring resistance (e.g.
63 desiccation-resistant stages) or resilience (e.g. plurivoltine taxa). This may cause a shift in community
64 composition, whereby either only a subset of tolerant species from the original pool will persist *in situ* or
65 less adapted species will be replaced by a new set of species more tolerant to flow intermittency. These
66 two processes, often acting hand-in-hand, determine variations in nestedness and turnover components of
67 β -diversity, respectively, and may ultimately result in biotic homogenisation of intermittent reaches and a
68 reduction of the total taxonomic diversity at the regional scale (Socolar et al. 2016).

69 As anticipated, variations in taxonomic composition further reflect changes in the organization of functional
70 space [functional diversity (FD); Petchey & Gaston, 2002], due to the selection of species based on their

71 biological traits related to physiology, behaviour, feeding strategy or dispersal (MacLean & Beissinger,
72 2017). A reduction in total diversity of traits and their dispersion should translate in a functional
73 homogenization of aquatic communities due to the depletion of less tolerant species, i.e. environmental
74 filtering (e.g. Elias et al., 2015; Leigh et al., 2016; Brown et al., 2018). Therefore, a nuanced understanding
75 of the underlying mechanisms shaping the structure of lotic communities in a flow intermittency regime is
76 best achieved by simultaneously looking at multiple spatial scales and diversity measures (Bruno et al.,
77 2019; Van Looy et al., 2019). While similar comprehensive studies have been performed in Mediterranean
78 streams, where flow intermittency is part of their natural hydrological cycle (García-Roger et al., 2013),
79 little is known in other biogeographic areas where streams are changing from perennial to intermittent
80 regimes as a result of anthropogenic activities, therefore generating novel community trajectories (Bogan
81 et al. 2015).

82 Alpine streams are expected to be particularly affected by flow intermittency, because they are fed
83 at least partially by snowfields and glaciers, which are rapidly disappearing (Gobiet et al., 2014). Moreover,
84 they host specialized organisms adapted to a narrow range of environmental conditions (McGregor et al.
85 1995). In these lotic systems, summer flow cessation periods are increasing, especially at the mid-latitudes,
86 as they are warming at a rate two to three times faster than the global average (Hansen et al., 2005;
87 Pederson et al., 2011). The rapid shrinkage of glaciers and snowpack (Rauscher et al., 2008) anticipates the
88 hydrological peak in spring and induces modifications in both thermal and hydrological regimes as well as in
89 sediment transport (Brighenti et al., 2019). Moreover, mountainous streams are suffering the heavy flow
90 regulation, hydropower and increased water abstraction aimed to fulfill local human needs e.g., agricultural
91 use, drinking water and energy production (Wohl, 2006; López-Rodríguez et al., 2019). The combined
92 effects of both climate change and anthropogenic disturbance are changing mountain streams from
93 perennial to temporary systems (Bruno et al., 2019), alternating non-flow events and rewetting phases
94 (Fenoglio et al., 2010), bringing new challenges to the conservation and management of these ecosystems.
95 In this context, periodic flow intermittency represents a major threat for stream macroinvertebrates in
96 mountainous lotic ecosystems (e.g. Doretto et al., 2018; Piano et al., 2019a; Piano et al., 2019b), whose
97 resistance and resilience abilities are expected to be limited compared to their counterpart in the

98 Mediterranean region, where the drying phase is a natural part of the stream hydrological cycle (Leigh et
99 al., 2016; Vannucchi et al., 2013). Investigating the effects of flow intermittency on Alpine benthic
100 invertebrate communities is therefore pivotal to understand how this phenomenon is changing their
101 communities and to predict potential threats in the near future.

102 We here investigated the effects of flow intermittency on Alpine benthic invertebrate communities
103 using a hierarchical sampling design, wherein a permanent reach, located upstream, and intermittent
104 reach, located downstream, were selected within 13 Alpine streams in SW-Alps recently facing summer
105 flow intermittency. Within each reach we sampled macroinvertebrates in seven patches differing in near-
106 bed conditions. Following modern standards in community ecology analyses (e.g. Pavoine et al., 2011;
107 Jarzyna & Jetz, 2018), we estimate multiple facets of taxonomic and functional diversity at both reach and
108 basin scale to evaluate how hydrological disturbance affects the assemblage composition of benthic
109 invertebrate communities. Specifically, we tested whether: (i) flow intermittency causes taxa loss at local
110 and/or at regional scale; (ii) communities in intermittent sites represent either a subset of taxa usually
111 inhabiting permanent reaches or they are composed by different taxa; and (iii) changes in taxonomic
112 composition are mirrored in the functional diversity due to the filtering of taxa with particular combinations
113 of traits. By assuming that flow intermittency is a major cause of species loss (e.g. Datry et al., 2017;
114 Samways et al., 2020) and considering that the overall pool of species in Alpine streams is expected to be
115 mainly composed by taxa not adapted to drying (e.g. Doretto et al., 2018, 2020a; Piano et al., 2019a), we
116 made the following hypotheses: (i) taxonomic diversity should decrease in intermittent sites at both reach
117 and basin; (ii) variation in taxa composition among permanent and intermittent reaches is best explained
118 by taxa loss (i.e., nestedness) than by taxa replacement (i.e. turnover); and (iii) loss of taxa displaying
119 particular trait combinations should directly lead to a reduction of functional diversity and a general
120 rearrangement of the total functional space.

121 **MATERIALS AND METHODS**

122 **Sampling design**

123 This study was conducted in 13 low to mid-order streams, experiencing dewatering events since
124 2011 in multiple reaches. The study streams were selected based on our previous knowledge (Fenoglio et
125 al., 2007) integrated with available historical hydrological data provided by the local Environmental
126 Protection Agency (ARPA). The study streams are all located in the hydroecoregion of SW-Alps (HER 4,
127 Piemonte, NW Italy; Wasson et al. 2007), thus they are characterized by comparable geology, climate,
128 topology and altitude (Moog et al., 2004). In each stream, we selected two 30 m long sampling sections,
129 differing in their hydrological regime: i) an upstream reach, with permanent water throughout the year
130 (hereinafter UP); and ii) a downstream reach, experiencing dewatering events during summer (hereinafter
131 DOWN). In the study area, flow intermittency events lasted on average two months since 2011, with the
132 riverbed almost completely dry for several kilometres (ARPA, 2013; personal observations). Stream drying
133 was caused by the reduction in precipitations, and consequently of snow accumulation, and the parallel
134 increase in water abstraction for human needs. UP and DOWN reaches were located within 10 km, in order
135 to reduce variations in environmental parameters (as demonstrated in Piano et al., 2019) and changes in
136 community composition attributable to geographical distance rather than differences in flow regimes. Both
137 permanent and intermittent sites were identified in the bottom of the valleys. Sampling site elevation was
138 on average 489 m a.s.l., ranging from 307 and 656 m, and permanent and intermittent reaches within the
139 same stream differed on average of 70 m in their elevation (min = 19 m; max = 155 m). Sampling was
140 performed in April 2017 under moderate flow conditions ($Q_{\text{mean}} = 3.98 \pm 4.56 \text{ m}^3\text{s}^{-1}$) occurring in both
141 reaches, approximately six months after water flow resumption in the DOWN reach (Hydrological bulletins,
142 www.arpa.piemonte.it).

143 **Data collection**

144 In each sampling site, we selected seven sampling patches representative of the different
145 conditions of flow velocity, water depth and substrate composition occurring within each section (7
146 samples x 2 reaches x 13 streams = 182 samples). In each patch, we measured flow velocity at 0.05 m from
147 the bottom and water depth with a current meter (Hydro-bios Kiel). We used a gravelometer to estimate
148 percentages of different substratum sizes based on the Wentworth's grain size classification (Wentworth,

149 1922): boulders (> 256 mm), cobbles (64–256 mm), gravel (2–64 mm), and fine sediment (< 2 mm). One
150 benthic sample was collected in each sampling patch, using a Surber sampler (250 μm mesh size; 0.062m²
151 area), and preserved in 75% ethanol (see Doretto et al., 2020a).

152 In the laboratory, macroinvertebrates were sorted from the substrate, counted and identified to
153 the family or genus level according to Campaioli et al. (1994, 1999). Each taxon was then assigned to: i) a
154 functional feeding group (FFG) based on its morpho-behavioral mechanisms of food acquisition (shredders,
155 scrapers, predators, collector-gatherers, and filterers); ii) a biological group based on its life-history traits
156 (size, life-cycle, respiration, reproduction, and locomotion); and iii) an ecological group based on its
157 ecological requirements (transversal and longitudinal distribution, preferred substrate and current
158 velocity). FFGs were assigned based on the classification provided by Merritt et al. (2017), whereas
159 biological and ecological groups were assigned based on the classification proposed by Usseglio-Polatera et
160 al. (2000), who grouped together taxa showing similar combinations of life-history traits (biological groups)
161 and ecological preferences (ecological groups). We preferred this approach as it allowed us to synthesize all
162 the most important life-history and ecological traits within few categories.

163 **Data analysis**

164 We conducted all analyses in R (R Core Team, 2019).

165 Taxonomic diversity. In order to test our first hypothesis (H1), we partitioned the total taxonomic richness
166 into its underlying components both within and among reaches. First, we decomposed the total diversity of
167 each stream (γ_w) into its within-reach diversity components (hereafter “reach scale”), namely average local
168 taxa richness of the seven patches within a reach (α_w) and the variation among these communities (β_w) (Fig.
169 1). We therefore obtained 13 values of each diversity component, one for each stream. In a second step,
170 we calculated the among-reach diversity components (hereafter “basin scale”), namely: i) average local
171 diversity (α_a), obtained by averaging the total taxa richness of each reach; ii) total diversity (γ_a), calculated
172 as the total number of taxa observed at the basin scale; and iii) the variation in taxa richness (β_a) among
173 reaches (Fig. 1). We therefore obtained a single value for each diversity component. Variation in taxa
174 composition among local communities (β -diversity) was calculated with the additive formula as the

175 difference between γ - and α -diversity (Anderson et al., 2011). Therefore, variation in taxa composition
176 within reaches (β_w) is calculated as the difference between γ_w and α_w , while variation in taxa composition
177 among reaches (β_a) is calculated as the difference between γ_a and α_a . Being calculated with this formula, β -
178 diversity expresses the number of species that accumulates at regional level as a result of differences
179 between sites at local level. Differences in taxa richness among permanent and intermittent reaches at the
180 reach scale was tested with a paired t-test. On the other hand, differences in taxa richness at basin scale
181 were tested with null modelling, by permuting samples over the two hydrological categories as only a single
182 value for permanent and intermittent reaches was produced. Significant differences between streams can
183 thus be obtained by comparing the observed value of the taxa richness to the null expectation where
184 hydrological categories (permanent vs intermittent) are randomly permuted across the sampling sites
185 (Legendre & Legendre, 1998).

186 In order to test our second hypothesis (H2), we decomposed total beta diversity, here intended as the
187 dissimilarity between communities in UP and DOWN sites (β_{total}) into its turnover (β_{repl}) and nestedness
188 (β_{rich}) components with the function *beta* in the BAT package (Cardoso et al., 2020). We decomposed beta
189 diversity within UP and DOWN reaches and we thus obtained 13 values of total beta diversity, turnover and
190 nestedness for UP and DOWN sites, respectively, and we tested for differences between UP and DOWN
191 sites with a paired t-test.

192 Functional diversity. In order to test our third hypothesis, we examined variation in functional
193 diversity between benthic invertebrate communities in UP vs DOWN using functional n -dimensional
194 hypervolumes (Mammola & Cardoso, 2020) as implemented in the BAT package (Cardoso et al., 2020). We
195 constructed the community functional hypervolume for each reach with the *hypervolume* package (Blonder
196 et al., 2014; Blonder, 2019). Since functional traits of benthic invertebrates here considered are categorical,
197 we used the approach proposed by Carvalho & Cardoso (2018) to incorporate categorical variables in the
198 hypervolume estimation procedure. We applied a Gower dissimilarity measure to the complete trait matrix
199 and extracted orthogonal morphological axes through Principal Coordinate Analysis (PCoA). We retained
200 the first three PCoA axes (43% Variance explained) to construct hypervolumes using a gaussian kernel
201 estimator (Blonder et al., 2018) and a default bandwidth for each axis. In turn, we estimated functional α

202 diversity, functional dispersion and functional evenness based on the approach by Mammola & Cardoso
203 (2020). To further inspect functional turnovers among permanent and intermittent reaches, we estimated
204 functional β -diversity following Carvalho & Cardoso (2018). This approach computes a pairwise
205 decomposition of the overall differentiation among kernel hypervolumes into two components: the
206 replacement of space between hypervolumes (β -replacement) and net differences between the amount of
207 space enclosed by each hypervolume (β -richness) (see Cardoso et al., 2014 and Mammola & Cardoso, 2020
208 for further details). As we had seven values for each reach within each flow category, we tested for
209 significant differences between UP and DOWN reaches using linear mixed models (LMMs), with the flow
210 intermittency category (UP vs DOWN) as independent variable and stream identity as a random factor. We
211 included the latter to account for spatial dependence of the data (sampling reaches nested within streams)
212 (Zuur et al., 2009).

213 To investigate which combination of traits is selected in UP and DOWN sites, we evaluated the
214 functional contribution of each taxon (observation) within communities to the functional space and
215 originality of each taxon within both UP and DOWN reaches. Taxa with high values of contribution provide
216 rare combinations of traits within the community thus contributing the most to the overall functional
217 diversity of the community (Pavoine et al., 2005; Violle et al., 2017), whereas taxa with high originality
218 values are the ones contributing the most in determining the functional dispersion of the community
219 (Mammola & Cardoso, 2020).

220

221 **RESULTS**

222 **Taxonomic diversity**

223 We observed a significant decline in taxa diversity from UP to DOWN reaches both within and
224 among reaches (Fig. 2a). When considering diversity within reaches, we could detect a significant decrease
225 in all diversity components, indicating that flow intermittency reduces both local (α_w : $t = 4.85$, $P < 0.001$)
226 and total (γ_w : $t = 6.06$, $P < 0.001$) diversity within reaches from UP to DOWN sites, as well as taxa variation
227 among habitat patches within reaches (β_w : $t = 4.26$, $P = 0.001$). When considering diversity among reaches,

228 we could detect a significant decline in both local (α_d : $P < 0.001$) and total (γ_d : $P < 0.001$) diversity
229 components from UP to DOWN sites, but not in terms of taxa variation (β_d : $P = 0.333$).

230 By partitioning total β -diversity, here intended as the dissimilarity among community matrices,
231 into its turnover and nestedness components, we observed a higher contribution of nestedness than
232 turnover in both UP and DOWN sites (UP: $\beta_{total} = 0.69$, $\beta_{repl} = 0.19$, $\beta_{rich} = 0.50$; DOWN: $\beta_{total} = 0.66$, $\beta_{repl} =$
233 0.21 , $\beta_{rich} = 0.45$), although no significant differences between UP and DOWN sites were observed
234 (turnover: $t = -0.507$, $P = 0.621$; nestedness: $t = 1.05$, $P = 0.313$). This suggests that the contribution of the
235 two components is equal in the two hydrological levels (Fig. 2b).

236

237 **Functional diversity**

238 Results of the LMMs showed that all functional metrics significantly differ between UP and DOWN
239 reaches (Tab. 1). α -diversity, dispersion and evenness were significantly lower in DOWN than in UP reaches
240 (Fig. 3). When considering β -diversity, we observed significantly higher values in pairwise comparisons
241 between communities in DOWN reaches, whereas communities in UP reaches were in general more similar
242 to one another, suggesting trait overdispersion in sites experiencing flow intermittency. In DOWN reaches
243 the β -richness component was more preponderant than in UP reaches, whereas the β -replacement was
244 higher in UP reaches. When evaluating the contribution of each taxon, we inferred that Tipulidae (Diptera)
245 was the taxon contributing the most to the functional space in UP reaches, and *Perla* (Plecoptera, Perlidae)
246 in DOWN reaches. In other words, Tipulidae contributed with unique traits in UP sites, while *Perla*
247 contributed with unique traits in DOWN sites.

248

249 **DISCUSSION**

250 According to our results, flow intermittency is causing a decline in macroinvertebrate taxonomic
251 richness at both reach and basin scale, suggesting that this phenomenon, which is expected to increase in
252 frequency in the near future in Alpine streams (Brighenti et al., 2019), will represent a threat for Alpine
253 stream biodiversity.

254 Decomposing the total diversity into its taxonomic components allowed us to obtain a more
255 detailed view on which of the two main mechanisms, i.e. species filtering (alpha diversity) and biotic
256 homogenization (beta diversity components) (Socolar et al. 2016), and at which spatial scale, is affecting
257 macroinvertebrate assemblages. We could highlight that the overall decline in macroinvertebrate richness
258 is mainly due to local diversity reduction among and within reaches. In particular, at the reach scale, local
259 macroinvertebrate communities are impoverished and become more homogeneous. Conversely, we did
260 not record a significant difference in taxonomic variation among reaches between permanent and
261 intermittent sites. These results are partially supported by García-Roger (2013), who found that variation in
262 intermittent streams during the wet season is accounted by both the microhabitat (in systems with pools
263 during the dry season) and the stream (in systems where the riverbed completely dries during the dry
264 season) scale. The lack of taxonomic variation among reaches observed in our study might indicate that the
265 biotic homogenization of macroinvertebrate communities caused by flow intermittency, which is a
266 relatively new phenomenon in the investigated streams, has not become pervasive yet. However, this
267 result should be interpreted with caution as we here identified our taxa at the family to genus levels,
268 preventing to infer whether a homogenization process is ongoing at species level too.

269 When analysing the contribution of nestedness and turnover components, we could detect that
270 variation among local communities is mainly due to richness difference in both permanent and intermittent
271 reaches, while taxa replacement seemingly played a minor role, in accordance with the fact we sampled
272 during the base flow regime. In fact, β -diversity should be low during non-drought phases and dominated
273 by nestedness as organisms experience favourable high hydrological connectivity between sites (Larned et
274 al., 2010). Conversely, the turnover component of dissimilarity increases with increasing intermittency (
275 Tornés & Ruhì, 2013; Piano et al., 2017; Aspin et al., 2018). Although we did not record significant
276 differences in the relative contribution of these two components in permanent and intermittent reaches,
277 we observed a slightly higher contribution of turnover in intermittent than permanent sites. Despite some
278 contrasting results found in literature (Datry et al., 2014), this is in agreement with Rolls et al. (2016) who
279 showed a dominance of turnover processes in the dissimilarity partitioning on macroinvertebrates and fish
280 subjected to intermittent flow. This may be due to the environmental filter imposed by physical and

281 chemical alterations associated with stream drying—lentification with the creation of isolated pools, where
282 organisms experience oxygen reduction and increase in nutrient concentration, fine sediment deposition
283 and biotic interactions (Lake, 2003)—that causes the replacement of the most sensitive taxa by tolerant
284 ones (Dolédec et al., 2017).

285 Functional diversity patterns showed that reaches experiencing flow intermittency are less
286 functionally diverse than permanent ones, with communities in intermittent sites showing lower diversity,
287 variability and a less homogeneous distribution of functional traits. Opposite results were obtained by
288 Belmar et al. (2019) in Mediterranean streams, where they did not observe any significant effect of flow
289 intermittence on functional diversity. These different outcomes may indicate that flow intermittency exerts
290 a stronger environmental pressure on benthic invertebrate communities in mountainous lotic ecosystems
291 than in Mediterranean streams. In fact, the overall pool of macroinvertebrates in Alpine areas lacks
292 functional traits to cope with flow intermittency, which are present in their counterpart in the
293 Mediterranean region, where the drying phase is a natural part of the stream hydrological cycle (Leigh et
294 al., 2016; Vannucchi et al., 2013). On the other hand, the analysis of functional β -diversity showed how
295 intermittent reaches experience trait overdispersion, and this is mainly due to a net difference in the
296 amount of space enclosed by the trait space rather than replacements of traits. In other words,
297 communities in intermittent streams are composed by taxa that show a subset of the entire pool of
298 functional traits, but this subset is different depending on the reach considered. Similar results were
299 obtained by Dolédec et al. (2017), as they observed that Mediterranean taxa, which have to cope with
300 higher hydrological variability, may perform more diverse functional roles than temperate taxa and
301 intermittent streams show higher functional variability than permanent systems within the Mediterranean
302 area. This may be due to the fact that intermittent sites are colonized by species with traits adapted to flow
303 intermittence even during baseflow conditions leading to the coexistence of taxa displaying a wide range of
304 traits. The variation in trait composition among reaches can be explained by the high variability of lotic
305 systems experiencing flow intermittency, which in turn affect the community assemblages through
306 stochastic extinctions and colonization (Datry et al., 2016).

307 The analysis of taxa contribution to the functional diversity allowed us to highlight which trait
308 combination is less typical in permanent and intermittent reaches. We observed that Tipulidae shows
309 unique traits for the environmental conditions in permanent sites, whereas *Perla* shows unique traits for
310 environmental conditions in intermittent sites, confirming the results that we obtained in related studies
311 (Doretto et al., 2018; Doretto et al., 2020b; Piano et al., 2019a). In other words, Tipulidae and *Perla* traits
312 are atypical (and possibly maladaptive) for permanent and intermittent stretches, respectively. Tipulidae is
313 a monovoltine, crawler taxon, living in slow flowing waters, such as side-arms or near the banks, whose
314 larvae have a dormancy phase, whereas *Perla* is a large-sized, monovoltine, crawler taxon, with aquatic
315 respiration, preferring oligotrophic, cold, fast-flowing waters (Usseglio-Polatera et al., 2000; Merritt et al.,
316 2017). Large semivoltine taxa with long larval life are expected to be unable to survive in periodically dry
317 environments, while fast growing, plurivoltine organisms are adapted to complete their life cycle in less
318 than one year, emerging just before the loss of surface water (Corbet et al., 2006; López-Rodríguez et al.,
319 2009a, b; Bonada & Dolédec, 2018). Small body size is another trait that increases the survival in
320 intermittent reaches as reduced dimensions allow fast development and population growth (Bonada et al.
321 2007). The shift from rheophilous to lentic taxa has also been recorded in literature (e.g. Acuña et al., 2005;
322 Pace et al., 2013; Chessmann, 2015), as flow intermittency is usually associated with lentication and
323 consequent oxygen reduction (Lake, 2003), which may compromise the survival of many taxa, especially in
324 mountainous lotic ecosystems where organisms have to be adapted to high flow velocities and the
325 pronounced oligotrophy (Fenoglio et al., 2010). In addition, it should be noted that, compared to other
326 exclusively lotic organisms, Tipulidae larvae can survive even in terrestrial habitats if characterized by
327 sufficient moisture (Pritchard, 1983), thus being naturally tolerant to intermittent regimes.

328 Overall, our results demonstrate that benthic invertebrate biodiversity is threatened at both local
329 and regional scale, whereas the biotic homogenization process is still ongoing. We can therefore suggest
330 that macroinvertebrate communities are currently facing an ongoing “mediterraneization process”, which
331 is still in progress, and the results of the analyses of functional diversity support this hypothesis. In fact, a
332 more in-depth analysis of their functional diversity revealed that Alpine streams are currently facing the
333 extinction of rheophilous organisms, typical of fast-flowing, well-oxygenated waters, which are gradually

334 replaced by lentic taxa, with life-traits that confer adaptations to flow intermittency. As the frequency and
335 magnitude of anthropogenic and climatic droughts in the mountain areas are predicted to increase
336 (Brighenti et al., 2019), further studies should be performed in the next future to prevent the loss of
337 specialized benthic species, their functional diversity, and the key role they play in the riverine ecological
338 network.

339

340 **Acknowledgements**

341 We thank M. Apostolo, R. Bolpagni, M. C. Bruno, G. Burgazzi, C. Garetto, A. Laini, D. Melchio, D.
342 Morandini, D. Nizzoli and B. Palmia for their assistance and contribution during the field and laboratory
343 activities. Marco Baltieri (ATAAI - Associazione Tutela Ambienti Acquatici e Ittiofauna) is greatly
344 acknowledged for his help in the identification of temporary sampling stretches. This work was realized
345 within the framework of the PRIN NOACQUA “Risposte di comuNità e processi ecOsistemici in corsi
346 d'ACQUA soggetti a intermittenza idrologica” - code 201572HW8F, funded by the Italian Ministry of
347 Education, University and Research and is part of the research fellowship “Aquatic invertebrate
348 communities as sentinels of climate change in Italian Alpine streams” funded by Fondazione CRT and of the
349 activities of ALPSTREAM, a research center financed by FESR, Interreg Alcotra 2014-2020, EcO Project of the
350 Piter Terres Monviso.

351

352 **Data Availability Statement**

353 The data that support the findings of this study are available from the corresponding author upon
354 reasonable request.

355

356 **Conflict of Interest Statement**

357 The authors declare no conflict of interest.

358

359

REFERENCES

- 360 Acuña, V., Muñoz, I., Giorgi, A., Omella, M., Sabater, F., & Sabater, S. (2005). Drought and post-
361 drought recovery cycles in an intermittent Mediterranean stream: structural and functional aspects. *Journal*
362 *of the North American Benthological Society*, **24(4)**, 919-933. <https://doi.org/10.1899/04-078.1>
- 363 Agenzia Regionale per la Protezione dell'Ambiente (2013) Idrologia in Piemonte nel 2012. Regione
364 Piemonte 26 pp.
- 365 Anderson, M. J., Crist, T. O., Chase, J. M., Vellend, M., Inouye, B. D., Freestone, A. L., ... Swenson, N.
366 J. (2011). Navigating the multiple meanings of β diversity: a roadmap for the practicing ecologist. *Ecology*
367 *Letters*, **14(1)**, 19-28. <https://doi.org/10.1111/j.1461-0248.2010.01552.x>
- 368 Aspin, T. W., Matthews, T. J., Khamis, K., Milner, A. M., Wang, Z., O'callaghan, M. J., & Ledger, M. E.
369 (2018). Drought intensification drives turnover of structure and function in stream invertebrate
370 communities. *Ecography*, **41(12)**, 1992-2004. <https://doi.org/10.1111/ecog.03711>
- 371 Belmar, O., Bruno, D., Guareschi, S., Mellado-Díaz, A., Millán A., & Velasco, J. (2019). Functional
372 responses of aquatic macroinvertebrates to flow regulation are shaped by natural flow intermittence in
373 Mediterranean streams. *Freshwater Biology* **64(5)**, 1064-1077. <https://doi.org/10.1111/fwb.13289>
- 374 Blonder, B. (with contribution of Harris, D. J.) (2019). hypervolume: High Dimensional Geometry
375 and Set Operations Using Kernel Density Estimation, Support Vector Machines, and Convex Hulls. R package
376 version 2.0.12. <https://CRAN.R-project.org/package=hypervolume>
- 377 Blonder, B., Lamanna, C., Violle, C., & Enquist, B. J. (2014). The n-dimensional hypervolume. *Global*
378 *Ecology and Biogeography*, **23(5)**, 595–609. <https://doi.org/10.1111/geb.12146>
- 379 Blonder, B., Morrow, C. B., Maitner, B., Harris, D. J., Lamanna, C., Violle, C., ... & Kerkhoff, A. J.
380 (2018). New approaches for delineating n-dimensional hypervolumes. *Methods in Ecology and Evolution*,
381 **9(2)**, 305-319. <https://doi.org/10.1111/2041-210X.12865>

382 Bogan, M. T., Boersma, K. S., & Lytle, D. A. (2015). Resistance and resilience of invertebrate
383 communities to seasonal and suprasedasonal drought in arid-land headwater streams. *Freshwater Biology*,
384 **60**(12), 2547-2558. <https://doi.org/10.1111/fwb.12522>

385 Bogan, M. T., Hwan, J. L., Cervantes-Yoshida, K., Ponce, J., & Carlson, S. M. (2017). Aquatic
386 invertebrate communities exhibit both resistance and resilience to seasonal drying in an intermittent
387 coastal stream. *Hydrobiologia*, **799**(1), 123-133. <https://doi.org/10.1007/s10750-017-3205-4>

388 Bonada, N., Rieradevall, M., & Prat, N. (2007). Macroinvertebrate community structure and
389 biological traits related to flow permanence in a Mediterranean river network. *Hydrobiologia*, **589**(1), 91-
390 106. <https://doi.org/10.1007/s10750-007-0723-5>

391 Bonada, N., & Dolédec, S. (2018). Does the Tachet trait database report voltinism variability of
392 aquatic insects between Mediterranean and Scandinavian regions?. *Aquatic Sciences*, **80**(1), 7.
393 <https://doi.org/10.1007/s00027-017-0554-z>

394 Brighenti, S., Tolotti, M., Bruno, M. C., Wharton, G., Pusch, M. T., & Bertoldi, W. (2019). Ecosystem
395 shifts in Alpine streams under glacier retreat and rock glacier thaw: A review. *Science of the Total*
396 *Environment*, **675**, 542-559. <https://doi.org/10.1016/j.scitotenv.2019.04.221>

397 Brown, L. E., Khamis, K., Wilkes, M., Blaen, P., Brittain, J. E., Carrivick, J. L., ... Hainie, S. (2018).
398 Functional diversity and community assembly of river invertebrates show globally consistent responses to
399 decreasing glacier cover. *Nature Ecology & Evolution*, **2**(2), 325–333. <https://doi.org/10.1038/s41559-017-0426-x>

401 Bruno, D., Belmar, O., Maire A., Morel, A., Dumont, B., & Datry, T. (2019). Structural and functional
402 responses of invertebrate communities to climate change and flow regulation in alpine catchments. *Global*
403 *Change Biology*, **25**, 1612-1628. <https://doi.org/10.1111/gcb.14581>

404 Campaioli, S., Ghetti, P. F., Minelli A., & Ruffo, S. (1994). Manuale per il riconoscimento dei
405 macroinvertebrati delle acque dolci italiane (Vol. I). Trento, IT: Provincia Autonoma di Trento.

406 Campaioli, S., Ghetti, P. F., Minelli A., & Ruffo, S. (1999). Manuale per il riconoscimento dei
407 macroinvertebrati delle acque dolci italiane (Vol. II). Trento, IT: Provincia Autonoma di Trento.

408 Cardoso, P., Rigal, F., Carvalho, J. C., Fortelius, M., Borges, P. A. V., Podani, J., & Schmera, D. (2014).
409 Partitioning taxon, phylogenetic and functional beta diversity into replacement and richness difference
410 components. *Journal of Biogeography*, **41**(4), 749–761. <https://doi.org/10.1111/jbi.12239>

411 Cardoso, P., Mammola, S., Rigal, F., & Carvalho, J. C. (2020). BAT: Biodiversity Assessment Tools. R
412 package version 2.0.0. <https://CRAN.R-project.org/package=BAT>

413 Carvalho, J. C., & Cardoso, P. (2018). Decomposing the causes for niche differentiation between
414 species using hypervolumes. *BioRxiv*, 485920. <https://doi.org/10.1101/485920>

415 Chessmann, B. C. (2015). Relationships between lotic macroinvertebrate traits and responses to
416 extreme drought. *Freshwater Biology*, **60**(1), 50-63. <https://doi.org/10.1111/fwb.12466>

417 Datry, T., Larned, S. T., & Tockner, K. (2014). Intermittent rivers: a challenge for freshwater ecology.
418 *BioScience*, **64**(3), 229-235. <https://doi.org/10.1093/biosci/bit027>

419 Datry, T., Moya, N., Zubieta, J., & Oberdorff, T. (2016a). Determinants of local and regional
420 communities in intermittent and perennial headwaters of the Bolivian Amazon. *Freshwater Biology*, **61**(8),
421 1335-1349. <https://doi.org/10.1111/fwb.12706>

422 Datry, T., Bonada, N., & Heino, J. (2016b). Towards understanding the organisation of
423 metacommunities in highly dynamic ecological systems. *Oikos*, **125**(2), 149-159.
424 <https://doi.org/10.1111/oik.02922>

425 Dolédec, S., Tilbian, J., & Bonada, N. (2017). Temporal variability in taxonomic and trait
426 compositions of invertebrate assemblages in two climatic regions with contrasting flow regimes. *Science of*
427 *the Total Environment*, **599**, 1912-1921. <https://doi.org/10.1016/j.scitotenv.2017.05.057>

428 Doretto, A., Piano, E., Falasco, E., Fenoglio, S., Bruno, M. C., & Bona, F. (2018). Investigating the role
429 of refuges and drift on the resilience of macroinvertebrate communities to drying conditions: an
430 experiment in artificial streams. *River Research and Applications*, **34**(7), 777-785.
431 <https://doi.org/10.1002/rra.3294>

432 Doretto, A., Bo, T., Bona, F., & Fenoglio, S. (2020a). Efficiency of Surber net under different flow and
433 substrate conditions: insights for macroinvertebrates sampling and river biomonitoring. *Knowledge and
434 Management of Aquatic Ecosystems*, 421, 10. <https://doi.org/10.1051/kmae/2020001>

435 Doretto, A., Bona, F., Falasco, E., Morandini, D., Piano, E., & Fenoglio, S. (2020b). Stay with the flow:
436 How macroinvertebrate communities recover during the rewetting phase in Alpine streams affected by an
437 exceptional drought. *River Research and Applications*, **36**(1), 91-101. <https://doi.org/10.1002/rra.3563>

438 Elias, C. L., Calapez, A. R., Almeida, S. F., & Feio, M. J. (2015). From perennial to temporary streams:
439 an extreme drought as a driving force of freshwater communities' traits. *Marine and Freshwater Research*,
440 **66**(5), 469-480. <https://doi.org/10.1071/MF13312>

441 Fenoglio, S., Bo, T., Cucco, M., & Malacarne, G. (2007). Response of benthic invertebrate
442 assemblages to varying drought conditions in the Po river (NW Italy). *Italian Journal of Zoology*, **74**(2), 191-
443 201. <https://doi.org/10.1080/11250000701286696>

444 Fenoglio, S., Bo, T., Cucco, M., Mercalli, L., & Malacarne, G. (2010). Effects of global climate change
445 on freshwater biota: A review with special emphasis on the Italian situation. *Italian Journal of Zoology*,
446 **77**(4), 374-383. <https://doi.org/10.1080/11250000903176497>

447 García-Roger, E. M., Sánchez-Montoya, M. D. M., Cid, N., Erba, S., Karaouzas, I., Verkaik, I., ... &
448 Prat, N. (2013). Spatial scale effects on taxonomic and biological trait diversity of aquatic
449 macroinvertebrates in Mediterranean streams. *Fundamental and Applied Limnology/Archiv für
450 Hydrobiologie*, **183**(2), 89-105. <https://doi.org/10.1127/1863-9135/2013/0429>

451 Gobiet, A., Kotlarski, S., Beniston, M., Heinrich, G., Rajczak, J., & Stoffel, M. (2014). 21st century
452 climate change in the European Alps—a review. *Science of the Total Environment*, **493**, 1138-1151.
453 <https://doi.org/10.1016/j.scitotenv.2013.07.050>

454 Hansen, J., Nazarenko, L., Ruedy, R., Sato, M., Willis, J., Del Genio, A., ..., Tausnev, N. (2005). Earth's
455 energy imbalance: Confirmation and implications. *Science*, **308**(5727), 1431-1435.
456 <https://doi.org/10.1126/science.1110252>

457 Heino, J., Virkkala, R., & Toivonen, H. (2009). Climate change and freshwater biodiversity: detected
458 patterns, future trends and adaptations in northern regions. *Biological Reviews*, **84**(1), 39-54.
459 <https://doi.org/10.1111/j.1469-185X.2008.00060.x>

460 Jarzyna, M. A., & Jetz, W. (2018). Taxonomic and functional diversity change is scale dependent.
461 *Nature Communication*, **9**(1), 1-8. <https://doi.org/10.1038/s41467-018-04889-z>

462 Lake P.S. (2003). Ecological effects of perturbation by drought in flowing waters. *Freshwater
463 Biology*, **48**, 1161-1172. <https://doi.org/10.1046/j.1365-2427.2003.01086.x>

464 Larned, S. T., Datry, T., Arscott, D. B., & Tockner, K. (2010). Emerging concepts in temporary-river
465 ecology. *Freshwater Biology*, **55**(4), 717-738. <https://doi.org/10.1111/j.1365-2427.2009.02322.x>

466 Legendre, P., & Legendre, L. (1998). Numerical Ecology, Volume 24. Developments in
467 Environmental Modelling.

468 Leigh, C., & Datry, T. (2017). Drying as a primary hydrological determinant of biodiversity in river
469 systems: A broad-scale analysis. *Ecography*, **40**(4), 487-499. <https://doi.org/10.1111/ecog.02230>

470 Leigh, C., Bonada, N., Boulton, A. J., Hugueny, B., Larned, S. T., Vander Vorste, R., & Datry, T. (2016).
471 Invertebrate assemblage responses and the dual roles of resistance and resilience to drying in intermittent
472 rivers. *Aquatic Sciences*, **78**(2), 291-301. <https://doi.org/10.1007/s00027-015-0427-2>

473 López-Rodríguez, M. J., Tierno de Figueroa, J. M., Fenoglio, S., Bo, T., & Alba-Tercedor, J. (2009a).
474 Life strategies of 3 Perlodidae species (Plecoptera) in a Mediterranean seasonal stream in southern Europe.
475 *Journal of the North American Benthological Society*, **28**(3), 611-625. <https://doi.org/10.1899/08-105.1>

476 López-Rodríguez M. J., Tierno de Figueroa, J. M., & Alba-Tercedor, J. (2009b). The life history of
477 *Serratella ignita* (Poda, 1761)(Insecta: Ephemeroptera) in a temporary and permanent Mediterranean
478 stream. *Aquatic Sciences*, **71**(2), 179-188. <https://doi.org/10.1007/s00027-009-9211-5>

479 López-Rodríguez, M. J., Muñoz, C. M., Ripoll-Martín, E., & de Figueroa, J. M. T. (2019). Effect of
480 shifts in habitats and flow regime associated to water diversion for agriculture on the macroinvertebrate
481 community of a small watershed. *Aquatic Ecology*, **53**(3), 1-13. [https://doi.org/10.1007/s10452-019-09703-](https://doi.org/10.1007/s10452-019-09703-6)
482 6

483 MacLean, S. A., & Beissinger, S. R. (2017). Species' traits as predictors of range shifts under
484 contemporary climate change: A review and meta-analysis. *Global Change Biology*, **23**(10), 4094–4105.
485 <https://doi.org/10.1111/gcb.13736>

486 Mammola, S., & Cardoso, P. (2020). Functional diversity metrics using kernel density n-dimensional
487 hypervolumes. *Methods in Ecology and Evolution*, early view.

488 McGregor, G., Petts, G. E., Gurnell, A. M., & Milner, A. M. (1995). Sensitivity of alpine stream
489 ecosystems to climate change and human impacts. *Aquatic Conservation: Marine and Freshwater*
490 *Ecosystems*, **5**, 233–247. <http://dx.doi.org/10.1002/aqc.3270050306>

491 Merritt, R. W., Cummins, K. W., & Berg, M. B. (2017). Trophic relationships of macroinvertebrates.
492 In *Methods in Stream Ecology, Volume 1 (Third Edition)* (pp. 413-433).

493 Milner, A. M., Khamis, K., Battin, T. J., Brittain, J. E., Barrand, N. E., Füreder, L., ... & Hodson, A. J.
494 (2017). Glacier shrinkage driving global changes in downstream systems. *Proceedings of the National*
495 *Academy of Sciences*, **114**(37), 9770–9778. <https://doi.org/10.1073/pnas.1619807114>

496 Moog, O., Schmidt-Kloiber, A., Ofenböck, T., & Gerritsen, J. (2004). Does the ecoregion approach
497 support the typological demands of the EU 'Water Framework Directive'? In: Hering, D., Verdonschot, P. F.
498 M., Moog, O., & Sandin, L. (eds). Integrated Assessment of Running Waters in Europe. Developments in
499 Hydrobiology, vol 175. Springer, Dordrecht, pp 21-33. https://doi.org/10.1007/978-94-007-0993-5_2

500 Pace, G., Bonada, N., & Prat, N. (2013). Long-term effects of climatic–hydrological drivers on
501 macroinvertebrate richness and composition in two Mediterranean streams. *Freshwater Biology*, **58(7)**,
502 1313-1328. <https://doi.org/10.1111/fwb.12129>

503 Pavoine, S., & Bonsall, M. B. (2011). Measuring biodiversity to explain community assembly: a
504 unified approach. *Biological Reviews*, **86(4)**, 792–812. <https://doi.org/10.1111/j.1469-185X.2010.00171.x>

505 Pavoine, S., Ollier, S., & Dufour, A.-B. (2005). Is the originality of a species measurable? *Ecology*
506 *Letters*, **8(6)**, 579–586. doi:10.1111/j.1461-0248.2005.00752.x

507 Pederson, G. T., Gray, S. T., Ault, T., Marsh, W., Fagre, D. B., Bunn, A. G., ..., Graumlich, L. J. (2011).
508 Climatic controls on the snowmelt hydrology of the northern Rocky Mountains. *Journal of Climatology*,
509 **24(6)**, 1666-1687. <https://doi.org/10.1175/2010JCLI3729.1>

510 Petchey, O. L., & Gaston, K. J. (2002). Functional diversity (FD), species richness and community
511 composition. *Ecology Letters*, **5(3)**, 402–411. <https://doi.org/10.1046/j.1461-0248.2002.00339.x>

512 Piano, E., Falasco, E., & Bona, F. (2017). How does water scarcity affect spatial and temporal
513 patterns of diatom community assemblages in Mediterranean streams?. *Freshwater Biology*, **62(7)**, 1276-
514 1287. <https://doi.org/10.1111/fwb.12944>

515 Piano, E., Doretto, A., Falasco, E., Fenoglio, S., Gruppuso, L., Nizzoli, D., Viaroli P., & Bona, F.
516 (2019a). If Alpine streams run dry: the drought memory of benthic communities. *Aquatic Sciences*, **81(2)**,
517 32. <https://doi.org/10.1007/s00027-019-0629-0>

518 Piano, E., Doretto, A., Falasco, E., Gruppuso, L., Fenoglio, S., & Bona, F. (2019b). The role of
519 recurrent dewatering events in shaping ecological niches of scrapers in intermittent Alpine streams.
520 *Hydrobiologia*, **841**(1), 177-189. <https://doi.org/10.1007/s10750-019-04021-2>

521 Pritchard, G. (1983). Biology of Tipulidae. *Annual Review of Entomology*, **28**(1), 1-22.

522 R Core Team (2019). R: A language and environment for statistical computing. R Foundation for
523 Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.

524 Rauscher, S. A., Pal, J. S., Diffenbaugh, N. S., & Benedetti, M.M. (2008). Future changes in
525 snowmelt-driven runoff timing over the western US. *Geophysical Research Letters*, **35**(16).
526 <https://doi.org/10.1029/2008GL034424>

527 Rolls R. J., Heino J. & Chessman B. C. (2016). Unravelling the joint effects of flow regime, climatic
528 variability and dispersal mode on beta diversity of riverine communities. *Freshwater Biology*, **61**(8), 1350-
529 1364. <https://doi.org/10.1111/fwb.12793>

530 Samways, M. J., Barton, P. S., Birkhofer, K., Chichorro, F., Deacon, C., Fartmann, T., ... & Hill, M. J.
531 (2020). Solutions for humanity on how to conserve insects. *Biological Conservation*, 108427.

532 Scheffer, M., Carpenter, S., Foley, J. A., Folke, C., & Walker, B. (2001). Catastrophic shifts in
533 ecosystems. *Nature*, **413**(6856), 591-596.

534 Socolar, J. B., Gilroy, J. J., Kunin, W. E., & Edwards, D. P. (2016). How should beta-diversity inform
535 biodiversity conservation? *Trends in Ecology and Evolution*, **31**(1), 67-80.
536 <https://doi.org/10.1016/j.tree.2015.11.005>

537 Tornés E. & Ruhì A. (2013). Flow intermittency decreases nestedness and specialization of diatom
538 communities in Mediterranean rivers. *Freshwater Biology*, **58**, 2555-2566.
539 <https://doi.org/10.1111/fwb.12232>

540 Usseglio-Polatera, P., Bournaud, M., Richoux, P., & Tachet, H. (2000). Biological and ecological traits
541 of benthic freshwater macroinvertebrates: relationships and definition of groups with similar traits.
542 *Freshwater Biology*, 43(2), 175-205. <https://doi.org/10.1046/j.1365-2427.2000.00535.x>

543 Vannucchi, P. E., López-Rodríguez, M. J., de Figueroa, J. M. T., & Gaino, E. (2013). Structure and
544 dynamics of a benthic trophic web in a Mediterranean seasonal stream. *Journal of Limnology*, 72(3), 51.
545 <https://doi.org/10.4081/jlimnol.2013.e51>

546 Van Looy, K., Tonkin, J. D., Flourey, M., Leigh, C., Soininen, J., Larsen, S., ... & Datry, T. (2019). The
547 three Rs of river ecosystem resilience: Resources, recruitment, and refugia. *River Research and*
548 *Applications*, 35(2), 107-120. <https://doi.org/10.1002/rra.3396>

549 Violle, C., Thuiller, W., Mouquet, N., Munoz, F., Kraft, N. J. B., Cadotte, M. W., ... Mouillot, D. (2017).
550 Functional Rarity: The Ecology of Outliers. *Trends in Ecology and Evolution*, 32(5), 356–367.
551 <https://doi.org/10.1016/j.tree.2017.02.002>

552 Wasson, J. G., Chandesris, A., Bautista, A. G., Pella, H., & Villeneuve, B. (2007). REBECCA,
553 Relationships between ecological and chemical status of surface waters. European Hydro-Ecoregions. EU
554 6th Framework Programme. Cemagref Institute, Lyon, France, 43 pp.

555 Whitehead, P. G., Wilby, R. L., Battarbee, R. W., Kernan, M., & Wade, A. J. (2009). A review of the
556 potential impacts of climate change on surface water quality. *Hydrological Science Journal*, 54(1), 101-123.
557 <https://doi.org/10.1623/hysj.54.1.101>

558 Wilhite, D. A. (2000). Drought as a natural hazard: concepts and definitions. Published in *Drought: A*
559 *Global Assessment*, Vol. I, edited by Donald A. Wilhite, chap. 1, pp. 3–18 (London: Routledge, 2000)

560 Wohl, E. (2006). Human impacts to mountain streams. *Geomorphology*, 79(3-4), 217-248.
561 <https://doi.org/10.1016/j.geomorph.2006.06.020>

562 Zuur A. F., Ieno E. N., Walker N. J., Saveliev A. A. & Smith G. M. (2009). Mixed effect models and
563 extensions in ecology with R. Springer, Berlin: 574 pp.

564

Tables

565

566

567

568

569

Table 1. Mean values with standard deviation in brackets of functional diversity measures in permanent (UP) and intermittent (DOWN) sites, and estimated parameters (Est), with standard errors (SE) in brackets, and p-values (P) of LMMs performed on functional metrics (FD = functional α -diversity; Fdis = functional dispersion; Feve = functional evenness) to test for differences between UP and DOWN sites. All comparisons are significant.

Functional measures	UP	DOWN	BEst \pm SE	P
FD	0.20 (\pm 0.10)	0.13 (\pm 0.13)	-0.071 (\pm 0.014)	< 0.001
Fdis	0.53 (\pm 0.04)	0.49 (\pm 0.06)	-0.053 (\pm 0.006)	< 0.001
Feve	0.17 (\pm 0.06)	0.13 (\pm 0.09)	-0.039 (\pm 0.014)	0.007
β -diversity (β -replacement + β - richness)	0.63 (\pm 0.14)	0.74 (\pm 0.15)	0.119 (\pm 0.003)	< 0.001
β -replacement	0.25 (\pm 0.16)	0.19 (\pm 0.18)	-0.062 (\pm 0.004)	< 0.001
β -richness	0.38 (\pm 0.24)	0.56 (\pm 0.29)	0.181 (\pm 0.006)	< 0.001

570

571 **Figure captions**

572 Figure 1. Schematic overview of the calculated diversity components (α_w = local diversity within
573 reaches; α_a = local diversity among reaches; β_w = variation among local communities within reaches; β_a =
574 variation among local communities among reaches; γ_w = total diversity within reaches; γ_a = total diversity
575 among reaches) to test the effect of flow intermittency (UP = orange; DOWN = light blue) within- (reach
576 scale) and among stretches (basin scale).

577 Figure 2. a) diversity partitioning of total diversity in the study area into its within- and among
578 reaches components in UP and DOWN reaches (α_w = dark grey; β_w = medium grey; β_a = light grey); b)
579 decomposition of β -diversity into its nestedness (dark grey) and turnover (light grey) components in in UP
580 and DOWN reaches.

581 Figure 3. a–c) Functional richness (a), dispersion (b) and evenness (c) of benthic invertebrate
582 communities in permanent (UP) and intermittent (DOWN) reaches; d–f) Density of functional β -diversity
583 values for pairwise comparison of communities in permanent (UP) and intermittent (DOWN) reaches. Total
584 functional β -diversity (d) is split in two components: β -replacement (e) is turnover in functional
585 composition explained by replacement of space between hypervolumes, and β -richness (f) is the turnover
586 in functional composition explained by net differences between the amount of space enclosed by each
587 hypervolume (Carvalho & Cardoso, 2018; Mammola & Cardoso, 2020).