

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Is vitamin B12 serum level involved in the spread of *Helicobacter pylori*?

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1717381> since 2019-11-24T12:10:54Z

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Is vitamin B12 serum level involved in the spread of *Helicobacter pylori*?

Ramy Ibrahim Kamal Jouness¹, Marco Astegiano²

¹ Chair of Gastroenterology, University of Turin, Turin, Italy

²Department of Gastroenterology and Hepatology, San Giovanni Antica Sede Hospital, Turin, Italy

Key words: vitamin B12; *Helicobacter pylori*; gastritis.

Address reprint requests: Department of Gastroenterology and Hepatology, Città della Salute e della Scienza, Molinette Hospital, SGAS, Via Cavour 31, 10123 Torino, Italy

Dear Sir,

the relationship between *Helicobacter pylori* (*H.pylori*) and the deficiency of important and vital elements, including iron, vitamins and other micronutrients has been reported in a great number of articles worldwide.¹ The main pathogenetic mechanism responsible for these manifestations is the modification of the gastric environment that affects the absorption of both nutrients and drugs.²

In a recent interesting study of Moosazadeh et al., an important relationship between the prevalence of *H.pylori* and serum levels of vitamin B12 in women was found, showing that the prevalence of this infection was significantly higher in patients with serum lower levels of vitamin B12, compared to individuals with normal levels. The authors concluded that “vitamin B12 serum level is a determining factor in the spread of *H.pylori* and there is an inverse relationship between the two variables”.³

Apparently, there is no consistent plausibility to support the former part of this conclusion. Literature data have reported that *H.pylori* has an important role in the pathogenesis of vitamin B12 deficiency and pernicious anemia.⁴ Various studies showed that a result of *H.pylori*-induced gastritis is the damage of parietal cells which are important for the production of intrinsic factor, essential for vitamin B12 absorption, and eradication of the infection has been shown to result in improved blood levels of vitamin B12.⁵ It was also demonstrated that antacid drugs used by infected symptomatic subjects and the modification of the intragastric pH caused by *H.pylori* are the main factors of malabsorption of vitamin B12. Annibale *et al.* described the presence of *H.pylori* -related gastritis as the unique pathological finding in 57.1% of patients with macrocytic anemia caused by B12 deficiency.⁶

In conclusion, it is clear from all data mentioned above that an inverse relationship between *H.pylori* and B12 serum levels is a real fact, but evidences from literature and clinical experience show a fundamental role of *H.pylori*, as an aetiologic factor, in determining vitamin B12 levels instead of the role of B12 vitamin in the spread of *H.pylori*, as Moosazadeh et al. concluded in their study.³

References

1. Franceschi F, Annalisa T, Teresa DR, Giovanna D, Ianiro G, Franco S et al. Role of *Helicobacter pylori* infection on nutrition and metabolism. World J Gastroenterol 2014;20:12809-17.
2. Adriani A, Repici A, Hickman I, Pellicano R. *Helicobacter pylori* infection and respiratory diseases: actual data and directions for future studies. Minerva Med 2014;105:1-8.

3. Moosazadeh M, Shahbazi N, Mahmoodi G, Abdi Joubari H, Mahdavi SA, Babania Rostam FZ et al. The relationship between the prevalence of *Helicobacter pylori* and serum levels of vitamin B12 in women 10-49 years. *Minerva Gastroenterol Dietol* 2015;61:131-5.
4. Perez-Perez GI. Role of *Helicobacter pylori* infection in the development of pernicious anemia. *Clin Infect Dis* 1997;25:1-39.
5. Kaptan K, Beyan C, Ural AU, Cetin T, Aycu F, Gulsen M et al. *Helicobacter pylori*: is it a novel causative agent in Vitamin B12 deficiency? *Arch Intern Med* 2000;160:12-6.
6. Annibale B, Capurso G, Delle Fave G. Consequences of *Helicobacter pylori* infection on the absorption of micronutrients. *Dig Liver Dis* 2002;34 (Suppl 2):S72-S77.