

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Evidence of genetic monogamy in the lemur Indri (*Indri indri*)

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1704214> since 2020-12-22T10:43:16Z

Published version:

DOI:10.1002/ajp.22993

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



Evidence of genetic monogamy in the lemur Indri (Indri indri).

Journal:	<i>American Journal of Primatology</i>
Manuscript ID	AJP-18-0248.R2
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Bonadonna, Giovanna; Universita degli Studi di Torino, Dept. Life Science and Systems Biology Torti, Valeria; University of Torino, Dept. Life Sciences and Systems Biology De Gregorio, Chiara; Universita degli Studi di Torino, Life Science and Systems Biology Valente, Daria; Universita degli Studi di Torino, Life Science and Systems Biology Randrianarison, Rose Marie; GERP (Groupe d'etude et de recherche sur les primates du Madagascar) Pozzi, Luca; University of Texas San Antonio, Anthropology Gamba, Marco; Universita degli Studi di Torino Giacoma, Cristina; Universita degli Studi di Torino, Life Science and Systems Biology
Indicate which taxonomic group was the subject of your study (select all that apply or type another option)::	Prosimians, Indri indri
Keywords:	Extra Pair Paternity, Extra Pair Copulation, Primates, Genetic Monogamy, Indri

SCHOLARONE™
Manuscripts

1
2
3 **1 Evidence of genetic monogamy in the lemur Indri (*Indri indri*).**

4
5
6 **2 *A study based on paternity analysis and history of indri family groups in the Maromizaha***

7
8 **3 *New Protected Area, Madagascar***

9
10
11
12
13
14 5 Giovanna Bonadonna¹, Valeria Torti¹, Chiara De Gregorio¹, Daria Valente¹, Rose Marie

15
16 6 Randrianarison,² Luca Pozzi³, Marco Gamba¹, Cristina Giacomini¹

17
18
19 7 ¹ Department of Life Sciences and Systems Biology, University of Torino, Via Accademia

20
21
22 8 Albertina 13, 10123 Torino, Italy

23
24
25 9 ² GERP (Group d'Etude et de Recherche sur les Primates de Madagascar), Fort Duchesne,

26
27
28 10 Antananarivo 101, Madagascar

29
30
31 11 ³ Department of Anthropology, The University of Texas at San Antonio, One UTSA Circle, San

32
33
34 12 Antonio, TX 78249, United States

35
36 **13 Abstract**

37
38
39 14 Monogamy is a rare strategy among mammals but relatively common among primates.

40
41
42 15 The study of evolution of monogamy in mammals and primates is lacking empirical studies

43
44
45 16 that assess the relationship between a pair-living social organization and genetic

46
47
48 17 monogamy. Sexual or genetic monogamy can only be assessed by performing molecular

49
50
51 18 analyses and investigating rates of Extra-Pair Paternity (EPP). Studying the occurrence of

52
53
54 19 EPP can provide valuable insights into reproductive strategies and their adaptive value.

55
56
57 20 The indri is a pair-living primate that lives in stable groups. Their social units are composed

58
59
60 21 of the reproductive pair and up to four more individuals, but extra-pair copulation (EPC)

1
2
3 22 can occur. This raises the question of whether this event may or may not lead to EPP.
4
5 23 Here, we investigated whether a pair-living social organization corresponds to genetic
6
7 24 monogamy in indris (*Indri indri*). We analyzed the paternity of 12 offspring from 7 pairs
8
9 25 using a set of six microsatellite loci on fecal samples (mean number of alleles 11.7 ± 1.8
10
11 26 (mean \pm SD). We found that in 92% of cases the genetic profile of the offspring matched
12
13 27 the paired male of the group for all the loci considered. In the only case of paternity
14
15 28 mismatch, the paternity assignment remained inconclusive. Our results show that in indri
16
17 29 genetic monogamy is the norm and support the hypothesis that pair-living social
18
19 30 organization is associated with low EPP rate. Also, our results are in contrast with the
20
21 31 hypothesis of infertility as a reason to engage in EPC for this species.
22
23
24
25
26
27

28 32 Keywords: Extra-Pair Copulation, Extra-Pair Paternity, primates, genetic monogamy, indri
29
30
31

32 33 **Research Highlights**

- 34 34 • In 92% of cases, the paired male of the group did not have any locus mismatch
35
36 35 with the offspring. In the only case of paternity mismatch, we were not able to
37
38 36 assign the sire identity.
- 39
40 37 • Our finding suggest that genetic monogamy is the norm in the indri, although EPC
41
42 38 can occasionally occur.
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

39 Introduction

40 Monogamy is a mating system with a social and a genetic component, both linked to the
41 social organization of a species. A social organization commonly associated with a
42 monogamous mating system is pair-living, defined as a permanent and continuous
43 association of one adult female and one adult male in space and time (Fuentes, 2000;
44 Kappeler & van Schaik, 2002).

45 Behavioral and genetic studies on pair-living species of birds and mammals
46 demonstrate that monogamy is a much more flexible and variable mating system in
47 respect to what was previously thought (Kleiman, 1977; Black, 1996; Tecot, Singletary &
48 Eadie, 2015), and different levels of monogamy can be recognized depending on the level
49 of analysis: a pair-living social organization does not necessarily imply sexual exclusivity
50 between partners (*sexual monogamy*), nor that all offspring are sired by the social father
51 (*genetic monogamy*) (Reichard, 2003). While behavioral studies can help in describing a
52 pair-living system (Kappeler & van Schaik, 2002), DNA analyses of the offspring and their
53 parents are necessary to provide evidence of genetic monogamy, which usually refers
54 more strictly to the mating system (Reichard, 2003). Genetic fingerprints of individuals
55 from wild populations can provide important insights into the relationship between pair-
56 living social organization and genetic monogamy, as well as a better understanding of the
57 evolution of this mating system in primates and mammals in general.

58 In the existing literature, two terms have been used to define evidence of offspring
59 sired by a different male than the social father: Extra Pair Paternity (EPP) and Extra Group
60 Paternity (EGP), referring to offspring fathered by a male outside the reproductive pair or

1
2
3 61 the group, respectively. EPP and EGP are interchangeable when considering pair-living
4
5 62 monogamous species (Isvaran & Clutton-Brock, 2007), as in our study; we decided to
6
7
8 63 apply the term EPP because we focus on the reproductive pair of the group.
9

10
11 64 Birds are the typical example of a taxon in which more than 90% of species were
12
13 65 considered sexually monogamous because of their pair-living organization (Lack, 1968).
14
15
16 66 However, molecular analysis have confirmed that EPP is widespread across birds (Griffith,
17
18 67 Owens & Thuman, 2002). Contrary to birds, pair living is very rare among mammals (3%)
19
20
21 68 (Kleiman, 1977), and, when present, it is more frequently related to genetic monogamy
22
23 69 (Lukas & Clutton-Brock, 2012, 2013). According to a study including 22 mammal species
24
25
26 70 generally considered to have a monogamous mating system, the variation of EPP among
27
28 71 species seems to be related to the social structure, with pair-living species having the
29
30
31 72 lowest rates of EPP (Cohas & Allainé, 2009). The strength of pair bonding has also been
32
33 73 suggested to possibly play a role (Cohas & Allainé, 2009; Clutton-Brock & Isvaran, 2006).
34
35

36 74 Among mammals, primates include the highest frequency of monogamous species
37
38 75 (29%; Lukas & Clutton-Brock, 2013), usually organized in pair-living groups with immature
39
40
41 76 or even mature offspring (Reichard & Boesch, 2003; Kappeler & van Schaik, 2002). Few
42
43
44 77 studies have investigated the genetic paternity in putatively monogamous primates (not
45
46 78 necessarily pair-living as we defined it) and some of them show that EPP occurs, including
47
48 79 in tarsiers (*Tarsius lariang*, Driller, Perwitasari-Farajallah, Zischler & Merker, 2009), lemurs
49
50
51 80 (*Hapalemur griseus alaotrensis*, Nievergelt, Mutschler, Feistner & Woodruff, 2002;
52
53 81 *Cheirogaleus medius*, Fietz et al. 2000; *Phaner furcifer*, Schülke, Kappeler & Zischler, 2004),
54
55
56 82 gibbons (*Hylobates lar*, Barelli et al. 2013; *Nomascus gabriellae*, Kenyon, Roos, Binh &
57
58
59
60

1
2
3 83 Chivers, 2011). To date, Azara's night monkey (*Aotus azarae*) and Muller's Bornean gibbon
4
5 84 (*Hylobates muelleri*) are the only two primate species in which no evidence of EPP was
6
7
8 85 found; however, in the latter case, only four offspring were tested (reviewed in Lambert,
9
10 86 Sabol & Solomon, 2018).

11
12
13 87 The occurrence of EPP has consequences on individuals' reproductive success. Several
14
15 88 non-exclusive hypotheses have been proposed to explain the presence of EPP, such as to
16
17 89 guard against infertility of the social mate, to increase the genetic diversity of the
18
19 90 offspring with consequences at population level, to provide "good genes" for the
20
21 91 offspring, and to gain direct benefit given by access to resources (Birkhead & Møller, 1992;
22
23 92 Griffith et al., 2002; Møller & Thornhill, 1998). However, there is still a lack of empirical
24
25 93 data to support these hypotheses (Griffith et al., 2002; Akçay & Roughgarden, 2007),
26
27 94 especially among primates.

28
29
30
31
32
33 95 The occurrence of EPP is directly related to the possibility of engagement in Extra-Pair
34
35 96 Copulation (EPC). A social context with easy access to potential mates within and between
36
37 97 groups increases the probability of EPCs and, as a consequence, can lead to higher rates of
38
39 98 EPP (Cohas & Allainé, 2009). In pair-living species, the opportunities of engaging in EPC are
40
41 99 lower because of more efficient control of the partner over potential mates. In territorial
42
43 100 species, the overlap between territories and the rate of intergroup encounters can
44
45 101 influence the chances of being in contact and interacting with potential mates, as
46
47 102 suggested in gibbons (*Hylobates lar*, Reichard & Sommer, 1997).

48
49
50
51 103 It seems that there is a relationship between genetic monogamy and the strength of
52
53 104 the pair bond (Huck, Fernandez-Duque, Babb, Schurr, 2014), and recent reviews proposed
54
55
56
57
58
59
60

1
2
3 105 the hypothesis that social organization and mate guarding are the main factors influencing
4
5 106 a monogamous mating system in primates (Lambert, Sabol & Solomon, 2018, Reichard
6
7
8 107 2018). In this study, we investigate the genetic relationship among reproductive pairs,
9
10 108 potential sires, and offspring to explore whether pair living is associated with genetic
11
12
13 109 monogamy in the indris (*Indri indri*).

14
15
16 110 Bonadonna and colleagues (2014) reported the first observation of EPC in indris
17
18 111 between a neighboring paired female and a paired male within the same population of
19
20
21 112 this study. This event raised the question of whether EPC may lead to EPP; our study aims
22
23 113 to address this question with subsequent eventual reconsideration of the monogamous
24
25
26 114 mating system of this species at the genetic level.

27
28
29 115 The indri is a diurnal lemur living in family groups composed of up to six individuals,
30
31 116 usually consisting of the reproductive pair and their offspring (Pollock, 1975, Bonadonna
32
33 117 et al., 2014). The pair bonding in indris is remarkable. The reproductive pair is the basal
34
35
36 118 unit of a group (Pollock, 1975) and we have records of individuals being together for at
37
38
39 119 least a decade (unpublished data).

40
41
42 120 This species can be considered a slow breeder. The gestation period reported for
43
44 121 indris is 157 days (Godfrey, Samonds, Jungers, Sutherlan, Irwin, 2004) and another study
45
46 122 reported 176 days (Weir, 2014) with interbirth intervals of two to three years (Weir,
47
48
49 123 2014), females give births to one offspring at a time between May and July, and
50
51 124 individuals presumably become potentially reproductive at the age of three, when they
52
53
54 125 start singing (unpublished data). Group members tend to be cohesive, and usually the
55
56 126 individuals of the reproductive pair stay within a distance between 0 and 13 m from each

1
2
3 127 other (Pollock, 1975). Each group occupies a stable and exclusive territory and intergroup
4
5 128 encounters are very rare (Bonadonna et al. 2017). The indri is one of the few singing
6
7
8 129 primates and, besides having a function in signaling territory occupancy and defense (Torti
9
10 130 et al., 2013), the song also plays an important role in the mating and social system of this
11
12 131 species. The temporal and spectral characteristics of the song can signal the pair-bond
13
14 132 strength to conspecifics (Gamba et al., 2016). Furthermore, the individual-specific features
15
16 133 of the indri song and acoustic similarities between fathers and sons can play a role in pair
17
18 134 formation, dispersal and avoidance of inbreeding (Torti et al. 2017).
19
20
21
22

23 135 Using fecal samples collected in Maromizaha between 2009 and 2015, we obtained
24
25 136 individual genetic fingerprints and examined the genetic relationships between sires and
26
27 137 offspring. In the presence of an observed EPC (Bonadonna et al., 2014), we hypothesized
28
29 138 that the indri population we studied would show a certain degree of EPP during successive
30
31 139 years and across different groups. Previous studies on gibbons (*H. lar*), demonstrated that
32
33 140 EPC may frequently occur after group encounters (Reichard, 1995). A more recent work
34
35 141 on the same population studied by Reichard has found an EPP rate of 9.5% on a total of 42
36
37 142 offspring (Bartlett, Light & Brockelman, 2016). Our prediction is that we would find a
38
39 143 limited occurrence of EPP in indris, which present a pair-living social organization where
40
41 144 EPC occurs.
42
43
44
45
46
47
48

49 145 **Methods**

50 51 52 146 *Study Site*

53
54
55 147 We conducted the study on a wild population of indris (*I. indri*) in the New Protected
56
57 148 Area (NAP) of Maromizaha (18°56'49" S – 48°27'53" E). We accessed the forest from the
58
59
60

1
2
3 149 village of Anevoka (on the RN2), 147 km east of Antananarivo in the Alaotra Mangoro
4
5 150 Region (Province of Toamasina), central-eastern Madagascar. Maromizaha (1880 ha) is a
6
7 151 primary and secondary mid-altitude (800-1200m) evergreen rainforest, and it is part of
8
9 152 the ecological corridor of Ankeniheny-Zahamena (Fig. 1). The study included an area of
10
11 153 approximately 140 ha of continuous rainforest.
12
13
14
15

16 154 *Subjects and Sample Collection*

17
18 155 Nine indri groups have been habituated to human presence and have been the subject
19
20 156 of ongoing etho-ecological studies in Maromizaha since 2009. Seven out of nine
21
22 157 habituated groups were included in this study (Fig. 1).
23
24
25

26 158 We updated group composition and demographic records year by year, identifying
27
28 159 each individual using natural marks. The population density in the study area was $27.7 \pm$
29
30 160 4.0 (mean \pm SD) individuals and 8.4 ± 1.0 (mean \pm SD) groups per km², comparable to
31
32 161 other populations inhabiting different sites (Bonadonna et al. 2017; Pollock 1975).
33
34
35

36 162 Each group was composed of a reproductive pair plus one to three individuals. These
37
38 163 additional indris are usually the pair's offspring; however, a case of immigration of an
39
40 164 individual into another group was observed in another forest (GB and VT, unpublished
41
42 165 data) (Table 1). We never observed copulation attempts outside the mating season, which
43
44 166 ranges between December and February (Pollock 1975, GB VT pers. obs.). The
45
46 167 reproductive pairs remained stable over time, except for one takeover in 2015, after the
47
48 168 death of the injured resident male (CDG, pers. obs.) (Table 1).
49
50
51
52
53

54 169 For DNA analyses, we included 26 individuals from seven groups: seven parental pairs
55
56 170 and 12 offspring. Unfortunately, not all the individuals that were part of a group during
57
58
59
60

1
2
3 171 our study were included in DNA analyses, either because we did not have samples
4
5 172 available, or because DNA extraction did not succeed (Table 1).
6
7

8
9 173 Between 2011 and 2015, we obtained at least two fecal samples for each animal, each
10
11 174 corresponding to a different defecation. We collected fecal samples immediately after
12
13 175 defecation to avoid individual misidentification during the sampling process. Feces were
14
15 176 put in 20ml labeled tubes filled with RNAlater® Ambion (Nsubuga et al. 2004) wearing
16
17 177 disposable gloves and stored at room temperature in the field and at 4°C once transferred
18
19 178 to the lab. We conducted all the genetic analyses at the New York University Molecular
20
21 179 Anthropology Laboratory.
22
23
24

25 26 180 *DNA Extraction* 27

28
29 181 Genomic DNA was extracted from feces using the QIAamp DNA® Stool Mini Kit
30
31 182 (Qiagen®, Hilden, Germany). To maximize the amount of DNA extracted we slightly
32
33 183 modified the manufacturer's protocol (QIAamp DNA Stoll Handbook 04/2010): we used
34
35 184 300 mg stool instead of 180-220 mg; we added 35 µL of proteinase K rather than 25 µL
36
37 185 and incubated at 70°C for 30 minutes, instead of 10 minutes, during the DNA purification
38
39 186 phase. We then applied 75 µL Buffer AE on the QIAamp membrane rather than 200 µL for
40
41 187 the first DNA elution and incubate the spin column with Buffer AE at room temperature
42
43 188 for 15 minutes instead of one minute. We used the same QIAamp membrane to obtain a
44
45 189 second DNA elution applying 50 µL of Buffer AE and incubate for 15 minutes at room
46
47 190 temperature.
48
49
50
51
52

53
54 191 For the samples collected in the field during 2014, DNA purification was conducted
55
56 192 using the automated robotic workstation QIAcube HT supported by the software
57
58
59
60

1
2
3 193 QIAxtractor 4.17.1 (Qiagen®) setting the protocol for QXT Liquid DNA V1. The preparation
4
5
6 194 of the samples required a bath at 70°C for at least five minutes of the 2.0 mL tubes
7
8 195 containing 300 mg of smashed feces and 1.6 mL of Buffer ASL. Afterwards, tubes were
9
10 196 centrifuged at maximum speed (13000 RPM) for 10 minutes. 200 µL of supernatant were
11
12 197 then transferred to separate wells of the QIAextractor lysis plate before starting the run.
13
14 198 At the end of the run, we obtained 70 µL of DNA elution for each sample. We stored the
15
16 199 extracted DNA at 4° C for immediate use.
17
18
19
20

21 *Microsatellite Genotyping*

22
23 201 We initially identified a set of 10 microsatellite marker loci as potentially variable in
24
25 202 indris (Zaonarivelo et al. 2007). Polymorphism of each locus was tested using monolocus
26
27 203 PCRs. Four loci failed to provide amplification products, and we therefore chose to use the
28
29 204 six loci among the ten tested that provided good quality amplification products for
30
31 205 multiplex PCRs (Table 2). The 5' end forward primer of each locus was labeled with a
32
33 206 fluorescent dye (FAM, HEX) to analyze simultaneously loci of similar allelic size. PCR
34
35 207 amplification was carried out in 10 µL reaction volume containing 2 µL DNA template, 5 µL
36
37 208 Multiplex PCR Master Mix (Qiagen®), 0.1 µL of each primer, and two µL RNase-free water.
38
39 209 The cycle conditions included a pre-incubation step at 95 °C for 15 min. We then
40
41 210 performed 50 cycles with denaturation at 94 °C for 30 s, annealing at 54°C or 60 °C
42
43 211 (depending on the locus, Table 2) for 90 s. The first extension phase was at 72 °C for 60 s;
44
45 212 the final extension phase was at 60 °C for 30 min.
46
47
48
49
50
51
52

53 213 We separated PCR products by electrophoresis using a 48 capillary ABI 3730 DNA
54
55 214 Analyzer (Applied Biosystems) for allele size estimates. We mixed 1 µL of PCR product with
56
57
58
59
60

1
2
3 215 6.85 μ L HiDi formamide (Applied Biosystems) and 0.15 μ L Genescan 500-ROX size
4
5
6 216 standard (Applied Biosystems). We carried out automated allele calling using the
7
8 217 GENEMAPPER 3.7 software (Applied Biosystems). We subsequently confirmed by eye all
9
10 218 the allele calls and checked for consistency across replicate PCRs of the same sample or
11
12
13 219 from the same individual. To minimize possible genotyping errors due to allelic dropout,
14
15 220 we repeated independent PCRs for each locus depending on whether the individual
16
17 221 resulted in being heterozygote (minimum three replicates) or homozygote (minimum five
18
19 222 replicates) for a certain locus.
20
21
22

23 223 We used the software CERVUS 3.0 to calculate observed and expected heterozygosity
24
25 224 to test deviation from Hardy-Weinberg equilibrium (HWE), and to estimate null allele
26
27 225 frequency for each locus.
28
29
30

31 226 *Paternity test and assignment*

32
33 227 We identified 14 paired individuals (7 males, 7 females), and 12 potential offspring.
34
35 228 We included the reproductive males of the neighboring groups as potential sires for each
36
37 229 potential offspring. Unpaired neighboring males exceeding the age of 3 at the time of
38
39 230 habituation were also included as potential sires because: (i) we could not exclude that
40
41 231 they had potentially mated with a female in the study population before or during the
42
43 232 sampling period; and (ii) the youngest indri we observed forming a group was 3 and half
44
45 233 years old. We also considered the presence of potential sires from the unsampled
46
47 234 neighboring groups. In the simulation, we included one unsampled potential sire for each
48
49 235 neighboring non-habituated group.
50
51
52
53
54
55
56
57
58
59
60

1
2
3 236 Ten out of the 12 offspring were born during the 2009-2015 study period in
4
5
6 237 Maromizaha; two were already present in the groups at the beginning of the habituation,
7
8 238 and we could not know whether they were born in the group or immigrated. We
9
10 239 considered these individuals as both potential offspring and potential sires. For these two
11
12
13 240 indris, all the reproductive males were also included as potential fathers. For the single
14
15 241 case of offspring and social father mismatch, we included each individual qualified as
16
17
18 242 potential sire as a candidate father. The average number of potential sires for the 12
19
20 243 offspring was 7.3 ± 2.2 (mean \pm SD) and the potential sires sampled were 4.1 ± 2.2 (mean
21
22
23 244 \pm SD).

245 Based on microsatellite genotypes, we ran parentage analyses using CERVUS 3.0
246 (Kalinowski, Taper, & Marshall, 2007). This program compares likelihood ratios (LOD
247 scores) of all candidate fathers, assigning or excluding paternity to the most likely parent
248 using statistical criteria generated by computer simulation. A true parent has a positive
249 LOD score; on the contrary, a negative LOD score indicates that the potential father is
250 unrelated to the offspring. The program takes genotyping errors and the presence of close
251 relatives into account (Marshall et al. 1998; Jones & Ardren, 2003); we applied a mistyping
252 rate of 0.01. This procedure ensures the running of blind analyses.

253 We set the confidence levels of paternity assignment at 90% (relaxed level) and 95%
254 (strict level). The critical value corresponding to a strict or relaxed confidence level was
255 given by the delta LOD score between the first and the second potential father,
256 automatically calculated by CERVUS. We considered the assignment of the father given a
257 known mother, to include the genotype of the mother when matching the genotype of

1
2
3 258 potential sires to offspring. This considers the case in which the parent pairs are mutually
4
5
6 259 exclusive.

7
8
9 260 We assigned paternity in the cases of complete genotypes matching between the
10
11 261 potential father and offspring. We excluded the paternity in cases of negative LOD scores
12
13 262 and more than one allelic mismatch between the offspring and the potential father. We
14
15
16 263 assigned EPP in cases of paternity exclusion for the social father; if the paternity was also
17
18 264 excluded for all the other possible sampled sires, we defined the assignment of EPP as
19
20
21 265 inconclusive.

22
23
24 266 *Ethical standards*

25
26
27 267 During the study, we did not have any physical contact with the animals. All fecal samples
28
29 268 were collected from the ground immediately after defecation. The research was
30
31 269 authorized by the “Ministère de l’Environnement et des Forêts” (MEF) of Madagascar.
32
33
34 270 Research permits: N° 243/ 09/ MEF/ SG/ DGF/ DCB.SAP/ SLRSE, N° 118/ 10/ MEF/ SG/
35
36 271 DGF/ DCB.SAP/ SCBSE; N° 293/ 10/ MEF/ SG/ DGF/ DCB.SAP/ SCB, N° 274/ 11/ MEF/ SG/
37
38 272 DGF/ DCB.SAP/ SCB, N°245/12/MEF/SG/DGF/DCB.SAP/SCB,
39
40
41 273 N°066/14/MEF/SG/DGF/DCB.SAP/ SCB. We adhered to applicable international, national,
42
43 274 and/or institutional guidelines for the study on animals and non-human primates,
44
45
46 275 including the American Society of Primatologist (ASP) Principle for the Ethical Treatment
47
48 276 of non-human Primates, and the European Union directive guidelines for the study on
49
50
51 277 animals and non-human primates (Directive 2010/63/EU). The study did not require an
52
53
54 278 IACUC approval.

Results*Genotypes and Identity of Individuals*

We included all the 26 individuals in the analyses. For three indris of the group 8MZ we failed to amplify the locus 67HDZ55, and for one of them (Jonah) also the locus 67HDZ62. We obtained confirmed genotype of 23 individuals for the locus 67HDZ55, 25 for the locus 67HDZ62, and 26 for all the four remaining loci (Table 3).

The number of alleles per locus varied between nine and 14, observed heterozygosity (HO) ranged from 0.840 to 0.962 and expected heterozygosity (HE) ranged between 0.814 and 0.928. None of the six loci showed significant deviation from the Hardy-Weinberg equilibrium (HWE) after Bonferroni correction (Table 3). Even if in some cases the presence of null allele frequency is close to the suggested threshold of 0.05, we included all the loci in the analyses because the level of heterozygosity observed was high and all known offspring-maternal pairs were free of possible homozygous mismatches for those loci. The combined non-exclusion probability of identity over six loci was 8.482×10^{-10} . Therefore, it is unlikely that the set of loci failed to differentiate between two randomly-selected individuals.

We found two cases of loci mismatch between mother and offspring. In one case (Locus 67HDZ55) the offspring resulted to be homozygous for this locus after six independent replicates (group 3MZ; genotypes: mother "Mena" 313-327; offspring "Blague" 330-330). In the other case (locus 67HDZ25), the mother was homozygous after eight independent replicates (group 4MZ; genotypes: mother "Eva" 224-224; offspring "Hendri" 226-228). In both cases the offspring matched with the mother for all the other

1
2
3 301 five loci, and all the six loci matched with the social father; therefore, we suggest that this
4
5 302 incompatibility is related to the estimated error rate of 0.0733 and of 0.0728 for the loci
6
7
8 303 67HDZ55 and 67HDZ25, respectively. It is possible that known parent mismatches at one
9
10 304 or more loci are due to the presence of null alleles (Kalinowski, Taper & Marshall, 2007),
11
12 305 especially in the case of homozygous individuals; the high likelihood of parentage given by
13
14 306 the other loci and the consistent match with both parents make it acceptable to consider
15
16 307 the known mothers as the true parents. The same methodological approach would have
17
18 308 been applied in the case of one locus mismatch with the fathers.
19
20
21
22
23

24 309 *Paternity*

25
26 310 Paternity based on CERVUS at a confidence level of 95% given a known mother was
27
28 311 assigned to a total of 11 individuals out of 12 (92% against an expected assignment of 61%
29
30 312 of the total offspring). We found a lower percentage of unassigned paternity than
31
32 313 expected, 8% versus 39%. None of the assigned paternity presented any mismatch with
33
34 314 the offspring for all the six loci considered, and the assigned fathers were consistently the
35
36 315 reproductive males of the group (Table 4).
37
38
39
40

41 316 Only one out of 12 offspring tested (Tsiky, group 6MZ) showed a mismatch with the
42
43 317 social father (Zokybe, group 6MZ) for all the six loci, and a negative LOD score. We have
44
45 318 also found that the LOD score was negative for all the potential fathers tested. However,
46
47 319 the reproductive female (Befotsy, group 6MZ) had no mismatch with the offspring (Table
48
49 320 4). Tsiky was born in the summer of 2010, the year after the habituation of the group;
50
51 321 Zokybe was the paired male of the group at the time of habituation (in September 2009)
52
53 322 and was the paired male at the time of the reproductive season (starting in December
54
55
56
57
58
59
60

1
2
3 323 2009); Tsiky was born in the summer of 2010, the first birth season after the habituation
4
5 324 of the group; therefore, we can consider the mismatch as a case of EPP with inconclusive
6
7
8 325 paternity assignment. This evidence indicates an EPP rate of 8% in our sample.
9

10
11 326 The most closely related individual to the indri with unassigned paternity was Hendri,
12
13 327 a potential reproductive male of the neighboring group 4MZ, having two loci mismatches
14
15
16 328 even when considering the genotype of the mother. However, it is unlikely that Hendri
17
18 329 can be the father: Hendri and Tsiky are heterozygotes for all the loci considered,
19
20
21 330 eliminating the possibility of allelic dropout at the mismatching loci; furthermore, the LOD
22
23 331 score was negative, excluding Hendri as a potential sire for Tsiky.
24
25

26 332 It must be noted that an unsampled individual was in the group 6MZ during the year
27
28 333 of conception and left the group the year after the birth of Tsiky (Table 1). If the male that
29
30 334 left the group was one of the offspring, we would exclude the possibility that he was a
31
32
33 335 potential sire, and we would hypothesize that the sire would be an unknown male.
34
35
36 336 Unfortunately, our samples did not allow us to test this hypothesis, leaving the paternity
37
38 337 assignment inconclusive.
39
40

41
42 338 The male (Emilio) involved in the EPC observed in 2011 (Bonadonna et al. 2014) with
43
44 339 the female of the group 3MZ (Mena) was not the sire of the individual (Blague) born after
45
46 340 the EPC; in fact, it was the resident and paired male of the group 3MZ (Ratsy) that was
47
48
49 341 found to be Blague's father (Table 4).
50
51
52
53
54
55
56
57
58
59
60

1
2
3 342 **Discussion**
4
5

6 343 Our study investigates the genetic relationship between offspring, social parents and
7
8
9 344 potential sires in indris, within the same population where an EPC was observed
10
11 345 (Bonadonna et al. 2014). The results we presented in this paper provide new insights into
12
13
14 346 the mating system of this pair-living primate.
15

16
17 347 The paternity analysis revealed that all genetic profiles of the offspring matched with
18
19 348 the social fathers, except for one case. Because in this case the paternity remained
20
21
22 349 inconclusive, we cannot exclude our hypothesis that EPC may lead to EPP in indris, but we
23
24 350 have to weigh this phenomenon across our study groups. In fact, we confirmed genetic
25
26
27 351 monogamy for 11 out of the 12 cases we tested. The 8% mismatch with the paired males
28
29 352 we found in indris is lower than the high rate of EPP found by Fietz and colleagues (2000)
30
31 353 in the nocturnal dwarf lemur (*C. medius*, 44%), but comparable with the study of Barelli et
32
33
34 354 al. (2013) reporting rates varying from 8.5% to 10% for white handed gibbons (*H. lar*).
35

36
37 355 Our results agree with the low rate of EPP in monogamous and constantly associated
38
39 356 pairs predicted by van Schaik & Kappeler (2003) and Clutton-Brock & Isvaran (2006). Our
40
41
42 357 study also supports the hypothesis of Cohas & Allainé (2009), who pointed out the
43
44 358 important role of pair-living social organization in ensuring paternity exclusivity, in
45
46
47 359 addition to pair-bonding strength and mate guarding.
48
49

50 360 White-handed gibbons share with indris several characteristics: they usually have a
51
52 361 stable reproductive pair per group, and they show exclusive territoriality (Brockelman et
53
54
55 362 al., 1998). They also share the emission of loud songs that, in both species, have a role in
56
57
58
59
60

1
2
3 363 signaling pair-bonding strength (Geissmann & Orgendilger 2000; Gamba et al., 2016). Long
4
5 364 distance calls can also broadcast information to potential mates (Torti et al., 2013, 2017;
6
7
8 365 Gamba et al., 2016). Each group exchanges information about individuals' sex, status, and
9
10 366 genetic relatedness throughout the emission of loud songs way beyond the limit of a
11
12 367 territory (Torti et al., 2017). In this way, groups are not isolated units and individuals can
13
14 368 communicate with one another without having visual or physical contact (Giacoma et al.,
15
16 369 2010; Gamba et al., 2016; Bonadonna et al. 2017). Such a system would allow regulating
17
18 370 inter-group dynamics including the possibility of engaging in EPC (Bonadonna et al., 2014),
19
20 371 and therefore providing a certain degree of flexibility in the mating system of this species.
21
22
23
24
25

26 372 The takeover of paired individuals is rare and, according to our observations, the case
27
28 373 reported for the group 3MZ in Maromizaha was the result of a conflict between males
29
30 374 (CDG, pers. obs.). We suggest that both pair-bonding strength and social structure play a
31
32 375 role in the flexibility of a monogamous mating system, as we found in indris.
33
34
35

36 376 It has been suggested that once a monogamous mating system is established, low
37
38 377 rates of EPP are related to intensive male care (Huck et al., 2014). In the indri, males cover
39
40 378 the primary role in territorial defense (Pollock 1975), which can be considered an indirect
41
42 379 form of parental care (Brockelman, 1975; Kleiman, 1977). The territorial defense is part of
43
44 380 the resource defense strategy: the male can guard the access to the female, and both the
45
46 381 female and offspring have access to resources (Clutton-Brock, 1991; Møller & Thornhill,
47
48 382 1998). This idea agrees with the hypothesis that parental care is an indirect evolutionary
49
50 383 consequence of mate guarding (Huck et al., 2014). There is no more direct form of
51
52
53
54
55
56
57
58
59
60

1
2
3 384 parental care by males in indris apart from the fact that they may occasionally transfer
4
5 385 infants from a branch to another while the mother is feeding (Torti, pers. comm.).
6
7

8
9 386 As suggested by Bonadonna and colleagues (2014), EPC can be a female strategy to
10
11 387 enhance male guarding. Given the fact that the EPC reported for indris has been observed
12
13 388 at the beginning of the mating season, it could have increased the male's guarding
14
15 389 behavior and also reduced the chances of EPP. An enhanced guarding of the paired male
16
17 390 would also improve resource monopolization, especially during the mating season
18
19 391 (Brotherton & Komers, 2003).
20
21
22
23

24 392 We can also indirectly assume that EPP and the occurrence of EPCs can be
25
26 393 underestimated. The direct and indirect benefits for females adopting differential
27
28 394 reproductive strategies in indris are not easily identifiable. Interestingly, the genetic
29
30 395 profile of the individual born after the breeding season in which we observed the EPC
31
32 396 suggests that the female successfully reproduced with her social partner during the same
33
34 397 year. Thus, our findings suggest that the female was not likely to engage in EPC to avoid
35
36 398 partner infertility (Palombit, 1994; Brotherton & Komers, 2003), as hypothesized for
37
38 399 white-handed gibbons (in which two out of three EPPs were found in the same pair-living
39
40 400 group, Barelli et al., 2013).
41
42
43
44
45

46 401 Our results excluded that the current paired male of the group 6MZ (Zokybe) sired the
47
48 402 oldest individual of the offspring (Tsiky), representing the only case of EPP in our study,
49
50 403 but he did not present any loci mismatch with the individual born in 2014 (Hira). We
51
52 404 consider two possible scenarios to explain our results. In the first scenario, the genetic
53
54 405 father of the individual with unassigned paternity might have preceded a male takeover,
55
56
57
58
59
60

1
2
3 406 which would explain the mismatch of the paired male as the sire of the offspring. Partner
4
5
6 407 take-over in the indri can occur. This fact would configure a forced “divorce” as reported
7
8 408 for brown titi monkeys (*Callicebus brunneus*; Lawrence, 2007), and owl monkeys (*Aotus*
9
10 409 *azarae*; Fernandez-Duque & Huck, 2013), and it agrees with the serial monogamy model
11
12 410 proposed by Fernandez-Duque & Huck (2013). This first scenario seems unlikely because
13
14
15 411 Zokybe was reported as the paired male during the reproductive season that preceded the
16
17
18 412 birth of Tsiky despite the changes in the group composition. In the second, and more likely
19
20 413 scenario, there was no takeover of the reproductive male because Zokybe was the paired-
21
22 414 male of the female at the time of conception. This scenario is in accordance with our
23
24
25 415 hypothesis that partner infertility is not a primary cause of EPC and EPP in indris because
26
27 416 Zokybe was likely the father of the younger offspring (no evidence of allelic mismatch).
28
29

30 417 Although we were not able to assign the paternity to an individual, based on our
31
32
33 418 results, we can draw some conclusion about the mating system in indris. According to our
34
35
36 419 findings and observations, individuals that reach maturity can either leave or remain in
37
38 420 the group. We found that two resident males (group 4MZ and 8MZ), both over the
39
40
41 421 reproductive age, were the offspring of the reproductive pairs. Furthermore, we never
42
43 422 observed a copulation event between a mother and her potential reproductive offspring,
44
45 423 and finally we did not find any case of offspring sired by an older brother. Therefore, we
46
47
48 424 did not find any evidence to support the hypothesis that adults prior dispersal are likely
49
50 425 candidates for siring offspring.
51
52

53 426 The floating of individuals without a territory, delayed dispersal and immigration have
54
55
56 427 an important role in the dynamics of the monogamous mating system in pair-living
57
58
59
60

1
2
3 428 primates (Porter, Grote, Fernandez-Duque & Di Fiore, 2017; Jacobs, Frankel, Rice, Kiefer &
4
5 429 Bradley, 2018). Unfortunately, there are few studies and data available on the dispersal
6
7
8 430 pattern and dynamics of the social organization in the indri. The role of these factors in
9
10 431 the social and mating system of this species need further investigation. However, even if
11
12 432 in one case we could not assign the paternity, in all the other cases we found no loci
13
14 433 mismatches between social fathers and offspring. In addition, we did not find any case of
15
16 434 ambiguity among potential fathers, making our results consistent.
17
18
19
20

21 435 Genetic analyses can reveal a level of plasticity in the mating system of species
22
23 436 considered monogamous (Díaz-Muñoz & Bales 2016). Although our study is limited to a
24
25 437 relatively small sample size, it is indicative that genetic monogamy seems to be the norm
26
27 438 in indris, with an EPP rate comparable to other pair-living monogamous primates.
28
29 439 Furthermore, it is unlikely that a female would reproduce with another male in the group
30
31 440 other than her partner. This situation would exclude that the social organization of indris
32
33 441 may change from pair-living to polyandrous as it has been found in primates with a similar
34
35 442 behavioral ecology, such as the gibbons (Barelli, Heistermann, Boesch, Reichard, 2008).
36
37
38
39
40

41 443 In conclusion, our study contributes to the scanty literature of genetic studies of wild
42
43 444 monogamous, pair-living primates. We found that the model of pair-living social
44
45 445 organization fits this species although it might not be consistently associated with genetic
46
47 446 monogamy despite the low EPP rate. However, our results were not fully conclusive about
48
49 447 paternity assignment, requiring further investigations on the genetic structure of the
50
51 448 population, and the dispersal dynamics. Future studies should focus on providing further
52
53
54
55
56
57
58
59
60

1
2
3 449 insight into the mechanisms involved in the maintenance of socially monogamous systems
4
5 450 and their genetic variability.
6
7
8

9 451 **Acknowledgments**

10
11
12 452 This research was supported by the University of Turin and the ACP Science and
13
14
15 453 Technology Programme of the ACP Group of States, with the financial assistance of the
16
17
18 454 European Union, through the Projects BIRD (Biodiversity Integration and Rural
19
20 455 Development; no. FED/2009/217077) and SCORE (Supporting Cooperation for Research
21
22 456 and Education; Contract no. ACP RPR 118 # 36). Part of the field research was supported
23
24
25 457 by grants from the Parco Natura Viva – Garda Zoological Park, PCI (Primate Conservation
26
27 458 Inc.), and scholarship granted by the project UNI.COO (UNITO for International
28
29
30 459 Cooperation). We are particularly grateful to Todd Disotell and the New York University
31
32 460 Molecular Anthropology Laboratory, Department of Anthropology, for hosting and
33
34
35 461 supporting GB in performing all the genetic analyses. We thank the *Ministère de*
36
37 462 *l'Environnement et des Forêts* (MEF) and Madagascar National Parks for granting the
38
39
40 463 research permits. We are also grateful to GERP for allowing us to collect data in the forest
41
42 464 under its management. We thank all the field assistants, research guides (Gilbert, Naivo,
43
44 465 Zafison) and the students that worked in the field. Lanto and Mamatin for their help and
45
46
47 466 logistic support. We thank R. Lovelace, and the member of Monmouth University Writing
48
49 467 Group for proofreading the manuscript. We would like to thank the two anonymous
50
51
52 468 reviewers for their suggestions and comments that helped improving the quality of this
53
54 469 manuscript. The authors have no conflict of interest to declare.
55
56
57
58
59
60

470 **References**

- 471 Akçay, E., & Roughgarden, J. (2007). Extra-pair paternity in birds: review of the genetic
472 benefits. *Evolutionary Ecology Research*, 9, 855-868.
- 473 Barelli, C., Heistermann, M., Boesch, C., & Reichard, U. H. (2008) Mating patterns and
474 sexual swellings in pair-living and multimale groups of wild white-handed gibbons,
475 *Hylobates lar*. *Animal Behaviour*, 75, 991-1001. DOI: 10.1016/j.anbehav.2007.08.012
- 476 Barelli, C., Matsudaira, K., Wolf, T., Roos, C., Heistermann, M., Hodges, K., ... Reichard,
477 U. H. (2013). Extra-pair paternity confirmed in wild white-handed gibbons. *American*
478 *Journal of Primatology*, 75, 1185-95. DOI: 10.1002/ajp.22180
- 479 Bartlett, T. Q., Light, L. E., & Brockelman, W. Y. (2016). Long-term home range use in
480 white-handed gibbons (*Hylobates lar*) in Khao Yai National Park, Thailand. *American*
481 *Journal of Primatology*, 78, 192-203. DOI: 10.1002/ajp.22492
- 482 Black, J.M. (1996). *Partnerships in Birds: The Study of Monogamy: The Study of*
483 *Monogamy*. Oxford, UK: Oxford University Press.
- 484 Birkhead, T.R., & Møller, A.P. (1992). *Sperm competition in birds: evolutionary causes*
485 *and consequences*. London, UK: Academic Press.
- 486 Bonadonna, G., Torti, V., Randrianarison, R. M., Martinet, N., Gamba, M., & Giacoma,
487 C. (2014). Behavioral correlates of extra-pair copulation in *Indri indri*. *Primates*, 55,
488 119-23. DOI: 10.1007/s10329-013-0376-0

- 1
2
3 489 Bonadonna, G., Torti, V., Sorrentino, V., Randrianarison, R. M., Zaccagno, M., Gamba,
4
5
6 490 M., ... Giacomini, C. (2017). Territory exclusivity and intergroup encounters in the indris
7
8 491 (Mammalia: Primates: Indridae: *Indri indri*) upon methodological tuning. *The European*
9
10 492 *Zoological Journal*, 84, 238-251. DOI: 10.1080/24750263.2017.1318184
11
12
13 493 Brockelman, W.Y. (1975) Competition, the fitness of offspring, and optimal clutch size.
14
15
16 494 *American Naturalist*, 109: 677-699.
17
18
19 495 Brockelman, W. Y., Reichard, U., Treesucon, U., & Raemaekers, J. J. (1998). Dispersal,
20
21 496 pair formation and social structure in gibbons (*Hylobates lar*). *Behavioral Ecology and*
22
23 497 *Sociobiology*, 42, 329-39. DOI: 10.1007/s002650050445
24
25
26
27 498 Brotherton, P. N., & Komers, P. E. (2003). Mate guarding and the evolution of social
28
29 499 monogamy in mammals. In *Monogamy: mating strategies and partnerships in birds,*
30
31 500 *humans and other mammals.* (pp. 42-58). Cambridge, UK: Cambridge University Press
32
33
34
35 501 Brotherton, P. N., & Komers, P. E. (2003). Mate guarding and the evolution of social
36
37 502 monogamy in mammals. In U.H. Reichard & C. Boesch (Eds.), *Monogamy: Mating*
38
39 503 *strategies and partnerships in birds, humans and other mammals* (pp. 42-58).
40
41
42 504 Cambridge, UK: Cambridge University Press
43
44
45 505 Clutton-Brock, T.H. (1991). *The evolution of parental care.* Princeton, NJ: Princeton
46
47 506 University Press.
48
49
50
51 507 Clutton-Brock, T. H., & Isvaran, K. (2006). Paternity loss in contrasting mammalian
52
53 508 societies. *Biology Letters*, 2, 513-516. DOI: 10.1098/rsbl.2006.0531
54
55
56
57
58
59
60

- 1
2
3 509 Cohas, A., & Allainé, D. (2009). Social structure influences extra-pair paternity in
4
5 510 socially monogamous mammals. *Biology letters*, 5, 313-316 DOI:
6
7 511 10.1098/rsbl.2008.0760
8
9
10
11 512 Díaz-Muñoz, S. L., & Bales, K. L. (2016). "Monogamy" in primates: variability, trends,
12
13 513 and synthesis: introduction to special issue on primate monogamy. *American Journal*
14
15 514 *of Primatology*, 78(3), 283-287.
16
17
18
19 515 Driller, C., Perwitasari-Farajallah, D., Zischler, H., & Merker, S. (2009). The social
20
21 516 system of Lariang tarsiers (*Tarsius lariang*) as revealed by genetic
22
23 517 analyses. *International journal of primatology*, 30(2), 267-281.
24
25
26
27 518 Fernandez-Duque, E., & Huck, M. (2013). Till Death (Or an Intruder) Do Us Part:
28
29 519 Intrasexual-Competition in a Monogamous Primate. *PLoS ONE*, 8(1): e53724. DOI:
30
31 520 10.1371/journal.pone.0053724
32
33
34
35 521 Fietz, J., Zischler, H., Schwegk, C., Tomiuk, J., Dausmann, K. H., & Ganzhorn, J. U.
36
37 522 (2000). High rates of extra-pair young in the pair-living fat-tailed dwarf lemur,
38
39 523 *Cheirogaleus medius*. *Behavioral Ecology and Sociobiology*, 49, 8-17. DOI:
40
41 524 10.1007/s002650000269
42
43
44
45 525 Fuentes, A. (2000). Hylobatid communities: changing views on pair bonding and social
46
47 526 organization in hominoids. *American Journal of Physical Anthropology*, 113(S31), 33-
48
49 527 60. DOI: 10.1002/1096-8644(2000)43:31+<33::AID-AJPA3>3.0.CO;2-D
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 528 Gamba, M., Favaro, L., Torti, V., Sorrentino, V., & Giacoma, C. (2011). Vocal tract
4
5 529 flexibility and variation in the vocal output in wild indris, *Bioacoustics*, 20, 251-265.
6
7
8 530 DOI: 10.1080/09524622.2011.9753649
9
10
11 531 Gamba, M., Torti, V., Estienne, V., Randrianarison, R. M., Valente, D., Rovara, P., ... &
12
13 532 Giacoma, C. (2016). The indris have got rhythm! Timing and pitch variation of a
14
15 533 primate song examined between sexes and age classes. *Frontiers in Neuroscience*, 10,
16
17 534 249. DOI: 10.3389/fnins.2016.00249
18
19
20
21 535 Garber, P. A., Porter, L. M., Spross, J., & Di Fiore, A. (2016). Tamarins: Insights into
22
23 536 monogamous and non-monogamous single female social and breeding systems.
24
25 537 *American Journal of Primatology*, 78, 298-314. DOI: 10.1002/ajp.22370
26
27
28
29 538 Geissmann, T., & Orgeldinger, M. (2000). The relationship between duet songs and
30
31 539 pair bonds in siamangs, *Hylobates syndactylus*. *Animal Behaviour*, 60, 805-809. DOI:
32
33 540 10.1006/anbe.2000.1540
34
35
36
37 541 Giacoma, C., Sorrentino, V., Rabarivola, C., & Gamba, M. (2010). Sex differences in the
38
39 542 song of *Indri indri*. *International Journal of Primatology*, 31, 539-551. DOI:
40
41 543 10.1007/s10764-010-9412-8
42
43
44
45 544 Godfrey, L. R., Samonds, K. E., Jungers, W. L., Sutherland, M. R. & Irwin, M. T.
46
47
48 545 (2004). Ontogenetic correlates of diet in Malagasy lemurs. *American Journal of*
49
50 546 *Physical Anthropology*, 123, 250–276.
51
52
53
54
55
56
57
58
59
60

- 1
2
3 547 Goossens, B., Graziani, L., Waits, L. P., Farand, E., Magnolon, S., Coulon, J., ... & Allainé,
4
5 548 D. (1998). Extra-pair paternity in the monogamous alpine marmot (*Marmota*
6
7
8 549 *marmota*): the roles of social setting and female mate choice. *Behavioral Ecology and*
9
10 550 *Sociobiology*, 59, 597-605. DOI: 10.1007/s002650050492
11
12
13 551 Griffith, S. C., Owens, I. P., & Thuman, K. A. (2002). Extra pair paternity in birds: a
14
15
16 552 review of interspecific variation and adaptive function. *Molecular ecology*, 11, 2195-
17
18 553 2212. DOI: 10.1046/j.1365-294X.2002.01613.x
19
20
21 554 Huck, M., Fernandez-Duque, E., Babb, P., & Schurr, T. (2014). Correlates of genetic
22
23
24 555 monogamy in socially monogamous mammals: insights from Azara's owl monkeys.
25
26 556 *Proceedings of the Royal Society of London B*, 281, 20140195. DOI:
27
28 557 10.1098/rspb.2014.0195
29
30
31
32 558 Isvaran, K., Clutton-Brock, T. (2007). Ecological correlates of extra-group paternity in
33
34 559 mammals. *Proceedings of the Royal Society of London B: Biological Sciences*, 274, 219-
35
36 560 224. DOI: 10.1098/rspb.2006.3723
37
38
39
40 561 Jacobs, R.L., Frankel, D.C., Rice, R.J., Kiefer, V.J., & Bradley, B.J. (2018). Parentage
41
42 562 complexity in socially monogamous lemurs (*Eulemur rubriventer*): Integrating genetic
43
44
45 563 and observational data. *American Journal of Primatology*, 80(2), e22738. DOI:
46
47 564 10.1002/ajp.22738
48
49
50 565 Jones, A. G., & Ardren, W. R. (2003). Methods of parentage analysis in natural
51
52 566 populations. *Molecular Ecology*, 12, 2511–2523. DOI: 10.1046/j.1365-
53
54
55 567 294X.2003.01928.x
56
57
58
59
60

- 1
2
3 568 Kalinowski, S. T., Taper, M. L., & Marshall, T. C. (2007). Revising how the computer
4
5
6 569 program CERVUS accommodates genotyping error increases success in paternity
7
8 570 assignment. *Molecular Ecology*, 16, 1099-1106. DOI: 10.1111/j.1365-
9
10 571 294X.2007.03089.x
11
12
13 572 Kappeler, P. M., & van Schaik, C. P. (2002). Evolution of primate social systems.
14
15
16 573 *International Journal of Primatology*, 23, 707-740. DOI: 10.1023/A:101552083
17
18
19 574 Kenyon, M., Roos, C., Binh, V. T., & Chivers, D. (2011). Extrapair paternity in golden-
20
21 575 cheeked gibbons (*Nomascus gabriellae*) in the secondary lowland forest of Cat Tien
22
23
24 576 National Park, Vietnam. *Folia Primatologica*, 82(3), 154-164.
25
26
27 577 Kohn, M., Knauer, F., Stoffella, A., Schröder, W., & Pääbo, S. (1995). Conservation
28
29 578 genetics of the European brown bear-a study using excremental PCR of nuclear and
30
31 579 mitochondrial sequences. *Molecular Ecology*, 4, 95-104. DOI: 10.1111/j.1365-
32
33 580 294X.1995.tb00196.x
34
35
36
37 581 Kleiman, D.G. (1977). Monogamy in mammals. *The Quarterly Review of Biology*. 52,
38
39 582 39-69. DOI: 10.1086/409721
40
41
42
43 583 Lack, D. L. (1968). *Ecological adaptations for breeding in birds*. London, UK: Methuen.
44
45
46 584 Lambert, C. T., Solomon, N. G., & Sabol, A. C. (2018) Genetic Monogamy in Socially
47
48 585 Monogamous Mammals Is Primarily Predicted by Multiple Life History Factors: A
49
50 586 Meta-Analysis. *Frontiers in Ecology and Evolution*, 6:139. doi:
51
52 587 10.3389/fevo.2018.00139
53
54
55
56
57
58
59
60

- 1
2
3 588 Lawrence, J.M. (2007). *Understanding the pair bond in brown titi monkeys (Callicebus*
4
5 589 *brunneus): male and female reproductive interests*. Doctoral dissertation. Columbia
6
7
8 590 University.
- 9
10
11 591 Lukas, D., & Clutton-Brock, T. (2012). Cooperative breeding and monogamy in
12
13 592 mammalian societies. *Proceedings of the Royal Society B: Biological Sciences*, 279,
14
15 593 2151-2156. DOI: 10.1098/rspb.2011.2468
- 16
17
18
19 594 Lukas, D., & Clutton-Brock, T. (2013). The evolution of social monogamy in mammals.
20
21 595 *Science*, 341, 526-530. DOI: 10.1098/rspb.2011.2468
- 22
23
24 596 Marshall, T. C., Slate, J. B. K. E., Kruuk, L. E. B., & Pemberton, J. M. (1998). Statistical
25
26 597 confidence for likelihood-based paternity inference in natural populations. *Molecular*
27
28 598 *ecology*, 7, 639-655. DOI: 10.1046/j.1365-294x.1998.00374.x
- 29
30
31
32 599 Nsubuga, A. M., Robbins, M. M., Roeder, A. D., Morin, P. A., Boesch, C., & Vigilant, L.
33
34 600 (2004). Factors affecting the amount of genomic DNA extracted from ape faeces and
35
36 601 the identification of an improved sample storage method. *Molecular Ecology*, 13,
37
38 602 2089-2094. 10.1111/j.1365-294X.2004.02207.x
- 39
40
41
42
43 603 Nievergelt, C. M., Mutschler, T., Feistner, A. T., & Woodruff, D. S. (2002). Social system
44
45 604 of the Alaotran gentle lemur (*Haplemur griseus alaotrensis*): genetic characterization
46
47 605 of group composition and mating system. *American Journal of Primatology*, 57, 157-
48
49 606 176. DOI: 10.1002/ajp.10046
- 50
51
52
53 607 Palombit, R. A. (1994). Extra-pair copulations in a monogamous ape. *Animal*
54
55 608 *Behaviour*, 47, 721-723.

- 1
2
3 609 Pollock, J.I. (1975). *The social behaviour and ecology of Indri indri*. Doctoral
4
5
6 610 dissertation, University of London.
7
8
9 611 Porter, A. M., Grote, M. N., Isbell, L. A., Fernandez-Duque, E., & Di Fiore, A. (2017).
10
11 612 Delayed dispersal and immigration in equatorial sakis (*Pithecia aequatorialis*): Factors
12
13 613 in the transition from pair-to group-living. *Folia Primatologica*, 88, 11-27. DOI:
14
15
16 614 10.1159/000464147
17
18
19 615 Reichard, U. (1995). Extra-pair copulations in a monogamous gibbon (*Hylobates*
20
21 616 *lar*). *Ethology*, 100, 99-112. 10.1111/j.1439-0310.1995.tb00319.x
22
23
24 617 Reichard, U. (2003). Monogamy: past and present. In U. Reichard & C. Boesch (Eds.)
25
26 618 *Monogamy: Mating strategies and partnerships in birds, humans and other mammals*
27
28 619 (pp. 3-25). Cambridge, UK: Cambridge University Press.
29
30
31
32 620 Reichard, U. H. (2018). Monogamy in Primates. In W. Trevathan, M. Cartmill, D.
33
34 621 Dufour, C. Larsen, D. ORourke, K. Rosenberg and K. Strier (Eds.), *The International*
35
36 622 *Encyclopedia of Biological Anthropology*. John Wiley and Sons, Inc.
37
38 623 doi:10.1002/9781118584538.ieba0326
39
40
41
42
43 624 Sommer, V., & Reichard, U. (1997). Group encounters in wild gibbons (*Hylobates lar*):
44
45 625 agonism, affiliation, and the concept of infanticide. *Behaviour*, 134, 1135-1174. DOI:
46
47 626 10.1163/156853997X00106
48
49
50
51 627 Schülke, O., Kappeler, P. M., & Zischler, H. (2004). Small testes size despite high extra-
52
53 628 pair paternity in the pair-living nocturnal primate *Phaner furcifer*. *Behavioral Ecology*
54
55 629 *and Sociobiology*, 55, 293-301. DOI: 10.1007/s00265-003-0709-x
56
57
58
59
60

- 1
2
3 630 Tecot, S. R., Singletary, B., & Eadie, E. (2016). Why “monogamy” isn't good enough.
4
5 631 *American Journal of Primatology*, 78, 340-354. DOI: 10.1002/ajp.22412
6
7
8 632 Torti, V., Gamba, M., Rabemananjara, Z. H., & Giacoma, C. (2013). The songs of the
9
10 633 indris (Mammalia: Primates: Indridae): contextual variation in the long-distance calls of
11
12 634 a lemur. *Italian Journal of Zoology*, 80, 596-607. DOI: 10.1080/11250003.2013.845261
13
14
15
16 635 Torti, V., Bonadonna, G., De Gregorio, C., Valente, D., Randrianarison, R. M., Friard, O.,
17
18 636 ... & Giacoma, C. (2017). An intra-population analysis of the indris' song dissimilarity in
19
20 637 the light of genetic distance. *Scientific Reports*, 7(1), 10140. DOI: 10.1038/s41598-017-
21
22 638 10656-9
23
24
25
26 639 van Schaik, C.P., Kappeler, P.M. (2003). The evolution of social monogamy in primates.
27
28 640 In U. Reichard & C. Boesch (Eds.) *Monogamy: Mating strategies and partnerships in*
29
30 641 *birds, humans and other mammals* (pp. 59-80). Cambridge, UK: Cambridge University
31
32 642 Press.
33
34
35
36 643 Weir, J.S. (2014). *Infant Development and Maternal Strategies in the Two Largest*
37
38 644 *Lemurs: The Diademed Sifaka (Propithecus diadema) and the Indri (Indri indri)*.
39
40 645 Doctoral dissertation. University of Victoria.
41
42
43
44
45 646 Zaonarivelo, J. R., Sommer, J. A., Shore, G. E., McGuire, S. M., Engberg, S. E.,
46
47 647 Brenneman, R. A., & Loid, E. E. (2007). Isolation and characterization of 20
48
49 648 microsatellite marker loci from the Indri (*Indri indri*) genome. *Molecular Ecology Notes*,
50
51 649 7, 25-28. DOI: 10.1111/j.1471-8286.2006.01451.x
52
53
54
55
56
57
58
59
60

Table 1 Annual group composition of the seven indri groups, since the habituation to 2015, including name and sex of the individuals. Gray cells: presence of an individual in the group. *Italic* individuals: genetic data not available. **Bold** names: offspring born in the group during the study period.

Group	Individual	Sex	2009	2010	2011	2012	2013	2014	2015
1MZ	Bevolo (R)	Female							
	Jery (R)	Male							
	<i>Fotsy (>3[†])</i>	<i>Male</i>							
	Maintso	Female							
	Berthy	Female							
	Akora	<i>Unknown</i>							
2MZ	Soa (R)	Female							
	Max (R)	Male							
	Kinga	<i>Unknown</i>							
	Fanihy	Female							
	Afo	<i>Unknown</i>							
3MZ	Mena (R)	Female							
	Ratsy (R)	Male							
	<i>Mahagaga (R)</i>	<i>Male</i>							
	<i>Tsara (>3[†])</i>	<i>Male</i>							
	Zandry	Female							
	Blague	Male							
	Faly	<i>Female</i>							
	Laro	<i>Unknown</i>							
<i>Tonga (>3[†])</i>	<i>Female</i>								
4MZ	Eva (R)	Female							
	Koto (R)	Male							
	Hendri (>3 [†])	Male							
	Gibet	Male							
	Nic	<i>Unknown</i>							
6MZ	Befotsy (R)	Female							
	Zokybe (R)	Male							
	<i>Marco (>3[†])</i>	<i>Male</i>							
	<i>Guido (>3[†])</i>	<i>Male</i>							
	Tsiky	Male							
	Hira	<i>Unknown</i>							
8MZ	Bemasoandro (R)	Female							
	Jonah (R)	Male							
	Cesar (>3)	Male							
	Zafy	Male							
	Mika	<i>Unknown</i>							

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

654

9MZ	Sissie (R)	Female		
	Emilio (R)	Male	EPC‡	
	Njaka	<i>Male</i>		
	Ovy	Male		
	Dosy	<i>Unknown</i>		

For Peer Review

655 **Table 2** Microsatellite Loci Applied with Respective Primers, Annealing Temperatures and Number of PCR (Torti et al. 2017)

656

Locus	Forward primer	Reverse Primer	Repeat motif	Annealing temp. (°C)	Number of PCR cycles	Size range (bp)
67HDZ25	GGACCCTAATTCA AATATCACCTC	GGCATTCTACT CCAGGTTGG	(CA)16	54	50	218-253
67HDZ62	AGCCCTTTCTCTC AATGCC	CCTTCTTTGTTAT CTTTCTGCATC	(GT)21	54	50	203-217
67HDZ18	GGACTGGTAGAT TTCTGGGTTTAG	CAGCCACTCCAA TGCAAAG	(CA)7C(CA)15	60	50	164-190
67HDZ55	TCAGGAGTTGGG ACCAGGG	ATGAAGGGATG GAGGTGGG	(GT)18	60	50	312-334
67HDZ180	TCCCCTCCTCAGT CATTCTC	CGTGAAGCTCGT GTGTATGG	(CA)17	60	50	113-136
67HDZ39	CAGAGCCAGGGT TCAAATTC	TTGTCTTTTCTGC CACTGTAGG	(CA)11	60	50	148-162

657 **Table 3** Per Locus Summary of six Microsatellites Markers for *Indri indri*

Locus	# Individuals	# Alleles	Observed Heterozygosity	Expected Heterozygosity	Hardy Weinberg Equilibrium p-value	Allele Null Probability
67HDZ25	26	13	0.923	0.876	0.1090	-0.0487
67HDZ62	25	9	0.840	0.814	0.4591	-0.0290
67HDZ18	26	11	0.962	0.883	0.8488	-0.0529
67HDZ55	23	14	0.913	0.928	0.7909	-0.0042
67HDZ180	26	11	0.846	0.876	0.8869	0.0091
67HDZ39	26	12	1.000	0.908	0.1046	-0.0589

658 **Table 4** Paternity analyses results of 12 offspring and respective known and potential parents of seven indri groups. For the potential
 659 fathers we included the number of loci mismatch and the sign of the LOD score (positive or negative). . We indicated if the paternity was

Group	Y of Birth	Offspring	Mother (# mismatches)	Potential Sires Sampled		Assigned Paternity		
				Paired Male (# mismatches, LOD score sign)	Neighboring Male(s) Genotyped (# mismatches, LOD score sign)	Paired Male	EPC Male [†]	Inconclusive Assignment
1MZ	2010	Maintso	Bevolo (0)	Jery (0, +)	Ratsy (6, -), Zokybe (6, -), Cesar (4, -)	x		
1MZ	2012	Berthy	Bevolo (0)	Jery (0, +)	Ratsy (5, -), Zokybe (6, -)	x		
2MZ	2012	Fanihy	Soa (0)	Max (0, +)	Ratsy (4, -)	x		
3MZ	2010	Zandry	Mena (0)	Ratsy (0, +)	Cesar (3, -), Jery (4, -), Max (4, -), Emilio (5, -)	x		
3MZ	2012	Blague	Mena (1)	Ratsy (0, +)	Jery (5, -), Max (5, -), Emilio (6, -)	x	excluded	
4MZ	N/A	Hendri	Eva (1)	Koto (0, +)	Jery (3, -), Emilio (4, -), Max (4, -), Jonah (4, -), Zokybe (5, -), Cesar (5, -), Ratsy (6, -)	x		
4MZ	2012	Gibet	Eva (0)	Koto (0, +)	Hendri (4, -), Zokybe (6, -)	x		
6MZ	2010	Tsiky	Befotsy (0)	Zokybe (6, -)	Hendri (2, -), Ratsy (4, -), Cesar(4, -), Koto (5, -), Jery (5, -), Max (5, -), Emilio (6, -)			x
6MZ	2014	Hira	Befotsy (0)	Zokybe (0, +)	Koto (2, -), Hendri (3, -), Jery (5, -)	x		
8MZ	N/A	Cesar	Bemasoandro (0)	Joanh (0, +)	Jery (2, -), Ratsy (2, -), Max (2, -), Zokybe (3, -), Koto(4, -) Emilio (4, -), Hendri (5, -)	x		
8MZ	2012	Zafy	Bemasoandro (0)	Joanh (0, +)	Cesar (3, -)	x		
9MZ	2013	Ovy	Sissie (0)	Emilio (0, +)	Ratsy (6, -)	x		
Percentage of paternity assignement						92%	0%	8%

660 assigned to the paired male or remained inconclusive, we specified if the male involved in the extra pair-copulation was excluded as sire.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

For Peer Review

† EPC male: There is only one case of EPC reported in indris (Bonadonna et al. 2014). Emilio is the male that participated in the EPC event observed during the mating season that preceded the birth of Blague.

1
2
3 662 **Figure 1.** Study site and indri groups included in the study
4
5
6 663 Study site located in Center-Eastern Madagascar (A), in the Maromizaha forest,
7
8 664 accessible from the village Anevoka (B). The multipurpose center (18°58'34" S –
9
10 665 48°27'53" E) was used as base-camp (C). Box C is showing the spatial distribution of
11
12 666 the indri groups included in the genetic analysis (solid green polygons), and the
13
14 667 schematic representation of unsampled neighboring groups (gray striped polygons,
15
16 668 GX1 to GX7). The uppercase names within each studied group are the social fathers;
17
18 669 the lowercase names are the offspring genotyped. "*" denotes the only individual that
19
20 670 resulted with inconclusive paternity assignment. Original satellite images (Google Earth
21
22 671 - Image © 2018 CNES / Airbus; downloaded on June 27, 2018) have been graphically
23
24 672 simplified and adjusted with GIMP 2.10.2. Map of the indri territories created with
25
26 673 ArcGIS® 10.2 (ESRI).
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Research Highlights

- In 92% of cases, the paired male of the group did not have any loci mismatch with the offspring. In the only case of paternity mismatch, we were not able to assign the sire identity.
- Our findings suggest that genetic monogamy is the norm in indri, although EPC can occasionally occur.

For Peer Review




Figure 1. Study site and indri groups included in the study

Study site located in Center-Eastern Madagascar (A), in the Maromizaha forest, accessible from the village Anevoka (B). The multipurpose center ($18^{\circ}58'34''$ S – $48^{\circ}27'53''$ E) was used as base-camp (C). Box C is showing the spatial distribution of the indri groups included in the genetic analysis (solid green polygons), and the schematic representation of unsampled neighboring groups (gray striped polygons, GX1 to GX7).

The uppercase names within each studied group are the social fathers; the lowercase names are the offspring genotyped. "*" denotes the only individual that resulted with inconclusive paternity assignment. Original satellite images (Google Earth - Image © 2018 CNES / Airbus; downloaded on June 27, 2018) have been graphically simplified and adjusted with GIMP 2.10.2. Map of the indri territories created with ArcGIS® 10.2 (ESRI).

127x124mm (300 x 300 DPI)



Spatial distribution of the indri groups included in the genetic and paternity analyses (solid green polygons), and the schematic representation of unsampled neighboring groups (gray striped polygons, GX1 to GX7).

The uppercase names within each studied group are the social fathers; the lowercase names are the offspring tested. "*" denotes the only individual we found with inconclusive paternity assignment.

12x16mm (300 x 300 DPI)