

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Estrogens Inhibit Amyloid- $\beta$ -Mediated Paired Helical Filament-Like Conformation of Tau Through Antioxidant Activity and miRNA 218 Regulation in hTau Mice**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1765727> since 2021-01-03T15:08:42Z

*Published version:*

DOI:10.3233/jad-200707

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

5 Michela Guglielmino<sup>a</sup>, Giusti Manassero<sup>a</sup>, Valeria Vascraveo<sup>a</sup>, Marika Venezia<sup>a</sup>  
6 Tabaton<sup>c</sup> and Elena Tamagno<sup>a,b,\*</sup>  
7 <sup>a</sup>*Department of Neuroscience, University of Torino, Torino, Italy*  
8 <sup>b</sup>*Neuroscience Institute of Cavalieri Ottolenghi Foundation (NICO), University of Torino, C*  
9 *Italy*  
10 <sup>c</sup>*Unit of Geriatric Medicine, Department of Internal Medicine and Medical Specialties (D*  
11 *Genova, Genova, Italy*

Accepted 15 July 2020

## 12 **Abstract.**

13 **Background:** The risk of developing Alzheimer's disease as well as its progression and severity are kn  
14 men and women, and cognitive decline is greater in women than in men at the same stage of disease an  
15 at least in part on estradiol levels.

16 **Objective:** In our work we found that biological sex influences the effect of amyloid- $\beta_{42}$  ( $A\beta_{42}$ ) mono  
17 tau conformational change.

18 **Methods:** In this study we used transgenic mice expressing the wild-type human tau (hTau) whic  
19 intraventricular (ICV) injections of  $A\beta$  peptides in nanomolar concentration.

20 **Results:** We found that  $A\beta_{42}$  produces pathological conformational changes and hyperphosphorylat  
21 male or ovariectomized female mice but not in control females. The treatment of ovariectomized fe  
22 replacement protects against the pathological conformation of tau and seems to be mediated by antiox  
23 as the ability to modulate the expression of miRNA 218 linked to tau phosphorylation.

24 **Conclusion:** Our study indicates that factors as age, reproductive stage, hormone levels, and the inte  
25 factors should be considered in women, in order to identify the best appropriate therapeutic approach in pr  
26 impairment.

27 **Keywords:** Alzheimer's disease, antioxidants, estradiol, miRNA, tau protein

## 28 **INTRODUCTION**

29 Hallmarks of Alzheimer's disease (AD) are the  
30 accumulation of amyloid- $\beta$  ( $A\beta$ ) peptides in amy-

loid plaques, and the aggregation  
form neurofibrillary tangles.

$A\beta$  derives from the amyloid  
precursor ( $A\beta$ PP) through  $\beta$  site APP  
(BACE) 1 and  $\gamma$ -secretase proc  
ates multiple C-termini, most en  
and 42.  $A\beta_{42}$  aggregates more  
bly than  $A\beta_{40}$  through sequentia

---

\*Correspondence to: Elena Tamagno PhD, Neuroscience Insti-  
tute Cavalieri Ottolenghi, University of Torino, Regione Gonzole  
10, 10043 Orbassano, Torino, Italy. Tel.: +390116707764; E-mail:  
elena.tamagno@unito.it

53 1) were subjected to intravitreal (ICV) injections of A $\beta$  peptides in nanomolar concentration. We discovered that A $\beta$ <sub>42</sub> monomers, but not oligomers: 54 1) produce paired helical filament-like conformation of tau protein, and 2) induce two phosphorylated epitopes which are not present in normal tau (Ser396 and 55 Ser422) through the activation of GSK3 $\beta$ , JNK, and ERK 1/2 kinases [4]. 56 57 58 59 60

61 Recent epidemiological studies showed that two-thirds of AD patients are women [5], and this 62 fact cannot be attributed only to their higher life expectancy. In this connection, the loss of estradiol 63 might be one of the factors leading to declining cognitive function in women [6]. 64 65 66

67 Of note, we found that oxidative stress, together with important oxidative stress-related risk factors 68 related to AD, such as hypoxia, hyperglycemia, and hypercholesterolemia, are potential causes of 69 the increased BACE1 activity [7]. In AD, estrogen neuroprotective activity is exerted at multiple 70 levels. Preclinical data showed that, in addition to their action against neuroinflammation and oxidative 71 stress, estrogens are able to influence both the main players of neurodegeneration, A $\beta$ , and tau 72 [8]. 73 74 75 76 77

78 In this paper, we pursue the hypothesis that biological sex influences the effect of A $\beta$ <sub>42</sub> monomers 79 on pathological tau conformational change. Our data revealed that A $\beta$ <sub>42</sub> monomers produce the 80 pathological conformational changes and hyperphosphorylation of tau protein in male or ovariectomized 81 female mice but not in control female. The treatment of ovariectomized females with estradiol replacement 82 protects against the pathological conformation of tau. 83 The hypothesized protective mechanism is mediated 84 both by their antioxidant activity and by their ability 85 to modulate the expression of miRNA 218 linked to tau phosphorylation. 86 87 88 89 90

53 5) , mouse tau gene (forward CCCACCTGTAAC-3', reverse GTATGTCCACCC-3'), and di 54 (forward 5'-CAGGCTTTGAA reverse 5'-TGAACCTGTGGG 55 3'). Mice were maintained on 129/SvJae/C57BL/6 background 56 were kept on a 12 h light/dark and water available *ad libitum*. 57 procedures on live animals were under the supervision of a licensed veter 58 to: 1) European Communities (November 24, 1986; 86/609/ Ministry of Health and University 59 tutional guidelines on animal welfare on Care and Protection of living animals 60 experimental or other scientific purposes (authorization No. 17/2010-B, June 30, 61 *hoc* Ethical Committee of the University (http://www.unito.it/unitoWAR/pagine/ricerca1/Ricerca\_comitato1). 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90

Two groups of 2-month-old male mice were treated for 3 h ( $n=60$ ) with O<sub>2</sub>/N<sub>2</sub>O anesthesia, hTau mice were injected with A $\beta$  peptides or saline. The sites used for injection were: anteroposterior lateral, 1.2 mm relative to Bregma; anteroposterior medial, 1.7 mm from the dural surface, validated by injecting one mouse with saline (1  $\mu$ l).

### *Ovariectomy*

Two groups of 2-month-old female mice underwent bilateral ovariectomy (OVX) or sham-operated groups. The bilateral ovariectomized female mice were exposed to a skin incision in the dorsal skin and n

151 centrifuged with sterile double distilled water,  
152 centrifuged at 10000 g for 10 min to remove possi-  
153 bly aggregates and then intraventricularly injected.  
154 The quality of A $\beta$  preparations was controlled using  
155 atomic force microscopy (AFM). AFM was car-  
156 ried out on a Multimode AFM with a Nanoscope  
157 V system operating in Tapping Mode using stan-  
158 dard antimony(n)-doped Si probes (T: 3.5–4.5 mm,  
159 L: 115–135 mm, W: 30–40 mm, f<sub>0</sub>:313–370 kHz, k:  
160 20–80 N/m) (Bruker). The scan rate was tuned pro-  
161 portionally to the area scanned and was kept in the  
162 0.5–1.2 Hz range. The sample was then diluted to  
163 5  $\mu$ M with PBS, and 50  $\mu$ l of solution was spot-  
164 ted onto a freshly cleaved muscovite mica disk and  
165 incubated for 5 min. The disk was then washed with  
166 ddH<sub>2</sub>O and dried under a gentle nitrogen stream.  
167 Samples were analyzed with the Scanning Probe  
168 Image Processor (SPIP Version 5.1.6 released April  
169 13, 2011) data analysis package (Nanoscience Instru-  
170 ments, Phoenix, AZ, USA). SPIP software was used  
171 to analyze the distribution of the molecular assem-  
172 blies of the different populations in terms of height  
173 and diameter, as previously described [10]. Our con-  
174 trols were hTau mice ICV injected with saline. The  
175 experiments were done four weeks after ovariecto-  
176 my. After two weeks from ovariectomy surgery, one  
177 group of female mice was subjected to a daily sub-  
178 cutaneous injection of 17 $\beta$ -estradiol (E<sub>2</sub>) for three  
179 weeks (1  $\mu$ g/Kg) [11]. Animals were allowed to  
180 recover for at least three weeks before experiments  
181 were performed and subsequently were subjected to  
182 intracerebroventricular injection of A $\beta$ <sub>42</sub> monomers  
183 (200 nM) or saline and sacrificed after 3 h.

#### 184 *Antibodies and immunoblot analysis*

185 Immunoblot analysis was performed using the  
186 following antibodies: MC1 (kind gift from Dr.

150 mM NaCl, 1 mM EGTA, 1 mM PMSF, phosphatase and protease inhibitors), then centrifuged at 10,000 g for 10 min to isolate soluble proteins. Supernatant (supernatant solution) were collected and incubated with sarkosyl (5% final concentration) at 4°C. The sarkosyl mixtures were centrifuged in Beckman SW 55 Ti rotor for 1 h at 4°C. Pellets were resuspended in sample buffer to obtain sarkosyl-soluble lysates (20  $\mu$ g) were run on 3–15% gradient PAGE gel (Invitrogen) and transferred to nitrocellulose membrane. Blots were probed with anti-A $\beta$  (1:1000) (5% no fat milk) and incubated with primary antibodies. Peroxidase-conjugated secondary antibodies were incubated at room temperature (RT) and developed using ECL-Plus Forte Western substrate (Wako Pure Chemical Industry, Japan). Densitometric values were normalized to  $\beta$ -actin.

#### *Total antioxidant capacity*

To evaluate the antioxidant capacity of brain tissues, we performed the total antioxidant capacity (TAC) dosage kit (ab65329, Abcam) was performed on total extracts according to the manufacturing protocol.

#### *Quantitative determination of 17 $\beta$ -estradiol*

To quantify E2 levels, blood samples were obtained from mice. To perform the measurement, we used a commercially available ELISA kit (E2 EIA kit, catalog # ADI-901-174) according to the manufacturer's protocol.




Fig. 1. Intracerebroventricular treatment with  $A\beta_{42}$  causes a change in tau conformation only in male or ovariectomized female mice. Representative western blot of brain samples from control (saline) and treated  $A\beta_{42}$  peptides for ICV male and female mice were probed with conformational tau antibody (MC1) and a total tau antibody (Tau 5) for detection. Some female mice were subjected to ovariectomy and served as loading control. Densitometric quantification shows an increase of the total protein level of both MC1 and Tau 5 in ovariectomized female mice injected or not with  $A\beta_{42}$ . The data are mean  $\pm$  standard error of the mean (SEM); \* $p < 0.05$  versus control; § $p < 0.05$  versus OVX by one-way ANOVA followed by Bonferroni post test  $n = 5$ .

233 *MicroRNA isolation and quantitative real time*  
 234 *PCR*

235 MicroRNA was isolated from brains of female  
 236 mice using the MagMAXmirVana kit and according  
 237 to manufacturer's protocol (Applied Biosystems,  
 238 Foster City, CA, USA). Subsequently, cDNA syn-  
 239 thesis was performed using the TaqMan Mi-croRNA  
 240 Reverse Transcription Kit (Applied Biosystems,  
 241 Foster City, CA, USA) and a RT-primer pool con-

taining microRNA-specific stem  
 miR218 (Mature miRNA Sequence: GAUCUAACCAUGU) and for snRNA U6 (Mature snRNA Sequence: GTGCTCGCTTCGACTAAAATTGGAACGATACAGCATGGCCCCTGCGCAAGGATTCGTGAAGCGTTCCATATTTT).

Each qPCR contained 1.3  $\mu$ L of 1  $\mu$ L 20X TaqMan MicroRNA Assay and 0.3  $\mu$ L 2X TaqMan Universal PCR Master




Fig. 2. The conformational change mediated by  $A\beta_{42}$  is induced by protein hyperphosphorylation. A) Representative extracts from control (saline) and treated  $A\beta_{42}$  peptides for ICV male and female hTau mice using antibodies specific for pathological tau phosphorylation sites such as AT8, pS422, and pS396. Some female mice were subjected to ovariectomy as loading control. Densitometric analysis shows a significant increase of total protein levels of AT8, pS422, and pS396 in ovariectomized female mice. B–D) Representative western blot of brain extracts from control (saline) and treated male and female hTau mice using pJNK (B), pERK1/2 (C), and pGSK3 $\beta$  (D). Some female mice were subjected to ovariectomy as loading control. Densitometric analysis shows a significant increase of total protein levels of all the three kinases in ovariectomized female mice. The data are mean  $\pm$  standard error of the mean (SEM);  $\xi p < 0.05$ ;  $**p < 0.01$  versus OVX by one-way ANOVA followed by Bonferroni post test  $n = 5$ .

264 All values were presented as mean  $\pm$  standard error  
265 of the mean (SEM). Means were compared by one or  
266 two-way analysis of variance (ANOVA) with Bon-  
267 ferroni as a *post-hoc* test. Values of  $*p < 0.05$  were  
268 considered significant,  $**p < 0.01$  very significant,  
269 and  $***p < 0.001$  extremely significant.

## 270 RESULTS

271 *Intracerebroventricular treatment with A $\beta$ <sub>42</sub>*  
272 *causes a change in tau conformation only in*  
273 *male or ovariectomized female mice*

274 Figure 1 shows that, as previously demonstrated,  
275 the intracerebroventricular injection of 200 nM A $\beta$ <sub>42</sub>  
276 in male hTau mice is able to determine a pathologi-  
277 cal conformational change of the tau protein. The  
278 same treatment is able to determine this effect only  
279 in female mice after ovariectomy. The same result  
280 was obtained by evaluating the total tau protein lev-  
281 els. As can be seen, treatment with A $\beta$ <sub>42</sub> significantly  
282 increases total tau levels in male (1.5-fold increase)  
283 and ovariectomized female mice (1.5-fold increase)  
284 with respect to control female mice injected or not  
285 with A $\beta$ <sub>142</sub>.

286 *The conformational change mediated by A $\beta$ <sub>42</sub> is*  
287 *induced by protein hyperphosphorylation*

288 To understand if the conformational change of tau  
289 was due to hyperphosphorylation, we measured the  
290 phosphorylation of some specific sites related to the  
291 pathology by western blotting. Phosphorylation lev-  
292 els were studied through the use of the AT8 antibody  
293 (which recognizes the Ser 202/Thr 205 epitopes), the  
294 antibody S396 and S422. These are all phosphoryla-  
295 tion sites closely associated with disease progression.

(+3-fold increase). A similar result was obtained  
measuring the nuclear levels of p-tau, which increase in the same groups  
times compared to the controls (Fig. 2).

*Estradiol hormone therapy protects female mice against*  
*A $\beta$ <sub>42</sub>-mediated tau conformational change*

To confirm that the presence of estradiol is involved in the different effect experimentally observed with A $\beta$ <sub>42</sub> on the pathological change of tau, groups of female mice injected or not were subcutaneously treated with estradiol (E<sub>2</sub>) (1  $\mu$ g/kg) and fed with a soybean diet for 4 weeks. We first tested the validity of the method for measuring the levels of estradiol in the blood of the different groups. As can be seen, oophorectomy significantly decreases circulating estradiol levels. E<sub>2</sub> treatment determines an increase in estradiol levels that becomes significantly higher with respect to control females (+100%) (Fig. 3). Figure 4 shows that treatment with estradiol completely blocks the pathological conformational change of total tau mediated by A $\beta$ <sub>42</sub> in ovariectomized females. Then, to further confirm the effect of estradiol, we studied the insolubilization of tau with sarkosyl detergent technique, and the presence of a tau band at approximately 75 kDa was revealed with Tau 46 antibody. Figure 4B shows that A $\beta$ <sub>42</sub> in ovariectomized female, with estradiol blocks the aggregation of tau (Fig. 4B).

*Estradiol therapy protects female mice against*  
*A $\beta$ <sub>42</sub>-mediated tau hyperphosphorylation*

Figure 5 shows that enrichment of tau in the nucleus is also followed by complete phosphorylation of tau in the nucleus.

Fig. 3. Estradiol hormone therapy significantly increases E<sub>2</sub> levels. Groups of female mice, ovariectomized or not, were subcutaneously treated with E<sub>2</sub> (1 μg/kg) and fed with a soy-free diet for three weeks. To test the validity of the treatment, we measured E<sub>2</sub> levels in serum of our experimental groups. We observed that the ovariectomy induces a significant decrease in hormone levels, whereas the treatment protects the decrease of E<sub>2</sub> levels, that significantly increased with respect to controls. The data are mean ± standard error of the mean (SEM); \**p* < 0.05; \*\**p* < 0.01 versus control by one-way ANOVA followed by Bonferroni post test *n* = 6.

344 mediated phosphorylation, after oophorectomy, of  
345 pathology-related sites (Fig. 5A). As expected, the  
346 kinases involved in the phosphorylation of the above  
347 sites do not appear induced; thus, as observed in  
348 Fig. 5B, the levels of nuclear pJNK, pERK, and  
349 pGSK3β are absolutely comparable to the levels of  
350 the control females.

351 *Estradiol hormone therapy protects male hTau*  
352 *mice against Aβ<sub>42</sub>-mediated tau conformational*  
353 *change and hyperphosphorylation*

354 To further confirm the protective role of estradiol  
355 on the pathological conformational change of  
356 tau and its hyperphosphorylation, we also treated  
357 hTau male mice with estradiol, following the same  
358 protocol as the females. As can be seen in Fig. 6,  
359 the treatment with estradiol is able to completely  
360 protect both the conformational change of tau as  
361 revealed by the significant increase of the band  
362 revealed with MC1 antibody as well as its hyper-  
363 phosphorylation revealed by using AT8 antibody  
364 that recognizes Ser202/Thr205 phospho-epitopes  
365 (Fig. 6).

and the simultaneous intracerebral  
tion of Aβ<sub>42</sub> induces a further d  
parameter (-70%). Treatment with  
the drop-in antioxidant capacity b  
to control values (Fig. 6A).

Finally, we measured levels  
recent discoveries demonstrate th  
tors are able to modulate the expres  
involved in tau phosphorylation [  
has been found that increase of m  
level of target protein tyrosine p  
consequent enhancement of tau p

We observed that levels of miR2  
higher in ovariectomized female m  
with Aβ<sub>42</sub>, whereas the E2 treatm  
a total protection of the miRNA .  
It is interesting to note that the m  
male hTau mice treated with Aβ  
obtained in females after oophorec

## DISCUSSION

In our work we pursued the hyp  
influences the effect of Aβ<sub>42</sub> mono  
cal tau conformational change. Our  
Aβ<sub>42</sub> produced pathological conf  
and hyperphosphorylation of tau  
or ovariectomized female mice  
female. The risk of developing  
progression and severity are know  
different in men and women [14  
the drop-in estrogen and the path  
by data indicating that early me  
the risk of developing dementia  
cognitive decline is greater in w  
at the same stage of disease and  
correlated with estrogen levels [1  
the therapeutic potential of estro






Fig. 4. Estradiol hormone therapy protects against A $\beta$ <sub>42</sub>-mediated tau conformational change. A) Representative samples from control (saline) and treated A $\beta$ <sub>42</sub> peptides for ICV female hTau mice using a conformational tau antibody (Tau 5) for detection. Some female mice were subjected to ovariectomy and/or to E<sub>2</sub> (1  $\mu$ g/kg) and 4 weeks.  $\beta$ -actin served as loading control. Densitometric quantification shows an increase of the total protein level of tau in ovariectomized female mice injected or not with A $\beta$ <sub>42</sub>; the treatment with estradiol completely protects the control from the increase of tau protein. B) Representative western blot of insoluble tau fraction by sarkosyl detergent technique (saline) and treated A $\beta$ <sub>42</sub> peptides for ICV female hTau using Tau 46 antibody for detection.  $\beta$ -actin served as loading control. To show the effect of estradiol on the aggregation of tau protein, we showed a band at approximately 75 kDa molecular weight revealed with Tau 46 antibody. The data are mean  $\pm$  standard error of the mean (SEM); \* $p$  < 0.05; \*\* $p$  < 0.05 versus OVX by one-way ANOVA followed by Bonferroni post test  $n$  = 5.

415 active in the last years and recently it has taken new  
 416 life, thus finding new therapeutic approaches for AD  
 417 is one of the most important challenges of modern  
 418 medicine. Numerous experimental evidences have

shown that estrogens have protective effects against the  
 induction of neuroinflammation and oxidative stress  
 in animal models of AD [16–18]. The encouraging  
*in vitro* results have been largely confirmed in clinical trials




Fig. 5. Estradiol hormone therapy protects against A $\beta$ <sub>42</sub>-mediated tau hyperphosphorylation. A) Representative western blots from control (saline) and treated A $\beta$ <sub>42</sub> peptides for ICV female hTau mice using antibodies specific for the detected phosphorylation sites such as AT8, pS422, and pS396. Some female mice were subjected to ovariectomy and/or to E<sub>2</sub> diet for three weeks.  $\beta$ -actin served as loading control. Densitometric analysis shows an increase of total protein levels of pS396 in ovariectomized female mice injected with or not with A $\beta$ <sub>42</sub>; the treatment with estradiol completely protects the levels of tau protein. B–D) Representative western blot of brain extracts from control (saline) and treated A $\beta$ <sub>42</sub> peptides for ICV female hTau mice using antibodies specific for pJNK (B), pERK1/2 (C), and pGSK3 $\beta$  (D). Some female mice were subjected to ovariectomy and/or to E<sub>2</sub> (1  $\mu$ M) diet for three weeks.  $\beta$ -actin served as loading control. Densitometric analysis shows an increase of total protein levels of pJNK, pERK, and pGSK3 $\beta$  in ovariectomized female mice injected with or not with A $\beta$ <sub>42</sub>; the treatment with estradiol completely protects the activity of these proteins. The data are mean  $\pm$  standard error of the mean (SEM); \* $p$  < 0.05; \*\* $p$  < 0.01 versus control; <sup>§</sup> $p$  < 0.05 versus OVX followed by Bonferroni post test  $n$  = 5.




Fig. 6. Estradiol hormone therapy protects male mice against  $A\beta_{42}$ -mediated tau conformational change as well as tau phosphorylation. Representative western blot of brain samples from control (saline) and treated  $A\beta_{42}$  peptides for ICV male hTau mice with MC1 and a specific antibody for tau pathological phosphorylation (AT8) for detection. Some male mice were subjected to a soy-free diet for three weeks.  $\beta$ -actin served as loading control. Densitometric quantification shows an increase in phosphorylation of both MC1 and AT8 in male mice injected or not with  $A\beta_{42}$ ; the treatment with estradiol completely protects the control mice from the increase of tau phosphorylation. The data are mean  $\pm$  standard error of the mean (SEM); \*\* $p < 0.01$  versus control followed by Bonferroni post test  $n = 5$ .

423 clinical trials had shown that the use of replacement  
 424 therapy significantly increased the risk of dementia  
 425 and cognitive decline [19, 20]. These studies sug-  
 426 gested that patients aged 65 and over, already in  
 427 menopause for a long time, did not represent an  
 428 adequate experimental group, because replacement  
 429 therapy is indicated for women who have just gone  
 430 through menopause, and suggested the presence of a  
 431 therapeutic window useful for this type of therapeu-  
 432 tic approach [21]. More recently, emerging evidence  
 433 suggests that protein tau could be a potential target for  
 434 estrogens. It has been demonstrated that  $17\beta$  estro-  
 435 diol promotes tau dephosphorylation *in vitro* in rat  
 436 cortical neurons and SH-SY5Y neuronal cells [22].  
 437 Other authors confirmed these results showing that  
 438 E2 prevent the phosphorylation of tau in an estro-  
 439 gen receptor-mediated and dose-dependent manner  
 440 [23]. *In vivo* studies have shown that estrogenic treat-  
 441 ment increases GSK3 $\beta$  phosphorylation in Ser 9/21,  
 442 a site that inactivates the kinase activity, protecting  
 443 the phosphorylation of pathological sites related to

disease progression [24]. Moreover, these results suggest that estrogens exert their neuroprotective effects through their alpha-type receptor signaling pathway with insulin-like growth factor 1 receptor by incorporating itself into a macromolecular complex that includes phosphoinositide-dependent kinase and protein kinase A (AKT). The activation of this signal pathways leads to an inhibition of GSK3 $\beta$  activity and therefore to a reduction in tau phosphorylation [25]. Our results confirm the neuroprotective effect of estrogens on the pathological hyperphosphorylation of tau mediated by  $A\beta$  after intracerebroventricular injection with  $A\beta_{42}$  in male mice. From our results, it cannot be concluded if estradiol acts as an antioxidant or as a neuroprotective agent, therefore, in a mode independent of its antioxidant activity or modulates at the receptor level the downstream pathways. We showed that it is not sufficient that, under  $A\beta$  treatment, hyperphosphorylation of tau is reduced; these results suggest a cellular mechanism of estradiol.




Fig. 7. Estradiol hormone therapy protects against oxidative stress and downregulate miRNA 218. A) Evaluation of antioxidant capacity in female mice subjected to ovariectomy and or to E<sub>2</sub> (1 µg/kg) and soy-free diet for three weeks. We show that ovariectomy significantly decreases the antioxidant capacity respect to control females, whereas the treatment with estradiol completely protects the increase. B) miRNA 21 levels capacity in female mice subjected to ovariectomy and or to E<sub>2</sub> (1 µg/kg) and soy-free diet for three weeks. We show that ovariectomy significantly increases the levels of miRNA 218 with respect to controls female, the treatment with estradiol completely protects the increase. C) Evaluation of the total antioxidant capacity in male mice treated or not with Aβ<sub>42</sub>. We show that Aβ<sub>42</sub> significantly decreases the antioxidant capacity respect to control males. D) miRNA 21 levels capacity in male mice treated or not with Aβ<sub>42</sub> significantly decreases miRNA level respect to control males. \**p* < 0.05; \*\**p* < 0.01 versus control; §*p* < 0.05 versus control. ANOVA followed by Bonferroni post test *n* = 5.

465  
466  
467  
468  
469

A lot of protective mechanisms relating to estrogens have been described in the literature [26]. Most recent studies have revealed that estrogens exert an antioxidant action not only by direct chemical neutralization of reactants, but also by modulating the

expression of antioxidant enzymes and levels of biological reducing agents [27]. In experimental models, we found that the decrease in antioxidant levels was followed by a decrease in antioxidant activity and that this event

with important oxidative stress-related risk factors related to AD such as hypoxia, hyperglycemia, and hypercholesterolemia, are potential causes of the increased BACE1, the crucial enzyme for A $\beta$  production, activity [7].

Recently emerging evidence suggests that estrogens are involved in regulation of microRNAs in many pathological conditions [28]. Rao et al. showed that estradiol regulates particular target miRNAs in a specific tissue and age manner in ovariectomized rats [29]. Furthermore, the deprivation of estrogens caused the progressive loss of regulation of the miRNA, leading to a lack of regulation even after the reintroduction of the estrogens [30]. In our work we focused our attention on the miRNA 218, because it is implicated in the phosphorylation of tau upon estrogen receptor (ER)  $\alpha$  and  $\beta$  activation. There are two known ERs, usually referred to as ER $\alpha$  and ER $\beta$ , and both are widely distributed in the brain [31]. In the brain of patients with AD, both ER $\alpha$  and ER $\beta$  are defective. Mitochondrial ER $\beta$  is reduced in the frontal cortex of female patients with AD [32], and the alternative splicing of ER $\alpha$  mRNA is decreased in the AD brain especially in female patients [33]. Moreover, in the hippocampus of AD patients the ER $\alpha$ -expressing neurons are reduced [34], whereas ER $\beta$  immunoreactivity is increased [35]. These findings indicate a potential role of these two receptors in the pathogenesis of AD. Then it has been reported that the neuroprotection against A $\beta$  toxicity by estrogens requires the expression of both receptors and the activation of mitogen-activated protein kinase pathway [36]. Specifically, Xiong et al. demonstrated opposite effects of these two receptors on tau phosphorylation. ER $\alpha$  overexpression increased miRNA 218 expression and the hyperphosphorylation of tau, whereas ER $\beta$  decreased miRNA 218 expression and tau phosphorylation [13]. Interestingly, a number of miRNA

We thank Mrs. Claudia Blengini for her assistance.  
Authors' disclosures available at [www.j-alz.com/manuscript-disclosure](http://www.j-alz.com/manuscript-disclosure)

## REFERENCES

- [1] Hubin E, van Nuland NA, Broese van Groenou MI, et al. (2014) Transient dynamics of A $\beta$  in the brain in Alzheimer's disease. *Cell Mol Life Sci* **368**, 1039-1050.
- [2] Nimmrich V, Ebert U (2009) Is Alzheimer's disease a presynaptic failure? Synaptic dysfunction and oligomeric beta-amyloid. *Rev Neurosci* **22**, 1-24.
- [3] Puzzo D, Privitera L, Fa' M, Stanzani A, Aziz F, Sakurai M, Ribe EM, Caccamo A, SS, Palmeri A, Arancio O (2011) Estrogen is necessary for hippocampal synaptic plasticity. *Ann Neurol* **69**, 819-830.
- [4] Manassero G, Guglielmotto M, Colombo L, Salmona M, Perry G, Olanow CW, Tabaton M (2016) Beta-amyloid oligomers, produce PHF-like protein. *Aging Cell* **15**, 914-923.
- [5] Beam CR, Kaneshiro C, Jang JY, Lee J, NL, Gatz M (2018) Differences between sexes in incidence rates of dementia and Alzheimer's disease. *Alzheimers Dis* **64**, 1077-1083.
- [6] Laws KR, Irvine K, Gale TM (2010) Cognitive impairment in Alzheimer's disease. *Chiatry* **6**, 54-65.
- [7] Tamagno E, Guglielmotto M, Morbelli E, et al. (2012) Amyloid- $\beta$  production: Mitogenic stress and BACE1. *Neurotox Res* **22**, 1-10.
- [8] Merlo S, Spampinato SF, Sortino M, et al. (2010) Alzheimer's disease: Still an attractive target for drug development from early clinical results. *Alzheimers Dis* **51**, 51-58.
- [9] Andorfer C, Kress Y, Espinoza M, Barde YA, Duff K, Davies P (2003) Estrogen and aggregation of tau in mice expressing tau isoforms. *J Neurochem* **86**, 582-592.
- [10] Messa M, Colombo L, del Favero S, Cagnotto A, Rossi A, Morbin M, Del Giudice M, Salmona M (2014) The peculiar role of

- 599  
600 [16] Villa A, Vegeto E, Poletti A, Maggi A (2016) Estrogens, neuroinflammation, and neurodegeneration. *Endocrinol Rev* **37**, 372-402.
- 601  
602  
603 [17] Khan M, Ullah R, Rehman SU, Shah SA, Saeed K, Muhammad T, Park HY, Jo MH, Choe K, Rutten BPF, Kim MO (2019) 17 $\beta$ -estradiol modulates SIRT1 and halts oxidative stress-mediated cognitive impairment in a male aging mouse model. *Cells* **8**, 928.
- 604  
605  
606  
607  
608 [18] Pandey R, Shukla P, Anjum B, Gupta HP, Pal S, Arjaria N, Gupta K, Chattopadhyay N, Sinha RA, Bandyopadhyay S (2020) Estrogen deficiency induces memory loss via altered hippocampal HB-EGF and autophagy. *J Endocrinol* **244**, 53-70.
- 609  
610  
611  
612  
613 [19] Rapp SR, Espeland MA, Shumaker SA, Henderson VW, Brunner RL, Manson JE, Gass MLS, Stefanick ML, Lane DS, Hays J, Johnson KC, Coker LH, Dailey M, Bowen D, WHIMS Investigators (2003) Effect of estrogen plus progestin on global cognitive function in postmenopausal women: The Women's Health Initiative Memory Study: A randomized controlled trial. *JAMA* **289**, 2663-2672.
- 614  
615  
616  
617  
618  
619 [20] Shumaker SA, Legault C, Rapp SR, Thal L, Wallace RB, Ockene JK, Hendrix SL, Jones BN 3rd, Assaf AR, Jackson RD, Kotchen JM, Wassertheil-Smoller S, Wactawski-Wende J; WHIMS Investigators (2003) Estrogen plus progestin and the incidence of dementia and mild cognitive impairment in postmenopausal women: The Women's Health Initiative Memory Study: A randomized controlled trial. *JAMA* **289**, 2651-2662.
- 620  
621  
622  
623  
624  
625  
626  
627  
628 [21] Henderson VW (2014) Alzheimer's disease: Review of hormone therapy trials and implications for treatment and prevention after menopause. *J Steroid Biochem Mol Biol* **142**, 99-106.
- 629  
630  
631  
632 [22] Alvarez-de-la-Rosa M, Silva I, Nilsen J, Pérez MM, García-Segura LM, Avila J, Naftolin F (2010) Estradiol prevents neural tau hyperphosphorylation characteristic of Alzheimer's disease. *Ann N Y Acad Sci* **1052**, 210-224.
- 633  
634  
635  
636 [23] Zhang Z, Simpkins JW (2010) Okadaic acid induces tau phosphorylation in SH-SY5Y cells in an estrogen-preventable manner. *Brain Res* **1345**, 176-181.
- 637  
638
- [29] Rao YS, Mott NN, Wang Y, Chung SH (2007) MicroRNAs in the aging female brain: A potential mechanism for age-specific estrogen effects. *Neurobiol Aging* **28**, 2795-2806.
- [30] Rao YS, Shults CL, Pinceti E, Pak J, Chung SH (2008) Ovarian hormone deprivation alters the expression of estradiol on microRNA expression in the mouse hypothalamus. *Oncotarget* **6**, 3696-3706.
- [31] Perez SE, Chen EY, Mufson EJ (2005) Estrogen receptor alpha and beta immunoreactivity in the postnatal rat brain. *Brain Res Dev* **23**, 139-148.
- [32] Long J, He P, Shen Y, Li R (2012) Mitochondrial dysfunction in the female brain: Deficiency of estrogen receptor alpha. *Neurobiol Aging* **30**, 545-558.
- [33] Ishunina TA, Swaab DF (2012) Decreased expression of estrogen receptor- $\alpha$  mRNA in the aging female brain. *Neurobiol Aging* **33**, 286-296.
- [34] Hu XY, Qin S, Lu YP, Ravid R, Zhang Z (2012) Decreased estrogen receptor- $\alpha$  expression in hippocampal neurons in relation to hyperphosphorylation of tau in Alzheimer patients. *Acta Neuropathol* **123**, 101-110.
- [35] Savaskan E, Olivieri G, Meier F, Rapp SR, Mufson EJ (2001) Hippocampal estrogen receptor- $\alpha$  immunoreactivity is increased in Alzheimer's disease. *Neurobiol Aging* **22**, 113-119.
- [36] Fitzpatrick JL, Mize CB, Wade CE, Mufson EJ, RA, Dorsa DM (2002) Estrogen-mimetic treatment against beta-amyloid toxicity requires estrogen receptor alpha or beta and a specific signaling pathway. *J Neurochem* **82**, 674-682.
- [37] Mathew LK, Skuli N, Mucaj V, Lee J, Imtiyaz HZ, Zhang Z, Davuluri R, Zhang Z, Lal P, Lathia JD, Rich JN, Keith B, Mufson EJ (2014) miR-218 opposes a critical signaling pathway in mesenchymal glioblastoma. *Proc Natl Acad Sci U S A* **111**, 291-296.