

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

## Demoralization and associated features in an Italian sample of end-of-life cancer patients

**This is a pre print version of the following article:**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1777400> since 2021-03-04T11:26:25Z

*Published version:*

DOI:10.1200/JCO.2019.37.15\_suppl.e23004

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

# Meeting Abstracts

[Home](#)[Search Abstracts](#)[Browse Abstracts](#)[Meeting Home](#)

## Demoralization and associated features in an Italian sample of end-of-life cancer patients.

Sub-category:  
End-of-Life Care

Category:  
Symptoms and Survivorship

Meeting:  
2019 ASCO Annual Meeting

Abstract No:  
e23004

Citation:  
J Clin Oncol 37, 2019 (suppl; abstr e23004)

Author(s): Mario Airoldi, Andrea Bovero, Rossana Botto, Beatrice Adriano, Marta Opezzo, Valentina Tesio, Riccardo Torta; 2nd Medical Oncology Division, A. O. Città della Salute e della Scienza di Torino, Torino, Italy; Psycho-oncology Unit, Department of Neuroscience, University of Turin, Turin, Italy; Department of Neuroscience, University of Turin, Torino, Italy; Department of Psychology, University of Turin, Turin, Italy

### Abstract Disclosures

#### Abstract:

**Background:** Demoralization is an expression of cancer-related existential distress that emerges from feelings of loss and changes linked to a life-threatening clinical condition. It is characterized by subjective incompetence, sense of failure, hopelessness, loss of purpose/meaning and low self-esteem. The study aimed to explore prevalence, latent dimensions and associated features of demoralization in a sample of end-of-life cancer patients. **Methods:** Participants were recruited for a cross-sectional assessment. For each patient, demoralization, anxiety, depression, physical symptoms, pain, spiritual well-being, and dignity were evaluated through validated rating scales and personal and medical data were gathered by a clinician. Final sample consisted of 235 end-of-life cancer patients with a Karnofsky Performance Status (KPS) lower than 50 and a life expectancy of few weeks. **Results:** High levels of demoralization occurred in 22.6% (n = 53) of patients. Sixty-four participants (27.2%) had low demoralization and 50.2% (n = 118) had moderate demoralization. Emotional Distress and Inability to Cope, Loss of Purpose and Meaning, Worthlessness, Sense of Failure and Dysphoria were the five latent dimensions of demoralization emerged from the factor analysis. Demoralization was significantly associated with depression, anxiety, dignity, spiritual well-being and the physical symptoms except for nausea and breathing problems. **Conclusions:** Demoralization levels seem to be higher in this distinctive population than in advanced cancer patients. This could be due to the fact that end-of-life cancer patients are in a severe clinical condition and nearing death. The emerged demoralization dimensions could be five forms of expression of the existential distress typical of this illness phase. The considerable number of patients suffering from demoralization strengthen the need for psychological interventions in order to reduce the existential distress at the end of life, focusing on finding meaning and detecting spiritual concerns.

#### ► Other Abstracts in this Sub-Category:

1. Randomized trial of a hospice video decision aid for patients with advanced cancer and their caregivers.

Meeting: 2019 ASCO Annual Meeting Abstract No: 11513 First Author: Areej El-Jawahri  
Category: Symptoms and Survivorship - End-of-Life Care

2. Early mortality after resection of locally advanced rectal cancer in elderly United States patients.

Meeting: 2019 ASCO Annual Meeting Abstract No: 11529 First Author: Helmhne M. Sineshaw  
Category: Symptoms and Survivorship - End-of-Life Care

3. Is there a shift in use of subacute rehabilitation (SAR) instead of hospice referral since immunotherapy became available?

Meeting: 2019 ASCO Annual Meeting Abstract No: 11530 First Author: Jonathan Yeh  
Category: Symptoms and Survivorship - End-of-Life Care

More...