

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

## Geomatics and Epidemiology: Associating Oxidative Stress and Greenness in Urban Areas

**This is a pre print version of the following article:**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1781336> since 2021-03-22T12:16:24Z

*Published version:*

DOI:10.1016/j.envres.2021.110999

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

# Environmental Research

## Geomatics and Epidemiology: Associating Oxidative Stress and Greenness in Urban Areas

--Manuscript Draft--

<b>Manuscript Number:</b>	
<b>Article Type:</b>	Research paper
<b>Section/Category:</b>	Environmental Health & Risk Assessment
<b>Keywords:</b>	Urban Vegetation; Remote Sensing; Public Health; Isoprostane
<b>Corresponding Author:</b>	Samuele De Petris University of Torino ITALY
<b>First Author:</b>	Samuele De Petris
<b>Order of Authors:</b>	Samuele De Petris Giulia Squillacioti, Dr. Roberto Bono, Prof. Enrico Borgogno-Mondino, Prof.
<b>Abstract:</b>	<p>Green spaces may benefit human health mainly by mitigating noise and air pollution, promoting physical or social activities and improving mental health. Based on the influence that green space exposure seems to exert on Public Health and using a multidisciplinary approach, we investigated, the association between oxidative stress (OS) and green exposure in children. Overall, 207 subjects (10-13 yrs.) living in Torino (NW- Italy) were involved in this study. Each participant provided a urinary sample, used to quantify a reliable OS biomarker (15-F2t-IsoP), and their residence addresses, used for geocoding. Green exposure was characterised by calculating i) the Soil Adjusted Vegetation Index (SAVI) within fixed buffers around each participant's home, using remotely-sensed data; ii) Tree Map accounting for evergreen/broadleaf species; iii) The percentage of green cover (PGC). Significant negative correlation (Pearson's <math>r = -0.758</math>, <math>p &lt; 0.001</math>) between PGC and 15-F2t-IsoP was found. Greater SAVI was associated with lower OS (Pearson's <math>r = -0.717</math>, <math>p &lt; 0.001</math>). Noticeably, evergreens seemed to determine a significant OS reduction compared to broadleaves (slope = -0.12 and -0.02, respectively; Warton-test <math>F = 12.48</math>, <math>p = 0.0011</math>). Finally, a spatial distribution of 15-F2t-IsoP estimates map, overlying with 2011 Census Data on same-aged dwellers of Torino, was generated. Predictive models accounting for green spaces influence on OS can be useful tool derived from geomatic employ in Public Health field. Future developments of such a multidisciplinary approach should be considered in urban planning and policy-makers decisions to better define priority zones to requalify in urban settings.</p>
<b>Suggested Reviewers:</b>	Andrea Lessio andrea.lessio@ithaca.polito.it Alessio Calantropio alessio.calantropio@polito.it Piero Boccoardo piero.boccoardo@polito.it Eija Parmes eija.parmes@vtt.fi Carmen de Keijzer carmen.dekeijzer@isglobal.org

**UNIVERSITY OF TURIN, ITALY**

Department of Agricultural, Forest and Food Sciences  
Largo P. Braccini 2, 10095, Grugliasco (TO), Italy

October 14<sup>th</sup>, 2020Dear Editor of *Environmental Research*,

Please find attached the manuscript “*Geomatics and Epidemiology: Associating Oxidative Stress and Greenness in Urban Areas*”. In my view, the relevance of the results can be summarised as follows. Firstly, the evidence that urban green spaces are able to influence oxidative stress in youths living in urban areas. Secondly, evergreens show a greater impact on this association compared to broadleaf species. Thirdly, the analysis on spatial distribution of oxidative stress can be a useful tool derived from GIS and remote sensing employ in Public Health, serving urban planning and policy-makers decisions to define potential priority zones to requalify in urban settings. Further key information about the presented manuscript:

- the manuscript is an original work, has not been previously published, and is not under consideration for publication elsewhere;
- the participation of all human subjects did not occur prior their informed consent was obtained;
- all authors have disclosed any potential competing interest regarding the submitted article;
- all authors have read the manuscript, agreeing on the submission to *Environmental Research*, and accepting the responsibilities for the manuscript’s contents;
- all authors have read and approved the paper and it has not been published previously nor is it being considered by any other peer-reviewed journal;

The multidisciplinary approach of this manuscript involves: biosphere, anthroposphere, atmosphere. In my point of view, this manuscript may deserve Your considerations because it fits some of subject areas included in the journal:

- Environmental risks assessment and management
- Air pollution quality and human health
- Risks and public health
- Environmental management and policy
- Environmental risks assessment and management

Please find below some suggestions for potential reviewers, as requested.

1. Andrea Lessio - andrea.lessio@ithaca.polito.it
2. Alessio Calantropio - alessio.calantropio@polito.it
3. Piero Boccardo - piero.boccardo@polito.it
4. Eija Parmes - eija.parmes@vtt.fi.
5. Carmen de Keijzer - carmen.dekeijzer@isglobal.org

Hoping that the manuscript may fulfil the scientific standards of *Environmental Research*.

Best Regards,  
Samuele De Petris

Contacts:

e-mail: samuele.depetris@unito.it

1. Oxidative stress and green exposure association was investigated in children
2. Green exposure was analyzed by geomatic techniques
3. Spectral vegetation index and tree census data were used to map oxidative stress
4. Evergreens trees determine a significant oxidative stress reduction
5. Geomatics can support urban planning to improve public health

## Geomatics and Epidemiology: Associating Oxidative Stress and Greenness in Urban Areas

DE PETRIS SAMUELE<sup>A\*</sup>, SQUILLACIOTI GIULIA<sup>B\*</sup>, BONO ROBERTO<sup>B</sup>, BORGOGNO-MONDINO ENRICO<sup>A</sup>

<sup>A</sup>Department of Agriculture, Forest and Food sciences, University of Torino, 10095 Grugliasco (TO), Italy; [samuele.depétris@unito.it](mailto:samuele.depétris@unito.it) (S.D.P.); [enrico.borgogno@unito.it](mailto:enrico.borgogno@unito.it) (E.B.M.)

<sup>B</sup>Department of Public Health and Pediatrics, University of Torino, 10121 Torino, Italy; [giulia.squillacioti@unito.it](mailto:giulia.squillacioti@unito.it) (G.S.); [roberto.bono@unito.it](mailto:roberto.bono@unito.it) (R.B.)

\*these two authors equally contributed to the conceptualization and execution of the manuscript

**Corresponding author:** Samuele De Petris, [samuele.depétris@unito.it](mailto:samuele.depétris@unito.it)

**Declarations of interest:** none.

## Geomatics and Epidemiology: Associating Oxidative Stress and Greenness in Urban

### Areas

**Abstract:** Green spaces may benefit human health mainly by mitigating noise and air pollution, promoting physical or social activities and improving mental health. Based on the influence that green space exposure seems to exert on Public Health and using a multidisciplinary approach, we investigated, the association between oxidative stress (OS) and green exposure in children. Overall, 207 subjects (10-13 yrs.) living in Torino (NW- Italy) were involved in this study. Each participant provided a urinary sample, used to quantify a reliable OS biomarker (15-F2t-IsoP), and their residence addresses, used for geocoding. Green exposure was characterised by calculating i) the Soil Adjusted Vegetation Index (SAVI) within fixed buffers around each participant's home, using remotely-sensed data; ii) Tree Map accounting for evergreen/broadleaf species; iii) The percentage of green cover (PGC). Significant negative correlation (*Pearson's r* = -0.758, *p* < 0.001) between PGC and 15-F2t-IsoP was found. Greater SAVI was associated with lower OS (*Pearson's r* = -0.717, *p* < 0.001). Noticeably, evergreens seemed to determine a significant OS reduction compared to broadleaves (slope = -0.12 and -0.02, respectively; *Warton-test F* = 12.48, *p* = 0.0011). Finally, a spatial distribution of 15-F2t-IsoP estimates map, overlying with 2011 Census Data on same-aged dwellers of Torino, was generated. Predictive models accounting for green spaces influence on OS can be useful tool derived from geomatic employ in Public Health field. Future developments of such a multidisciplinary approach should be considered in urban planning and policy-makers decisions to better define priority zones to requalify in urban settings.

**Keywords:** Urban Vegetation, Remote Sensing, Public Health, Isoprostane

## 1. Introduction

Green spaces are thought to benefit human health mainly through i) mitigation of noise and air pollution exposure ii) promotion of outdoor exercise and social activities iii) improvement of mental health. Participation in physical activity is significantly higher for people living close to parks, or green areas, determining a reduction of overweight, obesity and, in general, a lower body mass index (Bell et al., 2008). Physical activity is also showing a positive impact on health by reducing risk of diabetes (Dalton et al., 2016) and cardiovascular diseases (Fong et al., 2018; Pereira et al., 2012). Green space exposure proves to positively affect mental health and social engagement by reducing stress (Gong et al., 2016; Markevych et al., 2017), depressive symptoms (Garipey et al., 2015), aggressive behaviour (Younan et al. 2016), and contributing in decelerating the cognitive decline (de Keijzer et al., 2018). Moreover, maternal exposure to green space is positively associated with greater new-born's birthweight; this is mainly due to environmental pollution mitigation, opportunity of socialisation, and to exercise in contact with nature.

Studies addressing greenness and respiratory health possible association are still inconsistent. Some authors referred about a positive effect of higher exposures to greenness finding risk reduction of asthma and respiratory symptoms (Lovasi et al., 2008; Sbihi et al., 2015; Squillacioti et al., 2019b). On the other hand, some authors found adverse effect of green space exposure on asthma, rhinitis and respiratory health in general (Fuertes et al., 2014; Parmes et al., 2020).

Based on the influence that green space exposure seems to exert on Public Health, we speculated that greenness might also be involved in oxidative stress (OS) induction.

OS is a pre-pathological condition characterised by an imbalance between pro-oxidant and anti-oxidant species, in favour of pro-oxidants. OS does not represent a pathology

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

itself (Sies, 2015), but a risky condition related to the pathophysiological mechanisms behind several health impairments, such as Cardio Vascular Diseases (CVDs), diabetes and respiratory diseases. Furthermore, it is susceptible to those environmental risk factors able to act as pro-oxidants (e.g. air pollutants and tobacco smoke)(Milne et al., 2015).

At the meantime, OS may be influenced by individual characteristics of subjects like obesity, health *status*, tobacco smoking exposure and exercise training (Nikolaidis et al., 2011; van 't Erve et al., 2017). Within this context, we focused on the potential association between OS and urban green spaces, being this type of knowledge still lacking (Woo et al., 2009; Yeager et al., 2018). We operated by a multidisciplinary approach aiming at investigating strength of this potential association, specifically evaluating usefulness of geomatic tools in epidemiological studies. Many epidemiological studies investigating the relationship between green space and health benefits are already based on remotely-sensed data. Specifically, the Normalised Difference Vegetation Index (NDVI) is widely used in epidemiological research as spectral metric (Fong et al., 2018). NDVI does not provide any information about species characterisation (Fong et al., 2018). Few works, alternatively, have tried to characterise exposure with reference to land cover maps (Egorov et al., 2017; Parmes et al., 2020; Tsai et al., 2019) or tree *census* data, with the aim of taking into account type and shape of green spaces, or vegetation fraction cover (Browning and Rigolon, 2018; Lovasi et al., 2013; Pilat et al., 2012).

## 2. Materials

### 2.1. Study Area



1 The study area corresponds to the municipality of Torino (NW Italy, E: 396027 N:  
2 4991913 reference frame WGS84 UTM32N), sizing about 130 Km<sup>2</sup> (Fig.1). In 2019  
3 green areas covered 43% of the whole city, addressing Torino as one of the most  
4 relevant greenest cities (Baycan-Levent and Nijkamp, 2009; Li et al., 2015). With  
5 reference to green patches, the following metrics were computed: minimum and  
6 maximum patch size are 0.015 and 108 ha, respectively; patch size mode is 0.63 ha; 25<sup>th</sup>  
7 percentile is 0.38 ha; 75<sup>th</sup> percentile is 3.76 ha.




Figure 1 – Torino area localization (Reference frame: WGS84 UTM32N).

## 2.2. Multispectral Data

Multispectral imagery are widely used to detect and characterise vegetation in urban contexts, permitting to locally map vegetated areas (Mudele and Gamba, 2019; Rosina and Kopecká, 2016). We assumed that the maximum of vegetation vigour, in the area, occurs in the summer period (June-July, Zhou et al. 2016; De Petris S. et al. 2019;


1 Borgogno-Mondino, Sarvia, and Gomarasca 2019). Consequently, a Sentinel-2 Level  
2 2A image (S2), acquired by the Copernicus Sentinel-2 MSI (Multi Spectral Instrument)  
3 sensor on 14<sup>th</sup> June 2016, was obtained from the Theia CNES geoportal  
4 (<https://www.Theia-land.fr/en/product/sentinel-2-surface-reflectance/>, Baetens,  
5 Desjardins, and Hagolle 2019; Revel et al. 2019)). In summer time, vegetation-related  
6 biomass expresses its maximum in Torino. This assumption also relies on the observed  
7 inter-annual phenological behaviour of vegetation in urban context. Since  
8 anthropogenic green areas show no expansion dynamics, if compared with natural ones,  
9 and vegetation density does not change significantly, due to the human-management  
10 (i.e. pruning, mowing), biomass shows inter-annual flat behaviour (Li et al., 2017). S2  
11 Level 2A data are supplied already calibrated in at-the-ground-reflectance. Nominal  
12 radiometric accuracy is about 0.01 reflectance units (European Space Agency/Centre  
13 National d’Eudes Spatiales (CNES), 2019). Technical features S2 image are reported in  
14 Table 1. It is worth to remind that the minimum mapping unit from S2 imagery is 100  
15 m<sup>2</sup>; this size was retained appropriate if compared with the average dimension of green  
16 areas in Torino.

17 [ Here Table 1 ]

### 18 2.3.Epidemiological Data

19 Available epidemiologic sample refers to an on-going cohort whose subjects were  
20 recruited in 2010 as part of a research project funded by the Piedmont Regional Council  
21 focusing on the effects of environmental factors. This cross-sectional study involved  
22 207 healthy children (10–13 years old) from secondary schools located in Torino. All  
23 subjects gave the assent to participate along with their parents, who signed an informed  
24 consent. Since the study involved human subjects, the study protocol required the Ethics

1 Committee approval (protocol number 826/13/08). Parents filled out a questionnaire to  
2 provide general information and their home address; participants provided a sample of  
3 urine that was used for quantification of a reliable biomarker of OS, namely 15-F2t-  
4 urine that was used for quantification of a reliable biomarker of OS, namely 15-F2t-  
5 isoprostane (15-F2t-IsoP). At this concern, 15-F2t-IsoP was quantified to measure OS  
6 by a specific enzyme-linked immunosorbent assay (ELISA) kit (Oxford, MI, USA),  
7 according to the manufacturer's instructions. To achieve better accuracy each sample  
8 was diluted 1:4, as reported in previous works (Romanazzi et al., 2013; Squillacioti et  
9 al., 2019a). Urinary creatinine (crea) was quantified in each urinary sample in order to  
10 normalise urinary excretion rate and dilution. 15-F2t-IsoP was finally referred as ng/mg  
11 crea. Collected home addresses (Fig. 2) were used to geocode each subject by means of  
12 MMQGis QGIS plugin; positioning accuracy achieved by MMQGis is declared lower  
13 than 25m (Cetl et al., 2018).



54 *Figure 2 – Position (accuracy < 25 m) of sampled subjects (207) children aged between 10 and 13) within the study area.*

55  
56 *Reference frame is WGS84 UTM 32N.*

## 2.4. Auxiliary data

Auxiliary data, provided by institutional geoportals, were supplied in vector format.

A Tree Map (TM), updated at 2019 with a nominal scale 1:1000 was obtained as point layer from the Torino Geoportal. TM contains more than 160,000 individuals divided in evergreen and broadleaf species (Fig. 3). Main evergreen genera are: *Abies spp.*, *Picea spp.*, *Pinus spp.*, *Thuja spp.*, *Cedrus spp.*, *Cupressus spp.*, *Ilex spp.*, *Magnolia spp.* representing the 5% of the trees; while the broadleaf genera are: *Platanus spp.*, *Tilia spp.*, *Aesculus spp.*, *Celtis spp.*, *Acer spp.*, *Ulmus spp.*, *Carpinus spp.*, representing the 95%.

Census data (*CD*, 2011) were obtained from the ISTAT (Italian Statistics Institute) geoportal; *CD* has a nominal scale of 1:5000 and contains census sections (polygons) and the correspondent features of surveyed people, included the number of children aged between 10 and 13 years. *CD* was used to describe the spatial distribution of potential targets that was retained a driving factor to address future urban policies eventually aimed at mitigating OS effects.




Figure 3 - Tree Map (TM) providing position of trees within the study area. It was obtained from Torino Geoportal. Nominal scale is 1:1000 and reference frame WGS84 UTM 32N.

### 3. Methods and data processing

Spatial analyses were operated by *SAGA GIS* vs.7.1 and *QGIS* vs. 3.4.12; statistical analyses were performed using *Past* vs. 3.6.6 (Hammer et al., 2001).

#### 3.1. Detection and characterization of urban vegetation

In urban contexts soil background makes remotely sensed pixels not pure, i.e. mixed (Small and Lu, 2006; Song, 2005). In fact, background exerts considerable influence on the average pixel spectrum, thus influencing the accuracy of detection of green areas. To minimize such an effect while mapping vegetation within urban environments (Huete, 1988), the Soil Adjusted Vegetation Index (SAVI) was calculated by eq. 1:

$$SAVI = \frac{1.5 (\rho_{NIR} - \rho_{RED})}{\rho_{NIR} + \rho_{RED} + 0.5} \quad (1)$$

where  $\rho_{NIR}$  and  $\rho_{RED}$  are S2 band 4 (665 nm) and band 8 (840 nm), respectively. Not-vegetated areas (buildings, streets etc.) were masked out by SAVI thresholding: pixel showing SAVI values  $< 0.45$  were labelled as not-vegetated (Borgogno-Mondino et al., 2016; Burgan, 1993; Gao, 1996; Ormsby et al., 1987; Zhang et al., 2003). A map showing only vegetated areas was, therefore, generated (Fig.4), making possible to obtain preliminary information about spatial distribution of biomass in the area. Biomass is well-known to be related to spectrally derived vegetation indices (SAVI included).




Figure 4 – SAVI map obtained by S2 image. Not-vegetated pixels were masked out by SAVI thresholding (SAVI  $< 0.45$ ). Reference frame is WGS84 UTM32N.

### 3.2. Green Exposure Assessment

1  
2 Many studies attested that green spaces may positively contribute to children's health  
3  
4 (McMorris et al., 2015; Thiering et al., 2016). Greenness positive effects are assumed to  
5  
6 decay with distance and increase with exposure time. Consequently, a first reasonable  
7  
8 approach to quantify greenness exposure level is the characterisation of vegetated areas  
9  
10 falling in a buffer zone surrounding subject' houses (James et al., 2015). Considering that  
11  
12 children aged between 10 and 13, they can be supposed having an independent mobility  
13  
14 pattern that ranges between 500 and 1000 m from their houses (Fagerholm and Broberg,  
15  
16 2011; Mavoia et al., 2011; Tillberg Mattsson, 2002). Consequently, buffers having a fixed  
17  
18 radius of 500 m ( $B_{500}$ ) were mapped around geocoded addresses. Different metrics, useful  
19  
20 to measure and characterise green space falling in each buffer, were then calculated.  
21  
22

23  
24  
25 With reference to the masked SAVI map pixels were counted falling in  $B_{500}$  and the  
26  
27 related SAVI mean value (mSAVI) computed. The percentage of green cover (PGC) was  
28  
29 also calculated comparing the vegetation fraction with the whole area of the buffer.  
30  
31 mSAVI was then multiplied by PGC, assuming the latter as a sort of weight. A new  
32  
33 metric, hereinafter called *Unitary SAVI* (uSAVI) was finally obtained to somehow  
34  
35 measure biomass density. Using TM, two more variables accounting for the number of  
36  
37 evergreen and broadleaf tree falling in  $B_{500}$  were calculated, namely NET (Number of  
38  
39 Evergreen Trees) and NBT (Number of Broadleaf Trees).  
40  
41  
42  
43  
44  
45  
46  
47

### 3.3. Statistical Analysis and Data Processing

48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000

1 was tested using the Pearson's linear correlation ( $r$ ) and the correspondent linear  
2 regression was calibrated by Ordinary Least Squares (OLS). Correlation was also tested  
3  
4 by  $r$  for the following relationship: 15-F2t-IsoP vs PGC, IsoP vs NET, and 15-F2t-IsoP  
5  
6 vs NBT. Obtained values showed, for all the tested relationships, a weak correlation (see  
7  
8 Results and Discussion section). Nevertheless, an envelope function ( $Ev$ ) of these  
9  
10 distributions was considered bounding observations clouds.  $Ev$  can be interpreted as the  
11  
12 upper limit beyond which none 15-F2t-IsoP values exist referred to actual vegetation  
13  
14 parameters values. It was computed by dividing the 15-F2t-IsoP range into 20  
15  
16 equiprobable classes corresponding to a class width of about 1.4 points of 15-F2t-IsoP  
17  
18 value, starting from 3.56 ng/mg crea. With reference to the defined classes the  
19  
20 correspondent 15-F2t-IsoP and uSAVI, PGC, NET, NBT maximum values were  
21  
22 calculated and compared by scatterplots. Linear associations relating 15-F2t-IsoP- $Ev$  with  
23  
24 the other variables, were tested using Pearson's coefficient and correspondent linear  
25  
26 regression modelled. Namely, NET- $Ev$  and NBT- $Ev$  slope values were compared  
27  
28 according to the Warton method (Warton et al., 2006) in order to evaluate if evergreen  
29  
30 trees determined a significant difference in OS reduction rate (i.e. slope value of bivariate  
31  
32 model) with respect to broadleaf trees. Given these relationships, in order to spatialise the  
33  
34 information generating a map of estimate of 15-F2t-IsoP- $Ev$ , authors only focused on  
35  
36  $uSAVI-Ev$  since, among the tested metrics, from an operational point of view, it can be  
37  
38 easily and globally computed from free satellite data. Consequently, the same procedure  
39  
40 could be applied everywhere in spite of the existence of more specific databases  
41  
42 containing more accurate information about local vegetation. Nevertheless, other useful  
43  
44 information is derived from the interpretation of relationship between 15-F2t-IsoP- $Ev$  and  
45  
46 PGC- $Ev$ , NET- $Ev$  and NB- $Ev$  for investigation purposes, solely. With these premises  
47  
48 the linear regression model relating 15-F2t-IsoP- $Ev$  to uSAVI- $Ev$  was adopted to  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



1 spatialise 15-F2t-IsoP-Ev estimates in the study area. For this purpose a 500x500 m  
2 squared graticule (G) was generated assuming this size consistent with a walking  
3 mobility of 10–15 minutes (Wolch et al., 2014). For each cell of G, the uSAVI maximum  
4 value was computed from the previously generated SAVI masked map. G was then  
5 rasterised to generate uSAVI<sup>G</sup>. A raster map of estimates of 15-F2t-IsoP-Ev (hereinafter  
6 called 15-F2t-IsoP-Ev<sup>G</sup>) was finally computed by grid calculation tools, implementing  
7 the previously calibrated linear regression relating uSAVI to 15-F2t-IsoP-Ev. An  
8 accuracy assessment of 15-F2t-IsoP-Ev<sup>G</sup> was operated by a “leave-one-out” procedure  
9 (Brovelli et al., 2008) to compute the Mean Absolute Error (MAE) (Willmott and  
10 Matsuura, 2005).

#### 26 **4. Results and Discussions**

27 Concerning outliers detection in the 15-F2t-IsoP dataset 4 individuals were found having a 15-  
28 F2t-IsoP value > 40 ng/mg crea and removed. With reference to the test concerning relationship  
29 between OS and uSAVI (Fig. 5) a Pearson’s linear correlation coefficient was found equal to -  
30 0.045,  $p > 0.05$ . A linear regression was calibrated by OLS estimation finding a slope value of  
31 -2.214,  $C.Is$  95% (upper bound =9.538 and lower bound=4.68,  $p > 0.05$ ).  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65





Figure 5– Linear regression model (red line, dark-grey are 95% Cis) of 15-F2t-IsoP (dependent variable) and uSAVI (independent variable). Box-plot represent respectively: minimum, Q1, median, Q3, maximum of both uSAVI (up box-plot) and 15-F2t-IsoP (right) value distributions without outliers.

It is worth to stress that some of the OLS regression assumptions were violated: residuals were not normally distributed (Shapiro-Wilk test) showing autocorrelation (Durbin-Watson test) and, more in general, both  $r$  and the OLS regression model showed that OS and uSAVI were poorly correlated. Such poorly correlation values were possibly due to the fact that OS is also related with other factors like air quality, personal behaviour (i.e. diet, smoking), individual pathology. Therefore, all these relationships influence the covariance and, as well, correlation coefficient. Concerning the other tested relationships (15-F2t-IsoP vs PGC, IsoP vs NET, and 15-F2t-IsoP vs NBT) the following values were found (Tab.2).

[ Here Table 2]

It is worth to remind that OS level differences are due to several risk factors, both individual and environmental (Fig. 6). Therefore, a modelling of 15-F2t-IsoP based on uSAVI does not have the aim of producing reliable absolute 15-F2t-IsoP estimates; conversely, it is expected to produce a general overview, enhancing relative differences between different areas.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Figure 6 – OS is influenced by many individual (left ellipse) and environmental factors (right ellipse). This generates high variability in OS values if ordinary regression-based approaches are used. (Green ellipse)

Interaction between OS and greenness. OS level can be influenced by green areas increasing physical activity or mitigating air pollution (Nowak, Crane, and Stevens 2006; McMorris et al. 2015)

Figure 7 graphically shows scatterplots relating 15-F2t-IsoP to uSAVI, PGC, NET and NB. Red lines define the upper bound (*envelope*) of 15-F2t-IsoP estimates, which class *maxima*, used to generate the new metrics 15-F2t-IsoP-Ev, uSAVI-Ev, PGC-Ev, NET\_Ev and NB-E, have to somehow represent. Classes used for maximum value computation are reported in Tab. 3.




Figure 7 – Scatterplots relating 15-F2t-IsoP and vegetation parameters. A linear envelope function (Red line) can be defined for each cloud defining the upper boundary of the estimates.

[Here Table 3]

Figures 8 (a-b) and 9, and table 4 report results of the bivariate models that relate 15-F2t-IsoP-Ev to PGC-Ev, 15-F2t-IsoP-Ev and uSAVI-Ev, respectively.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65




Figure 8 – (A) bivariate model between 15-F2t-IsoP-Ev and uSAVI-Ev; (B) Bivariate model between 15-F2t-IsoP-Ev and PGC-Ev. Pearson correlation coefficient ( $r$ ) was reported with related significance level. Dark-grey limits are 95% CIs.

A significant negative correlation was found between 15-F2t-IsoP-Ev and PGC-Ev ( $r = -0.758$ ,  $p < 0.001$ ) and between 15-F2t-IsoP-Ev and uSAVI-Ev ( $r = -0.717$ ,  $p < 0.001$ ). Results prove that higher vegetation cover seems to significantly reduce OS. In particular, if vegetation cover is composed by trees (i.e. high uSAVI), OS tends to decrease. With respect to evergreen and broadleaf tree *genera*, the correspondent regressions were tested separately (Figure 9).





Figure 9 – (Triangle) bivariate model between 15-F2t-IsoP-Ev and NET-Ev; (Circle) Bivariate model between 15-F2t-IsoP-Ev and NBT-Ev. Pearson correlation coefficient ( $r$ ) was reported with related significance level. Dark-grey limits are 95% CIs. In Bold are highlighted IsoP reduction rate (slope values) of NET and NBT. NET slope is steeper negative than NBT one and the two slopes are statistically different ( $p < 0.01$ ).

1 To test significance of slope of NET and NBT, the Warton test (Warton et al., 2006) was applied  
2 finding a  $F$  value = 12.48 ,  $p = 0.0011$ . It proved that NET and NBT bivariate model gain values  
3 were significantly different. In particular, NET determines a negative steeper 15-F2t-IsoP-Ev  
4 reduction rate six-time greater in respect of NBT, suggesting that evergreen trees determine a  
5 stronger positive effect on OS. Statistic parameters of tested relationships are reported in Table  
6  
7  
8  
9  
10  
11  
12 4.

13  
14  
15 [Here Table 4]  
16  
17

18 Concerning accuracy of estimates, MAE of 15-F2t-IsoP-Ev<sup>G</sup> was computed by a leave-one-out  
19 approach. It resulted equal to 5.26 15-F2t-IsoP ng/mg crea. Results proved that green areas  
20 where evergreen trees dominate OS is lower. Consequently, urban planners could exploit this  
21 information while designing new restoration/requalification actions, possibly promoting these  
22 interventions especially in those parts of the city having a higher density of potential targets  
23 (i.e. children). In order to exemplify spatial implications of the above-mentioned relationships  
24 a 15-F2t-IsoP-Ev<sup>G</sup> map was generated with a grid size of 10 m and coupled, by overlaying, with  
25 CD (Fig. 10). This map is intended to represent an operational tool to describe OS risk of 10-  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2 areas could be added, or old ones requalified, providing estimates of associated 15-F2t-IsoP-  
3 Ev<sup>G</sup> resulting from the adopted choices.  
4  
5



37  
38 *Figure 10 –Map shows spatial distribution of 15-F2t-IsoP-Ev<sup>G</sup> estimates coupled by overlaying with 2011*

39 *Census Data (children). It could be useful to urban planners to define priority zones where creating/*  
40 *requalifying green areas. Dotted rectangles are critical zones where high children density and high OS values*  
41 *are located. Reference frame is WGS84 UTM32N.*  
42  
43  
44  
45

## 46 **5. Conclusions**

47  
48 This study aimed at investigating and formalising the relationship between OS and  
49 greenness with reference to children living in Torino (NW Italy). Significant negative  
50 correlations (Pearson's  $r = -0.758$ ,  $p < 0.001$ ) between PGC-Ev and 15-F2t-IsoP-Ev were  
51 found indicating that a higher degree of green areas around children's houses determines a  
52 lower OS levels in children. One of the potential mechanisms underlying this association  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 could be related to the presence of accessible green areas potentially used to perform  
2 physical activity.  
3

4 Moreover, our results emphasised that local biomass plays a key role in OS reduction. In  
5 fact, higher values of uSAVI correspond to lower levels of OS (Pearson's  $r = -0.717$ ,  $p <$   
6  $0.001$ ). High biomass can be generally related to a strong presence of trees: green areas  
7 where trees density was higher proved to be significantly related to lower OS values. Trees  
8 are thought to provide positive effects to health by reducing exposure to particular matter  
9 ( $PM_{10}$  or  $PM_{2.5}$ ) or ozone (Fausto Manes et al., 2012; F. Manes et al., 2012; Nowak et al.,  
10 2006). In particular, evergreen trees seem to determine a significant 15-F2t-IsoP-Ev  
11 reduction rate in respect to broadleaves (NET slope =  $-0.12$ , NBT slope =  $-0.02$ , Warton-  
12 test  $F = 12.48$ ,  $p = 0.0011$ ). With respect to evergreen trees, this can be related to: (a) a  
13 higher content of cuticular resins, which have an important role in pollution absorption and  
14 mitigation (Marando et al., 2016; Sawidis et al., 2012); (b) a total leaf area greater than  
15 broadleaf that, considering the annual phenology of urban trees, provides larger absorption  
16 surfaces for pollutants. Furthermore, in northern Italy, especially in Po river watershed, air  
17 quality dramatically decreases in winter (Bigi and Ghermandi, 2014; Finardi and Pellegrini,  
18 2004) when only the evergreen species can provide the mitigation effect on air pollution.  
19 uSAVI and OS bivariate model (15-F2t-IsoP-Ev<sup>G</sup> map) can be used for future developments  
20 related to urban planning strategies and policy makers decisions, making possible to define  
21 priority zones where requalification or new greenspaces should be considered. These plans  
22 might take into account the here presented results according to three main strategies: (a)  
23 green cover improvement; (b) trees or shrubs (i.e. high biomass density cover) planting in  
24 place of grass; (c) evergreen trees selection rather than broadleaves.  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54

55 It is worth to remind that calibrated uSAVI-OS model describes the Ev function containing  
56 OS estimates, meaning that one can infer only about the upper limits of expected 15-F2t-  
57  
58  
59  
60  
61  
62  
63  
64  
65



1 IsoP, with no possibility of knowing its exact local value. This is mainly due to the high  
2 intrinsic variability in OS-induced factors (individual and environmental). Since this study  
3  
4 mainly refers to children, further works are expected to test the association between  
5  
6 greenness and OS in differently aged subjects and different geographic urban areas where  
7  
8 the sole intensity of urbanization can produce different level of OS (Squillaciotti et al.,  
9  
10 2020). Expectation is that this approach could promote the highest public health standards  
11  
12  
13 even with the best urban green design *criteria*.  
14  
15  
16  
17  
18  
19  
20

## 21 **References**

- 22  
23  
24 Baetens, L., Desjardins, C., Hagolle, O., 2019. Validation of Copernicus Sentinel-2 Cloud Masks  
25 Obtained from MAJA, Sen2Cor, and FMask Processors Using Reference Cloud Masks  
26 Generated with a Supervised Active Learning Procedure. *Remote Sensing* 11, 433.
- 27 Baycan-Levent, T., Nijkamp, P., 2009. Planning and management of urban green spaces in Europe:  
28 Comparative analysis. *Journal of Urban Planning and Development* 135, 1–12.
- 29 Bell, J.F., Wilson, J.S., Liu, G.C., 2008. Neighborhood greenness and 2-year changes in body mass  
30 index of children and youth. *American journal of preventive medicine* 35, 547–553.
- 31 Bigi, A., Ghermandi, G., 2014. Long-term trend and variability of atmospheric PM10 concentration in  
32 the Po Valley.
- 33  
34 Borgogno-Mondino, E., Lessio, A., Gomasasca, M.A., 2016. A fast operative method for NDVI  
35 uncertainty estimation and its role in vegetation analysis. *European Journal of Remote Sensing*  
36 49, 137–156.
- 37 Borgogno-Mondino, E., Sarvia, F., Gomasasca, M.A., 2019. Supporting Insurance Strategies in  
38 Agriculture by Remote Sensing: A Possible Approach at Regional Level, in: *International*  
39 *Conference on Computational Science and Its Applications*. Springer, pp. 186–199.
- 40 Brovelli, M.A., Crespi, M., Fratarcangeli, F., Giannone, F., Realini, E., 2008. Accuracy assessment of  
41 high resolution satellite imagery orientation by leave-one-out method. *ISPRS Journal of*  
42 *Photogrammetry and Remote Sensing* 63, 427–440.
- 43  
44 Browning, M., Rigolon, A., 2018. Do Income, Race and Ethnicity, and Sprawl Influence the  
45 Greenspace-Human Health Link in City-Level Analyses? Findings from 496 Cities in the  
46 United States. *International Journal of Environmental Research and Public Health* 15, 1541.  
47 <https://doi.org/10.3390/ijerph15071541>
- 48 Burgan, R.E., 1993. Monitoring vegetation greenness with satellite data. US Department of Agriculture,  
49 Forest Service, Intermountain Research Station.
- 50 Cetl, V., Kliment, T., Jogun, T., 2018. A comparison of address geocoding techniques—case study of the  
51 city of Zagreb, Croatia. *Survey Review* 50, 97–106.
- 52 Dalton, A.M., Jones, A.P., Sharp, S.J., Cooper, A.J.M., Griffin, S., Wareham, N.J., 2016. Residential  
53 neighbourhood greenspace is associated with reduced risk of incident diabetes in older people:  
54 a prospective cohort study. *BMC Public Health* 16, 1171. <https://doi.org/10.1186/s12889-016-3833-z>
- 55  
56  
57 de Keijzer, C., Tonne, C., Basagaña, X., Valentín, A., Singh-Manoux, A., Alonso, J., Antó, J.M.,  
58 Nieuwenhuijsen, M.J., Sunyer, J., Dadvand, P., 2018. Residential surrounding greenness and  
59  
60  
61  
62  
63  
64  
65

- cognitive decline: a 10-year follow-up of the Whitehall II cohort. *Environmental health perspectives* 126, 077003.
- 1  
2 De Petris S., Berretti R., Sarvia F., Borgogno-Mondino E., 2019. Precision arboriculture: a new  
3 approach to tree risk management based on geomatics tools, in: *SPIE Remote Sensing*, 2019.  
4 Presented at the Remote Sensing for Agriculture, Ecosystems, and Hydrology XXI, SPIE.  
5 <https://doi.org/10.1117/12.2532778>
- 6  
7 Egorov, A.I., Griffin, S.M., Converse, R.R., Styles, J.N., Sams, E.A., Wilson, A., Jackson, L.E., Wade,  
8 T.J., 2017. Vegetated land cover near residence is associated with reduced allostatic load and  
9 improved biomarkers of neuroendocrine, metabolic and immune functions. *Environmental*  
10 *Research* 158, 508–521. <https://doi.org/10.1016/j.envres.2017.07.009>
- 11 European Space Agency/Centre National d’Eudes Spatiales (CNES), 2019. 3rd Sentinel-2 Validation  
12 Team Meeting - 12-14 March 2019 - Toulouse, France . Toulouse, France, p. 30.
- 13 Fagerholm, N.C., Broberg, A., 2011. Mapping and characterising children’s daily mobility in urban  
14 residential areas in Turku, Finland. *Fennia-International Journal of Geography* 189, 31–46.
- 15 Finardi, S., Pellegrini, U., 2004. 6.13 SYSTEMATIC ANALYSIS OF METEOROLOGICAL  
16 CONDITIONS CAUSING SEVERE URBAN AIR POLLUTION EPISODES IN THE  
17 CENTRAL PO VALLEY.
- 18 Fong, K.C., Hart, J.E., James, P., 2018. A Review of Epidemiologic Studies on Greenness and Health:  
19 Updated Literature Through 2017, Current environmental health reports. NLM (Medline).  
20 <https://doi.org/10.1007/s40572-018-0179-y>
- 21  
22 Fuertes, E., Markevych, I., Berg, V., 2014. Greenness and allergies: evidence of differential associations  
23 in two areas in Germany. *J Epidemiol Community Health*. <https://doi.org/10.1136/jech>
- 24 Gao, B.-C., 1996. NDWI—A normalized difference water index for remote sensing of vegetation liquid  
25 water from space. *Remote sensing of environment* 58, 257–266.
- 26 Garipey, G., Kaufman, J.S., Blair, A., Kestens, Y., Schmitz, N., 2015. Place and health in diabetes: the  
27 neighbourhood environment and risk of depression in adults with Type 2 diabetes. *Diabetic*  
28 *Medicine* 32, 944–950. <https://doi.org/10.1111/dme.12650>
- 29  
30 Gong, Y., Palmer, S., Gallacher, J., Marsden, T., Fone, D., 2016. A systematic review of the relationship  
31 between objective measurements of the urban environment and psychological distress.  
32 *Environment International* 96, 48–57. <https://doi.org/10.1016/j.envint.2016.08.019>
- 33 Hammer, Ø., Harper, D.A., Ryan, P.D., 2001. PAST: paleontological statistics software package for  
34 education and data analysis. *Palaeontologia electronica* 4, 9.
- 35 Huete, A., 1988. Huete, AR A soil-adjusted vegetation index (SAVI). *Remote Sensing of Environment*.  
36 *Remote sensing of environment* 25, 295–309.
- 37 James, P., Banay, R.F., Hart, J.E., Laden, F., 2015. A review of the health benefits of greenness. *Current*  
38 *epidemiology reports* 2, 131–142.
- 39  
40 Li, F., Song, G., Liujun, Z., Yanan, Z., Di, L., 2017. Urban vegetation phenology analysis using high  
41 spatio-temporal NDVI time series. *Urban Forestry & Urban Greening* 25, 43–57.
- 42 Li, X., Zhang, C., Li, W., Ricard, R., Meng, Q., Zhang, W., 2015. Assessing street-level urban greenery  
43 using Google Street View and a modified green view index. *Urban Forestry & Urban Greening*  
44 14, 675–685.
- 45 Lovasi, G.S., O’Neil-Dunne, J.P.M., Lu, J.W.T., Sheehan, D., Perzanowski, M.S., MacFaden, S.W.,  
46 King, K.L., Matte, T., Miller, R.L., Hoepner, L.A., Perera, F.P., Rundle, A., 2013. Urban Tree  
47 Canopy and Asthma, Wheeze, Rhinitis, and Allergic Sensitization to Tree Pollen in a New York  
48 City Birth Cohort. *Environmental Health Perspectives* 121, 494–500.  
49 <https://doi.org/10.1289/ehp.1205513>
- 50  
51 Lovasi, G.S., Quinn, J.W., Neckerman, K.M., Perzanowski, M.S., Rundle, A., 2008. Children living in  
52 areas with more street trees have lower prevalence of asthma. *Journal of Epidemiology &*  
53 *Community Health* 62, 647–649. <https://doi.org/10.1136/jech.2007.071894>
- 54 Manes, F., Blasi, C., Salvatori, E., Capotorti, G., Galante, G., Feoli, E., Incerti, G., 2012. Natural  
55 vegetation and ecosystem services related to air quality improvement: tropospheric ozone  
56 removal by evergreen and deciduous forests in Latium (Italy). *Annali di Botanica* 2, 79–86.
- 57 Manes, Fausto, Incerti, G., Salvatori, E., Vitale, M., Ricotta, C., Costanza, R., 2012. Urban ecosystem  
58 services: tree diversity and stability of tropospheric ozone removal. *Ecological Applications* 22,  
59 349–360.  
60  
61  
62  
63  
64  
65

- 1 Marando, F., Salvatori, E., Fusaro, L., Manes, F., 2016. Removal of PM10 by forests as a nature-based  
2 solution for air quality improvement in the Metropolitan city of Rome. *Forests* 7, 150.
- 3 Markevych, I., Schoierer, J., Hartig, T., Chudnovsky, A., Hystad, P., Dzhambov, A.M., de Vries, S.,  
4 Triguero-Mas, M., Brauer, M., Nieuwenhuijsen, M.J., Lupp, G., Richardson, E.A., Astell-Burt,  
5 T., Dimitrova, D., Feng, X., Sadeh, M., Standl, M., Heinrich, J., Fuertes, E., 2017. Exploring  
6 pathways linking greenspace to health: Theoretical and methodological guidance.  
7 *Environmental Research* 158, 301–317. <https://doi.org/10.1016/j.envres.2017.06.028>
- 8 Mavoia, S., Oliver, M., Witten, K., Badland, H.M., 2011. Linking GPS and travel diary data using  
9 sequence alignment in a study of children's independent mobility. *International Journal of*  
10 *Health Geographics* 10, 64.
- 11 McMorris, O., Villeneuve, P.J., Su, J., Jerrett, M., 2015. Urban greenness and physical activity in a  
12 national survey of Canadians. *Environmental research* 137, 94–100.
- 13 Milne, G.L., Dai, Q., Roberts, L.J., 2015. The isoprostanes - 25 years later. *Biochimica et Biophysica*  
14 *Acta - Molecular and Cell Biology of Lipids* 1851, 433–445.  
15 <https://doi.org/10.1016/j.bbali.2014.10.007>
- 16 Mudele, O., Gamba, P., 2019. Mapping vegetation in urban areas using Sentinel-2, in: 2019 Joint Urban  
17 Remote Sensing Event (JURSE). IEEE, pp. 1–4.
- 18 Nikolaidis, M.G., Kyparos, A., Vrabas, I.S., 2011. F2-isoprostane formation, measurement and  
19 interpretation: The role of exercise. *Progress in Lipid Research* 50, 89–103.  
20 <https://doi.org/10.1016/J.PLIPRES.2010.10.002>
- 21 Nowak, D.J., Crane, D.E., Stevens, J.C., 2006. Air pollution removal by urban trees and shrubs in the  
22 United States. *Urban forestry & urban greening* 4, 115–123.
- 23 Ormsby, J.P., Choudhury, B.J., Owe, M., 1987. Vegetation spatial variability and its effect on vegetation  
24 indices. *International Journal of Remote Sensing* 8, 1301–1306.
- 25 Parmes, E., Pesce, G., Sabel, C.E., Baldacci, S., Bono, R., Brescianini, S., D'Ippolito, C., Hanke, W.,  
26 Horvat, M., Liedes, H., Maio, S., Marchetti, P., Marcon, A., Medda, E., Molinier, M., Panunzi,  
27 S., Pärkkä, J., Polańska, K., Prud'homme, J., Ricci, P., Snoj Tratnik, J., Squillacioti, G., Stazi,  
28 M.A., Maesano, C.N., Annesi-Maesano, I., 2020. Influence of residential land cover on  
29 childhood allergic and respiratory symptoms and diseases: Evidence from 9 European cohorts.  
30 *Environmental Research* 183, 108953. <https://doi.org/10.1016/j.envres.2019.108953>
- 31 Pereira, G., Foster, S., Martin, K., Christian, H., Boruff, B.J., Knuiman, M., Giles-Corti, B., 2012. The  
32 association between neighborhood greenness and cardiovascular disease: an observational  
33 study. *BMC Public Health* 12, 466. <https://doi.org/10.1186/1471-2458-12-466>
- 34 Pilat, M.A., McFarland, A., Snelgrove, A., Collins, K., Waliczek, T.M., Zajicek, J., 2012. The effect of  
35 tree cover and vegetation on incidence of childhood asthma in metropolitan statistical areas of  
36 Texas. *HortTechnology* 22, 631–637.
- 37 Revel, C., Lonjou, V., Marcq, S., Desjardins, C., Fougny, B., Coppolani-Delle Luche, C., Guillemint,  
38 N., Lacamp, A.-S., Lourme, E., Miquel, C., 2019. Sentinel-2A and 2B absolute calibration  
39 monitoring. *European Journal of Remote Sensing* 52, 122–137.
- 40 Romanazzi, V., Pirro, V., Bellisario, V., Mengozzi, G., Peluso, M., Pazzi, M., Bugiani, M., Verlatto, G.,  
41 Bono, R., 2013. 15-F2t isoprostane as biomarker of oxidative stress induced by tobacco smoke  
42 and occupational exposure to formaldehyde in workers of plastic laminates. *Science of The*  
43 *Total Environment* 442, 20–25. <https://doi.org/10.1016/j.scitotenv.2012.10.057>
- 44 Rosina, K., Kopecká, M., 2016. Mapping of urban green spaces using Sentinel-2A data: Methodical  
45 aspects, in: 6th International Conference on Cartography and GIS, Albena. Bulgarian  
46 Cartographic Association (in Print). pp. 562–568.
- 47 Sawidis, T., Krystallidis, P., Veros, D., Chettri, M., 2012. A study of air pollution with heavy metals in  
48 Athens city and Attica basin using evergreen trees as biological indicators. *Biological trace*  
49 *element research* 148, 396–408.
- 50 Sbihi, H., Tamburic, L., Koehoorn, M., Brauer, M., 2015. Greenness and Incident Childhood Asthma:  
51 A 10-Year Follow-up in a Population-based Birth Cohort. *American journal of respiratory and*  
52 *critical care medicine* 192, 1131–3. <https://doi.org/10.1164/rccm.201504-0707LE>
- 53 Sies, H., 2015. Oxidative stress: a concept in redox biology and medicine. *Redox biology* 4, 180–3.  
54 <https://doi.org/10.1016/j.redox.2015.01.002>

- 1 Small, C., Lu, J.W., 2006. Estimation and vicarious validation of urban vegetation abundance by spectral  
2 mixture analysis. *Remote Sensing of Environment* 100, 441–456.
- 3 Song, C., 2005. Spectral mixture analysis for subpixel vegetation fractions in the urban environment:  
4 How to incorporate endmember variability? *Remote Sensing of Environment* 95, 248–263.
- 5 Squillaciotti, G., Bellisario, V., Grignani, E., Mengozzi, G., Bardaglio, G., Dalmaso, P., Bono, R.,  
6 2019a. The Asti Study: The Induction of Oxidative Stress in A Population of Children  
7 According to Their Body Composition and Passive Tobacco Smoking Exposure. *International*  
8 *Journal of Environmental Research and Public Health* 16, 490.  
9 <https://doi.org/10.3390/ijerph16030490>
- 10 Squillaciotti, G., Bellisario, V., Grosso, A., Ghelli, F., Piccioni, P., Grignani, E., Corsico, A., Bono, R.,  
11 2020. Formaldehyde, Oxidative Stress, and FeNO in Traffic Police Officers Working in Two  
12 Cities of Northern Italy. *International Journal of Environmental Research and Public Health* 17,  
13 1655.
- 14 Squillaciotti, G., Bellisario, V., Levra, S., Piccioni, P., Bono, R., 2019b. Greenness Availability and  
15 Respiratory Health in a Population of Urbanised Children in North-Western Italy. *International*  
16 *Journal of Environmental Research and Public Health* 17, 108.  
17 <https://doi.org/10.3390/ijerph17010108>
- 18 Thiering, E., Markevych, I., Brüske, I., Fuertes, E., Kratzsch, J., Sugiri, D., Hoffmann, B., von Berg, A.,  
19 Bauer, C.-P., Koletzko, S., 2016. Associations of residential long-term air pollution exposures  
20 and satellite-derived greenness with insulin resistance in German adolescents. *Environmental*  
21 *health perspectives* 124, 1291–1298.
- 22 Tillberg Mattsson, K., 2002. Children's (in) dependent mobility and parents' chauffeuring in the town  
23 and the countryside. *Tijdschrift voor economische en sociale geografie* 93, 443–453.
- 24 Tsai, W.-L., Leung, Y.-F., McHale, M.R., Floyd, M.F., Reich, B.J., 2019. Relationships between urban  
25 green land cover and human health at different spatial resolutions. *Urban Ecosystems* 22, 315–  
26 324. <https://doi.org/10.1007/s11252-018-0813-3>
- 27 van 't Erve, T.J., Kadiiska, M.B., London, S.J., Mason, R.P., 2017. Classifying oxidative stress by F2-  
28 isoprostane levels across human diseases: A meta-analysis. *Redox Biology* 12, 582–599.  
29 <https://doi.org/10.1016/j.redox.2017.03.024>
- 30 Warton, D.I., Wright, I.J., Falster, D.S., Westoby, M., 2006. Bivariate line-fitting methods for allometry.  
31 *Biological reviews* 81, 259–291.
- 32 Willmott, C.J., Matsuura, K., 2005. Advantages of the mean absolute error (MAE) over the root mean  
33 square error (RMSE) in assessing average model performance. *Climate research* 30, 79–82.
- 34 Wolch, J.R., Byrne, J., Newell, J.P., 2014. Urban green space, public health, and environmental justice:  
35 The challenge of making cities 'just green enough.' *Landscape and urban planning* 125, 234–  
36 244.
- 37 Woo, J., Tang, N., Suen, E., Leung, J., Wong, M., 2009. Green space, psychological restoration, and  
38 telomere length. *Lancet (London, England)* 373, 299–300. [https://doi.org/10.1016/S0140-6736\(09\)60094-5](https://doi.org/10.1016/S0140-6736(09)60094-5)
- 39 Yeager, R., Riggs, D.W., DeJarnett, N., Tollerud, D.J., Wilson, J., Conklin, D.J., O'Toole, T.E.,  
40 McCracken, J., Lorkiewicz, P., Xie, Z., Zafar, N., Krishnasamy, S.S., Srivastava, S., Finch, J.,  
41 Keith, R.J., DeFilippis, A., Rai, S.N., Liu, G., Bhatnagar, A., 2018. Association Between  
42 Residential Greenness and Cardiovascular Disease Risk. *Journal of the American Heart*  
43 *Association* 7, e009117. <https://doi.org/10.1161/JAHA.118.009117>
- 44 Zhang, X., Friedl, M.A., Schaaf, C.B., Strahler, A.H., Hodges, J.C., Gao, F., Reed, B.C., Huete, A.,  
45 2003. Monitoring vegetation phenology using MODIS. *Remote sensing of environment* 84,  
46 471–475.
- 47 Zhou, D., Zhao, S., Zhang, L., Liu, S., 2016. Remotely sensed assessment of urbanization effects on  
48 vegetation phenology in China's 32 major cities. *Remote Sensing of Environment* 176, 272–  
49 281.

Table 1. Technical features of the S2 images obtained from Theia CNES provider (<https://www.Theia-land.fr/en/product/sentinel-2-surface-reflectance/>)

<b>Band ID</b>	<b>Central wavelength (nm)</b>	<b>Bandwidth (nm)</b>	<b>Nominal Geometric resolution by Theia CNES provider</b>
B2	490	98	10 m
B3	560	45	10 m
B4	665	38	10 m
B5	705	19	10 m (20 m native resolution)
B6	740	18	10 m (20 m native resolution)
B7	775	28	10 m (20 m native resolution)
B8	840	145	10 m
B8A	865	33	10 m (20 m native resolution)
B11	1610	143	10 m (20 m native resolution)
B12	2200	242	10 m (20 m native resolution)

Table 2 – Coefficients of linear regression models between 15-F2t-IsoP vs PGC, IsoP vs NET, and 15-F2t-IsoP vs NBT.

All dependent variables were poorly correlated with OS.

<b>Dependent variable</b>	<b><i>r</i></b>	<b><i>p</i>-value</b>	<b>R<sup>2</sup></b>	<b>Slope</b>	<b><i>p</i>(slope)</b>	<b>Slope (95% C.Is)</b>	<b>Intercept</b>
uSAVI	-0.045	0.520	0.002	-2.214	0.520	(-9.538, 4.685)	11.967
PGC	-0.046	0.519	0.002	-2.126	0.519	(-8.747, 4.819)	12.035
NBT	-0.028	0.691	0.001	-0.001	0.691	(-0.003, 0.002)	11.699
NET	-0.030	0.673	0.001	-0.005	0.673	(-0.025, 0.014)	11.523

*Table 3 – Twenty equi-probable classes of 15-F2t-IsoP (divided by about 1.4 ng/mg crea step) and vegetation parameters. In order to calculate the envelope function of each scatterplot, maximum values of equi-probable class were extracted. For each equi-probable class maximum value and coefficient of variation (%) were reported. Slash sign denotes only one value within equi-probable class.*

<b>15-F2t-IsoP</b>		<b>uSAVI</b>		<b>NBT</b>		<b>NET</b>		<b>PGC</b>	
<b>Max</b>	<b>CV%</b>	<b>Max</b>	<b>CV%</b>	<b>Max</b>	<b>CV%</b>	<b>Max</b>	<b>CV%</b>	<b>Max</b>	<b>CV%</b>
<b>3.56</b>	39.9	0.47	46.9	990	55.2	102	91.5	0.514	43.5
<b>4.93</b>	10.8	0.42	38.8	1874	58.9	89	84.3	0.487	34.6
<b>6.35</b>	7.7	0.44	37.5	1239	51.7	115	95.8	0.512	34.5
<b>7.52</b>	4.2	0.57	51.6	1016	37.6	113	94.9	0.586	46.5
<b>9.16</b>	5.5	0.63	46.2	1310	48.6	136	100.0	0.619	40.8
<b>10.55</b>	4.2	0.65	45.5	1262	45.8	244	99.4	0.616	39.9
<b>11.85</b>	3.2	0.44	40.3	1144	33.0	115	80.7	0.492	35.9
<b>13.35</b>	3.0	0.50	35.1	1124	30.2	165	91.4	0.497	31.2
<b>14.56</b>	3.2	0.42	45.8	1103	72.6	108	119.2	0.477	43.5
<b>16.05</b>	2.6	0.42	43.1	1144	41.8	115	79.6	0.473	41.1
<b>17.17</b>	1.2	0.44	60.6	923	60.0	117	167.9	0.517	57.6
<b>18.74</b>	2.2	0.47	46.1	1107	50.3	92	108.4	0.484	44.3
<b>20.23</b>	1.8	0.36	43.7	1163	55.5	79	105.8	0.426	42.6
<b>21.18</b>	1.4	0.42	36.7	1111	58.6	79	137.0	0.457	33.1
<b>22.88</b>	1.8	0.41	59.7	1118	88.9	76	95.8	0.471	60.3
<b>24.2</b>	2.7	0.56	20.7	892	2.2	77	51.3	0.575	13.5
<b>26.81</b>	/	0.41	/	995	/	101	/	0.469	/
<b>28.98</b>	0.7	0.34	69.5	1109	52.1	19	/	0.411	66.9
<b>34.34</b>	/	0.11	/	206	/	0	/	0.139	/
<b>39.65</b>	/	0.16	/	170	/	6	/	0.190	/

Table 4 – Bivariate relationships statistics of 15-F2t-IsoP-Ev and vegetation parameters. Significance levels: \*  $p$

$<0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$ .

<b>Proxy</b>	<b><math>r</math></b>	<b><math>R^2</math></b>	<b>Slope</b>	<b>Slope (95% C.I.s)</b>	<b>Intercept</b>
<b>PGC</b>	-0.758***	0.57	-62.75***	(-83.671 -34.792)	34.26
<b>uSAVI</b>	-0.717***	0.51	-0.01***	(-0.016, -0.005)	0.59
<b>NET</b>	-0.696***	0.48	-0.12***	(-0.192, -0.023)	30.02
<b>NBT</b>	-0.743***	0.55	-0.02***	(-0.031, -0.010)	39.17

**S.D.P.:** Conceptualization, Methodology, Software, Formal analysis, Writing - Original Draft; **G.S.:** Conceptualization, Methodology, Software, Formal analysis, Writing - Original Draft, Data curation; **R.B.:** Supervision, Project administration, Writing - Review & Editing; **E.B.M.:** Supervision, Project administration, Writing - Review & Editing



**Declaration of interests**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

**Declaration of interests**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.