



UNIVERSITÀ DEGLI STUDI DI TORINO

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Cytokine, chemokine, and growth factor profile of platelet-rich plasma

This is the author's manuscript
Original Citation:
Availability:
This version is available http://hdl.handle.net/2318/1557497 since 2017-02-21T12:18:47Z
Published version:
DOI:10.3109/09537104.2016.1143922
Terms of use:
Open Access
Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This is an author version of the contribution published on: Questa è la versione dell'autore dell'opera: Platelets. 2016 Mar 7:1-5. [Epub ahead of print]
Cytokine, chemokine, and growth factor profile of platelet-rich plasma. Mussano F, Genova T, Munaron L, Petrillo S, Erovigni F, Carossa S.

> *The definitive version is available at:* La versione definitiva è disponibile alla URL: http://www.ncbi.nlm.nih.gov/pubmed/26950533

Cytokine, chemokine and growth factor profile of Platelet Rich Plasma

Mussano F¹*, Genova T^{1,2}*, Munaron L^{2,3}, Petrillo S, Erovigni F¹, Carossa S¹

*Equally contributed to the paper

CIR Dental School, Department of Surgical Sciences, University of Turin, via Nizza
 230, 10126 Turin, Italy;

2. Department of Life Sciences and Systems Biology, University of Turin, via Accademia Albertina 13, 10123, Turin, Italy;

3. Centre for Nanostructured Interfaces and Surfaces (NIS), Turin, Italy;

4. Molecular Biotechnology Center, University of Turin, Turin, Italy.

A short title not exceeding 50 characters (including spaces):

Biomolecules within Platelet Rich Plasma Keywords (2-6):

Platelet Rich Plasma

Growth factors

Chemokine

Interleukines

Abstract (150-400 words unstructured)

During wound healing, biologically active molecules are released from platelets. The rationale of using Platelet Rich Plasma (PRP) relies on the concentration of bioactive molecules and subsequent delivery to healing sites. These bioactive molecules have been seldom simultaneously quantified within the same PRP preparation. In the present study, the flexible Bio-Plex system was employed to assess the concentration of a large range of cytokines, chemokines and growth factors in sixteen healthy volunteers so as to determine whether significant baseline differences may be found. Besides IL-1b, IL-1ra, IL-4, IL-6, IL-8, IL-12, IL-13, IL-17, INF-γ, TNF-α, monocyte chemoattractant protein-1 (MCP-1) (CCL-2), MIP-1a (CCL-3), RANTES (CCL-5), bFGF, PDGF, VEGF that were already quantified elsewhere, the authors reported also on the presence of IL-2, IL-5, IL-7, IL-9, IL-10, IL-15 G-CSF, GM-CSF, Eotaxin (CCL-11), CXCL10 chemokine (IP-10), MIP 1b (CCL-4). Among the most interesting results, it is convenient to mention the high concentrations of the HIV-suppressive and inflammatory cytokine RANTES and a statistically significant difference between males and females in the content of PDGF-BB. These data are consistent with previous reports pointing out that gender, diet and test system affect the results of platelet function in healthy subjects, but seem contradictory when compared to other quantification assays in serum and plasma.

The inconsistencies affecting the experimental results found in literature, along with the variability found in the content of bioactive molecules, urge further research, hopefully in form of Randomized Controlled Clinical Trials, in order to find definitive evidence of the efficacy of PRP treatment in various pathologic and regenerative conditions.

Introduction

Platelet-rich plasma (PRP) emerges amongst the most innovative autologous blood products used to enhance tissue healing and regeneration. Blood withdrawn from a patient's peripheral vein is centrifuged to concentrate platelets. PRP may be immediately used either as it is, or in combination with other biomaterials. For sake of precision, based on leukocytes and fibrin content, PRP products are subdivided as follows: pure platelet-rich plasma (P-PRP) also known as plasma rich in growth factors (PRGF), leukocyte- and platelet-rich plasma (L-PRP), pure platelet-rich fibrin (P-PRF), and leuko-cyte- and platelet-rich fibrin (L-PRF)¹. Usually in the form of gel or liquid, P-PRP and L-PRP have both a low-density fibrin network, while P-PRF and L-PRF, available only in the gel form, contain high-density fibrin network.

Platelet-derived growth factor (PDGF), epidermal growth factor (EGF), insulin-like growth factor (IGF-I), transforming growth factor β -I (TGF β -I), vascular endothelial growth factor (VEGF), hepatocyte growth factor (HGF), and basic fibroblast growth factor (bFGF) are just a few of the growth factors (GFs) present within the α -granules of platelets ^{2,3}. Such a wide range of GFs is one of the keys to understand the multifaceted roles of PRP, including proliferation and differentiation of cells belonging to the musculoskeletal and vascular systems ⁴. However, α -granules also contain different interleukins (ILs) and chemokines, such as IL-1 β , IL-8, and MIP-1-2-3 ^{5,6}, that are inflammation mediators able to stimulate cell chemotaxis and maturation.

Several studies ^{7–9} evaluated in vitro the effect of PRPs supporting their ability to improve proliferation and osteogenic activity of osteoblasts and mesenchymal stem cells. However, these effects were dependent on the PRP composition ¹⁰. Somehow data become controversial when in vivo settings are compared, since opposite outcomes are equally available in the scientific literature in favor ^{11–21} or against ^{19,22–27} the addition of PRP to biologic and synthetic graft materials for bone regeneration purposes. Interestingly, the association of PRP with mesenchymal stem cells showed mostly satisfying results ^{28,29} in the field. Recently, regenerative medicine and tissue engineering focused on the use of GFs ³⁰ and cell-based therapy to improve the quality and speed of healing, extending the PRP application from the traditional dental and maxillofacial field to the treatment of musculoskeletal injuries ³¹.

Currently, on the market, there exist more than 40 commercial systems able to concentrate whole blood into a platelet-rich substance. Such a relevant number of different preparation protocols should be taken into account whenever inconsistencies are found from comparing clinical outcomes reported in the literature ³². The quality of PRP could be affected by patient's heterogeneity in terms of age, gender, body mass index, comorbidities, healing capabilities, and different lifestyles ³³. In order to reduce at least the protocol dependent variability, the authors selected a closed preparation system, which, being automatic, limits the possibility of operator dependent errors and reduces the risk of microbial contamination, while processing the blood.

The aim of the present research is to assess the concentration of a large range of cytokines, chemokines and growth factors in a proper sample of healthy volunteers so as to determine whether significant baseline differences may be found.

Material and methods

Sample preparation

Sixteen healthy volunteers (mean age= 24.25 ± 4.96 years, age range: 20 to 40 years, 10 females and 6 males) were recruited as possible blood donors, based on their medical histories, at the Interdepartmental Research Center (IRC) Dental School of the University of Turin, between January 1st and February 1st 2015. The study protocol was approved by the Ethics Committee of the IRC. The informed consent was obtained. PRP samples were prepared with the CPunT System, (Eltek Group, Casale Monferrato, Italy). The CPunT Preparation System is a closed system medical device that consists of a main unit, which separates the PRP from other substances, and the appropriate disposable components (for detailed information please refer to supplementary materials). Briefly, the samples were processed according to the following protocol. First, blood was withdrawn and centrifuged at 1200 rpm for 11 minutes in an ELTEK centrifuge (Cat number 10714800), after adding ACD (Sodium Citrate / Citric Acid / Glucose SALF SpA, Italy). This centrifugation separated the buffy coat and the red blood cells from platelets and plasma, which were automatically pushed into a separate bag under the control of an optical sensor so as to prevent any buffy coat contaminations. The bag was subsequently positioned in the centrifuge basket for a second centrifugation (2000 rpm for 9 min). At this step, the Platelet Concentrate (PC) was sedimented on the bottom of the bag. Part of PPP (Poor Platelets Plasma), 4 mL, was removed through the accessory 10-mL syringe with needle. The bag with Platelet Concentrate and the remaining plasma were mixed gently, massaging the bag. Thus, the Platelet Rich Plasma (PRP) was obtained.

Detection of Interleukins, Chemokines and Growth Factors using Bio-Plex system

The PRP samples thus obtained were characterized by measuring the concentration of the following specific biomolecules: interleukin-1b, interleukin 1ra, interleukin-2, interleukin-4, interleukin-5, interleukin-6, interleukin-7, interleukin-8, interleukin-9, interleukin-10, interleukin-12, interleukin-13, interleukin-15, interleukin-17, granulocyte-colony stimulating factor (G-CSF), granulocyte-macrophage colony stimulating factor (GM-CSF), interferon-gamma (INF- γ), tumor necrosis factor- α (TNF- α), monocyte chemoattractant protein-1 (MCP-1), CXCL10 chemokine (IP-10), MIP-1a, MIP 1b, RANTES, eotaxin, platelet derived growth factor (PDGF), basic-fibroblastic growth factor (bFGF), vascular endothelial growth factor (VEGF). The flexible Bio-Plex system (Bio-Rad Laboratories, Hercules, CA, USA) was emplyed. All samples were analysed following the manufacturer's protocol. At least two independent repetitions in duplicate were made per sample. Concentrations of the analytes were expressed in pg/ml. A standard curve ranging on average from 0.15 pg/ml to 3700 pg/ml (High Photomultiplier Tube Setting -PMT setting) was prepared and then fitted by Bio-Plex Manager software.

Blood count

Platelets from every processed sample were counted with Neubauer Improved counting chamber (Marienfeld, Lauda-Königshofen, Germany).

Statistical Analysis

Red blood cells and platelet count were analyzed using Student t test. A descriptive analysis of Bio-Plex data was performed presenting data using means±standard error

mean (SEM). Differences between groups were analyzed using the two-way ANOVA with Tukey's multiple comparison test.

All the statistical analysis were performed using GraphPad Prism 6 software (GraphPad Software, Inc., La Jolla, CA, USA) and were conducted with a 0.05 level of significance.

Results

The blood processing work-flow used to get PRP is portrayed in Fig.1 along with the number of the platelets at the key steps of the procedure. The concentrations of the Cytokines (Fig.2), Chemokines (Fig.3) and Growth factors (Fig.4) detected within the PRPs of the 16 healthy volunteers are reported as means and are subdivided by gender. It is noteworthy that only PDGF differed in a statistically significant way between males and females.

Discussion

Platelets are fundamental factors of the clotting cascade that is central to the wound healing ³⁴. During this process, biologically active molecules are released from the α -granules ^{5,35}. Within the first 10 minutes after activation about 70% of the GFs are secreted reaching almost 100% within the first hour ³⁶. The rationale of the use of PRP is that it concentrates more platelets than the whole blood, allowing the delivery of bioactive GFs and molecules that promote tissue healing. To date, however, there is still little information available about chemokines, cytokines and growth factors simultaneously quantified within the same PRP preparation. This lack of knowledge may be due to the analytic techniques used, which are mostly ELISA-based kits ⁶.

To address the issue, the flexible Bio-Plex system was employed in the present study. Based on a capture sandwich immunoassay, this technology allowed for the simultaneous dosage of different biomolecules to be done in a single microplate well, as previously described ³⁷. Besides the biomolecules that had already been quantified in PRPs such as: IL-1b IL-1ra ^{38,39}, IL-4 ⁴⁰, IL-6 ³⁹, IL-8 ^{40,41}, IL-12 ⁶, IL-13 ⁴⁰, IL-17 ⁴⁰, INF- γ (detected contrary to ⁴⁰), TNF- α ^{39,40}, monocyte chemoattractant protein-1 (MCP-1) (CCL-2) ⁴², MIP-1a (CCL-3) ⁴², RANTES (CCL-5) ⁶, bFGF ⁶, PDGF ^{6,38,41}, VEGF ^{6,38,41} the authors reported also on the presence of IL-2, IL-5, IL-7, IL-9, IL-10, IL-15 G-CSF, GM-CSF, Eotaxin (CCL-11), CXCL10 chemokine (IP-10), MIP 1b (CCL-4).

Some considerations immediately emerge when analyzing the results here shown. Although RANTES (CCL5) plays an active role in recruiting leukocytes into inflammatory sites, its more refined modulatory and chemotactic properties have been already shown by El-Sharkawy et al ⁶. The high concentration of RANTES here detected is therefore consistent with the acceleration of healing and regenerative processes often favorably reported. Interestingly, from the data analysis, the only difference concerning possible gender variability dealt with the concentration of PDGF, which was higher in males, independently of the platelet count. This result may be partially discordant with the data published by Gomez et al.⁴³, but it is certainly consistent with Miller and colleagues ⁴⁴, who pointed out that gender, diet and test system affected the results of platelet function in healthy subjects. It is noteworthy that the values of PDGF-BB here described are quite similar to those reported previously ⁶, while the content of PDGF was respectively higher in serum ⁴⁵ and lower in plasma ⁴⁶ according to other studies. Contrary to Amable et al. ⁴⁰, INF-γ could be detected and quantified in non irrelevant amount. Also, other authors described more elevated PDGF concentrations in plasma than we could determine within PRPs. ⁴⁷

These inconsistencies affecting the experimental results found in literature may owe to countless factors such as the varied types of activating methods (CaCl2, thrombin, batroxobin, bovine thrombin, and thrombin added to CaCl2), the different sample volumes of blood (from 9 to 120 mL) or PRP (from 3 to 32 mL), the number of spins during centrifugation (1 or 2), and the platelet concentration (from 1x to 18x) ^{48,49}.

The widely recognized importance of PRP as an accessible source of growth factors is supported even by this study portraying the variety of bioactive molecules. Indeed, it must not be neglected that, besides the mitogenic and anabolic properties individually attributed to PDGF-BB, FGF-b, VEGF and other well known growth factors, synergistic effects may be taken into consideration also in case of the relatively low concentrations of signal molecules whose role is less characterized ⁵⁰. In conclusion, further research, hopefully in form of Randomized Controlled Clinical Trials, is required to find definitive evidence of the efficacy of PRP treatment in various pathologic and regenerative conditions.

Bibliography

1. Dohan Ehrenfest DM, Andia I, Zumstein MA, Zhang C-Q, Pinto NR, Bielecki T. Classification of platelet concentrates (Platelet-Rich Plasma-PRP, Platelet-Rich Fibrin-PRF) for topical and infiltrative use in orthopedic and sports medicine: current consensus, clinical implications and perspectives. Muscles, ligaments and tendons journal. 2014 [accessed 2015 Jun 13];4(1):3–9. http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4049647&tool=pmcentrez &rendertype=abstract

2. Pietrzak WS, Eppley BL. Platelet rich plasma: biology and new technology. The Journal of craniofacial surgery. 2005 [accessed 2015 Nov 17];16(6):1043–54. http://www.ncbi.nlm.nih.gov/pubmed/16327552

3. Mejia HA, Bradley JP. The Effects of Platelet-Rich Plasma on Muscle: Basic Scienceand Clinical Application. Operative Techniques in Sports Medicine. 2011 [accessed2015Nov17];19(3):149–153.http://www.sciencedirect.com/science/article/pii/S1060187211000372

4. Stiles CD. The molecular biology of platelet-derived growth factor. Cell. 1983 [accessed 2015 Nov 17];33(3):653–5. http://www.ncbi.nlm.nih.gov/pubmed/6307524

5. Nurden AT. Platelets, inflammation and tissue regeneration. Thrombosis and Haemostasis. 2011 [accessed 2015 Oct 16];105(Suppl. 1):S13–S33. http://www.ncbi.nlm.nih.gov/pubmed/21479340

6. El-Sharkawy H, Kantarci A, Deady J, Hasturk H, Liu H, Alshahat M, Van Dyke TE. Platelet-rich plasma: growth factors and pro- and anti-inflammatory properties. Journal of periodontology. 2007 [accessed 2015 Nov 17];78(4):661–9. http://www.ncbi.nlm.nih.gov/pubmed/17397313

7. Lu HH, Vo JM, Chin HS, Lin J, Cozin M, Tsay R, Eisig S, Landesberg R. Controlled delivery of platelet-rich plasma-derived growth factors for bone formation. Journal of biomedical materials research. Part A. 2008 [accessed 2015 Nov 17];86(4):1128–36. http://www.ncbi.nlm.nih.gov/pubmed/18181109

8. Man Y, Wang P, Guo Y, Xiang L, Yang Y, Qu Y, Gong P, Deng L. Angiogenic and osteogenic potential of platelet-rich plasma and adipose-derived stem cell laden alginate microspheres. Biomaterials. 2012 [accessed 2015 Nov 17];33(34):8802–11. http://www.ncbi.nlm.nih.gov/pubmed/22981779

9. Kawasumi M, Kitoh H, Siwicka KA, Ishiguro N. The effect of the platelet concentration in platelet-rich plasma gel on the regeneration of bone. The Journal of bone and joint surgery. British volume. 2008 [accessed 2015 Nov 17];90(7):966–72. http://www.ncbi.nlm.nih.gov/pubmed/18591611

10. Salamanna F, Veronesi F, Maglio M, Della Bella E, Sartori M, Fini M. New and emerging strategies in platelet-rich plasma application in musculoskeletal regenerative procedures: general overview on still open questions and outlook. BioMed research

international. 2015 [accessed 2015 Oct 1];2015:846045. http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4436449&tool=pmcentrez &rendertype=abstract

11. Mariano R, Messora M, de Morais A, Nagata M, Furlaneto F, Avelino C, Paula F, Ferreira S, Pinheiro M, de Sene JP. Bone healing in critical-size defects treated with platelet-rich plasma: a histologic and histometric study in the calvaria of diabetic rat. Oral surgery, oral medicine, oral pathology, oral radiology, and endodontics. 2010 [accessed 2015 Nov 17];109(1):72–8. http://www.ncbi.nlm.nih.gov/pubmed/19926499

12. Messora MR, Nagata MJH, Mariano RC, Dornelles RCM, Bomfim SRM, Fucini SE, Garcia VG, Bosco AF. Bone healing in critical-size defects treated with plateletrich plasma: a histologic and histometric study in rat calvaria. Journal of periodontal research. 2008 [accessed 2015 Nov 17];43(2):217–23. http://www.ncbi.nlm.nih.gov/pubmed/18302625

13. Simman R, Hoffmann A, Bohinc RJ, Peterson WC, Russ AJ. Role of platelet-richplasma in acceleration of bone fracture healing. Annals of plastic surgery. 2008[accessed2015Nov17];61(3):337–44.http://www.ncbi.nlm.nih.gov/pubmed/18724139

14. Kang Y-H, Jeon SH, Park J-Y, Chung J-H, Choung Y-H, Choung H-W, Kim E-S, Choung P-H. Platelet-rich fibrin is a Bioscaffold and reservoir of growth factors for tissue regeneration. Tissue engineering. Part A. 2011 [accessed 2015 Nov 2];17(3-4):349–59. http://www.ncbi.nlm.nih.gov/pubmed/20799908

15. Rai B, Oest ME, Dupont KM, Ho KH, Teoh SH, Guldberg RE. Combination of platelet-rich plasma with polycaprolactone-tricalcium phosphate scaffolds for segmental bone defect repair. Journal of biomedical materials research. Part A. 2007 [accessed 2015 Nov 17];81(4):888–99. http://www.ncbi.nlm.nih.gov/pubmed/17236215

16. Chang S-H, Hsu Y-M, Wang YJ, Tsao Y-P, Tung K-Y, Wang T-Y. Fabrication of
pre-determined shape of bone segment with collagen-hydroxyapatite scaffold and
autogenous platelet-rich plasma. Journal of materials science. Materials in medicine.2009[accessed2015Nov17];20(1):23–31.http://www.ncbi.nlm.nih.gov/pubmed/18651114

17. Chen J-C, Ko C-L, Shih C-J, Tien Y-C, Chen W-C. Calcium phosphate bone cementwith 10 wt% platelet-rich plasma in vitro and in vivo. Journal of dentistry. 2012[accessed2015Nov18];40(2):114–22.http://www.ncbi.nlm.nih.gov/pubmed/22101118

18. Messora MR, Nagata MJH, Fucini SE, Pola NM, Campos N, de Oliveira GC V, Bosco AF, Garcia VG, Furlaneto FAC. Effect of platelet-rich plasma on the healing of mandibular defects treated with fresh frozen bone allograft: a radiographic study in dogs. The Journal of oral implantology. 2014 [accessed 2015 Nov 18];40(5):533–41.

http://www.ncbi.nlm.nih.gov/pubmed/25295885

19. Mooren RECM, Merkx MAW, Bronkhorst EM, Jansen JA, Stoelinga PJW. The effect of platelet-rich plasma on early and late bone healing: an experimental study in goats. International journal of oral and maxillofacial surgery. 2007 [accessed 2015 Nov 17];36(7):626–31. http://www.ncbi.nlm.nih.gov/pubmed/17521885

20. Nagata M, Messora M, Okamoto R, Campos N, Pola N, Esper L, Sbrana M, Fucini S, Garcia V, Bosco A. Influence of the proportion of particulate autogenous bone graft/platelet-rich plasma on bone healing in critical-size defects: an immunohistochemical analysis in rat calvaria. Bone. 2009 [accessed 2015 Nov 18];45(2):339–45. http://www.ncbi.nlm.nih.gov/pubmed/19410024

21. Kanthan SR, Kavitha G, Addi S, Choon DSK, Kamarul T. Platelet-rich plasma (PRP) enhances bone healing in non-united critical-sized defects: a preliminary study involving rabbit models. Injury. 2011 [accessed 2015 Nov 17];42(8):782–9. http://www.ncbi.nlm.nih.gov/pubmed/21329922

22. Nikolidakis D, van den Dolder J, Wolke JGC, Jansen JA. Effect of platelet-rich plasma on the early bone formation around Ca-P-coated and non-coated oral implants in cortical bone. Clinical oral implants research. 2008 [accessed 2015 Nov 17];19(2):207–13. http://www.ncbi.nlm.nih.gov/pubmed/18067601

23. Cinotti G, Corsi A, Sacchetti B, Riminucci M, Bianco P, Giannicola G. Bone ingrowth and vascular supply in experimental spinal fusion with platelet-rich plasma. Spine. 2013 [accessed 2015 Nov 17];38(5):385–91. http://www.ncbi.nlm.nih.gov/pubmed/22885836

24. Hatakeyama M, Beletti ME, Zanetta-Barbosa D, Dechichi P. Radiographic and histomorphometric analysis of bone healing using autogenous graft associated with platelet-rich plasma obtained by 2 different methods. Oral surgery, oral medicine, oral pathology, oral radiology, and endodontics. 2008 [accessed 2015 Nov 17];105(1):e13–8. http://www.ncbi.nlm.nih.gov/pubmed/18155595

25. Torres J, Tamimi FM, Tresguerres IF, Alkhraisat MH, Khraisat A, Lopez-Cabarcos E, Blanco L. Effect of solely applied platelet-rich plasma on osseous regeneration compared to Bio-Oss: a morphometric and densitometric study on rabbit calvaria. Clinical implant dentistry and related research. 2008 [accessed 2015 Nov 17];10(2):106–12. http://www.ncbi.nlm.nih.gov/pubmed/18462207

26. Giovanini AF, Deliberador TM, Gonzaga CC, de Oliveira Filho MA, Göhringer I, Kuczera J, Zielak JC, de Andrade Urban C. Platelet-rich plasma diminishes calvarial bone repair associated with alterations in collagen matrix composition and elevated CD34+ cell prevalence. Bone. 2010 [accessed 2015 Nov 17];46(6):1597–603. http://www.ncbi.nlm.nih.gov/pubmed/20206725

27. Broggini N, Hofstetter W, Hunziker E, Bosshardt DD, Bornstein MM, Seto I, Weibrich G, Buser D. The influence of PRP on early bone formation in membrane

protected defects. A histological and histomorphometric study in the rabbit calvaria. Clinical implant dentistry and related research. 2011 [accessed 2015 Nov 17];13(1):1–12. http://www.ncbi.nlm.nih.gov/pubmed/20156229

28. Zhong W, Sumita Y, Ohba S, Kawasaki T, Nagai K, Ma G, Asahina I. In vivo comparison of the bone regeneration capability of human bone marrow concentrates vs. platelet-rich plasma. PloS one. 2012 [accessed 2015 Nov 17];7(7):e40833. http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3395629&tool=pmcentrez &rendertype=abstract

29. Kasten P, Beverungen M, Lorenz H, Wieland J, Fehr M, Geiger F. Comparison of platelet-rich plasma and VEGF-transfected mesenchymal stem cells on vascularization and bone formation in a critical-size bone defect. Cells, tissues, organs. 2012 [accessed 2015 Nov 17];196(6):523–33. http://www.ncbi.nlm.nih.gov/pubmed/22796828

30. Anitua E, Sánchez M, Orive G. The importance of understanding what is plateletrich growth factor (PRGF) and what is not. Journal of shoulder and elbow surgery / American Shoulder and Elbow Surgeons ... [et al.]. 2011 [accessed 2015 Nov 17];20(1):e23–4; author reply e24. http://www.ncbi.nlm.nih.gov/pubmed/21050777

31. Kaux J-F, Janssen L, Drion P, Nusgens B, Libertiaux V, Pascon F, Heyeres A, Hoffmann A, Lambert C, Le Goff C, et al. Vascular Endothelial Growth Factor-111 (VEGF-111) and tendon healing: preliminary results in a rat model of tendon injury. Muscles, ligaments and tendons journal. 2014 [accessed 2015 Nov 17];4(1):24–8. http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4049645&tool=pmcentrez &rendertype=abstract

32. Kaux J-F, Le Goff C, Seidel L, Péters P, Gothot A, Albert A, Crielaard J-M. [Comparative study of five techniques of preparation of platelet-rich plasma]. Pathologie-biologie. 2011 [accessed 2015 Nov 17];59(3):157–60. http://www.ncbi.nlm.nih.gov/pubmed/19481375

33. Evanson JR, Guyton MK, Oliver DL, Hire JM, Topolski RL, Zumbrun SD, McPherson JC, Bojescul JA. Gender and age differences in growth factor concentrations from platelet-rich plasma in adults. Military medicine. 2014 [accessed 2015 Nov 17];179(7):799–805. http://www.ncbi.nlm.nih.gov/pubmed/25003868

34. Wroblewski AP, Mejia HA, Wright VJ. Application of Platelet-Rich Plasma to Enhance Tissue Repair. Operative Techniques in Orthopaedics. 2010 [accessed 2015 Nov 18];20(2):98–105.

http://www.sciencedirect.com/science/article/pii/S1048666609001463

35. Nurden AT, Nurden P, Sanchez M, Andia I, Anitua E. Platelets and wound healing. Frontiers in bioscience: a journal and virtual library. 2008 [accessed 2015 Nov 8];13:3532–48. http://www.ncbi.nlm.nih.gov/pubmed/18508453

36. Marx RE. Platelet-rich plasma (PRP): what is PRP and what is not PRP? Implantdentistry.2001[accessed2015Nov18];10(4):225–8.

http://www.ncbi.nlm.nih.gov/pubmed/11813662

37. Sacerdote P, Mussano F, Franchi S, Panerai AE, Bussolati G, Carossa S, Bartorelli A, Bussolati B. Biological components in a standardized derivative of bovine colostrum. Journal of dairy science. 2013 [accessed 2015 Nov 18];96(3):1745–54. http://www.ncbi.nlm.nih.gov/pubmed/23332842

38. Oh JH, Kim W, Park KU, Roh YH. Comparison of the Cellular Composition and Cytokine-Release Kinetics of Various Platelet-Rich Plasma Preparations. The American journal of sports medicine. 2015 Oct 15 [accessed 2015 Nov 18]. http://www.ncbi.nlm.nih.gov/pubmed/26473014

39. Kaur D, Sharma RR, Marwaha N. Defining an appropriate leucoreduction strategy by serial assessment of cytokine levels in platelet concentrates prepared by different methods. Asian journal of transfusion science. [accessed 2015 Nov 18];9(1):31–5. http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4339928&tool=pmcentrez &rendertype=abstract

40. Amable PR, Carias RBV, Teixeira MVT, da Cruz Pacheco I, Corrêa do Amaral RJF, Granjeiro JM, Borojevic R. Platelet-rich plasma preparation for regenerative medicine: optimization and quantification of cytokines and growth factors. Stem cell research & therapy. 2013 [accessed 2015 Nov 18];4(3):67. http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3706762&tool=pmcentrez &rendertype=abstract

41. Kang J, Hur J, Kang J-A, Yun J-Y, Choi J-I, Ko SB, Lee C-S, Lee J, Han J-K, Kim HK, et al. Activated platelet supernatant can augment the angiogenic potential of human peripheral blood stem cells mobilized from bone marrow by G-CSF. Journal of molecular and cellular cardiology. 2014 [accessed 2015 Nov 18];75:64–75. http://www.ncbi.nlm.nih.gov/pubmed/25016235

42. Galliera E, Corsi MM, Banfi G. Platelet rich plasma therapy: inflammatory molecules involved in tissue healing. Journal of biological regulators and homeostatic agents. [accessed 2015 Nov 18];26(2 Suppl 1):35S–42S. http://www.ncbi.nlm.nih.gov/pubmed/23648197

43. Gómez LA, Escobar M, Peñuela O. Standardization of a Protocol for Obtaining Platelet Rich Plasma from blood Donors; a Tool for Tissue Regeneration Procedures. Clinical laboratory. 2015 [accessed 2015 Nov 18];61(8):973–80. http://www.ncbi.nlm.nih.gov/pubmed/26427141

44. Miller CH, Rice AS, Garrett K, Stein SF. Gender, race and diet affect platelet function tests in normal subjects, contributing to a high rate of abnormal results. British journal of haematology. 2014 [accessed 2015 Nov 18];165(6):842–53. http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4477706&tool=pmcentrez &rendertype=abstract

45. Kleiner G, Marcuzzi A, Zanin V, Monasta L, Zauli G. Cytokine levels in the serum

of healthy subjects. Mediators of inflammation. 2013 [accessed 2015 Nov 18];2013:434010.

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3606775&tool=pmcentrez &rendertype=abstract

46. Biancotto A, Feng X, Langweiler M, Young NS, McCoy JP. Effect of anticoagulants on multiplexed measurement of cytokine/chemokines in healthy subjects. Cytokine. 2012 [accessed 2015 Nov 18];60(2):438–46. http://www.sciencedirect.com/science/article/pii/S1043466612002153

47. Ashwood P, Krakowiak P, Hertz-Picciotto I, Hansen R, Pessah I, Van de Water J. Elevated plasma cytokines in autism spectrum disorders provide evidence of immune dysfunction and are associated with impaired behavioral outcome. Brain, behavior, and immunity. 2011 [accessed 2015 Sep 7];25(1):40–5. http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2991432&tool=pmcentrez &rendertype=abstract

48. Anitua E, Sánchez M, Orive G, Andía I. The potential impact of the preparation rich in growth factors (PRGF) in different medical fields. Biomaterials. 2007 [accessed 2015 Nov 18];28(31):4551–60. http://www.ncbi.nlm.nih.gov/pubmed/17659771

49. Castillo TN, Pouliot MA, Kim HJ, Dragoo JL. Comparison of growth factor and platelet concentration from commercial platelet-rich plasma separation systems. The American journal of sports medicine. 2011 [accessed 2015 Nov 18];39(2):266–71. http://www.ncbi.nlm.nih.gov/pubmed/21051428

50. Honda Y, Ding X, Mussano F, Wiberg A, Ho C-M, Nishimura I. Guiding the osteogenic fate of mouse and human mesenchymal stem cells through feedback system control. Scientific reports. 2013 [accessed 2015 Feb 24];3:3420. http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3851880&tool=pmcentrez &rendertype=abstract