

Impact of asymptomatic COVID-19 patients in global surgical practice during the COVID-19 pandemic

Editor

The rapid spread of COVID-19 has changed the global surgical care¹⁻³. Patients infected with COVID-19 may present without typical symptoms, and such asymptomatic patients may potentially trigger in-hospital outbreaks by transmitting the disease to health care providers and other hospitalized patients^{4,5}. Further, asymptomatic COVID-19 patients have worse postoperative outcomes with an unexpectedly

high morbidity and mortality, reaching 20.5 per cent deaths⁶. However, we do not have objective global data on this issue. In an attempt to clarify the current global surgical practice under the COVID-19 pandemic particularly focusing on the preoperative screening of asymptomatic COVID-19 patients, we conducted a cross-sectional online survey on surgical practice. The survey was administered to surgeons worldwide through international surgical societies, social media and personal contacts during 2 April to 8 April. The results were analyzed by country's cumulative deaths

number by 8 April, 2020 (high risk, >5000; intermediate risk, 100-5000; low risk, <100) (<https://covid19.who.int/>).

The survey was completed by a total of 936 centres in 71 countries, involving 5 high risk countries (330 centres), 20 intermediate risk countries (242 centres) and 46 low risk countries (364 centres). Table 1 summarizes the results of the survey. The majority (73.8 per cent) of the centers performed preoperative COVID-19 testing based on symptoms or suspicious radiologic findings. Universal COVID-19 testing for every patient was performed in only 16.6

Table 1 Preoperatively asymptomatic COVID-19 patients by the countries' risk group					
	Overall 936	Countries by risk group			P Value
		High risk 330	Int. risk 242	Low risk 364	
Testing policies before surgery. No. (%)^a					
Everyone	155 (16.6)	91 (27.6)	43 (17.8)	21 (5.8)	<.001
All oncologic patients	42 (4.5)	24 (7.3)	14 (5.8)	4 (1.1)	<.001
All emergency patients	73 (7.8)	41 (12.4)	22 (9.1)	10 (2.7)	<.001
Symptomatic or suspicious radiological features	691 (73.8)	214 (64.8)	183 (75.6)	294 (80.8)	.03
Experienced preoperatively asymptomatic patients tested positive after surgery. No. (%)					
Yes	257 (27.5)	170 (51.5)	66 (27.3)	21 (5.8)	<.001
No	679 (72.5)	160 (48.5)	176 (72.7)	343 (94.2)	
Experienced preoperatively asymptomatic patients tested positive after surgery with symptoms. No. (%)					
Yes	231 (24.7)	158 (47.9)	55 (22.7)	18 (4.9)	<.001
No	705 (75.3)	172 (52.1)	187 (77.3)	346 (95.1)	
Experienced preoperatively asymptomatic patients tested positive after surgery without symptoms. No. (%)					
Yes	157 (16.8)	108 (32.7)	39 (16.1)	10 (2.7)	<.001
No	779 (83.2)	222 (67.3)	203 (83.9)	354 (97.3)	
Preventive measures changed due to the asymptomatic patients? (2nd survey) No. (%)^{a,b}					
Yes, PPE use	40 (26.8)				
Yes, PPE availability	23 (15.4)				
Yes, preoperative patient testing	55 (36.9)				
Yes, staff testing	26 (17.4)				
Yes, staff rotation	16 (10.7)				
Yes, COVID-19 dedicated area	54 (36.2)				
Yes, stop elective surgeries	31 (20.8)				
No	27 (18.1)				
Did you observe a worse surgical outcome related to asymptomatic COVID-19 patient? (2nd survey) No. (%)^b					
Yes	46 (30.9)				
No/I don't know	103 (69.1)				

Int: intermediate. ^aPercentages do not add up to 100 because of the multiple choice question. ^bThe second survey respondents are the centers that answered 'yes' in the question 'Experienced preoperatively asymptomatic patients tested positive after surgery' ($n = 149$). The data are not analyzed by countries' risk group.

per cent of the centers, and the proportion was less than 30 per cent even in severely affected countries. Asymptomatic COVID-19 patients who tested positive postoperatively were reported from 27.5 per cent of the centres, with higher rates in higher risk countries (high risk, 51.5 per cent; intermediate risk, 27.3 per cent; low risk, 5.8 per cent; $p < 0.001$). To further clarify the clinical impact of asymptomatic COVID-19 patients, the second survey was sent to 218 centres that experienced such patients and had contact information, and 149 responded. The median number of asymptomatic COVID-19 patients testing positive after surgery per centre was 2 (range, 1 to 20). After experiencing the first asymptomatic COVID-19 patients, most centers (81.9 per cent) changed testing policies and/or preventive measures in surgical practice. Worse surgical outcomes in asymptomatic COVID-19 patients were reported from 30.9 per cent of the centres.

To our knowledge, this is one of the largest international surveys on COVID-19 in the field of surgery. Acknowledging the biases caused by unintended selection of centres, unequal number of centres per country and uneven geographical coverage, this large survey captured the global surgical practice under the COVID-19 pandemic and highlighted the threat of asymptomatic patients. Asymptomatic COVID-19 patients who were tested positive postoperatively may represent insufficient preoperative screening in the current surgical practice. The clinical impact of asymptomatic COVID-19 patients for surgeons is evident by the fact that the detection of such asymptomatic patients resulted in the change of testing policies and/or preventive measures in most of the centres. We strongly believe that in the coming phase of pandemic, during which many medical centres will resume elective surgeries, a call for action in surgical departments for more intensive screening programs is needed in global plans.

Acknowledgement

We would like to thank all the collaborators of “S-COVID Collaborative

Group” (listed separately) for their contribution to the study; the surgical societies (European Society of Surgical Oncology, Latin American Society of Surgical Oncology, Russian Society of Colorectal Surgeons and Società Italiana di Chirurgia Colo-Rettale) for distributing the survey; Dr. Krishn Khanna for English editing; and all the surgeons who translated the original questionnaire into 13 languages and shared the project in their countries: A. Frontali, J. Carvas, L. Sanchez-Guillen, C. Blajut, M. Sbah, E. Karbovnichaya, E. Samadov, N. Buchs, M. Ninkovic, T. Araki, G. Gallo, A. Sturiale. The study was registered with an analysis plan on ClinicalTrials.gov (NCT04344197).

V. Bellato^{1*}, T. Konishi^{2*},
G. Pellino^{3,4}, Y. An⁵, A. Piciocchi⁶,
B. Sensi¹, L. Siragusa¹, K. Khanna⁷,
B.M. Pirozzi¹, M. Franceschilli¹,
M. Campanelli¹, S. Efetov⁸, G.S. Sica¹
and on behalf of S-COVID
Collaborative Group

*Vittoria Bellato and Tsuyoshi Konishi contributed equally to this work as the co-first authors. Contributors of ‘S-COVID Collaborative Group’ are listed in Supporting Information.

¹Department of Surgery, Minimally Invasive Unit, Università degli Studi di Roma “Tor Vergata”, Rome, Italy,

²Department of Gastroenterological Surgery, Cancer Institute Hospital of the Japanese Foundation for Cancer Research, Tokyo, Japan, ³Department of Advanced Medical and Surgical Sciences, Università degli Studi della Campania “Luigi Vanvitelli”, Naples, Italy, ⁴Department of

Colorectal Surgery, Vall d’Hebron University Hospital, Barcelona, Spain,

⁵Department of General Surgery, Beijing Friendship Hospital, Capital Medical University, Beijing, China, ⁶GIMEMA Foundation, Rome, Italy, ⁷Department of Orthopaedic surgery, Rush university

medical center, Chicago, IL, and ⁸Department of Surgery, I.M. Sechenov First Moscow State Medical University, Moscow, Russia

DOI: 10.1002/bjs.11800

- 1 COVIDSurg Collaborative. Global guidance for surgical care during the COVID-19 pandemic. *Br J Surg* 2020; <https://doi.org/10.1002/bjs.11646> [Epub ahead of print].
- 2 Soreide K, Hallet J, Matthews JB, Schnitzbauer AA, Line PD, Lai PBS *et al*. Immediate and long-term impact of the COVID-19 pandemic on delivery of surgical services. *Br J Surg* 2020; <https://doi.org/10.1002/bjs.11670> [Epub ahead of print].
- 3 Spinelli A, Pellino G. COVID-19 pandemic: perspectives on an unfolding crisis. *Br J Surg* 2020; <https://doi.org/10.1002/bjs.11627> [Epub ahead of print].
- 4 Di Marzo F, Sartelli M, Cennamo R, Toccafondi G, Coccolini F, La Torre G, *et al*. Recommendations for general surgery activities in a pandemic scenario (SARS-CoV-2). *Br J Surg* 2020; <https://doi.org/10.1002/bjs.11652> [Epub ahead of print].
- 5 Mowbray NG, Ansell J, Horwood J, Cornish J, Rizkallah P, Parker A *et al*. Safe management of surgical smoke in the age of COVID-19. *Br J Surg* 2020; <https://doi.org/10.1002/bjs.11679> [Epub ahead of print].
- 6 Lei S, Jiang F, Su W, Chen C, Chen J, Mei W *et al*. Clinical characteristics and outcomes of patients undergoing surgeries during the incubation period of COVID-19 infection. *EClinicalMedicine* 2020; <https://doi.org/10.1016/j.eclinm.2020.100331b> [Epub ahead of print].

Supporting information

Additional supporting information can be found online in the Supporting Information section at the end of the article.