

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Rhythm in dyadic interactions

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1786997> since 2021-04-22T08:49:46Z

Published version:

DOI:10.1098/rstb.2020-0337

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

RHYTHM IN DYADIC INTERACTIONS

Koen de Reus^{1,2*#}, Masayo Soma^{3*}, Marianna Anichini^{4,5}, Marco Gamba⁶, Marianne de Heer Kloots⁷,
Miriam Lense⁸, Julia Hyland Bruno⁹, Laurel Trainor¹⁰, Andrea Ravignani¹

¹Comparative Bioacoustics Group, Max Planck Institute for Psycholinguistics, Nijmegen, The Netherlands

²Artificial Intelligence Lab, Vrije Universiteit Brussel, Brussels, Belgium

³Department of Biology, Hokkaido University, Sapporo, Japan

⁴BRAIN/Neurosciences and Cognitive Sciences, Hanse-Wissenschaftskolleg (Institute for Advanced Study), Delmenhorst, Germany

⁵Division Animal Physiology & Behaviour, Department of Neuroscience, School of Medicine and Health Sciences, University of Oldenburg, Oldenburg, Germany

⁶Department of Life Sciences and Systems Biology, University of Turin, Turin, Italy

⁷Institute for Advanced Study, University of Amsterdam, Amsterdam, The Netherlands

⁸Department of Otolaryngology-Head and Neck Surgery, Vanderbilt University Medical Center, Nashville, TN, USA

⁹Center for Science and Society, Columbia University, New York, NY, USA

¹⁰Department of Psychology, Neuroscience and Behaviour, McMaster University, Hamilton, Ontario, Canada

* These authors share first authorship

Corresponding authors: kdereus95@gmail.com, andrea.ravignani@mpi.nl

Abstract

This review paper discusses *rhythmic* interactions and distinguishes them from *non-rhythmic* interactions. We report on communicative behaviours in social and sexual contexts, as found in dyads of humans, non-human primates, non-primate mammals, birds, anurans and insects. We discuss observed instances of rhythm in dyadic interactions, identify knowledge gaps and propose suggestions for future research. We find that most studies on rhythmicity in interactive signals mainly focus on one modality (acoustic or visual) and suggest more work should be performed on multimodal signals. Although the social functions of interactive rhythms have been fairly well described, developmental research on rhythms used to regulate social interactions is still lacking. Future work should also focus on identifying the exact timing mechanisms involved. Rhythmic signalling behaviours are widespread and critical in regulating social interactions across taxa, but many questions remain unexplored. A multidisciplinary, comparative cross-species approach may help provide answers.

Key words: animal communication, dyadic interaction, rhythm, signal timing, synchrony, multimodality

Introduction

Animals rely on effective signal transfer for communication with conspecifics. The sender must produce a clear signal that can be readily detected by the receiver [1]. A variety of sensory systems evolved to accommodate the production and reception of signals, and over time, these systems were fine-tuned by selective evolutionary pressures leading to remarkable species-specific adaptations (e.g. 3-ossicle middle ear in mammals adapted for acoustic transmission of high frequency signals [2]). Signals can be produced in several modalities, including acoustic, visual, chemical, tactile, etc.; each modality has its own (psycho)physical limitations that impose constraints on communication. For example, acoustic signals are effective for long-range information transfer but are affected by signal attenuation and degradation, especially at higher frequencies [3]. In the visual modality (e.g. signed languages), receivers are required to be in close contact with the signaller to accurately discriminate the signals because obstacles can block them. Regardless of the modality in which signalling occurs, *interactive* signalling will henceforth refer to a communicative exchange during which signals are transferred, back and forth, over a short time scale.

Timing is key in interactive signalling behaviour [4-6]. Individuals take turns in human conversation and entrain to the rhythms of other players while making music [7, 8]. Individually timed behaviour in groups can lead to impressive collective phenomena such as synchronous movements to musical rhythms at concerts [9], cricket choruses [10] and the bioluminescent flashing of fireflies [11]. Unfortunately, natural interactions involving many agents are often difficult to study under controlled conditions. Hence, for practical reasons, most studies on interactive rhythms minimise the number of individuals by focusing on *dyadic* interactions. Some species can achieve a high degree of temporal coordination, and even synchronisation, during conspecific interactions [10, 12-14]. Therefore, we ask: how does rhythm shape dyadic interactions and how do dyadic interactions shape rhythm?

Here, we review stable interactions in dyads across species, modalities and contexts. This review covers humans and nonhuman animals - namely insects, anurans, birds, non-human primates and non-primate mammals - interacting mainly in the acoustic and visual modalities, and in social and sexual contexts. We (i) begin by proposing widely applicable definitions for relevant concepts (see Table 1); (ii) highlight rhythmic dyadic interactions in each group; and (iii) discuss the existing behaviours in light of Tinbergen's 4 questions while proposing suggestions for future research.

We first need to distinguish between what constitutes a rhythmic versus non-rhythmic interaction. A *rhythmic interaction* is one where two conspecifics adjust their individual timing to each other to create temporal regularities that facilitate the interaction. The rhythmic structure that emerges can be measured and quantified along sequential and temporal dimensions [15]. For instance, in piano duets and dyadic finger tapping experiments, a statistical association, either synchronous and/or asynchronous, can be established between the temporal intervals of the two players [16, 17]. Based on the concepts described in Table 1, any interaction that requires timing adjustment between two conspecifics such as dyadic synchrony and turn-taking is considered rhythmic in this paper. A *non-rhythmic interaction* is one where there is no timing adjustment between the members of the dyad. Despite this, individual rhythms can still play an important role in regulating the interaction. For example, male northern elephant seals (*Mirounga angustirostris*) can recognise the rhythmic structure of their rivals' vocalisations; depending on the individual's status within the colony, males will either ignore or move away from the vocalising male [18]. In the absence of a temporal relationship between the signals of two conspecifics, both alternation and antiphony are considered non-rhythmic. In the supplement, readers can find examples of both interactions for all animal clades discussed in this review (Table S1) and a discussion on non-rhythmic interactions.

Table 1. Definitions of terms relating to dyadic interactions.

Term	Definition
<i>Signals</i>	‘Traits that (1) change another organism’s behaviour while benefitting the sender, that (2) are evolved for this function, and that (3) have their effects through the evolved response of the receiver’ ([19], p.1011)
<i>Rhythm</i>	‘Pattern of time intervals between the onset of events’ ([20], p.165)
<i>Rhythmic interaction</i>	An interaction where two conspecifics adjust their timing behaviour to each other to create temporal regularities that facilitate the interaction
<i>Entrainment</i>	‘Spatiotemporal coordination resulting from rhythmic responsiveness to a perceived rhythmic signal’ ([21], p.5)
<i>Duets</i>	‘Joint [...] displays where two [partners] coordinate their [signals] with a degree of temporal precision’ [22]. Traditionally, duets occur in the acoustic modality, but we believe the use of the term should be extended to all other communicative modalities.
<i>Synchrony</i>	‘Precise coincidence of events in time’ ([23], p.158)
<i>Turn-taking</i>	‘Orderly exchange of purely communicative signals or behaviours (e.g. peek-a-boo games in humans) between individuals characterized by principles for the coordination of turn transfer, which result in observable temporal regularities’ ([24], p.2). We subscribe to the idea that turn-taking is mostly rhythmic in its temporal dimension [6].
<i>Alternation</i>	‘Where the regularly repeating signals of two [...] individuals are broadcast such that they do not occur at the same time’ ([25], p.4)
<i>Antiphony</i>	‘When [...] two animals transmit sounds among themselves in response to preceding signals’ ([26], p.155)

Humans

Humans are a highly social species and perform many types of duets; here we focus on two: speech and music. In the rhythmic domain, speech generally involves turn-taking, whereas music is typically performed in synchrony, although people can chant speech together and call-and-response is used in music. Such behaviours involve complex social interactions and serve to communicate information, express emotion, and socially bond [27, 28]. These acoustic behaviours are performed with others in complex behavioural patterns involving cues including gesture, touch, body sway and dance, but also facial expressions and gaze direction [29-32].

Infancy and childhood

Human infants cannot survive without caregivers for a considerable period, and their early development and learning is done in a social context, often primarily with the mother [27]. Within this social context, movement behaviours [33], vocal behaviours [34], gaze [35], autonomic functioning [36] and hormone expression [37] are coordinated in time between infants and their caregivers. This early temporal coordination may enable infants to (i) regulate physiological and behavioural processes for survival [37], (ii) develop self-regulation [38], (iii) differentiate the emergence of their self from others [27], (iv) begin building social and empathetic relations with others [39], and (v) learn speech and music through continual reciprocal adjustments [40].

Caregivers provide rhythmic input to their infants in the forms of singing, patting, and rocking, which helps infants to regulate their states [41]. The periodicities found in rhythms of infant-directed singing provide a context for bidirectional entrainment between infants and caregivers, which can be seen in the coordination of autonomic and brain responses between mothers and infants [35, 42, 43]. Synchronous movement to music during infancy also has important social consequences. 14-month-old infants who are bounced in synchrony to music with an experimenter are much more likely to subsequently help that experimenter compared to infants who are bounced out of sync [39, 44], suggesting that the origins of empathy and friendship may be found in coordinated rhythmic behaviours.

While the bidirectionality of interactions begins in infancy, caregivers initially have a broader role in structuring and scaffolding the interaction [45, 46]. As children develop their advancing language, social and cognitive skills support and expand their active participation in complex, rhythmic verbal and nonverbal interactions [45]. In the verbal domain, children become increasingly proficient at dialog and conversation [45, 47]. Active turn-taking during conversation, which reflects partners' attunement to each other, is associated with children's language and social skill development [48, 49]. Children who experience more appropriate turn-taking styles during conversations with their parents are more liked by their peers [50], and also exhibit more appropriate turn-taking with their peers even when the peers are novel acquaintances [51]. In the nonverbal domain, unfamiliar child peers who actively engage in coordinated rhythmic movement activities together show greater cooperation [52] and perceive that they are closer and more similar to each other, potentially due to increased intentional communication [53].

Adulthood

In adults, entrainment during dyadic conversation is observed through temporal adaptations within each individual's speech pattern that occur over multiple timescales [54]. The precise timing involved in conversational entrainment suggests the presence of underlying rhythmic processes that allow for accurate timing predictions [55]. During a conversation, dyads converge their individual speech rhythms as measured via speech rate, prosody, and respiratory movement [56-58], as well as turn-taking timing (e.g., by minimizing silent gaps and avoiding overlap) [55, 59, 60]. The degree of entrainment in conversation is impacted by the interaction context (e.g. friendly versus unfriendly [54]). Rhythms are crucial in these interactions as their temporal regularity enables prediction and sampling of environmental stimuli [61].

Coordination in both musical and conversational duets relies on visual and auditory cues [62, 63] but, in contrast to speech which largely involves turn taking, musical interactions typically involve synchrony. While the two interacting musicians may sing or play different pitches, their outputs must fit together both rhythmically and harmonically to create a single meaningful joint performance. The

high real-time temporal demands of duet music-making requires continual adjustments, anticipation and prediction (if an individual reacts to rather than anticipates their partner's output, they will be late and not in sync with them) [8]. The continual adaptation of two people tapping together can be seen in 'lag-1 correlations', whereby on a given tap, the individual who is slightly ahead will slow down on their next tap and the individual who is slightly behind will speed up on their next tap [64]. Studies of string quartets also show mutual adjustments of timing, with some quartets being more leader-driven and others more egalitarian [65-67]. However, asynchronies increase when one musician in a duet is replaced with a recording [68] and when tempo preferences of two musicians are divergent [69], highlighting the importance of bidirectional coordination. Musicians convey their upcoming intentions through body sway, similarly to how people use hand gestures when they speak. Findings from string quartets show that the body sway of one musician predicts the upcoming body sway of another musician [70]. In short, the better overall communication is among group members, the higher rated is the quality of their performance [29]. Music affords an ideal context in which people can synchronise their movements, and when a person experiences even a short period of synchronisation with another person, it has social consequences, leading to increased liking, cooperation and trust [71, 72]. These social consequences may explain why music-making in groups is universal across human societies and likely an evolutionary adaptation [73].

Nonhuman primates

Dyadic interactions among primates are essential to determine hierarchies within stable groups [74], and establish alliances and partnerships [75]. In several monkey and ape species, where vocal communication follows precise temporal and social rules, strong social bonds shape interactive signalling patterns [76, 77]. For instance, bonobos respond preferentially to conspecifics with whom they have close bonds, and take turns to avoid overlap when vocalising with those [78]. Unfortunately, rhythmic features have generally been understudied in primate communication, with few exceptions.

Primates may form long-lasting, socially monogamous pairs with opposite-sex conspecifics. Tarsiidae, Callicebinae, Hylobatidae and Indriidae include the main species of pair living primates producing duets [79, 80]. These pair-living primates face the need to defend a reduced home range [81] and males within these pairs struggle to protect mating exclusivity [82]. In both gibbons and indris, duets serve to inform neighbouring groups about the occupation of a territory and to defend it during group encounters [83]. In indris, duetting mediates group cohesiveness (see panel C of Figure 1, [84]), while in gibbons it can inform about the presence of particular predators [85]. Duetting in siamang pairs (*Hylobates syndactylus*) is associated with a rhythmic swinging from branch to branch, which helps maintain entrained vocal displays between males and females [86]. Depending on the species, simultaneous singing or avoiding song overlap advertises the strength of a bond and may predict both behavioural coordination and grooming rates [87].




Figure 1. Rhythm and social interactions in indris. Panel A shows a spectrogram of a vocal interaction between two males from the same family group that compete for food; the adult male emits a low-pitched grunt [88] followed by a kiss and a wheeze from the younger male [89]. Panel C shows a spectrogram of part of an unusual solo song by a male indri. Panel E shows a spectrogram of part of a duet by a pair of indris ('cohesion song'). After dispersing within a territory, they emit a particular song type to regroup in a particular location [84]. Red shading denotes the fundamental frequency of the female's calls, and blue shading denotes the fundamental frequency of the male's units. Panels B, D and F are sketches of the animals represented in the adjacent spectrograms.

Non-primate mammals

Rhythmic interactions in other mammalian species allow individuals to identify conspecifics and maintain affiliations with other group members. We summarise findings of interactive exchanges in animal groups inhabiting different physical and socio-ecological environments including cetaceans, pinnipeds, bats, rodents, elephants, antelopes, and meerkats.

Cetacean vocal interactions span alternation and synchrony. Bottlenose dolphins exchange signature whistles for individual recognition and group cohesion [90, 91]. These exchanges are closely coordinated in time such that acoustic overlap is minimal (silent gaps < 1 s) [92]. Male bottlenose dolphins also engage in coordinated cooperative interactions when coercing females by synchronising their threat vocalisations and matching tempos [93].

In pinnipeds, mother-pup recognition is a common dyadic interaction, where mothers and pups use vocal cues to identify and localise each other in the breeding colony, though species differ in whether mother, pup or both individuals vocalise [94, 95]. Harbour seals may also adjust their call timing and call asynchronously relative to conspecific vocalisations [96]. Within a breeding colony, this turn-taking strategy may avoid acoustic overlap and make individual pup calls more conspicuous.

Several bat species engage in antiphonal exchanges of stereotyped calls, also for mother-pup recognition [97-99]. Moreover, adult white-winged vampire bats (*Diademus youngi*) respond to contact

calls in a duet-like fashion by temporally coordinating their reply (silent gaps < 500 ms), potentially to monitor the spatial positions of conspecifics [100].

Few studies reported on rhythmicity in rodent dyadic communication. In Alston's singing mice (*Scotinomys teguina*), depending on the social context, males adjust their signal timing to sing in turns and avoid acoustic overlap (silent gaps around 500 ms) [101]. Naked mole rats (*Heterocephalus glaber*) produce soft chirps antiphonally between two or more individuals to identify their social status and maintain affiliations (silent gaps < 400 ms) [102]. Middle East blind mole rats (*Spalax ehrenbergi*) communicate using vibratory signals by engaging in alternating head-drumming duets (silent gaps < 2 s) [103].

Data on interactive temporal coordination in dyads is limited for remaining mammal species. Female African elephants (*Loxodonta africana*) use antiphonal rumbling sequences between group members to maintain social distance [104]. An African antelope, the klipspringer (*Oreotragus oreotragus*), engages in alarm call duetting and calls are produced in alternation, with female calls closely following those of males [105]. Finally, meerkats (*Suricata suricatta*) avoid overlapping conspecifics in low-conflict group sunning calls by vocalising in turns [106]. Here, group turn-taking is an outcome of dyadic interactions between group members and mechanistically relies on two alternating processes: call inhibition and call excitation.

Birds

Duets, which require a certain degree of temporal coordination (see Figure 1E), are also widespread across birds, especially in mating contexts. Vocal or dance duets occur in 18-20% of all avian species [107], which accounts for mostly song-duetting species among songbirds and dance-duetting species among non-songbirds (but see [108, 109]). Avian duets are generally performed by paired partners or prospective mates, and serve different functions including mate-guarding, joint resource defence, and/or mutual courtship [110, 111].

Some barbet species (Capitonidae) [112] and the magpie lark (*Grallina cyanoleuca*) [113] perform multimodal duets, where the two birds simultaneously coordinate vocalisations and body movements. For singing, species-specific duet rules are well documented and show interspecific variation, even among closely related species. For example, in Thryothorus wrens, some species produce perfect antiphonal duets without overlap, while others sing in synchrony [114]. In avian song duetting, each of the sexes often produces a particular phrase at the precise onset or offset of its partner's singing, making it a perfectly timed collective display [115]. In contrast, rhythmicity in dance duets is still relatively unexplored. For example, in the red-crowned crane (*Grus japonensis*), males and females exhibit sequences of multiple dance elements for their joint display [116], but it is not known whether the paired cranes precisely time and synchronise their dancing with each other.

The degree to which two individuals coordinate their signalling during a duet varies even among dyads of the same species. Although temporal coordination can greatly influence reproductive fitness as better-coordinated signals are more effective for securing mates or reproductive resources [117], not every pair will be able to achieve the same degree of coordination. In fact, high level coordination can only be achieved by pairs with longer partnerships [118]. Indeed, newly established pairs of canebrake wrens (*Cantorchilus zeledoni*) improve the coordination of their duets over time, suggesting that learning enables fine duetting [119].

Multimodal signal coordination in dyadic interactions is not restricted to species known for their duetting. The zebra finch has been intensively studied and disproportionately so relative to other songbirds, but lacks duetting as only males sing [120]. Despite this, mated pairs display tight temporal coordination in both visual and auditory modalities. Zebra finches form strong dyads with a lifelong mate [118] where coordinated behaviours serve as an honest signal of the pair's coalition quality. Indeed, the strength of the bond in zebra finch pairs predicts the degree of coordination of movements [121] and call exchanges [122], which mechanistically rely on predictive timing [123]. Pairs use such vocal exchanges to maintain a joint behavioural state [124] and coordinate shared parenting duties [125].

Anurans

Dyadic interactions in anurans occur in mating contexts, where same-sex duets serve to compete for access to mates and opposite-sex duets serve to attract potential mates [126]. Anurans show a precedence effect, where calls of leading individuals are preferred when two identical calls are presented in close succession [25], but exceptions exist [127]. Moreover, calls that alternate in precise phase relationships are preferred by females of some species (e.g. the midwife toad *Alytes obstetricans*) [127]. Precise patterns of signal timing largely influence mate choice and are thus under strong sexual selection in anurans [126].

Males typically form large groups and produce loud alternating advertisement calls to attract females for mating [128]. Instances of non-random call timing in males were studied in larger groups, but also in dyads [129]. In duetting male pairs of the Neotropical toad (*Rhinella ocellata*), males call in alternation and avoid overlap with each other [130]. Moreover, the call delay of the responding male varies depending on the call duration of the male that initiated the interaction. This alternation pattern shows a high degree of temporal coordination within bouts. Call timing adjustment between two males of the European tree frog (*Hyla arborea*) can result in near-perfect antiphonal calling with mean relative phase angles distributed around 180° [131]. The onset of vocalisations can even be adjusted based on the distance separating individuals, as evidenced in bullfrogs (*Rana catesbeiana*) [132]. Bullfrog males call more frequently following calls of distant neighbours than of those nearby ones; nearby neighbours may inhibit the vocal response of the focal individual. In sum, males try to avoid signal overlap in an effort to increase their conspicuousness by paying selective attention to their close neighbours and vary their call timing accordingly [131]. However, precise alternation of vocalisations is not always the norm. A few species signal using overlapped calls. For example, males of the American toad (*Bufo americanus*) signal in synchrony or near-perfect synchrony [12]. This timing strategy may help reduce predation risk [133] and/or increase the chorus' audibility in order to attract more females to the breeding area [129].

Even though they are rarely described, duets between males and females do occur at the beginning of courtship. In the South African clawed frog (*Xenopus laevis*), receptive females produce a vocalisation named 'call rapping', which is composed of a rapid series of loud clicks, in response to the male's advertising call [134]. The female call spurs the male to move towards the sound source and produce an intense bout of calling within 1-2 s of the female's signal. A possible explanation for the evolution of female courtship vocalisations in anurans could be found by looking at similarities in terms of breeding biology [134].

Insects

Observations of non-random timing suggest that insects can adjust the onset of their signals. Temporal relationships of interactive signals are described for the acoustic modality in bush-crickets,

grasshoppers, mosquitos and flies, for the vibratory modality in planthoppers, leafhoppers, stoneflies, stink bugs and lacewings, and for the bioluminescent modality in fireflies [10, 135]. Generally, males initiate insect duets and females respond with a fixed latency relative to the male signal [135]. The temporal pattern of the duet and the time window of the female reply are species-specific [136], hence allowing species recognition [135]. The duration and complexity of the male signal affects the reply latency of the female [137]. The longer and more complex the male signal is, the more time is needed to process the information encoded in the call and assess the male's quality. However, the latency of the reply may also depend on the female's readiness to mate [137].

In the leafhopper *Aphrodes makarovi*, sexual communication and mate recognition are mediated via species-specific and sex-specific vibrational signals. During duetting, the females' response overlaps with the last portion of the male's signal [138]. Males can only evaluate the non-overlapped part of the female call, and consequently adjust their signal period length to the duration of the females' reply to obtain a longer non-overlapped segment of the females' call [139]. While competing for access to females in the presence of rivals, male broad-winged bush katydids (*Scudderia pistillata*) produce an acoustic tick that mimics the female's tick signal and response timing [140]. This indicates to the female when the call has concluded and also confounds eavesdropping males by making it difficult for them to accurately move towards the true female's sound source [137]. Not only does this reduce the risk of competing males interrupting the established duet, but it also increases the coordination of the duet itself.

Some insects perform multimodal duets. For temporal coordination to occur in duets of the fruit fly *Drosophila virilis*, females need to detect the male's sound cues and be in close physical proximity [141]. During courtship, males tap the female's abdomen and lick the genitalia in a precisely timed manner to coordinate the duet. Females may choose to mate only with males that provide multiple timing cues during the duet. In the katydid *Onomarchus unnotatus*, male and female signals alternate in a clear phase relationship. Males produce a low-pitched call which receives a female vibrational signal in response [142] (see Figure S1). This unique multimodal duet may have evolved in response to predation; silent flying females were predated at a higher rate than calling males. With females using vibrational signals, the roles are now reversed: males must search for females by moving towards the vibration source and face increased risk of predation by bats.

Same-sex dyadic interactions in insects occur in highly competitive environments, hence the resulting temporal patterns are an outcome of competition between signalling males. In species where females prefer to mate with males leading the call sequence, males alternate calls to avoid losing their leader role [10, 25]. In tarbush grasshoppers (*Ligurotettix planum*), males engage in acoustic 'fights' to defend or conquer a mating territory. The male that cannot match the signalling rate and length of the signal of its opponent gives up the fight [143]. The aggressive calls emitted by competing males thus represent an honest signal of male fitness.

Table 2. Summary table of rhythmic interactive signalling in dyads across taxa.

Composition of dyad	Interaction context	Taxonomic group			
		<i>Mammals (including humans)</i>	<i>Birds</i>	<i>Anurans</i>	<i>Insects</i>
Kin	Parent-offspring	A, V	A, V	-	-
	Social group member	A, V	A, V	-	-
Non-kin	Social group member	A, V	A, V	-	-
	Male-female	A, V	A, V	A	A, V, T
	Male-male	A, V	A, V	A	A, V, T

Note. Although many signalling modalities have been proposed, the distinctions between them are often murky, hence, in the spirit of simplicity and to avoid confusion, we only include the main ones: auditory (A), visual (V) and tactile (T). Bioluminescent signals enhance visual perception [144] and are grouped under the visual modality. Vibratory signals are substrate-borne signals that are perceived using mechanoreceptors [145], similar to touch, hence they are considered part of the tactile modality. Cases where no empirical evidence was found are represented by (-).

Ultimate causes of dyadic interactions: function and phylogeny

Dyads can consist of mates, siblings, parent and offspring, or any two individuals from the same group. Dyadic interactions serve a variety of functions including social bonding, sharing emotion, establishing hierarchies and partnerships, mate-guarding, courtship, and joint resource defence, etc.

[14, 27, 28, 40, 74, 75, 110, 111]. Studying dyads means that experiments possibly neglect relevant group dynamics, especially in highly social species. However, it also enables researchers to study *semi-natural* behavioural interactions under more controlled experimental conditions.

In this review, we identified two settings in which dyadic behavioural interactions can occur: social and sexual. Helping behaviour, learning behaviour, and parent-offspring recognition were classified as occurring in social settings. Dyadic interactions in sexual settings mostly pertain to opposite-sex conspecifics that are potential mates or already form an established pair, but they can also involve same-sex conspecifics during intrasexual competition occurring during mate search and attraction [127]. Rhythmic behaviours have been reported for both contexts in mammals and birds, but in anurans and insects, they have only been reported for sexual settings.

Animals produce many different types of rhythms; hence they could entrain to conspecific rhythms in ways that can be difficult to observe. For instance, individual rhythms do not need to occur in the same modality to interact rhythmically (i.e. coordination of an infant's brain responses to its mother's singing [42]). Moreover, the dynamics of the interaction can be both unidirectional and bidirectional [29, 65-67]. In music, bidirectional interaction improves coordination while unidirectional interactions create larger asynchronies [68]. Does this finding in humans carry over to interactions in other species? One could test whether the degree of coordination is affected by unidirectionality in songbird species in which one of the sexes does not sing (e.g. the Java sparrow or zebra finch). It seems that several species overcome such asynchrony by signalling in more than one modality to achieve higher levels of temporal coordination.

Even though multimodal signals are widespread across numerous taxa [62, 86, 112, 113, 141], their modalities are often studied separately [146]. Rhythmic interactive signalling has been reported mainly for the acoustic and visual modalities (see Table 2). However, birds can communicate using vibratory signals [147], but avian rhythmic interactions in the tactile modality have not been described. Similarly, anurans have visual signals [148], but rhythmicity in this modality has not been described during conspecific interactions. Future work on interactive rhythms should investigate the unexplored modalities in these animal groups. In addition, analysing rhythmic patterns of multimodal signals during interactive communication will provide more insight into the functions of signal timing.

Proximate causes of dyadic interactions: ontogeny and mechanisms

A wide repertoire of signals is needed to become an active, grown-up participant in dyadic interactions and precise timing of signals requires extensive practice. The social environment during development plays a crucial role in shaping adult signals. Human children benefit from structured bidirectional interactions scaffolded by their caregivers [45, 46]. Similarly, young birds learn to participate in adult interactions. Male zebra finches learn their courtship song from their father, but can also receive feedback from other individuals within their social environment [149]. It has even been argued that learning of species-specific rhythms starts before birth, with possible long-term effects on vocal and social development [150]. The social context experienced during development modulates male signal features and female mate preferences in insects which will constrain the interaction [151]. Future work on dyadic interactions should accurately describe the social environment in which animals have been raised as it may influence both rhythm perception and production. Moreover, longitudinal studies should aim to follow developing individuals to investigate the effect of development on rhythmic interactions.

Coordinated signal exchanges in dyadic interactions show reply latencies ranging from the order of milliseconds to seconds [92, 100, 101]. The precise temporal coordination of behaviours must be regulated by the presence of underlying rhythmic processes, and many timing mechanisms proposed in the literature are based on the concept of coupled oscillators [55]. Human behaviours such as conversational entrainment [54] or the synchronous playing of music [8] rely on mechanisms that allow us to make accurate timing predictions about the onset of upcoming signals. Pinnipeds and birds also show some affinity with predictive timing. Similar to conversational turn-taking, seal pups may adapt the timing of their calls [96], and some songbirds attempt to simultaneously synchronise their vocalisations and body movements [112, 121, 122]. Insects are also capable of synchronous displays in larger groups, but the neural processes that regulate signal timing are different from mammals and birds. Instead of being able to predict the exact onsets of events, signalling in anurans and insects is reactive [152]. Similar observable and interactive behaviours can thus be produced using different mechanisms, but we do not know whether these mechanisms are learned and whether they change across modalities.

Conclusions

Rhythmic behaviours are widespread among animal clades, and crucial in regulating social interactions in dyads and larger groups. Unfortunately, in the temporal domain of animal communication, many species remain unexplored and several questions unanswered. Within species and across contexts, which features of signal timing are stable, and which are constrained and to what degree? How are timing mechanisms learned and do they change from one modality to the next? Lines of investigation that integrate ecological and neuroethological perspectives have begun to resituate rhythmic behaviours within animal communication systems (see [153]), but they have not been described in this paper. Such a multidisciplinary approach would allow researchers to design species-specific experiments to infer how rhythm functions during interactive signalling [154]. We thus strongly encourage future studies on interactive rhythmic behaviours in the hope of ultimately developing an integrative cross-species framework [4]. The comparative method could then provide crucial insights into the evolution and adaptive functions of interactive rhythmic behaviour across taxa [5, 6].

Data accessibility

This article has no additional data.

Competing interests

The authors declare no competing or financial interests.

Author contributions

K.R. conceived the manuscript. All authors contributed in writing and revising of the manuscript, and gave final approval for publication.

Funding

This work was supported by the Research Foundation Flanders (FWO) [G034720N] to Bart de Boer.; JSPS Grant-in-Aid for Young Scientist [16H06177] and for Scientific Research [20K06809] to M.S.; Junior Fellowship of the Hanse-Wissenschaftskolleg (Institute for Advanced Study) to M.A.; NIMH and NCCIH [R61MH123029], and NIDCD [R21DC016710] to M.L. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

Acknowledgments

We thank the organisers of the Lorentz workshop on Synchrony and Rhythmic Interaction: From Neurons to Ecology, the Lorentz Center, and the NIAS, ILLC and ANU.

Reference list

- [1] Dusenbery, D. B. 1992 *Sensory ecology: how organisms acquire and respond to information*. New York, WH Freeman.
- [2] Tucker, A. S. 2017 Major evolutionary transitions and innovations: the tympanic middle ear. *Philosophical Transactions of the Royal Society B: Biological Sciences* **372**, 20150483. (DOI:10.1098/rstb.2015.0483).
- [3] Wiley, R. H. & Richards, D. G. 1978 Physical constraints on acoustic communication in the atmosphere: Implications for the evolution of animal vocalizations. *Behav. Ecol. Sociobiol.* **3**, 69-94. (DOI:10.1007/BF00300047).
- [4] Anichini, M., de Heer Kloots, M. & Ravignani, A. 2020 Interactive rhythms in the wild, in the brain, and in silico. *Canadian journal of experimental psychology/Revue canadienne de psychologie experimentale* **74**, 170-175. (DOI:10.1037/cep0000224).
- [5] Ravignani, A., Bowling, D. & Fitch, W. T. 2014 Chorusing, synchrony, and the evolutionary functions of rhythm. *Front. Psychol.* **5**, 1118-1118. (DOI:10.3389/fpsyg.2014.01118).
- [6] Ravignani, A., Verga, L. & Greenfield, M. D. 2019 Interactive rhythms across species: the evolutionary biology of animal chorusing and turn-taking. *Ann. N. Y. Acad. Sci.* **1453**, 12-21. (DOI:10.1111/nyas.14230).
- [7] Nguyen, T., Schleihauf, H., Kayhan, E., Matthes, D., Vrtička, P. & Hoehl, S. 2020 Neural synchrony in mother–child conversation: Exploring the role of conversation patterns. *Soc. Cogn. Affect. Neurosci.* **2020**, 1-10. (DOI:10.1093/scan/nsaa079).
- [8] Keller, P. E., Novembre, G. & Hove, M. J. 2014 Rhythm in joint action: Psychological and neurophysiological mechanisms for real-time interpersonal coordination. pp. 20130394, Royal Society of London.
- [9] Trehub, S. E., Becker, J. & Morley, I. 2015 Cross-cultural perspectives on music and musicality. *Philosophical Transactions of the Royal Society B: Biological Sciences* **370**, 20140096. (DOI:10.1098/rstb.2014.0096).
- [10] Greenfield, M. D. 1994 Cooperation and conflict in the evolution of signal interactions. *Annu. Rev. Ecol. Syst.* **25**, 97-126. (DOI:10.1146/annurev.es.25.110194.000525).
- [11] Buck, J. 1988 Synchronous rhythmic flashing of fireflies. II. *Q. Rev. Biol.* **63**, 265-289. (DOI:10.1086/415929).
- [12] Grafe, T. U. 1999 A function of synchronous chorusing and a novel female preference shift in an anuran. *Proc. R. Soc. Lond. B. Biol. Sci.* **266**, 2331-2336. (DOI:10.1098/rspb.1999.0927).
- [13] Backwell, P., Jennions, M., Passmore, N. & Christy, J. 1998 Synchronized courtship in fiddler crabs. *Nature* **391**, 31-32. (DOI:10.1038/34076).
- [14] Hall, M. L. 2009 A Review of Vocal Duetting in Birds. *Adv Study Behav* **40**, 67-121. (DOI:10.1016/S0065-3454(09)40003-2).
- [15] Ravignani, A. & Norton, P. 2017 Measuring rhythmic complexity: A primer to quantify and compare temporal structure in speech, movement, and animal vocalizations. *Journal of Language Evolution* **2**, 4-19. (DOI:10.1093/jole/lzx002).
- [16] Goebel, W. & Palmer, C. 2009 Synchronization of timing and motion among performing musicians. *Music Perception* **26**, 427-438. (DOI:10.1525/mp.2009.26.5.427).
- [17] Loehr, J. D. & Palmer, C. 2011 Temporal coordination between performing musicians. *Q. J. Exp. Psychol.* **64**, 2153-2167. (DOI:10.1080/17470218.2011.603427).
- [18] Mathevon, N., Casey, C., Reichmuth, C. & Charrier, I. 2017 Northern Elephant Seals Memorize the Rhythm and Timbre of Their Rivals' Voices. *Curr. Biol.* **27**, 2352-2356.e2. (DOI:10.1016/j.cub.2017.06.035).
- [19] Stegmann, U. E. 2005 John Maynard Smith's notion of animal signals. *Biology and Philosophy* **20**, 1011-1025. (DOI:10.1007/s10539-005-9020-8).
- [20] van den Bosch der Nederlanden, C. M., Taylor, J. E. T. & Grahn, J. A. 2019 Neural basis of rhythm perception. (eds. M. Thaut & D. Hodges), pp. 165-186. Oxford, Oxford University Press.

- [21] Phillips-Silver, J., Aktipis, C. A. & Bryant, G. A. 2010 The ecology of entrainment: Foundations of coordinated rhythmic movement. *Music Perception* **28**, 3-14. (DOI:10.1525/mp.2010.28.1.3).
- [22] Farabaugh, S. M. 1982 The ecological and social significance of duetting. (eds. D. E. Kroodsma & E. H. Miller), pp. 85-124. New York, Academic Press.
- [23] Ravignani, A. 2017 Interdisciplinary debate: Agree on definitions of synchrony. *Nature* **545**, 158.
- [24] Pika, S., Wilkinson, R., Kendrick, K. H. & Vernes, S. C. 2018 Taking turns: Bridging the gap between human and animal communication. *Proceedings of the Royal Society B: Biological Sciences* **285**, 20180598. (DOI:10.1098/rspb.2018.0598).
- [25] Greenfield, M. D. 2005 Mechanisms and Evolution of Communal Sexual Displays in Arthropods and Anurans. *Advances in the Study of Behaviour* **35**, 1-62. (DOI:10.1016/S0065-3454(05)35001-7).
- [26] Yoshida, S. & Okanoya, K. 2005 Evolution of Turn-Taking: A Bio-Cognitive Perspective. *Cognitive Studies* **12**, 153-165.
- [27] Feldman, R. 2007 Parent–Infant Synchrony: Biological Foundations and Developmental Outcomes. *Current Directions in Psychological Science* **16**, 340-345. (DOI:10.1111/j.1467-8721.2007.00532.x).
- [28] Juslin, P. N. & Sloboda, J. A. 2013 *Music and Emotion*. pp. 583-645, Elsevier Inc.
- [29] Chang, A., Kragness, H. E., Livingstone, S. R., Bosnyak, D. J. & Trainor, L. J. 2019 Body sway reflects joint emotional expression in music ensemble performance. *Sci. Rep.* **9**, 1-11. (DOI:10.1038/s41598-018-36358-4).
- [30] Chauvigné, L. A. S., Walton, A., Richardson, M. J. & Brown, S. 2019 Multi-person and multisensory synchronization during group dancing. *Human Movement Science* **63**, 199-208. (DOI:10.1016/j.humov.2018.12.005).
- [31] Goldin-Meadow, S. 1999 The role of gesture in communication and thinking. *Trends Cogn. Sci.* **3**, 419-429. (DOI:10.1016/S1364-6613(99)01397-2).
- [32] Holler, J., Schubotz, L., Kelly, S., Hagoort, P., Schuetze, M. & Özyürek, A. 2014 Social eye gaze modulates processing of speech and co-speech gesture. *Cognition* **133**, 692-697. (DOI:10.1016/j.cognition.2014.08.008).
- [33] Tronick, E. Z. & Cohn, J. F. 1989 Infant-Mother Face-to-Face Interaction: Age and Gender Differences in Coordination and the Occurrence of Miscoordination. *Child Dev.* **60**, 92-92. (DOI:10.2307/1131074).
- [34] Beebe, B., Alson, D., Jaffe, J., Feldstein, S. & Crown, C. 1988 Vocal congruence in mother-infant play. *J. Psycholinguist. Res.* **17**, 245-259. (DOI:10.1007/BF01686358).
- [35] Leong, V., Byrne, E., Clackson, K., Georgieva, S., Lam, S. & Wass, S. 2017 Speaker gaze increases information coupling between infant and adult brains. *Proc Natl Acad Sci U S A* **114**, 13290-13295. (DOI:10.1073/pnas.1702493114).
- [36] Leclère, C., Viaux, S., Avril, M., Achard, C., Chetouani, M., Missonnier, S. & Cohen, D. 2014 Why Synchrony Matters during Mother-Child Interactions: A Systematic Review. *PLoS ONE* **9**, e113571. (DOI:10.1371/journal.pone.0113571).
- [37] Feldman, R. 2017 The Neurobiology of Human Attachments. *Trends Cogn. Sci.* **21**, 80-99. (DOI:10.1016/j.tics.2016.11.007).
- [38] Feldman, R., Greenbaum, C. W. & Yirmiya, N. 1999 Mother-infant affect synchrony as an antecedent of the emergence of self-control. *Dev. Psychol.* **35**, 223-231. (DOI:10.1037/0012-1649.35.1.223).
- [39] Trainor, L. J. & Cirelli, L. 2015 Rhythm and interpersonal synchrony in early social development. *Ann. N. Y. Acad. Sci.* **1337**, 45-52. (DOI:10.1111/nyas.12649).
- [40] Lytle, S. R. & Kuhl, P. K. 2018 Social interaction and language acquisition: Toward a neurobiological view. (eds. E. M. Fernández & H. S. Cairns), pp. 615-634, Wiley Blackwell.
- [41] Trehub, S. E., Ghazban, N. & Corbeil, M. 2015 Musical affect regulation in infancy. *Ann. N. Y. Acad. Sci.* **1337**, 186-192. (DOI:10.1111/nyas.12622).
- [42] Cirelli, L. K., Jurewicz, Z. B. & Trehub, S. E. 2020 Effects of maternal singing style on mother–infant arousal and behavior. *J. Cognit. Neurosci.* **32**, 1213-1220. (DOI:10.1162/jocn_a_01402).
- [43] Patel, A. D. 2010 *Music, language, and the brain*, Oxford University Press.
- [44] Cirelli, L. K., Trehub, S. E. & Trainor, L. J. 2018 Rhythm and melody as social signals for infants. *Ann. N. Y. Acad. Sci.* **1423**, 66-72. (DOI:10.1111/nyas.13580).

- [45] Harrist, A. W. & Waugh, R. M. 2002 Dyadic synchrony: Its structure and function in children's development. *Dev. Rev.* **22**, 555-592. (DOI:10.1016/S0273-2297(02)00500-2).
- [46] Maccoby, E. E. 2004 The role of parents in the socialization of children: An historical overview. (eds. R. D. Parke, P. A. Ornstein, J. J. Rieser & C. Zahn-Waxler), pp. 589-615, American Psychological Association.
- [47] Casillas, M. & Frank, M. C. 2017 The development of children's ability to track and predict turn structure in conversation. *Journal of Memory and Language* **92**, 234-253. (DOI:10.1016/j.jml.2016.06.013).
- [48] Tomasello, M. & Farrar, M. J. 1986 Joint Attention and Early Language. *Child Dev.* **57**, 1454. (DOI:10.2307/1130423).
- [49] Stanton-Chapman, T. L. & Snell, M. E. 2011 Promoting turn-taking skills in preschool children with disabilities: The effects of a peer-based social communication intervention. *Early Childhood Research Quarterly* **26**, 303-319. (DOI:10.1016/j.ecresq.2010.11.002).
- [50] Black, B. & Logan, A. 1995 Links between Communication Patterns in Mother-Child, Father-Child, and Child-Peer Interactions and Children's Social Status. *Child Dev.* **66**, 255-271. (DOI:10.1111/j.1467-8624.1995.tb00869.x).
- [51] Black, B. & Hazen, N. L. 1990 Social Status and Patterns of Communication in Acquainted and Unacquainted Preschool Children. *Dev. Psychol.* **26**, 379-387. (DOI:10.1037/0012-1649.26.3.379).
- [52] Kirschner Sebastian, S. & Tomasello, M. 2010 Joint music making promotes prosocial behavior in 4-year-old children. *Evol Hum Behav* **31**, 354-364. (DOI:10.1016/j.evolhumbehav.2010.04.004).
- [53] Rabinowitch, T.-C. & Knafo-Noam, A. 2015 Synchronous Rhythmic Interaction Enhances Children's Perceived Similarity and Closeness towards Each Other. *PLoS ONE* **10**, e0120878. (DOI:10.1371/journal.pone.0120878).
- [54] Abney, D. H., Paxton, A., Dale, R. & Kello, C. T. 2014 Complexity matching in dyadic conversation. *J. Exp. Psychol. Gen.* **143**, 2304-2315. (DOI:10.1037/xge0000021).
- [55] Wilson, M. & Wilson, T. P. 2005 An oscillator model of the timing of turn-taking. *Psychonomic Bulletin and Review* **12**, 957-968. (DOI:10.3758/BF03206432).
- [56] Buder, E. H., Warlaumont, A. S., Kimbrough, D. & Chorna, L. B. 2010 Dynamic Indicators of Mother-Infant Prosodic and Illocutionary Coordination. In *Speech Prosody* (pp. 6-9).
- [57] Schultz, B. G., O'Brien, I., Phillips, N., McFarland, D. H., Titone, D. & Palmer, C. 2016 Speech rates converge in scripted turn-taking conversations. *Appl. Psycholinguist.* **37**, 1201-1220. (DOI:10.1017/S0142716415000545).
- [58] McFarland, D. H. 2001 Respiratory Markers of Conversational Interaction. *J. Speech. Lang. Hear. Res.* **44**, 128-143. (DOI:10.1044/1092-4388(2001/012)).
- [59] O'Dell, M., Nieminen, T. & Lennes, M. 2012 Modeling turn-taking rhythms with oscillators. *Linguistica Uralica* **48**, 218-227. (DOI:10.3176/lu.2012.3.08).
- [60] Stivers, T., Enfield, N. J., Brown, P., Englert, C., Hayashi, M., Heinemann, T., Hoymann, G., Rossano, F., de Ruiter, J. P., Yoon, K.-E., et al. 2009 Universals and cultural variation in turn-taking in conversation. *Proceedings of the National Academy of Sciences* **106**, 10587. (DOI:10.1073/pnas.0903616106).
- [61] Haegens, S. & Zion Golumbic, E. 2018 Rhythmic facilitation of sensory processing: A critical review. *Neurosci Biobehav Rev* **86**, 150-165. (DOI:10.1016/j.neubiorev.2017.12.002).
- [62] Bishop, L. & Goebel, W. 2015 When they listen and when they watch: Pianists' use of nonverbal audio and visual cues during duet performance. *Music. Sci.* **19**, 84-110. (DOI:10.1177/1029864915570355).
- [63] Moran, N., Hadley, L. V., Bader, M. & Keller, P. E. 2015 Perception of 'Back-Channeling' Nonverbal Feedback in Musical Duo Improvisation. *PLoS ONE* **10**, e0130070. (DOI:10.1371/journal.pone.0130070).
- [64] Repp, B. H. & Su, Y. H. 2013 Sensorimotor synchronization: A review of recent research (2006-2012). *Psychonomic Bulletin and Review* **20**, 403-452. (DOI:10.3758/s13423-012-0371-2).
- [65] Gilboa, A. & Tal-Shmotkin, M. 2012 String quartets as self-managed teams: An interdisciplinary perspective. *Psychology of Music* **40**, 19-41. (DOI:10.1177/0305735610377593).
- [66] Timmers, R., Endo, S., Bradbury, A. & Wing, A. M. 2014 Synchronization and leadership in string quartet performance: a case study of auditory and visual cues. *Front. Psychol.* **5**, 645. (DOI:10.3389/fpsyg.2014.00645).

- [67] Wing, A. M., Endo, S., Bradbury, A. & Vorberg, D. 2014 Optimal feedback correction in string quartet synchronization. *J R Soc Interface* **11**, 20131125. (DOI:10.1098/rsif.2013.1125).
- [68] Demos, A. P., Carter, D. J., Wanderley, M. M. & Palmer, C. 2017 The Unresponsive Partner: Roles of Social Status, Auditory Feedback, and Animacy in Coordination of Joint Music Performance. *Front. Psychol.* **8**, 149. (DOI:10.3389/fpsyg.2017.00149).
- [69] Zamm, A., Wellman, C. & Palmer, C. 2016 Endogenous rhythms influence interpersonal synchrony. *J. Exp. Psychol. Hum. Percept. Perform.* **42**, 611-616. (DOI:10.1037/xhp0000201).
- [70] Chang, A., Livingstone, S. R., Bosnyak, D. J. & Trainor, L. J. 2017 Body sway reflects leadership in joint music performance. *Proc Natl Acad Sci U S A* **114**, E4134-E4141. (DOI:10.1073/pnas.1617657114).
- [71] Hove, M. J. & Risen, J. L. 2009 It's all in the timing: Interpersonal synchrony increases affiliation. *Soc. Cogn.* **27**, 949-960. (DOI:10.1521/soco.2009.27.6.949).
- [72] Wiltermuth, S. S. & Heath, C. 2009 Synchrony and cooperation. *Psychol. Sci.* **20**, 1-5. (DOI:10.1111/j.1467-9280.2008.02253.x).
- [73] Savage, P. E., Loui, P., Tarr, B., Schachner, A., Glowacki, L., Mithen, S. & Fitch, W. T. 2020 Music as a coevolved system for social bonding. *Behav. Brain Sci.*, 1-36. (DOI:10.1017/S0140525X20000333).
- [74] Cheney, D. L., Silk, J. B. & Seyfarth, R. M. 2016 Network connections, dyadic bonds and fitness in wild female baboons. *Royal Society Open Science* **3**, 160255. (DOI:10.1098/rsos.160255).
- [75] Silk, J. B., Alberts, S. C. & Altmann, J. 2003 Social Bonds of Female Baboons Enhance Infant Survival. *Science* **302**, 1231-1234. (DOI:10.1126/science.1088580).
- [76] Lemasson, A., Ouattara, K., Petit, E. J. & Zuberbühler, K. 2011 Social learning of vocal structure in a nonhuman primate? *BMC Evol. Biol.* **11**, 1-7. (DOI:10.1186/1471-2148-11-362).
- [77] Lemasson, A. & Hausberger, M. 2004 Patterns of vocal sharing and social dynamics in a captive group of Campbell's monkeys (*Cercopithecus campbelli campbelli*). *J Comp Psychol* **118**, 347-359. (DOI:10.1037/0735-7036.118.3.347).
- [78] Levréro, F., Touitou, S., Frédet, J., Nairaud, B., Guéry, J. P. & Lemasson, A. 2019 Social bonding drives vocal exchanges in Bonobos. *Sci. Rep.* **9**, 1-11. (DOI:10.1038/s41598-018-36024-9).
- [79] Geissmann, T. 2000 Gibbon songs and human music from an evolutionary perspective. (eds. N. Wallin, B. Merker & S. Brown), pp. 103-123. Cambridge, Massachusetts, MIT Press.
- [80] Torti, V., Bonadonna, G., De Gregorio, C., Valente, D., Randrianarison, R. M., Friard, O., Pozzi, L., Gamba, M. & Giacoma, C. 2017 An intra-population analysis of the indris' song dissimilarity in the light of genetic distance. *Sci. Rep.* **7**, 1-12. (DOI:10.1038/s41598-017-10656-9).
- [81] Bonadonna, G., Torti, V., Sorrentino, V., Randrianarison, R. M., Zaccagno, M., Gamba, M., Tan, C. L. & Giacoma, C. 2017 Territory exclusivity and intergroup encounters in the indris (Mammalia: Primates: Indridae: *Indri indri*) upon methodological tuning. *The European Zoological Journal* **84**, 238-251. (DOI:10.1080/24750263.2017.1318184).
- [82] Bonadonna, G., Torti, V., Gregorio, C., Valente, D., Randrianarison, R. M., Pozzi, L., Gamba, M. & Giacoma, C. 2019 Evidence of genetic monogamy in the lemur Indri (*Indri indri*). *Am. J. Primatol.* **81**, e22993. (DOI:10.1002/ajp.22993).
- [83] Gamba, M., Torti, V., Estienne, V., Randrianarison, R. M., Valente, D., Rovara, P., Bonadonna, G., Friard, O. & Giacoma, C. 2016 The Indris Have Got Rhythm! Timing and Pitch Variation of a Primate Song Examined between Sexes and Age Classes. *Front. Neurosci.* **10**, 249. (DOI:10.3389/fnins.2016.00249).
- [84] Torti, V., Gamba, M., Rabemananjara, Z. H. & Giacoma, C. 2013 The songs of the indris (Mammalia: Primates: Indridae): contextual variation in the long-distance calls of a lemur. *Ital. J. Zool.* **80**, 596-607. (DOI:10.1080/11250003.2013.845261).
- [85] Clarke, E., Reichard, U. H. & Zuberbühler, K. 2006 The Syntax and Meaning of Wild Gibbon Songs. *PLoS ONE* **1**, e73. (DOI:10.1371/journal.pone.0000073).
- [86] Badraun, J. C., Mootnick, A. R., Deaner, R. O., Agoramoorthy, G. & McNeese, K. M. 1998 Hand modulation of vocalization in Siamangs *Hylobates syndactylus*. *Int Zoo Yearb* **36**, 84-89. (DOI:10.1111/j.1748-1090.1998.tb02888.x).
- [87] Geissmann, T. & Orgeldinger, M. 2000 The relationship between duet songs and pair bonds in siamangs, *Hylobates syndactylus*. *Anim. Behav.* **60**, 805-809. (DOI:10.1006/anbe.2000.1540).

- [88] Maretti, G., Sorrentino, V., Finomana, A., Gamba, M. & Giacoma, C. 2010 Not just a pretty song: an overview of the vocal repertoire of Indri indri. *Journal of Anthropological Sciences* **88**, 151-165.
- [89] Valente, D., De Gregorio, C., Torti, V., Miaretsoa, L., Friard, O., Randrianarison, R. M., Giacoma, C. & Gamba, M. 2019 Finding Meanings in Low Dimensional Structures: Stochastic Neighbor Embedding Applied to the Analysis of Indri indri Vocal Repertoire. *Animals* **9**, 243. (DOI:10.3390/ani9050243).
- [90] Lilly, J. C. & Miller, A. M. 1961 Sounds emitted by the bottlenose dolphin. *Science* **133**, 1689-1693. (DOI:10.1126/science.133.3465.1689).
- [91] Caldwell, C. A. 1968 Vocalisation of native dolphins in small groups. *Science* **159**, 1121-1123.
- [92] Nakahara, F. & Miyazaki, N. 2011 Vocal exchanges of signature whistles in bottlenose dolphins (*Tursiops truncatus*). *J. Ethol.* **29**, 309-320. (DOI:10.1007/s10164-010-0259-4).
- [93] Moore, B. L., Connor, R. C., Allen, S. J., Krützen, M. & King, S. L. 2020 Acoustic coordination by allied male dolphins in a cooperative context. *Proceedings of the Royal Society B: Biological Sciences* **287**, 20192944. (DOI:10.1098/rspb.2019.2944).
- [94] Insley, S. J., Phillips, A. & Charrier, I. 2003 A review of social recognition in pinnipeds. *Aquat. Mamm.* **29**, 181-201. (DOI:10.1578/016754203101024149).
- [95] Charrier, I. 2020 Mother–Offspring Vocal Recognition and Social System in Pinnipeds. (eds. T. Aubin & N. Mathevon), pp. 231-246, Springer, Cham.
- [96] Ravnani, A. 2019 Timing of antisynchronous calling: A case study in a harbor seal pup (*Phoca vitulina*). *J Comp Psychol* **133**, 272-277. (DOI:10.1037/com0000160).
- [97] Esser, K.-H. & Schmidt, U. 2010 Mother-Infant Communication in the Lesser Spear-nosed Bat *Phyllostomus discolor* (Chiroptera, Phyllostomidae) - Evidence for Acoustic Learning. *Ethology* **82**, 156-168. (DOI:10.1111/j.1439-0310.1989.tb00496.x).
- [98] Balcombe, J. P. & McCracken, G. F. 1992 Vocal recognition in mexican free-tailed bats: do pups recognize mothers? *Anim. Behav.* **43**, 79-87. (DOI:10.1016/S0003-3472(05)80073-9).
- [99] De Fanis, E. & Jones, G. 1996 Allomaternal care and recognition between mothers and young in pipistrelle bats (*Pipistrellus pipistrellus*). *J. Zool.* **240**, 781-787. (DOI:10.1111/j.1469-7998.1996.tb05324.x).
- [100] Carter, G. G., Fenton, M. B. & Faure, P. A. 2009 White-winged vampire bats (*Diaemus youngi*) exchange contact calls. *Can J Zool* **87**, 604-608. (DOI:10.1139/Z09-051).
- [101] Okobi, D. E., Banerjee, A., Matheson, A. M. M., Phelps, S. M. & Long, M. A. 2019 Motor cortical control of vocal interaction in neotropical singing mice. *Science* **363**, 983-988. (DOI:10.1126/science.aau9480).
- [102] Yosida, S. & Okanoya, K. 2009 Naked Mole-Rat is Sensitive to Social Hierarchy Encoded in Antiphonal Vocalization. *Ethology* **115**, 823-831. (DOI:10.1111/j.1439-0310.2009.01677.x).
- [103] Rado, R., Levi, N., Hauser, H., Witcher, J., Alder, N., Intrator, N., Wollberg, Z. & Terkel, J. 1987 Seismic signalling as a means of communication in a subterranean mammal. *Anim. Behav.* **35**, 1249-1251. (DOI:10.1016/S0003-3472(87)80183-5).
- [104] Leighty, K. A., Soltis, J., Wesolek, C. M. & Savage, A. 2008 Rumble vocalizations mediate interpartner distance in African elephants, *Loxodonta africana*. *Anim. Behav.* **76**, 1601-1608. (DOI:10.1016/j.anbehav.2008.06.022).
- [105] Tilson, R. L. & Norton, P. M. 1981 Alarm Duetting and Pursuit Deterrence in an African Antelope. *The American Naturalist* **118**, 455-462. (DOI:10.1086/283840).
- [106] Demartsev, V., Strandburg-Peshkin, A., Ruffner, M. & Manser, M. 2018 Vocal Turn-Taking in Meerkat Group Calling Sessions. *Curr. Biol.* **28**, 3661-3666.e3. (DOI:10.1016/j.cub.2018.09.065).
- [107] Tobias, J. A., Sheard, C., Seddon, N., Meade, A., Cotton, A. J. & Nakagawa, S. 2016 Territoriality, Social Bonds, and the Evolution of Communal Signaling in Birds. *Frontiers in Ecology and Evolution* **4**, 74. (DOI:10.3389/fevo.2016.00074).
- [108] Malacarne, G., Cucco, M. & Camanni, S. 1991 Coordinated visual displays and vocal duetting in different ecological situations among western palearctic non-passerine birds. *Ethology Ecology and Evolution* **3**, 207-219. (DOI:10.1080/08927014.1991.9525369).
- [109] Soma, M. & Garamszegi, L. Z. 2015 Evolution of courtship display in Estrildid finches: dance in relation to female song and plumage ornamentation. *Frontiers in Ecology and Evolution* **3**, 4. (DOI:10.3389/fevo.2015.00004).

- [110] Hall, M. L. 2004 A review of hypotheses for the functions of avian duetting. *Behav. Ecol. Sociobiol.* **55**, 415-430. (DOI:10.1007/s00265-003-0741-x).
- [111] Langmore, N. E. 1998 Functions of duet and solo songs of female birds. *Trends in Ecology and Evolution* **13**, 136-140. (DOI:10.1016/S0169-5347(97)01241-X).
- [112] Short, L. L. & Horne, J. F. M. 1983 A Review of Duetting, Sociality and Speciation in Some African Barbets (Capitonidae). *The Condor* **85**, 323. (DOI:10.2307/1367069).
- [113] Reş, P. & Magrath, R. D. 2016 Multimodal duetting in magpie-larks: How do vocal and visual components contribute to a cooperative signal's function? *Anim. Behav.* **117**, 35-42. (DOI:10.1016/j.anbehav.2016.04.024).
- [114] Mann, N. I., Dingess, K. A., Barker, K. F., Graves, J. A. & Slater, P. J. B. 2009 A comparative study of song form and duetting in neotropical *Thryothorus* wrens. *Behaviour* **146**, 1-43. (DOI:10.1163/156853908X390913).
- [115] Brumm, H. & Slater, P. 2007 Animal Communication: Timing Counts. *Curr. Biol.* **17**, R521-R523. (DOI:10.1016/j.cub.2007.04.053).
- [116] Takeda, K. F., Hiraiwa-Hasegawa, M. & Kutsukake, N. 2019 Uncoordinated dances associated with high reproductive success in a crane. *Behav. Ecol.* **30**, 101-106. (DOI:10.1093/beheco/ary159).
- [117] Reş, P. 2018 Multimodal coordination enhances the responses to an avian duet. *Behav. Ecol.* **29**, 411-417. (DOI:10.1093/beheco/ary174).
- [118] Hall, M. L. & Magrath, R. D. 2007 Temporal coordination signals coalition quality. *Curr. Biol.* **17**, R406-R407.
- [119] Rivera-Cáceres, K. D., Quirós-Guerrero, E., Araya-Salas, M. & Searcy, W. A. 2016 Neotropical wrens learn new duet rules as adults. *Proceedings of the Royal Society B: Biological Sciences* **283**, 20161819. (DOI:10.1098/rspb.2016.1819).
- [120] Griffith, S. C. & Buchanan, K. L. 2010 The Zebra Finch: The ultimate Australian supermodel. *Emu* **110**, v-xii. (DOI:10.1071/MUv110n3_ED).
- [121] Prior, N. H., Smith, E., Dooling, R. J. & Ball, G. F. 2020 Familiarity Enhances Moment-to-Moment Behavioral Coordination in Zebra Finch (*Taeniopygia guttata*) Dyads. *J Comp Psychol* **134**, 135-148. (DOI:10.1037/com0000201).
- [122] D'Amelio, P. B., Trost, L. & ter Maat, A. 2017 Vocal exchanges during pair formation and maintenance in the zebra finch (*Taeniopygia guttata*). *Front Zool* **14**, 1-12. (DOI:10.1186/s12983-017-0197-x).
- [123] Benichov, J. I., Benezra, S. E., Vallentin, D., Globerson, E., Long, M. A. & Tchernichovski, O. 2016 The forebrain song system mediates predictive call timing in female and Male Zebra finches. *Curr. Biol.* **26**, 309-318. (DOI:10.1016/j.cub.2015.12.037).
- [124] Perez, E. C., Elie, J. E., Boucaud, I. C. A., Crouchet, T., Soulage, C. O., Soula, H. A., Theunissen, F. E. & Vignal, C. 2015 Physiological resonance between mates through calls as possible evidence of empathic processes in songbirds. *Horm. Behav.* **75**, 130-141. (DOI:10.1016/j.yhbeh.2015.09.002).
- [125] Boucaud, I. C. A., Perez, E. C., Ramos, L. S., Griffith, S. C. & Vignal, C. 2017 Acoustic communication in zebra finches signals when mates will take turns with parental duties. *Behav. Ecol.* **28**, 645-656. (DOI:10.1093/beheco/arw189).
- [126] Grafe, T. U. 2005 Anuran choruses as communication networks. (pp. 277-299).
- [127] Bosch, J. & Márquez, R. 2001 Call timing in male-male acoustical interactions and female choice in the midwife toad *Alytes obstetricans*. *Copeia* **2001**, 169-177. (DOI:10.1643/0045-8511(2001)001[0169:CTIMMA]2.0.CO;2).
- [128] Wells, K. D. & Schwartz, J. J. 2006 *The Behavioral Ecology of Anuran Communication*. pp. 44-86, Springer New York.
- [129] Wells, K. D. 1977 The social behaviour of anuran amphibians. *Anim. Behav.* **25**, 666-693. (DOI:10.1016/0003-3472(77)90118-X).
- [130] Caldwell, J. P. & Shepard, D. B. 2007 Calling Site Fidelity and Call Structure of a Neotropical Toad, *Rhinella Ocellata* (Anura: Bufonidae). *J. Herpetol.* **41**, 611-621. (DOI:10.1670/07-025.1).
- [131] Klump, G. M. & Gerhardt, H. C. 1992 Mechanisms and Function of Call-Timing in Male-Male Interactions in Frogs. (pp. 153-174, Springer US).

- [132] Boatright-Horowitz, S. L., Horowitz, S. S. & Simmons, A. M. 2000 Patterns of Vocal Interactions in a Bullfrog (*Rana catesbeiana*) Chorus: Preferential Responding to Far Neighbors. *Ethology* **106**, 701-712. (DOI:10.1046/j.1439-0310.2000.00580.x).
- [133] Tuttle, M. D. & Ryan, M. J. 1982 The role of synchronized calling, ambient light, and ambient noise, in anti-bat-predator behavior of a treefrog. *Behav. Ecol. Sociobiol.* **11**, 125-131. (DOI:10.1007/BF00300101).
- [134] Tobias, M. L., Viswanathan, S. S. & Kelley, D. B. 1998 Rapping, a female receptive call, initiates male-female duets in the South African clawed frog. *Proc Natl Acad Sci U S A* **95**, 1870-1875. (DOI:10.1073/pnas.95.4.1870).
- [135] Bailey, W. J. 2003 Insect duets: Underlying mechanisms and their evolution. *Physiol. Entomol.* **28**, 157-174. (DOI:10.1046/j.1365-3032.2003.00337.x).
- [136] Zimmermann, U., Rheinlaender, J. & Robinson, D. 1989 Journal of Sensory, Comparative Cues for male phonotaxis in the duetting bushcricket *Leptophyes punctatissima*. *Journal of Comparative Physiology A* **164**, 621-628.
- [137] Bailey, W. J. & Hammond, T. J. 2003 Duetting in insects does call length influence reply latency? *J. Zool.* **260**, 267-274. (DOI:10.1017/S0952836903003728).
- [138] Kuhelj, A., de Groot, M., Blejec, A. & Virant-Doberlet, M. 2015 The Effect of Timing of Female Vibrational Reply on Male Signalling and Searching Behaviour in the Leafhopper *Aphrodes makarovi*. *PLoS ONE* **10**, e0139020. (DOI:10.1371/journal.pone.0139020).
- [139] Kuhelj, A., de Groot, M., Blejec, A. & Virant-Doberlet, M. 2016 Sender-receiver dynamics in leafhopper vibrational duetting. *Anim. Behav.* **114**, 139-146. (DOI:10.1016/j.anbehav.2016.02.001).
- [140] Villarreal, S. M. & Gilbert, C. 2014 Male *Scudderia pistillata* katydids defend their acoustic duet against eavesdroppers. *Behav. Ecol. Sociobiol.* **68**, 1669-1675. (DOI:10.1007/s00265-014-1775-y).
- [141] LaRue, K. M., Clemens, J., Berman, G. J. & Murthy, M. 2015 Acoustic duetting in *Drosophila virilis* relies on the integration of auditory and tactile signals. *eLife* **4**, 1-23. (DOI:10.7554/eLife.07277).
- [142] Rajaraman, K., Godthi, V., Pratap, R. & Balakrishnan, R. 2015 A novel acoustic-vibratory multimodal duet. *J. Exp. Biol.* **218**, 3042-3050. (DOI:10.1242/jeb.122911).
- [143] Greenfield, M. D. & Minckley, R. L. 1993 Acoustic Dueling in Tarbush Grasshoppers: Settlement of Territorial Contests via Alternation of Reliable Signals. *Ethology* **95**, 309-326. (DOI:10.1111/j.1439-0310.1993.tb00480.x).
- [144] Jeng, M.-L. 2019 Biofluorescence in Terrestrial Animals, with Emphasis on Fireflies: A Review and Field Observation. IntechOpen.
- [145] Tuthill, J. C. & Wilson, R. I. 2016 Mechanosensation and Adaptive Motor Control in Insects. *Curr. Biol.* **26**, R1022-R1038. (DOI:10.1016/j.cub.2016.06.070).
- [146] Halfwerk, W., Varkevisser, J., Simon, R., Mendoza, E., Scharff, C. & Riebel, K. 2019 Toward Testing for Multimodal Perception of Mating Signals. *Frontiers in Ecology and Evolution* **7**, 124. (DOI:10.3389/fevo.2019.00124).
- [147] Ota, N., Gahr, M. & Soma, M. 2015 Tap dancing birds: The multimodal mutual courtship display of males and females in a socially monogamous songbird. *Sci. Rep.* **5**, 1-6. (DOI:10.1038/srep16614).
- [148] Grafe, T. U. & Wanger, T. C. 2007 Multimodal Signaling in Male and Female Foot-Flagging Frogs *Staurois guttatus* (Ranidae): An Alerting Function of Calling. *Ethology* **113**, 772-781. (DOI:10.1111/j.1439-0310.2007.01378.x).
- [149] Carouso-Peck, S., Menyhart, O., DeVoogd, T. J. & Goldstein, M. H. 2020 Contingent parental responses are naturally associated with zebra finch song learning. *Anim. Behav.* **165**, 123-132. (DOI:10.1016/j.anbehav.2020.04.019).
- [150] Lampen, J., McAuley, J. D., Chang, S. E. & Wade, J. 2019 Neural activity associated with rhythmicity of song in juvenile male and female zebra finches. *Behav. Processes* **163**, 45-52. (DOI:10.1016/j.beproc.2017.12.003).
- [151] Desjonquères, C., Maliszewski, J., Lewandowski, E. N., Speck, B. & Rodríguez, R. L. 2019 Social ontogeny in the communication system of an insect. *Anim. Behav.* **148**, 93-103. (DOI:10.1016/j.anbehav.2018.12.002).

- [152] Greenfield, M. D. 1994 Synchronous and alternating choruses in insects and anurans: Common mechanisms and diverse functions. *Integr. Comp. Biol.* **34**, 605-615. (DOI:10.1093/icb/34.6.605).
- [153] Hoffmann, S., Trost, L., Voigt, C., Leitner, S., Lemazina, A., Sagunsky, H., Abels, M., Kollmansperger, S., Maat, A. T. & Gahr, M. 2019 Duets recorded in the wild reveal that interindividually coordinated motor control enables cooperative behavior. *Nature Communications* **10**, 1-11. (DOI:10.1038/s41467-019-10593-3).
- [154] Henry, M., Cook, P. F., De Reus, K., Nityananda, V., Rouse, A. A. & Kotz, S. A. An ecological approach to measuring rhythm abilities across the animal kingdom. *Philosophical Transactions of the Royal Society B: Biological Sciences*.