

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Cosmic-Ray Anisotropies in Right Ascension Measured by the Pierre Auger Observatory

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1802045> since 2021-09-17T18:56:51Z

Published version:

DOI:10.3847/1538-4357/ab7236

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

COSMIC-RAY ANISOTROPIES IN RIGHT ASCENSION
MEASURED BY THE PIERRE AUGER OBSERVATORY

A. AAB⁷⁵, P. ABREU⁶⁷, M. AGLIETTA^{50,49}, I.F.M. ALBUQUERQUE¹⁹, J.M. ALBURY¹², I. ALLEKOTTE¹, A. ALMELA^{8,11}, J. ALVAREZ CASTILLO⁶³, J. ALVAREZ-MUÑIZ⁷⁴, G.A. ANASTASI^{58,49}, L. ANCHORDOQUI⁸², B. ANDRADA⁸, S. ANDRINGA⁶⁷, C. ARAMO⁴⁷, P.R. ARAÚJO FERREIRA³⁹, H. ASOREY⁸, P. ASSIS⁶⁷, G. AVILA^{9,10}, A.M. BADESCU⁷⁰, A. BAKALOVA³⁰, A. BALACEANU⁶⁸, F. BARBATO^{56,47}, R.J. BARREIRA LUZ⁶⁷, K.H. BECKER³⁵, J.A. BELLIDO¹², C. BERAT³⁴, M.E. BERTAINA^{58,49}, X. BERTOU¹, P.L. BIERMANN¹⁰⁰¹, T. BISTER³⁹, J. BITEAU³², A. BLANCO⁶⁷, J. BLAZEK³⁰, C. BLEVE³⁴, M. BOHÁČOVÁ³⁰, D. BONCIOLI^{53,43}, C. BONIFAZI²⁴, L. BONNEAU ARBELETCHÉ¹⁹, N. BORODAI⁶⁴, A.M. BOTTI^{8,37}, J. BRACK¹⁰⁰⁴, T. BRETZ³⁹, F.L. BRIECHELE³⁹, P. BUCHHOLZ⁴¹, A. BUENO⁷³, S. BUITINK¹⁴, M. BUSCEMI^{54,44}, K.S. CABALLERO-MORA⁶², L. CACCIANIGA⁵⁵, L. CALCAGNI⁴, A. CANCIO^{11,8}, F. CANFORA^{75,77}, I. CARACAS³⁵, J.M. CARCELLER⁷³, R. CARUSO^{54,44}, A. CASTELLINA^{50,49}, F. CATALANI¹⁷, G. CATALDI⁴⁵, L. CAZON⁶⁷, M. CERDA⁹, J.A. CHINELLATO²⁰, K. CHOI¹³, J. CHUDOBA³⁰, L. CHYTKA³¹, R.W. CLAY¹², A.C. COBOS CERUTTI⁷, R. COLALILLO^{56,47}, A. COLEMAN⁸⁸, M.R. COLUCCIA^{52,45}, R. CONCEIÇÃO⁶⁷, A. CONDORELLI^{42,43}, G. CONSOLATI^{46,51}, F. CONTRERAS^{9,10}, F. CONVENGA^{52,45}, C.E. COVAULT^{80,1007}, S. DASSO^{5,3}, K. DAUMILLER³⁷, B.R. DAWSON¹², J.A. DAY¹², R.M. DE ALMEIDA²⁶, J. DE JESÚS^{8,37}, S.J. DE JONG^{75,77}, G. DE MAURO^{75,77}, J.R.T. DE MELLO NETO^{24,25}, I. DE MITRI^{42,43}, J. DE OLIVEIRA²⁶, D. DE OLIVEIRA FRANCO²⁰, V. DE SOUZA¹⁸, J. DEBATIN³⁶, M. DEL RÍO¹⁰, O. DELIGNY³², N. DHITAL⁶⁴, A. DI MATTEO⁴⁹, M.L. DÍAZ CASTRO²⁰, C. DOBRIGKEIT²⁰, J.C. D'OLIVO⁶³, Q. DOROSTI⁴¹, R.C. DOS ANJOS²³, M.T. DOVA⁴, J. EBR³⁰, R. ENGEL^{36,37}, I. EPICOCO^{52,45}, M. ERDMANN³⁹, C.O. ESCOBAR¹⁰⁰², A. ETCHEGOYEN^{8,11}, H. FALCKE^{75,78,77}, J. FARMER⁸⁷, G. FARRAR⁸⁵, A.C. FAUTH²⁰, N. FAZZINI¹⁰⁰², F. FELDBUSCH³⁸, F. FENU^{58,49}, B. FICK⁸⁴, J.M. FIGUEIRA⁸, A. FILIPČIČ^{72,71}, M.M. FREIRE⁶, T. FUJII^{87,1005}, A. FUSTER^{8,11}, C. GALEA⁷⁵, C. GALELLI^{55,46}, B. GARCÍA⁷, A.L. GARCIA VEGAS³⁹, H. GEMMEKE³⁸, F. GESUALDI^{8,37}, A. GHERGHEL-LASCU⁶⁸, P.L. GHIA³², U. GIACCARI⁷⁵, M. GIAMMARCHI⁴⁶, M. GILLER⁶⁵, J. GLOMBITZA³⁹, F. GOBBI⁹, G. GOLUP¹, M. GÓMEZ BERISSO¹, P.F. GÓMEZ VITALE^{9,10}, J.P. GONGORA⁹, N. GONZÁLEZ⁸, I. GOOS^{1,37}, D. GÓRA⁶⁴, A. GORGI^{50,49}, M. GOTTOWIK³⁵, T.D. GRUBB¹², F. GUARINO^{56,47}, G.P. GUEDES²¹, E. GUIDO^{49,58}, S. HAHN^{37,8}, R. HALLIDAY⁸⁰, M.R. HAMPTEL⁸, P. HANSEN⁴, D. HARARI¹, V.M. HARVEY¹², A. HAUNGS³⁷, T. HEBBEKER³⁹, D. HECK³⁷, G.C. HILL¹², C. HOJVAT¹⁰⁰², J.R. HÖRANDEL^{75,77}, P. HORVATH³¹, M. HRABOVSKÝ³¹, T. HUEGE^{37,14}, J. HULSMAN^{8,37}, A. INSOLIA^{54,44}, P.G. ISAR⁶⁹, J.A. JOHNSEN⁸¹, J. JURYSSEK³⁰, A. KÄÄPÄ³⁵, K.H. KAMPERT³⁵, B. KELHAUER³⁷, J. KEMP³⁹, H.O. KLAGES³⁷, M. KLEIFGES³⁸, J. KLEINFELLER⁹, M. KÖPKE³⁶, G. KUKEC MEZEK⁷¹, A. KUOTB AWAD³⁶, B.L. LAGO¹⁶, D. LAHURD⁸⁰, R.G. LANG¹⁸, M.A. LEIGUI DE OLIVEIRA²², V. LENOK³⁷, A. LETESSIER-SELVON³³, I. LHENRY-YVON³², D. LO PRESTI^{54,44}, L. LOPES⁶⁷, R. LÓPEZ⁵⁹, A. LÓPEZ CASADO⁷⁴, R. LOREK⁸⁰, Q. LUCE³⁶, A. LUCERO⁸, A. MACHADO PAYERAS²⁰, M. MALACARI⁸⁷, G. MANCARELLA^{52,45}, D. MANDAT³⁰, B.C. MANNING¹², J. MANSHANDEN⁴⁰, P. MANTSCH¹⁰⁰², A.G. MARIAZZI⁴, I.C. MARIS¹³, G. MARSELLA^{52,45}, D. MARTELLO^{52,45}, H. MARTINEZ¹⁸, O. MARTÍNEZ BRAVO⁵⁹, M. MASTRODICASA^{53,43}, H.J. MATHES³⁷, J. MATTHEWS⁸³, G. MATTHIAE^{57,48}, E. MAYOTTE³⁵, P.O. MAZUR¹⁰⁰², G. MEDINA-TANCO⁶³, D. MELO⁸, A. MENSNIKOV³⁸, K.-D. MERENDA⁸¹, S. MICHAL³¹, M.I. MICHELETTI⁶, L. MIRAMONTI^{55,46}, D. MOCKLER¹³, S. MOLLERACH¹, F. MONTANET³⁴, C. MORELLO^{50,49}, G. MORLINO^{42,43}, M. MOSTAFÁ⁸⁶, A.L. MÜLLER^{8,37}, M.A. MULLER^{20,1003,24}, S. MÜLLER³⁶, R. MUSSA⁴⁹, M. MUZIO⁸⁵, W.M. NAMASAKA³⁵, L. NELLEN⁶³, M. NICULESCU-OGLINZANU⁶⁸, M. NIECHCIOL⁴¹, D. NITZ^{84,1006}, D. NOSEK²⁹, V. NOVOTNY²⁹, L. NOŽKA³¹, A. NUCITA^{52,45}, L.A. NÚÑEZ²⁸, M. PALATKA³⁰, J. PALLOTTA², M.P. PANETTA^{52,45}, P. PAPENBREER³⁵, G. PARENTE⁷⁴, A. PARRA⁵⁹, M. PECH³⁰, F. PEDREIRA⁷⁴, J. PEKALA⁶⁴, R. PELAYO⁶¹, J. PEÑA-RODRIGUEZ²⁸, L.A.S. PEREIRA²⁰, J. PEREZ ARMAND¹⁹, M. PERLIN^{8,37}, L. PERRONE^{52,45}, C. PETERS³⁹, S. PETREIRA^{42,43}, T. PIEROG³⁷, M. PIMENTA⁶⁷, V. PIRRONELLO^{54,44}, M. PLATINO⁸, B. PONT⁷⁵, M. POTHAST^{77,75}, P. PRIVITERA⁸⁷, M. PROUZA³⁰, A. PUYLEART⁸⁴, S. QUERCHFELD³⁵, J. RAUTENBERG³⁵, D. RAVIGNANI⁸, M. REININGHAUS^{37,8}, J. RIDKY³⁰, F. RIEHN⁶⁷, M. RISSE⁴¹, P. RISTORI², V. RIZI^{53,43}, W. RODRIGUES DE CARVALHO¹⁹, J. RODRIGUEZ ROJO⁹, M.J. RONCORONI⁸, M. ROTH³⁷, E. ROULET¹, A.C. ROVERO⁵, P. RUEHL⁴¹, S.J. SAFFI¹², A. SAFTOIU⁶⁸, F. SALAMIDA^{53,43}, H. SALAZAR⁵⁹, G. SALINA⁴⁸, J.D. SANABRIA GOMEZ²⁸, F. SÁNCHEZ⁸, E.M. SANTOS¹⁹, E. SANTOS³⁰, F. SARAZIN⁸¹, R. SARMENTO⁶⁷, C. SARMIENTO-CANO⁸, R. SATO⁹, P. SAVINA^{52,45,32}, C. SCHÄFER³⁷, V. SCHERINI⁴⁵, H. SCHELER³⁷, M. SCHIMASSEK^{36,8}, M. SCHIMP³⁵, F. SCHLÜTER^{37,8}, D. SCHMIDT³⁶, O. SCHOLTEN^{76,14}, P. SCHOVÁNEK³⁰, F.G. SCHRÖDER^{88,37}, S. SCHRÖDER³⁵, S.J. SCIUTTO⁴, M. SCORNAVACCHE^{8,37}, R.C. SHELLARD¹⁵, G. SIGL⁴⁰, G. SILLI^{8,37}, O. SIMA^{68,1007}, R. ŠMÍDA⁸⁷, P. SOMMERS⁸⁶, J.F. SORIANO⁸², J. SOUCHARD³⁴, R. SQUARTINI⁹, M. STADELMAIER^{37,8}, D. STANCA⁶⁸, S. STANIČ⁷¹, J. STASIELAK⁶⁴, P. STASSI³⁴, A. STREICH^{36,8}, M. SUÁREZ-DURÁN²⁸, T. SUDHOLZ¹², T. SUOMIJÄRVI³², A.D. SUPANITSKY⁸, J. ŠUPÍK³¹, Z. SZADKOWSKI⁶⁶, A. TABOADA^{36,8}, O.A. TABORDA¹, A. TAPIA²⁷, C. TIMMERMANS^{77,75}, P. TOBISKA³⁰, C.J. TODERO PEIXOTO¹⁷, B. TOMÉ⁶⁷, G. TORRALBA ELIPE⁷⁴, A. TRAVAINI⁹, P. TRAVNICEK³⁰, C. TRIMARELLI^{53,43}, M. TRINI^{71,1007}, M. TUEROS⁴, R. ULRICH³⁷, M. UNGER³⁷, M. URBAN³⁹, L. VACLAVEK³¹, J.F. VALDÉS GALICIA⁶³, I. VALIÑO^{42,43}, L. VALORE^{56,47}, A. VAN VLIET⁷⁵, E. VARELA⁵⁹, B. VARGAS CÁRDENAS⁶³, A. VÁSQUEZ-RAMÍREZ²⁸, D. VEBERIČ³⁷, C. VENTURA²⁵, I.D. VERGARA QUISEPÉ⁴, V. VERZI⁴⁸, J. VICHA³⁰, L. VILLASEÑOR⁵⁹, J. VINK⁷⁹, S. VOROBIOV⁷¹, H. WAHLBERG⁴, A.A. WATSON¹⁰⁰⁰, M. WEBER³⁸, A. WEINDL³⁷, L. WIENCKE⁸¹, H. WILCZYŃSKI⁶⁴, T. WINCHEN¹⁴, M. WIRTZ³⁹, D. WITTKOWSKI³⁵, B. WUNDHEILER⁸, A. YUSHKOV³⁰, E. ZAS⁷⁴, D. ZAVRTANIK^{71,72}, M. ZAVRTANIK^{72,71}, L. ZEHNER⁷¹, A. ZEPEDA⁶⁰, M. ZIOLKOWSKI⁴¹, F. ZUCCARELLO^{54,44}

(THE PIERRE AUGER COLLABORATION)

¹Centro Atómico Bariloche and Instituto Balseiro (CNEA-UNCuyo-CONICET), San Carlos de Bariloche, Argentina

²Centro de Investigaciones en Láseres y Aplicaciones, CITEDEF and CONICET, Villa Martelli, Argentina

³Departamento de Física and Departamento de Ciencias de la Atmósfera y los Océanos, FCEyN, Universidad de Buenos Aires and CONICET,

- Buenos Aires, Argentina
- ⁴IFLP, Universidad Nacional de La Plata and CONICET, La Plata, Argentina
- ⁵Instituto de Astronomía y Física del Espacio (IAFE, CONICET-UBA), Buenos Aires, Argentina
- ⁶Instituto de Física de Rosario (IFIR) – CONICET/U.N.R. and Facultad de Ciencias Bioquímicas y Farmacéuticas U.N.R., Rosario, Argentina
- ⁷Instituto de Tecnologías en Detección y Astropartículas (CNEA, CONICET, UNSAM), and Universidad Tecnológica Nacional – Facultad Regional Mendoza (CONICET/CNEA), Mendoza, Argentina
- ⁸Instituto de Tecnologías en Detección y Astropartículas (CNEA, CONICET, UNSAM), Buenos Aires, Argentina
- ⁹Observatorio Pierre Auger, Malargüe, Argentina
- ¹⁰Observatorio Pierre Auger and Comisión Nacional de Energía Atómica, Malargüe, Argentina
- ¹¹Universidad Tecnológica Nacional – Facultad Regional Buenos Aires, Buenos Aires, Argentina
- ¹²University of Adelaide, Adelaide, S.A., Australia
- ¹³Université Libre de Bruxelles (ULB), Brussels, Belgium
- ¹⁴Vrije Universiteit Brussels, Brussels, Belgium
- ¹⁵Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, RJ, Brazil
- ¹⁶Centro Federal de Educação Tecnológica Celso Suckow da Fonseca, Nova Friburgo, Brazil
- ¹⁷Universidade de São Paulo, Escola de Engenharia de Lorena, Lorena, SP, Brazil
- ¹⁸Universidade de São Paulo, Instituto de Física de São Carlos, São Carlos, SP, Brazil
- ¹⁹Universidade de São Paulo, Instituto de Física, São Paulo, SP, Brazil
- ²⁰Universidade Estadual de Campinas, IFGW, Campinas, SP, Brazil
- ²¹Universidade Estadual de Feira de Santana, Feira de Santana, Brazil
- ²²Universidade Federal do ABC, Santo André, SP, Brazil
- ²³Universidade Federal do Paraná, Setor Palotina, Palotina, Brazil
- ²⁴Universidade Federal do Rio de Janeiro, Instituto de Física, Rio de Janeiro, RJ, Brazil
- ²⁵Universidade Federal do Rio de Janeiro (UFRJ), Observatório do Valongo, Rio de Janeiro, RJ, Brazil
- ²⁶Universidade Federal Fluminense, EEIMVR, Volta Redonda, RJ, Brazil
- ²⁷Universidad de Medellín, Medellín, Colombia
- ²⁸Universidad Industrial de Santander, Bucaramanga, Colombia
- ²⁹Charles University, Faculty of Mathematics and Physics, Institute of Particle and Nuclear Physics, Prague, Czech Republic
- ³⁰Institute of Physics of the Czech Academy of Sciences, Prague, Czech Republic
- ³¹Palacky University, RCPTM, Olomouc, Czech Republic
- ³²Institut de Physique Nucléaire d’Orsay (IPNO), Université Paris-Sud, Univ. Paris/Saclay, CNRS-IN2P3, Orsay, France
- ³³Laboratoire de Physique Nucléaire et de Hautes Energies (LPNHE), Universités Paris 6 et Paris 7, CNRS-IN2P3, Paris, France
- ³⁴Univ. Grenoble Alpes, CNRS, Grenoble Institute of Engineering Univ. Grenoble Alpes, LPSC-IN2P3, 38000 Grenoble, France, France
- ³⁵Bergische Universität Wuppertal, Department of Physics, Wuppertal, Germany
- ³⁶Karlsruhe Institute of Technology, Institute for Experimental Particle Physics (ETP), Karlsruhe, Germany
- ³⁷Karlsruhe Institute of Technology, Institut für Kernphysik, Karlsruhe, Germany
- ³⁸Karlsruhe Institute of Technology, Institut für Prozessdatenverarbeitung und Elektronik, Karlsruhe, Germany
- ³⁹RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany
- ⁴⁰Universität Hamburg, II. Institut für Theoretische Physik, Hamburg, Germany
- ⁴¹Universität Siegen, Fachbereich 7 Physik – Experimentelle Teilchenphysik, Siegen, Germany
- ⁴²Gran Sasso Science Institute, L’Aquila, Italy
- ⁴³INFN Laboratori Nazionali del Gran Sasso, Assergi (L’Aquila), Italy
- ⁴⁴INFN, Sezione di Catania, Catania, Italy
- ⁴⁵INFN, Sezione di Lecce, Lecce, Italy
- ⁴⁶INFN, Sezione di Milano, Milano, Italy
- ⁴⁷INFN, Sezione di Napoli, Napoli, Italy
- ⁴⁸INFN, Sezione di Roma “Tor Vergata”, Roma, Italy
- ⁴⁹INFN, Sezione di Torino, Torino, Italy
- ⁵⁰Osservatorio Astrofisico di Torino (INAF), Torino, Italy
- ⁵¹Politecnico di Milano, Dipartimento di Scienze e Tecnologie Aerospaziali, Milano, Italy

- ⁵²Università del Salento, Dipartimento di Matematica e Fisica “E. De Giorgi”, Lecce, Italy
- ⁵³Università dell’Aquila, Dipartimento di Scienze Fisiche e Chimiche, L’Aquila, Italy
- ⁵⁴Università di Catania, Dipartimento di Fisica e Astronomia, Catania, Italy
- ⁵⁵Università di Milano, Dipartimento di Fisica, Milano, Italy
- ⁵⁶Università di Napoli “Federico II”, Dipartimento di Fisica “Ettore Pancini”, Napoli, Italy
- ⁵⁷Università di Roma “Tor Vergata”, Dipartimento di Fisica, Roma, Italy
- ⁵⁸Università Torino, Dipartimento di Fisica, Torino, Italy
- ⁵⁹Benemérita Universidad Autónoma de Puebla, Puebla, México
- ⁶⁰Centro de Investigación y de Estudios Avanzados del IPN (CINVESTAV), México, D.F., México
- ⁶¹Unidad Profesional Interdisciplinaria en Ingeniería y Tecnologías Avanzadas del Instituto Politécnico Nacional (UPIITA-IPN), México, D.F., México
- ⁶²Universidad Autónoma de Chiapas, Tuxtla Gutiérrez, Chiapas, México
- ⁶³Universidad Nacional Autónoma de México, México, D.F., México
- ⁶⁴Institute of Nuclear Physics PAN, Krakow, Poland
- ⁶⁵University of Łódź, Faculty of Astrophysics, Łódź, Poland
- ⁶⁶University of Łódź, Faculty of High-Energy Astrophysics, Łódź, Poland
- ⁶⁷Laboratório de Instrumentação e Física Experimental de Partículas – LIP and Instituto Superior Técnico – IST, Universidade de Lisboa – UL, Lisboa, Portugal
- ⁶⁸“Horia Hulubei” National Institute for Physics and Nuclear Engineering, Bucharest-Magurele, Romania
- ⁶⁹Institute of Space Science, Bucharest-Magurele, Romania
- ⁷⁰University Politehnica of Bucharest, Bucharest, Romania
- ⁷¹Center for Astrophysics and Cosmology (CAC), University of Nova Gorica, Nova Gorica, Slovenia
- ⁷²Experimental Particle Physics Department, J. Stefan Institute, Ljubljana, Slovenia
- ⁷³Universidad de Granada and C.A.F.P.E., Granada, Spain
- ⁷⁴Instituto Galego de Física de Altas Enerxías (IGFAE), Universidade de Santiago de Compostela, Santiago de Compostela, Spain
- ⁷⁵IMAPP, Radboud University Nijmegen, Nijmegen, The Netherlands
- ⁷⁶KVI – Center for Advanced Radiation Technology, University of Groningen, Groningen, The Netherlands
- ⁷⁷Nationaal Instituut voor Kernfysica en Hoge Energie Fysica (NIKHEF), Science Park, Amsterdam, The Netherlands
- ⁷⁸Stichting Astronomisch Onderzoek in Nederland (ASTRON), Dwingeloo, The Netherlands
- ⁷⁹Universiteit van Amsterdam, Faculty of Science, Amsterdam, The Netherlands
- ⁸⁰Case Western Reserve University, Cleveland, OH, USA
- ⁸¹Colorado School of Mines, Golden, CO, USA
- ⁸²Department of Physics and Astronomy, Lehman College, City University of New York, Bronx, NY, USA
- ⁸³Louisiana State University, Baton Rouge, LA, USA
- ⁸⁴Michigan Technological University, Houghton, MI, USA
- ⁸⁵New York University, New York, NY, USA
- ⁸⁶Pennsylvania State University, University Park, PA, USA
- ⁸⁷University of Chicago, Enrico Fermi Institute, Chicago, IL, USA
- ⁸⁸University of Delaware, Department of Physics and Astronomy, Bartol Research Institute, Newark, DE, USA
-
- ¹⁰⁰⁰School of Physics and Astronomy, University of Leeds, Leeds, United Kingdom
- ¹⁰⁰¹Max-Planck-Institut für Radioastronomie, Bonn, Germany
- ¹⁰⁰²Fermi National Accelerator Laboratory, USA
- ¹⁰⁰³also at Universidade Federal de Alfenas, Poços de Caldas, Brazil
- ¹⁰⁰⁴Colorado State University, Fort Collins, CO, USA
- ¹⁰⁰⁵now at Hakubi Center for Advanced Research and Graduate School of Science, Kyoto University, Kyoto, Japan
- ¹⁰⁰⁶also at Karlsruhe Institute of Technology, Karlsruhe, Germany
- ¹⁰⁰⁷also at Radboud University Nijmegen, Nijmegen, The Netherlands

ABSTRACT

We present measurements of the large-scale cosmic-ray anisotropies in right ascension, using data

collected by the surface detector array of the Pierre Auger Observatory over more than 14 years. We determine the equatorial dipole component, \vec{d}_\perp , through a Fourier analysis in right ascension that includes weights for each event so as to account for the main detector-induced systematic effects. For the energies at which the trigger efficiency of the array is small, the ‘‘East-West’’ method is employed. Besides using the data from the array with detectors separated by 1500 m, we also include data from the smaller but denser sub-array of detectors with 750 m separation, which allows us to extend the analysis down to ~ 0.03 EeV. The most significant equatorial dipole amplitude obtained is that in the cumulative bin above 8 EeV, $d_\perp = 6.0_{-0.9}^{+1.0}\%$, which is inconsistent with isotropy at the 6σ level. In the bins below 8 EeV, we obtain 99% CL upper-bounds on d_\perp at the level of 1 to 3 percent. At energies below 1 EeV, even though the amplitudes are not significant, the phases determined in most of the bins are not far from the right ascension of the Galactic center, at $\alpha_{GC} = -94^\circ$, suggesting a predominantly Galactic origin for anisotropies at these energies. The reconstructed dipole phases in the energy bins above 4 EeV point instead to right ascensions that are almost opposite to the Galactic center one, indicative of an extragalactic cosmic ray origin.

1. INTRODUCTION

The distribution of cosmic-ray (CR) arrival directions is expected to provide essential clues to understanding the CR origin. Being charged particles, they are significantly deflected by the magnetic fields present in our galaxy (Haverkorn 2015) and, for those arriving from outside it, also by the extragalactic magnetic fields (Feretti et al. 2012). Since the deflections get smaller for increasing rigidities, it is only at the highest energies that one may hope to observe localized flux excesses associated with individual CR sources. On the other hand, as the energies lower and the deflections become large, the propagation eventually becomes diffusive and it is likely that only large-scale patterns, such as a dipolar flux modulation, may be detectable. However, the small amplitudes of these anisotropies make their observation quite challenging.

Due to the Earth’s rotation, cosmic-ray observatories running for long periods of time have an almost uniform exposure in right ascension. This enables them to achieve a high sensitivity to the modulation of the flux in this angular coordinate. In particular, for a dipolar cosmic-ray flux the first-harmonic modulation in right ascension provides a direct measurement of the projection of the dipole in the equatorial plane, \vec{d}_\perp . The possible sources of systematic uncertainties that could affect these measurements, such as those from remaining non-uniformities of the exposure or those related to the effects of atmospheric variations, can often be accounted for. Even when this is not possible, as can happen when the trigger efficiency of the array is small, methods that are insensitive to these systematic effects can be adopted to reconstruct \vec{d}_\perp , although they have a somewhat reduced sensitivity to the modulations. On the other hand, at low energies the number of events detected is large, what tends to enhance the statistical sensitivity of the measurements.

The projection of the dipole along the Earth rotation axis d_z can, in principle, be reconstructed from the study of the azimuthal modulation of the CR fluxes. This requires accounting in detail for the effects of the geomagnetic field on the air showers, which can affect the reconstruction of the CR energies in an azimuthally dependent way. Also, the presence of a tilt of the array can induce a spurious contribution to d_z . When the trigger efficiency of the array is small, these effects may lead to systematic uncertainties that cannot be totally corrected for, particularly given the azimuthal dependence of the trigger efficiency arising from the actual geometry of the surface detector array of the Pierre Auger Observatory. Due to these limitations, we will here restrict our analysis to the determination of \vec{d}_\perp through the study of the distribution in right ascension of the events recorded in different energy bins. We note that the determination of d_z for energies $E \geq 4$ EeV, for which that detector has full efficiency for zenith angles up to 80° , was discussed in detail in The Pierre Auger Collaboration (2015a, 2017a, 2018).

At $E \geq 8$ EeV, a significant first-harmonic modulation in right ascension, corresponding to an amplitude $d_\perp \sim 6\%$, has been detected by the Pierre Auger Observatory (The Pierre Auger Collaboration 2017a). The reconstructed direction of the three-dimensional dipole suggests a predominant extragalactic origin of the CR anisotropies at energies above 4 EeV, and the dipolar amplitudes obtained in different bins show a growing trend with increasing energies (The Pierre Auger Collaboration 2017a, 2018).

The phase in right ascension of the dipolar modulation of the flux determined above 8 EeV is $\alpha_d \simeq 100^\circ$. This is nearly opposite to the phases measured at PeV energies by IceCube and IceTop (IceCube Collaboration 2012, 2016), which lie not far from the Galactic center direction which is at $\alpha_{GC} = -94^\circ$. Also the KASCADE-Grande measurements, involving CR energies from few PeV up to few tens of PeV, lead to phases lying close to the right

ascension of the Galactic center, even though the measured amplitudes are not statistically significant (KASCADE-Grande Collaboration 2019).¹ All this is in agreement with the expectation that for energies above that of the knee of the CR spectrum, which corresponds to the steepening taking place at ~ 4 PeV, the outward diffusive escape of the CRs produced in the Galaxy should give rise to a dipolar flux component having its maximum not far from the Galactic center direction. Also at energies above few EeV, where the propagation would become more rectilinear, a continuous distribution of Galactic sources should give rise to a dipolar component not far from the GC direction (The Pierre Auger Collaboration 2018). Departures from these behaviors could however result if the CR source distribution is not symmetric with respect to the Galactic center (such as in the presence of a powerful nearby CR source), in the presence of drift motions caused by the regular Galactic magnetic field components (Ptuskin et al. 1993), or when the contribution from the extragalactic component becomes sizable. Note that the expected direction of a dipole of extragalactic origin will depend on the (unknown) distribution of the CR sources and on the effects of the deflections caused by the Galactic magnetic field, as was discussed in detail in The Pierre Auger Collaboration (2018).

The change from a Galactic CR origin towards a predominantly extragalactic origin is expected to take place somewhere above the knee. More precise measurements of the large-scale anisotropies, filling the gap between the IceCube/IceTop or KASCADE-Grande measurements and the dipole determined by the Pierre Auger Observatory above 8 EeV, should provide information about this transition. In fact, although at energies below 8 EeV the reported dipolar amplitudes are not significant, indications that a change in the phase of the anisotropies in right ascension takes place around few EeV are apparent in the Pierre Auger Observatory measurements (The Pierre Auger Collaboration 2011a, 2012, 2013; Sidelnik 2013; Al Samarai 2015). One has to keep in mind in this discussion that the energy at which the total anisotropy becomes of predominantly extragalactic origin may be different from the energy at which the CR flux becomes of predominantly extragalactic origin, since the intrinsic anisotropies of each component are likely different.

We present here an update of the measurements of the large-scale anisotropies that are sensitive to the equatorial component of a dipole, for the whole energy range from ~ 0.03 EeV up to ≥ 32 EeV, covering more than three decades of energy. The results above 4 EeV are an update of those presented in The Pierre Auger Collaboration (2018), including two more years of data, corresponding to an increase in the exposure by 20%. At lower energies, we provide a major update of the latest published results (Al Samarai 2015), with 50% more exposure for the SD1500 array and twice as much for the SD750 array. At energies below 2 EeV, possible systematic effects related to the reduced trigger efficiency could be significant. To study the modulation in right ascension in this regime we have then to resort to the “East-West” method, which has larger associated uncertainties but is not affected by most of the systematic effects (Nagashima et al. 1989; Bonino et al. 2011). At energies below 0.25 EeV, it proves convenient to use the data from the sub-array of detectors with 750 m spacing which, although being much smaller, can detect a larger number of events at these energies.

2. THE OBSERVATORY AND THE DATASET

The Pierre Auger Observatory (The Pierre Auger Collaboration 2015b), located near the city of Malargüe in western Argentina (at latitude 35.2° South), is the largest existing CR observatory. Its surface detector array (SD) consists of water-Cherenkov detectors having each one 12 tonnes of ultra-pure water viewed by three 9 inch phototubes. The main array, SD1500, consists of detectors distributed on a triangular grid with separations of 1,500 m that span an area of $3,000 \text{ km}^2$. A smaller sub-array, SD750, covers an area of 23 km^2 with detectors separated by 750 m, making it sensitive also to smaller CR energies. These arrays sample the secondary particles of the air showers reaching ground level. In addition, the fluorescence detector (FD) consists of 27 telescopes that overlook the SD array. The FD can determine the longitudinal development of the air showers by observing the UV light emitted by atmospheric nitrogen molecules excited by the passage of the charged particles of the shower. This fluorescence light can be detected during clear moonless nights, with a corresponding duty cycle of about 15% (The Pierre Auger Collaboration 2015b). The SD arrays have instead a continuous operation, detecting events with a duty cycle close to 100%. They also have a more uniform (and simpler to evaluate) exposure. This is why the studies of the large-scale anisotropies that we perform here are based on the much larger number of events recorded by the surface arrays.

For the SD1500 array, the dataset considered in this work includes events with energies above 0.25 EeV that were detected from 2004 January 1 up to 2018 August 31. For energies below 4 EeV, it includes events with zenith angles up to 60° , allowing coverage of 71% of the sky, and the quality trigger applied requires that all the six detectors

¹ Hints of anisotropies on smaller angular scales were also found recently in a reanalysis of KASCADE-Grande data (Ahlers 2019).

surrounding the one with the largest signal be active at the time the event is detected. For energies above 4 EeV, more inclined events can be reliably reconstructed (The Pierre Auger Collaboration 2014b) and hence the zenith-angle range is extended up to 80° , allowing coverage of 85% of the sky. Moreover, given that at these energies the number of detectors triggered by each shower is large (4 or more detectors for more than 99% of the events), we also include in this case events passing a relaxed trigger condition, allowing that one of the six detectors that are neighbors to the one with the largest signal be missing or not functioning, provided that the reconstructed shower core be contained inside a triangle of nearby active detectors (The Pierre Auger Collaboration 2017a). The integrated exposure of the array for $\theta \leq 60^\circ$ and using the strict trigger selection is $60,700 \text{ km}^2 \text{ sr yr}$, while that for $\theta \leq 80^\circ$ and relaxing the trigger is $92,500 \text{ km}^2 \text{ sr yr}$.

The CR arrival directions are reconstructed from the timing of the signals in the different triggered stations, and the angular resolution is better than 1.6° (The Pierre Auger Collaboration 2015b), so that it has negligible impact on the reconstruction of the dipole. The energies of the events with $\theta \leq 60^\circ$ are assigned in terms of the reconstructed signals at a reference distance from the shower core of 1000 m. They are corrected for atmospheric effects, accounting for the pressure and air density variations following The Pierre Auger Collaboration (2017b), as well as for geomagnetic effects, following The Pierre Auger Collaboration (2011b). The inclined events, whose signals are dominantly produced by the muonic component of the showers, have a negligible dependence on atmospheric variations, while geomagnetic effects are already taken into account in their reconstruction (The Pierre Auger Collaboration 2014b). Their energies are assigned in terms of the estimated muon number at ground level. The SD1500 array has full trigger efficiency for $E \geq 2.5 \text{ EeV}$ if one considers events with $\theta \leq 60^\circ$, and for $E \geq 4 \text{ EeV}$ for events with $\theta \leq 80^\circ$. The energies of the CRs are calibrated using the hybrid events measured simultaneously by the SD and FD detectors, in the regimes of full trigger efficiency. For lower energies, in which case we consider events with $\theta \leq 60^\circ$, the energy assignment is performed using the extrapolation of the corresponding calibration curve. The energy resolution for events with $\theta \leq 60^\circ$ is about 7% above 10 EeV, and degrades for lower energies, reaching about 20% at 1 EeV, while the systematic uncertainty in the energy scale is 14% (see Verzi (2019) for details). The more inclined events have an energy resolution of 19%, with a similar systematic uncertainty (The Pierre Auger Collaboration 2014b).

For energies below 0.25 EeV, and down to $\sim 0.03 \text{ EeV}$ (below which the trigger efficiency is tiny), we use the events from the denser and smaller SD750 array, since the accumulated statistics is larger. The dataset comprises events with zenith angles up to 55° detected from 2012 January 1 up to 2018 August 31. The trigger applied requires that all six detectors around the one with the largest signal be functioning and the associated exposure is $234 \text{ km}^2 \text{ sr yr}$. The energies are assigned in terms of the reconstructed signals at a reference distance from the shower core of 450 m. They are corrected for atmospheric effects following The Pierre Auger Collaboration (2017b). The SD750 array has full trigger efficiency for $E \geq 0.3 \text{ EeV}$ if one considers events with $\theta \leq 55^\circ$ (The Pierre Auger Collaboration 2015b). The energies are calibrated with hybrid events observed in the regime of full trigger efficiency and below that threshold the energy assignment is performed on the basis of the extrapolation of the corresponding calibration curve. At 0.3 EeV the energy resolution is about 18% (Coleman 2019).

3. THE ANALYSIS METHOD

The weighted first-harmonic analysis in the right ascension angle α , often referred to as Rayleigh analysis, provides the Fourier coefficients as

$$a = \frac{2}{\mathcal{N}} \sum_{i=1}^N w_i \cos \alpha_i, \quad b = \frac{2}{\mathcal{N}} \sum_{i=1}^N w_i \sin \alpha_i, \quad (1)$$

where the sums run over all N detected events. The weights w_i , which are of order unity, account for the effects of the non-uniformities in the exposure as a function of time, with the normalization factor being $\mathcal{N} \equiv \sum_i w_i$. The amplitude and phase of the first-harmonic modulation are given by $r = \sqrt{a^2 + b^2}$ and $\varphi = \arctan(b/a)$. The probability to obtain an amplitude larger than the one measured as a result of a fluctuation from an isotropic distribution is $P(\geq r) = \exp(-\mathcal{N}r^2/4)$. To obtain the weights, we permanently monitor the number of active unitary detector cells, corresponding to the number of active detectors that are surrounded by an hexagon of working detectors or, when considering the relaxed trigger condition above 4 EeV, we also account for detector configurations with only five active detectors around the central one. We obtain from this the exposure of the Auger Observatory in bins of right ascension of the zenith of the array, α^0 . This angle is given by $\alpha^0(t_i) \equiv 2\pi t_i/T_s \pmod{2\pi}$, with the origin of time being taken such that $\alpha^0(0) = 0$. The sidereal-time period, $T_s \simeq 23.934 \text{ h}$, corresponds to one extra cycle per year with respect to the solar frequency. The fraction of the total exposure that is associated to a given α^0 bin, taken to be of 1.25° width (5 minutes), is proportional to the total number of unitary cells in that bin, $N_{\text{cell}}(\alpha^0)$. The weights w_i account for the

relative variations of N_{cell} as a function of α^0 , i.e.

$$w_i = \left(\frac{N_{\text{cell}}(\alpha^0(t_i))}{\langle N_{\text{cell}} \rangle} \right)^{-1}, \quad (2)$$

with $\langle N_{\text{cell}} \rangle = 1/(2\pi) \int_0^{2\pi} d\alpha^0 N_{\text{cell}}(\alpha^0)$. Including these weights in the Fourier coefficients eliminates the spurious contribution to the amplitudes associated to the non-uniform exposure in right ascension.

We note that if one were to consider periods of only a few months, the resulting modulation of $N_{\text{cell}}(\alpha^0)$ could amount to an effect of a few percent on the modulation in right ascension of the event rates. However, after considering several years, the modulations that appear on shorter time scales tend to get averaged out, with the surviving effects being now typically at the level of about $\pm 0.5\%$. The effects of the tilt of the SD array ([The Pierre Auger Collaboration 2012](#)), which is inclined on average by $\sim 0.2^\circ$ towards $\phi \simeq -30^\circ$ (i.e. towards the South-East), can also be accounted for by adding an extra factor in the weights ([The Pierre Auger Collaboration 2018](#)). However, this is actually only relevant when performing the Fourier analysis in the azimuth variable ϕ , something we will not perform here.

When the triggering of the array is not fully efficient, there are additional systematic effects related to the interplay between the atmospheric effects in the air-shower development and the energy-dependent trigger efficiency. In particular, changes in the air density modify the Molière radius determining the lateral spread of the electromagnetic component of the showers. The fall-off of the signal at ground level is preferentially harder under hot weather conditions and steeper under cold ones. The detection efficiency of the SD is thus expected to follow these variations to some extent, being on average larger when the weather is hot than when it is cold. As a consequence, one could expect that, at energies below full trigger efficiency, a spurious modulation could appear at the solar frequency.

Moreover, we have found that the amplitude of the modulation of the rates at the antisidereal frequency, which is that corresponding to one cycle less per year than the solar frequency, suggests that spurious unaccounted effects become relevant below 2 EeV. In particular, the Fourier amplitude corresponding to the antisidereal time period $T_{\text{as}} = 24.066$ h in the bin [1, 2] EeV is $r = 0.005$. This has a probability of arising as a fluctuation of less than 0.1%. A non-negligible antisidereal amplitude could for instance appear in the presence of daily and seasonal systematic effects which are not totally accounted for. Since in this case comparable spurious amplitudes could be expected in the sidereal and antisidereal sidebands ([Farley & Storey 1954](#)), we only use the Rayleigh method described before in the bins above 2 EeV. We have checked that in the bins above 2 EeV the amplitudes at both the solar and antisidereal frequencies are consistent with being just due to fluctuations, so that there are no signs indicating that surviving systematic effects could be present at the sidereal frequency at these energies (see Table 1 in the Appendix).² Alternatively, one can use for the energies below 2 EeV the differential East-West (EW) method ([Bonino et al. 2011](#)), which is based on the difference between the counting rates of the events measured from the East sector and those from the West sector. Since the exposure is the same for events coming from the East and for those coming from the West³, and also the spurious modulations due to the atmospheric effects are the same in both sectors, the relative difference between both rates, $(E - W)/(E + W)$, is not sensitive to these experimental and atmospheric systematic effects. This allows one to reconstruct in a clean way the modulation of the rate itself, without the need to apply any correction but at the expense of a reduced sensitivity to the amplitude of the CR flux modulations.

In this approach ([Bonino et al. 2011](#)), the first-harmonic amplitude and phase are calculated using a slightly modified Fourier analysis that accounts for the subtraction of the Western sector from the Eastern one. The Fourier coefficients are defined as

$$a_{\text{EW}} = \frac{2}{N} \sum_{i=1}^N \cos(\alpha^0(t_i) - \xi_i), \quad b_{\text{EW}} = \frac{2}{N} \sum_{i=1}^N \sin(\alpha^0(t_i) - \xi_i), \quad (3)$$

where $\xi_i = 0$ for events coming from the East and $\xi_i = \pi$ for those coming from the West, so as to easily implement the subtraction of data from the two hemispheres.

In the case in which the dominant contribution to the flux modulation is purely dipolar, the amplitude $r_{\text{EW}} = \sqrt{a_{\text{EW}}^2 + b_{\text{EW}}^2}$ and phase $\varphi_{\text{EW}} = \arctan(b_{\text{EW}}/a_{\text{EW}})$ obtained with this method are related to the ones from the Rayleigh formalism through $r = \frac{\pi \langle \cos \delta \rangle}{2 \langle \sin \theta \rangle} r_{\text{EW}}$ and $\varphi = \varphi_{\text{EW}} + \pi/2$, where $\langle \cos \delta \rangle$ is the average of the cosine of the declination of the events and similarly $\langle \sin \theta \rangle$ is the average of the sine of their zenith angles ([Bonino et al. 2011](#)). The probability to obtain an amplitude larger than the one measured as a result of a fluctuation from an isotropic

² Given that, for events with zenith angles smaller than 60° , the trigger efficiency is larger than $\sim 95\%$ above 2 EeV, the efficiency related systematic effects are negligible above this threshold.

³ A possible tilt of the array in the East-West direction, giving just a constant term in the East-West rate difference, does not affect the determination of the first-harmonic modulation.




Figure 1. Reconstructed equatorial-dipole amplitude (left) and phase (right). The upper limits at 99% CL are shown for all the energy bins in which the measured amplitude has a chance probability greater than 1%. The gray bands indicate the amplitude and phase for the energy bin $E \geq 8$ EeV. Results from other experiments are shown for comparison (IceCube Collaboration 2012, 2016; KASCADE-Grande Collaboration 2019).

distribution is $P(\geq r_{EW}) = \exp(-Nr_{EW}^2/4)$.

The amplitude of the equatorial dipole component is related to the amplitude of the first-harmonic modulation through $d_{\perp} \simeq r/\langle \cos \delta \rangle$, and its phase α_d coincides with the first-harmonic phase φ .

4. RIGHT ASCENSION MODULATION FROM 0.03 EeV UP TO $E \geq 32$ EeV

In Table 1, we report the results for the reconstructed equatorial dipole in different energy bins, covering the range from ~ 0.03 EeV up to $E \geq 32$ EeV. The energies defining the boundaries of the bins are 2^n EeV, with $n = -5, -4, \dots, 4, 5$. As mentioned previously, the results are obtained from the study of the right ascension modulation using different methods and datasets. We use the weighted Rayleigh analysis in the energy bins above 2 EeV, for which the systematic effects associated with the non-saturated detector efficiency and to the effects related to atmospheric variations are well under control. When this is not the case, we report the results of the East-West method which, although having larger uncertainties, is quite insensitive to most sources of systematic effects in the right ascension distribution. For energies above 0.25 EeV, we report the results obtained with the data from the SD1500 array, while for lower energies we use the dataset from the SD750 array since, having a lower threshold, it leads to a larger number of events despite the reduced size of the array. In that case, given that the SD750 array is not fully efficient below 0.3 EeV, we just use the East-West method.

| | E [EeV] | E_{med} [EeV] | N | d_{\perp} [%] | $\sigma_{x,y}$ [%] | α_d [°] | $P(\geq d_{\perp})$ | d_{\perp}^{UL} [%] |
|-----------------------|-------------|------------------------|-----------|------------------------|--------------------|----------------|----------------------|-----------------------------|
| East-West (SD750) | 1/32 - 1/16 | 0.051 | 432,155 | $1.0^{+1.0}_{-0.4}$ | 0.91 | 112 ± 71 | 0.54 | 3.3 |
| | 1/16 - 1/8 | 0.088 | 924,856 | $0.6^{+0.6}_{-0.3}$ | 0.52 | -44 ± 68 | 0.50 | 2.0 |
| | 1/8 - 1/4 | 0.161 | 488,752 | $0.2^{+0.8}_{-0.2}$ | 0.63 | -31 ± 108 | 0.94 | 2.0 |
| East-West (SD1500) | 1/4 - 1/2 | 0.43 | 770,316 | $0.6^{+0.5}_{-0.3}$ | 0.48 | -135 ± 64 | 0.45 | 1.8 |
| | 1/2 - 1 | 0.70 | 2,388,467 | $0.5^{+0.3}_{-0.2}$ | 0.27 | -99 ± 43 | 0.20 | 1.1 |
| | 1 - 2 | 1.28 | 1,243,103 | $0.18^{+0.47}_{-0.02}$ | 0.35 | -69 ± 100 | 0.87 | 1.1 |
| Rayleigh (SD1500) | 2 - 4 | 2.48 | 283,074 | $0.5^{+0.4}_{-0.2}$ | 0.34 | -11 ± 55 | 0.34 | 1.4 |
| | 4 - 8 | 5.1 | 88,325 | $1.0^{+0.7}_{-0.4}$ | 0.61 | 69 ± 46 | 0.23 | 2.6 |
| | 8 - 16 | 10.3 | 27,271 | $5.6^{+1.2}_{-1.0}$ | 1.1 | 97 ± 12 | 2.3×10^{-6} | – |
| | 16 - 32 | 20.3 | 7,664 | $7.5^{+2.3}_{-1.8}$ | 2.1 | 80 ± 17 | 1.5×10^{-3} | – |
| | ≥ 32 | 40 | 1,993 | 13^{+5}_{-3} | 4.1 | 152 ± 19 | 5.3×10^{-3} | – |
| | ≥ 8 | 11.5 | 36,928 | $6.0^{+1.0}_{-0.9}$ | 0.94 | 98 ± 9 | 1.4×10^{-9} | – |

Table 1. Equatorial dipole reconstruction in different energy bins. Indicated are the median energies in each bin E_{med} , number of events N , amplitude of d_{\perp} , uncertainty $\sigma_{x,y} = \sigma/\langle \cos \delta \rangle$ of the components d_x or d_y , right ascension phase, probability to get a larger amplitude from fluctuations of an isotropic distribution and 99% CL upper limit on the amplitude.

For each energy bin, we report in Table 1 the number of events N , the amplitude d_{\perp} , the uncertainty $\sigma_{x,y}$ of

the components d_x or d_y , the right ascension phase of the dipolar modulation α_d , the chance probability $P(\geq d_\perp)$ and, when the measured amplitude has a probability larger than 1%, we also report the 99% CL upper limit on the amplitude of the equatorial dipole d_\perp^{UL} . The upper limits on the first-harmonic amplitude at a given confidence level CL (CL = 0.99 for 99% CL) are derived from the distribution for a dipolar anisotropy of unknown amplitude, marginalized over the dipole phase, requiring that

$$\int_0^{r^{\text{UL}}} dr \frac{r}{\sigma^2} \exp\left[-\frac{r^2 + s^2}{2\sigma^2}\right] I_0\left(\frac{rs}{\sigma^2}\right) = \text{CL}, \quad (4)$$

with $I_0(x)$ the zero-order modified Bessel function, s the measured amplitude and the dispersion being $\sigma = \sqrt{2/N}$ for the Rayleigh analysis while $\sigma = (\pi\langle\cos\delta\rangle/2\langle\sin\theta\rangle)\sqrt{2/N}$ for the East-West method. These bounds on the first-harmonic amplitude are then converted into the corresponding upper limit for the amplitude of the equatorial dipole using that $d_\perp^{\text{UL}} = r^{\text{UL}}/\langle\cos\delta\rangle$. For the uncertainties in the phase, we use the two-dimensional distribution marginalized instead over the dipole amplitude r (Linsley 1975). In Table 2 in the Appendix we also report the results obtained above 2 EeV with the East-West method, which are consistent with those obtained with the Fourier analysis in Table 1 but have larger uncertainties.

Fig. 1 shows the equatorial dipole amplitude (left panel) and phase (right panel) that were determined in all the energy bins considered, as reported in Table 1. Also shown are the results obtained by the IceCube, IceTop and KASCADE-Grande experiments in the 1–30 PeV range (IceCube Collaboration 2012, 2016; KASCADE-Grande Collaboration 2019). We also show the 99% CL upper limit d_\perp^{UL} in the cases in which the measured amplitude has more than 1% probability to be a fluctuation from an isotropic distribution. The results for the integral bin with $E \geq 8$ EeV, that was considered in The Pierre Auger Collaboration (2017a), is shown as a gray band.

A trend of increasing amplitudes for increasing energies is observed, with values going from $d_\perp \simeq 0.1\%$ at PeV energies, to $\sim 1\%$ at EeV energies and reaching $\sim 10\%$ at 30 EeV. Regarding the phases, a transition between values lying close to the right ascension of the Galactic center, $\alpha_d \simeq \alpha_{\text{GC}}$, towards values in a nearly opposite direction, $\alpha_d \simeq 100^\circ$, is observed to take place around a few EeV.




Figure 2. Components of the dipole in the equatorial plane for different energy bins above 0.25 EeV (left panel) and below 1 EeV (right panel). The horizontal axis corresponds to the component along the direction $\alpha = 0$ while the vertical axis to that along $\alpha = 90^\circ$. The radius of each circle corresponds to the 1σ uncertainty in d_x and d_y . The Galactic center direction is also indicated. The measurements from IceCube (IC) and IceTop (IT) at PeV energies are also indicated in the right panel (IceCube Collaboration 2012, 2016).

The overall behavior of the amplitudes and phases in the d_x – d_y plane is depicted in Fig. 2. The left panel includes the energies above 0.25 EeV while the right panel those below 1 EeV. In these plots, the right ascension α_d is the polar angle, measured anti-clockwise from the x-axis (so that $d_x = d_\perp \cos\alpha_d$ and $d_y = d_\perp \sin\alpha_d$). The circles shown have a radius equal to the 1σ uncertainties $\sigma_{x,y}$ in the dipole components $d_{x,y}$ (reported in the Table 1), effectively including $\sim 39\%$ of the two-dimensional confidence region. One can appreciate in this plot how the amplitudes decrease for

decreasing energies, and how the phases change as a function of the energy, pointing almost in the opposite direction of the Galactic center above 4 EeV and not far from it below 1 EeV.

The values of the anisotropy parameters obtained above are based, by construction, on the event content in the energy intervals under scrutiny. The finite resolution on the energies induces bin-to-bin migration of events. Due to the steepness of the energy spectrum, the migration happens especially from lower to higher energy bins. This influences the energy dependence of the recovered parameters. However, given that the size of the energy bins chosen here is much larger than the resolution, the migration of events remains small enough to avoid significant distortions for the recovered values above full efficiency. For instance, given the energy resolution of the SD1500 array (Verzi 2019) and assuming a dipole amplitude scaling as $E^{0.8}$, as was found in The Pierre Auger Collaboration (2018) to approximately hold above 4 EeV, the impact of the migrations remains below an order of magnitude smaller than the statistical uncertainties associated to the recovered parameters. In the energy range below full efficiency, additional systematic effects enter into play on the energy estimate. We note that forward-folding simulations of the response function effects into an injected anisotropy show that the recovered parameters are not impacted by more than their current statistical uncertainties. A complete unfolding of these effects is left for future studies. It requires an accurate knowledge of the response function of the SD arrays down to low energies, which is not available at the moment.

5. DISCUSSION AND CONCLUSIONS

We have updated the searches for anisotropies on large angular scales using the cosmic rays detected by the Pierre Auger Observatory. The analysis covered more than three orders of magnitude in energy, including events with $E \geq 0.03$ EeV and hence encompassing the expected transition between Galactic and extragalactic origins of the cosmic rays. This was achieved by studying the first-harmonic modulation in right ascension of the CR fluxes determined with the SD1500 and the SD750 surface detector arrays. This allowed us to determine the equatorial component of a dipolar modulation, \vec{d}_\perp , or eventually to set strict upper-bounds on it.

For the inclusive bin above 8 EeV, the first-harmonic modulation in right ascension leads to an equatorial dipole amplitude $d_\perp = 0.060_{-0.009}^{+0.010}$, which has a probability to arise by chance from an isotropic distribution of 1.4×10^{-9} , corresponding to a two-sided Gaussian significance of 6σ . The phase of the maximum of this modulation is at $\alpha_d = 98^\circ \pm 9^\circ$, indicating an extragalactic origin for these CRs. When splitting the bin above 8 EeV, as originally done in The Pierre Auger Collaboration (2018), one finds indications of an increasing amplitude with increasing energies, and the direction of the dipole suggests that it has an extragalactic origin in all the three bins considered. A growing dipole amplitude for increasing energies could for instance be associated with the larger relative contribution to the flux that arises at high energies from nearby sources, that are more anisotropically distributed than the integrated flux from the distant ones. A suppression of the more isotropic contribution from distant sources is expected to result from the strong attenuation of the CR flux that should take place at the highest energies as a consequence of their interactions with the background radiation (Greisen 1966; Zatsepin & Kuzmin 1966).

At energies below 8 EeV, none of the amplitudes are significant, and we set 99% CL upper bounds on d_\perp at the level of 1 to 3%. The phases measured in most of the bins below 1 EeV are not far from the direction towards the Galactic center. All this suggests that the origin of these dipolar anisotropies changes from a predominantly Galactic one to an extragalactic one somewhere in the range between 1 EeV and few EeV. The small size of the dipolar amplitudes in this energy range, combined with the indications that the composition is relatively light (The Pierre Auger Collaboration 2014a), disfavor a predominant flux component of Galactic origin at $E > 1$ EeV (The Pierre Auger Collaboration 2013). Models of Galactic CRs relying on a mixed mass composition, with rigidity dependent spectra, have been proposed to explain the knee (at ~ 4 PeV) and second-knee (at ~ 0.1 EeV) features in the spectrum (Candia et al. 2003). The predicted anisotropies depend on the details of the Galactic magnetic field model considered and, below 0.5 EeV, they are consistent with the upper bounds we obtained. An extrapolation of these models, considering that there is no cutoff in the Galactic component, would predict dipolar anisotropies at the several percent level beyond the EeV, in tension with the upper bounds in this range. The conflict is even stronger for Galactic models (Calvez et al. 2010) having a light CR composition that extends up to the ankle energy (at ~ 5 EeV). The presence of a more isotropic extragalactic component making a significant contribution already at EeV energies could dilute the anisotropy of Galactic origin, so as to be consistent with the bounds obtained. Note that even if the extragalactic component were completely isotropic in some reference frame, the motion of the Earth with respect to that system could give rise to a dipolar anisotropy through the Compton-Getting effect (Compton & Getting 1935). For instance, for a CR distribution that is isotropic in the CMB rest frame, the resulting Compton-Getting dipole amplitude would be about 0.6% (Kachelriess & Serpico 2006). This amplitude depends on the relative velocity and on the CR spectral slope, but not directly on the particle charge. The deflections of the extragalactic CRs caused by the Galactic magnetic

field are expected to further reduce this amplitude, and also to generate higher harmonics, in a rigidity dependent way, so that the exact predictions are model dependent. The Compton-Getting extragalactic contribution to the dipolar anisotropy is hence below the upper limits obtained.

More data, as well as analyses exploiting the discrimination between the different cosmic-ray mass components that will become feasible with the upgrade of the Pierre Auger Observatory currently being implemented (Castellina 2019), will be crucial to understand in depth the origin of the cosmic rays at these energies and to learn how their anisotropies are produced.

ACKNOWLEDGMENTS

The successful installation, commissioning, and operation of the Pierre Auger Observatory would not have been possible without the strong commitment and effort from the technical and administrative staff in Malargüe. We are very grateful to the following agencies and organizations for financial support:

Argentina – Comisión Nacional de Energía Atómica; Agencia Nacional de Promoción Científica y Tecnológica (ANPCyT); Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET); Gobierno de la Provincia de Mendoza; Municipalidad de Malargüe; NDM Holdings and Valle Las Leñas; in gratitude for their continuing cooperation over land access; Australia – the Australian Research Council; Brazil – Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq); Financiadora de Estudos e Projetos (FINEP); Fundação de Amparo à Pesquisa do Estado de Rio de Janeiro (FAPERJ); São Paulo Research Foundation (FAPESP) Grants No. 2010/07359-6 and No. 1999/05404-3; Ministério da Ciência, Tecnologia, Inovações e Comunicações (MCTIC); Czech Republic – Grant No. MSMT CR LTT18004, LO1305, LM2015038 and CZ.02.1.01/0.0/0.0/16.013/0001402; France – Centre de Calcul IN2P3/CNRS; Centre National de la Recherche Scientifique (CNRS); Conseil Régional Ile-de-France; Département Physique Nucléaire et Corpusculaire (PNC-IN2P3/CNRS); Département Sciences de l’Univers (SDU-INSU/CNRS); Institut Lagrange de Paris (ILP) Grant No. LABEX ANR-10-LABX-63 within the Investissements d’Avenir Programme Grant No. ANR-11-IDEX-0004-02; Germany – Bundesministerium für Bildung und Forschung (BMBF); Deutsche Forschungsgemeinschaft (DFG); Finanzministerium Baden-Württemberg; Helmholtz Alliance for Astroparticle Physics (HAP); Helmholtz-Gemeinschaft Deutscher Forschungszentren (HGF); Ministerium für Innovation, Wissenschaft und Forschung des Landes Nordrhein-Westfalen; Ministerium für Wissenschaft, Forschung und Kunst des Landes Baden-Württemberg; Italy – Istituto Nazionale di Fisica Nucleare (INFN); Istituto Nazionale di Astrofisica (INAF); Ministero dell’Istruzione, dell’Università e della Ricerca (MIUR); CETEMPS Center of Excellence; Ministero degli Affari Esteri (MAE); México – Consejo Nacional de Ciencia y Tecnología (CONACYT) No. 167733; Universidad Nacional Autónoma de México (UNAM); PAPIIT DGAPA-UNAM; The Netherlands – Ministry of Education, Culture and Science; Netherlands Organisation for Scientific Research (NWO); Dutch national e-infrastructure with the support of SURF Cooperative; Poland -Ministry of Science and Higher Education, grant No. DIR/WK/2018/11; National Science Centre, Grants No. 2013/08/M/ST9/00322, No. 2016/23/B/ST9/01635 and No. HARMONIA 5–2013/10/M/ST9/00062, UMO-2016/22/M/ST9/00198; Portugal – Portuguese national funds and FEDER funds within Programa Operacional Factores de Competitividade through Fundação para a Ciência e a Tecnologia (COMPETE); Romania – Romanian Ministry of Research and Innovation CNCS/CCCDI-UESFISCDI, projects PN-III-P1-1.2-PCCDI-2017-0839/19PCCDI/2018 and PN18090102 within PNCDI III; Slovenia – Slovenian Research Agency, grants P1-0031, P1-0385, I0-0033, N1-0111; Spain – Ministerio de Economía, Industria y Competitividad (FPA2017-85114-P and FPA2017-85197-P), Xunta de Galicia (ED431C 2017/07), Junta de Andalucía (SOMM17/6104/UGR), Feder Funds, RENATA Red Nacional Temática de Astropartículas (FPA2015-68783-REDT) and María de Maeztu Unit of Excellence (MDM-2016-0692); USA – Department of Energy, Contracts No. DE-AC02-07CH11359, No. DE-FR02-04ER41300, No. DE-FG02-99ER41107 and No. DE-SC0011689; National Science Foundation, Grant No. 0450696; The Grainger Foundation; Marie Curie-IRSES/EPLANET; European Particle Physics Latin American Network; and UNESCO.

APPENDIX

In Table 1 we report the amplitudes and probabilities obtained with the SD1500 array at the solar and antisidereal frequencies, in all bins above 2 EeV for which the Rayleigh analysis was applied at the sidereal frequency. One can see that all these amplitudes are consistent with being fluctuations, showing then no signs of remaining systematic effects. We also report in Table 2 the equatorial dipole amplitudes and phases obtained with the East-West method above 2 EeV, and compare them with the results for the same datasets that were obtained with the Rayleigh method (reported in Table 1). The inferred equatorial dipole amplitudes turn out to be consistent, although the statistical uncertainty obtained with the East-West method is larger by a factor $\pi\langle\cos\delta\rangle/2\langle\sin\theta\rangle$ (Bonino et al. 2011). Given

that above full trigger efficiency one has that $\langle \sin \theta \rangle \simeq 0.58$ when considering $\theta < 60^\circ$, as we do for $E < 4$ EeV, or $\langle \sin \theta \rangle \simeq 0.65$ when considering $\theta < 80^\circ$, as we do for $E \geq 4$ EeV, and that $\langle \cos \delta \rangle \simeq 0.78$ in both zenith ranges, the statistical uncertainties obtained in the East-West analysis are larger by a factor of about 2.1 than those obtained with the Rayleigh analysis for $\theta < 60^\circ$, or by a factor of about 1.9 for $\theta < 80^\circ$, as can be seen in Table 2.

| E [EeV] | N | solar | | antisidereal | |
|-----------|---------|---------------------|-------------|---------------------|-------------|
| | | r [%] | $P(\geq r)$ | r [%] | $P(\geq r)$ |
| 2 - 4 | 283,074 | $0.6^{+0.3}_{-0.2}$ | 0.07 | $0.5^{+0.3}_{-0.2}$ | 0.20 |
| 4 - 8 | 88,325 | $0.8^{+0.5}_{-0.3}$ | 0.24 | $0.5^{+0.5}_{-0.2}$ | 0.59 |
| 8 - 16 | 27,271 | $0.6^{+1.1}_{-0.2}$ | 0.79 | $0.5^{+1.1}_{-0.1}$ | 0.83 |
| 16 - 32 | 7,664 | $1.1^{+2.0}_{-0.3}$ | 0.79 | $3.1^{+1.9}_{-1.1}$ | 0.16 |
| ≥ 32 | 1,993 | $1.5^{+4.4}_{-0.1}$ | 0.90 | $1.3^{+4.6}_{-0.0}$ | 0.92 |
| ≥ 8 | 36,928 | $0.3^{+1.1}_{-0.0}$ | 0.93 | $1.0^{+0.8}_{-0.4}$ | 0.39 |

Table 1. Fourier amplitudes at the solar and antisidereal frequencies, and the probabilities to get larger values from statistical fluctuations of an isotropic distribution, for the different energy bins above 2 EeV.

| E [EeV] | N | East-West (SD1500) | | | | Rayleigh (SD1500) | | |
|-----------|---------|---------------------|--------------------|----------------|----------------------|---------------------|--------------------|----------------|
| | | d_\perp [%] | $\sigma_{x,y}$ [%] | α_d [°] | $P(\geq d_\perp)$ | d_\perp [%] | $\sigma_{x,y}$ [%] | α_d [°] |
| 2 - 4 | 283,074 | $0.2^{+0.9}_{-0.2}$ | 0.72 | -16 ± 167 | 0.94 | $0.5^{+0.4}_{-0.2}$ | 0.34 | -11 ± 55 |
| 4 - 8 | 88,325 | $1.7^{+1.3}_{-0.7}$ | 1.1 | 41 ± 38 | 0.33 | $1.0^{+0.7}_{-0.4}$ | 0.61 | 69 ± 46 |
| 8 - 16 | 27,271 | $6.4^{+2.3}_{-1.7}$ | 2.1 | 147 ± 18 | 8.3×10^{-3} | $5.6^{+1.2}_{-1.0}$ | 1.1 | 97 ± 12 |
| 16 - 32 | 7,664 | $9.3^{+4.5}_{-3.0}$ | 3.9 | 67 ± 24 | 5.8×10^{-2} | $7.5^{+2.3}_{-1.8}$ | 2.1 | 80 ± 17 |
| ≥ 32 | 1,993 | 25^{+9}_{-6} | 7.6 | 151 ± 17 | 4.1×10^{-3} | 13^{+5}_{-3} | 4.1 | 152 ± 19 |
| ≥ 8 | 36,928 | $6.6^{+2.0}_{-1.5}$ | 1.8 | 132 ± 15 | 8.6×10^{-4} | $6.0^{+1.0}_{-0.9}$ | 0.94 | 98 ± 9 |

Table 2. Equatorial dipole reconstruction above 2 EeV obtained using the East-West method. Indicated are the number of events, amplitude of d_\perp , uncertainty $\sigma_{x,y} = \sigma/(\cos \delta)$ of the components d_x or d_y , right ascension phase α_d and probability $P(\geq d_\perp)$ to get a larger amplitude from fluctuations of an isotropic distribution. For comparison we also include in the last two columns the values of d_\perp and α_d that were obtained in the Rayleigh analysis (reported in Table 1).

REFERENCES

- Ahlers, M. 2019, *ApJL*, 886, L18
Al Samarai, I. for the Pierre Auger Collaboration 2016, *PoS ICRC2015*, 372
Bonino, R. et al. 2011, *ApJ*, 738, 67
Candia, J., Mollerach, S. and Roulet, E. 2003, *JCAP*, 05, 003
Calvez, A., Kusenko, A. and Nagataki, S. 2010, *PhRvL*, 105, 091101
Castellina, A. for the Pierre Auger Collaboration 2019, *EPJ Web Conf.*, 210, 06002
Coleman, A. for the Pierre Auger Collaboration 2019, *PoS ICRC2019*, 225
Compton, A.H. and Getting, I.A. 1935, *PhRv*, 47, 817
Farley, F.J.M. and Storey, J.R. 1954, *Proc. Phys. Soc. A*, 67, 996
Feretti, L. et al. 2012, *A&A Rv*, 20, 54
Greisen, K. 1966, *PhRvL*, 16, 748
Haverkorn, M. 2015, *Astrophys. Space Sci. Library*, 407, 483
IceCube Collaboration 2012, *ApJ*, 746, 33
IceCube Collaboration 2016, *ApJ*, 826, 220
Kachelriess, M. and Serpico, P.D. 2006, *Phys. Lett. B*, 640, 225
KASCADE-Grande Collaboration 2019, *ApJ*, 870, 91
Linsley, J. 1975, *PhRvL*, 34, 1530
Nagashima, K. et al. 1989, *Il Nuovo Cimento C*, 12, 695
Ptuskin, V.S. et al. 1993, *A&A*, 268, 726
Sidelnik, I. for the Pierre Auger Collaboration 2013, *Proceeding of the 33rd ICRC*, arXiv:1307.5059
The Pierre Auger Collaboration 2011a, *Astropart. Phys.*, 34, 627
The Pierre Auger Collaboration 2011b, *JCAP*, 11, 022
The Pierre Auger Collaboration 2012, *ApJS*, 203, 34
The Pierre Auger Collaboration 2013, *ApJL*, 762, L13
The Pierre Auger Collaboration 2014a, *PhRvD*, 90, 122006
The Pierre Auger Collaboration 2014b, *JCAP*, 08, 019
The Pierre Auger Collaboration 2015a, *ApJ*, 802, 111
The Pierre Auger Collaboration 2015b, *NIM A*, 798, 172
The Pierre Auger Collaboration 2017a, *Science*, 357, 1266
The Pierre Auger Collaboration 2017b, *JINST*, 12, P02006
The Pierre Auger Collaboration 2018, *ApJ*, 868, 4
Verzi, V. for the Pierre Auger Collaboration 2019, *PoS ICRC2019*, 450
Zatsepin, G.T. and Kuzmin, V.A. 1966, *JETP Lett.*, 4, 78