

## ANATOMY/PHYSIOLOGY

## G-spot: Fact or Fiction?: A Systematic Review



Pedro Vieira-Baptista, MD,<sup>1,2,3</sup> Joana Lima-Silva, MD,<sup>1,2</sup> Mario Preti, MD,<sup>4</sup> Joana Xavier, MD,<sup>5</sup> Pedro Vendeira, PhD,<sup>6</sup> and Colleen K. Stockdale, MD<sup>7</sup>

## ABSTRACT

**Introduction:** The G-spot, a putative erogenous area in the anterior vaginal wall, is a widely accepted concept in the mainstream media, but controversial in medical literature.

**Aim:** Review of the scientific data concerning the existence, location, and size of the G-spot.

**Methods:** Search on Pubmed, Pubmed Central, Cochrane, clinicaltrials.gov and Google Scholar from inception to November 2020 of studies on G-spot's existence, location and nature. Surveys, clinical, physiological, imaging, histological and anatomic studies were included.

**Main Outcome Measure:** Existence, location, and nature of the G-spot.

**Results:** In total, 31 eligible studies were identified: 6 surveys, 5 clinical, 1 neurophysiological, 9 imaging, 8 histological/anatomical, and 2 combined clinical and histological. Most women (62.9%) reported having a G-spot and it was identified in most clinical studies (55.4% of women); in 2 studies it was not identified in any women. Imaging studies had contradictory results in terms of its existence and nature. Some showed a descending of the anterior vaginal wall, that led to the concept of clitorurethrovaginal complex. In anatomic studies, one author could systematically identify the G-spot, while another group did not find it. Studies on innervation of the vaginal walls did not systematically identify an area with richer innervation.

**Conclusion:** The different studies did systematically agree on the existence of the G-spot. Among the studies in which it was considered to exist, there was no agreement on its location, size, or nature. The existence of this structure remains unproved. **Vieira-Baptista P, Lima-Silva J, Preti M, et al. G-spot: Fact or Fiction?: A Systematic Review. Sex Med 2021;XX:XXXXXX.**

Copyright © 2021 The Authors. Published by Elsevier Inc. on behalf of the International Society for Sexual Medicine. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

**Key Words:** G-spot; Gräfenberg Spot; Orgasm; Sexual Function; Clitorurethrovaginal Complex

## INTRODUCTION

The possible existence of an erotogenic area in the anterior vaginal wall has been referred at least since the 11th century, or even earlier.<sup>1,2</sup>

In 1950 Gräfenberg published a seminal paper: “*The role of urethra in female orgasm*”. His main statement was that “*an erotic zone always could be demonstrated on the anterior wall of the vagina*

*along the course of the urethra*” and that this area swells with sexual stimulation, reaching its maximum at the end of the orgasm. Gräfenberg did not share previous points of view on the “need” of vaginal orgasms and, in fact, wrote that “*we can almost say that there is no part of the female body which does not give sexual response, the partner has only to find the erotogenic zones.*”<sup>3</sup>

In 1981, Addiego et al. published a case report of “female ejaculation” associated with an “*erotically sensitive spot, palpable*

Received April 8, 2021. Accepted August 10, 2021.

<sup>1</sup>Hospital Lusíadas Porto, Porto, Portugal;

<sup>2</sup>Lower Genital Tract Unit, Centro Hospitalar de São João, Porto, Portugal;

<sup>3</sup>LAP, a Unilabs Company, Porto, Portugal;

<sup>4</sup>Department of Surgical Sciences University of Torino, Torino, Italy;

<sup>5</sup>Gynecology Department, Centro Hospitalar de São João, Porto, Portugal;

<sup>6</sup>Urology Department, Saúde Atlântica - Clínica do Dragão, Porto, Portugal;

<sup>7</sup>Department of Obstetrics and Gynecology, University of Iowa Hospitals and Clinics, Iowa City, IA, USA

Copyright © 2021 The Authors. Published by Elsevier Inc. on behalf of the International Society for Sexual Medicine. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

<https://doi.org/10.1016/j.esxm.2021.100435>

through the anterior wall of the vagina.”<sup>4</sup> Honoring the paper published 3 decades before, they named that area “Gräfenberg spot.” Later it was abridged to “G-spot” by Ladas et al.<sup>5–7</sup>

The concept of “G-spot” soon gained popularity, especially in the mainstream media. Despite this widespread generalized acceptance, in the medical literature it is still shrouded in controversy, with the studies aimed to prove its existence or inexistence often potentially biased by the socio-cultural background.<sup>8</sup> In 2012, Ostrzenski dissected the anterior vaginal wall of cadaver and claimed to have found it.<sup>9</sup> Nevertheless, it was not the final evidence. For the scientific community the question of whether it exists or is a mere scientific unicorn remains: if it does, there will be surgical implications (identification in urogynecological approach of the anterior vaginal wall, surgeries aiming at increasing the function of the G-spot) and possible new approaches to treat female sexual dysfunction. This structure has even been attempted to be replicated in male-to-female transgenders.<sup>10</sup>

For the public in general, the idea of the existence of 2 types of orgasm brings back Freudian concepts – and puts pressure upon women who cannot achieve a “vaginal orgasm.” The G-spot, despite supposedly being stimulated through the vagina, seemed to somehow fill this gap in the rhetoric of female liberation.<sup>8</sup> Some authors developed the concept of “clitourethrovaginal complex,” referring to a descending of the anterior vaginal wall, that was reported in some imaging studies and which can take the focus of a specific area or anatomic structure.<sup>11–13</sup>

Independently of the existence of any specific structure or complex, any attempt to reduce the female orgasm to its mere stimulation will always fall short: several complex factors must be considered, including intimacy, the hormone milieu, previous experiences, cultural and religious beliefs, etc. The brain is undeniably the key player in terms of female orgasm.<sup>8</sup>

The objective of this review was to evaluate the available studies in humans (questionnaires, anatomical, histological, imaging or physiological) on the existence, location, and histological nature of the G-spot and its possible practical implications.

## MATERIAL AND METHODS

A Pubmed, Pubmed Central, Cochrane, clinicaltrials.gov and Google Scholar literature search was performed, from inception to November 2020, restricted to abstracts in English, French, Portuguese, Spanish, or Italian language. Only studies on humans were included.

The search string used was: (“G-spot” OR “Gräfenberg spot”). Studies referring to self-awareness, clinical, imaging, histological, neurophysiological or anatomical evidence concerning the G-spot were included in the review. All abstracts were checked by 2 of the authors for eligibility.

The study protocol development and the review were conducted in accordance with the Preferred Reporting Items for Systematic Reviews (PRISMA) (Figure 1).<sup>14</sup>

PROSPERO registration was not possible, as currently it does not accept registrations for scoping reviews, literature reviews or mapping reviews.

Due to the nature of the study, it was considered exempt from IRB approval.

## RESULTS

Of the 32 studies included, 6 concerned self-awareness (surveys), 5 digital/instrumental exploration of the vaginal walls, 9 dissection of cadavers/biopsies/evaluation of surgical samples, 9 imaging evaluation (7 ultrasound and 2 magnetic resonance imaging), and 1 neurophysiological evaluation. (Figure 2) Two studies involved both clinical and histological evaluation.<sup>15,16</sup>

### Self-Awareness Studies/Surveys

In the 6 studies of this type, women answered whether or not they believed they had a G-spot and/or more sensitive area in the vagina. Of 5072 participants, 3195 (62.9%) reported to have of such an area. In one study, 1245 also answered if they believed in the existence of an area of higher sensitivity in the vagina: 84.3% did and 65.9% reported having it (7.3% in the posterior vaginal wall).<sup>17</sup>

All studies except one<sup>18</sup> exclusively enrolled heterosexual women. Nearly half of the participants in the surveys were health care professionals.

One study asked if the G-spot was associated with ejaculation: 72.6% answered affirmatively.<sup>19</sup>

One study evaluated pairs of twins and found no evidence of a genetic factor.<sup>20</sup>

A recent study associated likelihood to report having a G-spot with higher education and higher Female Sexual Function Index (FSFI).<sup>21</sup> (Table 1)

### Clinical Evidence

The 7 studies that explored digital or instrumental exploration of the vaginal walls, searching for the G-spot involved 1842 women (1500 from 1 study<sup>15</sup>). The G-spot was identified in 55.4% (1020/1842) women. In 2 studies it was identified in all women ( $n = 74$ )<sup>23,24</sup> and in another 2 studies in none ( $n = 83$ ).<sup>25,26</sup>

The criteria for identification of the G-spot, depending on the study, could be the report of a more sensitive area and/or a bulging/swelling upon stimulation.

In most cases, digital stimulation was performed, using 1 or 2 fingers by one of the investigators (and the partner in one




Figure 1. Preferred Reporting Items for Systematic Reviews (PRISMA).




Figure 2. a) Number of studies, according to category, in which the authors concluded for or agains the existence of the G-spot; b) number of women, according to category in which some evidence or perception of the existence of the G-spot (regardless of the authors' conclusions).

**Table 1.** Studies on self-perception of the existence of the G-spot

Author	Country	Year	Type of study and population	N =	Age range	Results
Davidson J et al. <sup>17</sup>	USA	1989	<ul style="list-style-type: none"> <li>Anonymous questionnaire (55% answer rate)</li> <li>Membership lists of the 2 national organizations representing health and counseling professions</li> <li>Heterosexual women</li> <li>Six open-ended and 182 closed-form item</li> </ul>	1245	(mean 38)	<ul style="list-style-type: none"> <li>84.3% believed that a sensitive area exists in the vagina</li> <li>65.9% reported they had a more sensitive area in the vagina (55.1% in the anterior vaginal wall and 7.3% in the posterior wall)</li> </ul>
Darling C et al. <sup>19</sup>	USA	1990	<ul style="list-style-type: none"> <li>Anonymous questionnaire (mailed) (192 open-ended and closed-form items)</li> <li>Asked if they were aware of a "sensitive area located in the vagina"</li> <li>Professional women (health care)</li> </ul>	1230	22-82 (mean 38.4)	<ul style="list-style-type: none"> <li>65.9% reported having a G-spot (sensitive area)</li> <li>Stimulation of that area led to orgasm in 72.6%</li> <li>Overall, 40% reported ejaculating (82% if reported having a sensitive area in the vagina)</li> <li>Sensitive area associated with multiple orgasms</li> </ul>
Kratochvíl S <sup>22</sup> *	Czech Republic	1993	<ul style="list-style-type: none"> <li>Women treated for neurotic disorders (200) and health professional/councilor (100)</li> <li>Questionnaire</li> </ul>	300	Unknown	<ul style="list-style-type: none"> <li>Anterior wall slightly more sensitive than the posterior one</li> <li>One third reported effective stimulation in the depth of the vagina with cervical tapping.</li> <li>Stimulation in the area corresponding to the alleged G spot was acknowledged as effective by 10%–20%</li> </ul>
Burri AV et al. <sup>20</sup>	UK	2010	<ul style="list-style-type: none"> <li>Random exclusively heterosexual twins</li> <li>Postal self-administered questionnaire (TEIQue-SF; TIPI; frequency of orgasm)</li> <li>Questioned "Do you believe you have a so-called G-spot, a small area the size of a 20p coin on the front wall of your vagina that is sensitive to deep pressure?"</li> </ul>	1804 (902 pairs of twins)	22-83	<ul style="list-style-type: none"> <li>56% reported having a G-spot</li> <li>Women reporting always having an orgasm or having multiple orgasms with intercourse, or more satisfied with their relationship answered more often they had a G-spot</li> <li>Older women less likely to report it</li> <li>No evidence of a genetic factor</li> <li>"Extraversion" and "openness to new experience" associated with reporting having a G-spot</li> <li>Correlation with different factors (sexual behavior, relationship satisfaction, attitude towards sexuality) showed the assumption of the presence of a G-spot to be a pseudo-phenomenon</li> </ul>
Wang X et al. <sup>18</sup>	China	2012	<ul style="list-style-type: none"> <li>Convenience (online) sampling of WSW</li> <li>Asked if they were aware of the G-spot and if their partners deliberately tried to stimulate it</li> </ul>	184	19-46	<ul style="list-style-type: none"> <li>65.2% reported having had their G-spot stimulated</li> <li>49.2% reported bleeding after sex (vs 12.5% in those not reporting G-spot stimulation, <math>P &lt; .001</math>)</li> </ul>
Ellibeş Kaya A et al. <sup>21</sup>	Turkey	2018	<ul style="list-style-type: none"> <li>Cross sectional</li> <li>Pre-menopausal, sexually active women &gt;18 y</li> <li>Exclusion criteria included women who desired cosmetic genital surgery</li> <li>Asked participant if they "felt a coin-size sensitive area in the anterior vaginal wall at the time of finger or penis penetration or pressure"</li> <li>FSFI, FGSIS</li> </ul>	309	18-54	<ul style="list-style-type: none"> <li>51.5% believed the G spot exists</li> <li>19.7% indecisive/do not know</li> <li>Higher education levels correlated with reporting G spot existence</li> <li>Sexual dysfunction lower in those who reported its existence</li> <li>Higher mean score in the FSFI orgasm domain in women who reported the G spot exists</li> </ul>

FSFI = Female Sexual Function Index; FGSIS = Female Genital Self- Image scale; TEIQue-SF = Trait Emotional Intelligence Questionnaire- Short Form; TIPI = Ten-Item Personality Index; WSW = Women who have sex with women; \* access only to the abstract.

study<sup>26</sup>). In one study, a vibrator was used in cases in which “ejaculation was hard to induce.”<sup>23</sup>

As for location, it was reported to be localized “1 cm deep in the vaginal wall,”<sup>24</sup> “anterior vaginal wall”<sup>23</sup> or “connected to the hymen” (below the urethral meatus).<sup>15</sup> The 2 papers that concluded against the existence of the G-spot showed pleasurable spots in different areas of the vagina and in the cervix.<sup>25,26</sup>

Three papers reported on the swelling of a specific area upon stimulation: absent in one,<sup>25</sup> and always present in 2.<sup>23,24</sup>

Several biases could be identified, for instance, inclusion of a majority of sex workers,<sup>25</sup> or inclusion of only coital anorgasmic women<sup>26</sup> (both studies failed to provide evidence of the existence of the G-spot), which do not represent the general female population. Also, the context of a clinical study or having the stimulation induced by an investigator may hinder the ability to reach orgasm. (Table 2)

### Imaging Evidence

Imaging evaluation of the possible existence of the G-spot was performed in 9 studies: 2 using MRI (n = 23) and 7 using ultrasound (n = 116; one study did not report the number of involved subjects<sup>31</sup>).

One of the MRI studies involved women with midthoracic complete spinal cord injury (n = 2); brain PET-MRI showed activation of the region of the nucleus of the solitary tract. The authors concluded for the existence of a possible by-pass of the spinal cord, via vagus nerve and activated by vaginal stimulation.<sup>28</sup>

The other MRI study retrospectively evaluated MRIs performed for diverse clinical reasons (n = 21), assuming Ostrzensky’s putative location of the G-spot. They described a “G-spot complex” in 62% of women (in all women if vaginal gel had been used and in 3 of 11 if it was not used).<sup>39</sup>

The studies using ultrasound used either 2D or 3D flat or transvaginal probes to evaluate the clitoris and anterior vaginal wall, sometimes complemented with doppler evaluation.

One descriptive study reported a hyperechoic area between the clitoris and the vagina and two lateral hypoechoic areas (cavernous veins)<sup>31</sup>; in another one, the thickness of the urethrovaginal space was positively correlated with vaginal orgasm (specially for the distal segment).<sup>32</sup>

Battaglia et al. reported the existence of a gland-like structure (“female prostate”) in the urethrovaginal space, fed by small vessels in 2 studies (n = 87). Dimensions of these structures were described to be bigger in women who reported vaginal orgasms and to correlate with androgens’ levels and time since intercourse. The authors did not assume this area to be the G-spot.<sup>35,36</sup> A previous study (n = 5) reported the absence of glandular structures in that area.<sup>34</sup>

In 3 studies a close relationship between the root of the clitoris and the anterior vaginal wall during perineal

contraction and/or vaginal penetration and/or stimulation was described. In one of the studies, the authors assumed the root of the clitoris to be the G-spot<sup>34</sup>; Buisson et al. assumed the G-spot was the “clitourethrovaginal complex.”<sup>12,13</sup> The descent of these structures had previously been described by Foldes et al.<sup>31</sup> (Table 2)

### Neuro and Electrophysiological Evidence

Shafik et al. evaluated the electrical activity of the vagina (n = 24) and reported electrical waves (pacemaker) generated in the proximal vagina, and noted the intensity was responsive to vaginal pressure.<sup>30</sup> (Table 2)

### Anatomical and Histological Evidence

The histological nature of the G-spot was evaluated in 7 studies (biopsies/surgical specimens/autopsies), for a total of 244 women. In the largest study (n = 175) it was identified in 47.4% of women.<sup>16</sup> In this study, it was described as being composed of epithelial, glandular and erectile tissue.

In Ostrzenski’s first paper (dissection of one cadaver) the G-spot was described as a fibroconnective sac, containing erectile-like tissue (no histological examination performed).<sup>9</sup> Later, in 2 other papers he described it as a vein like structure, with ability to expand.<sup>43,38</sup> This description does not match the usual histology of erectile tissues (arteries, vascular shunts, venous sinusoids and a matrix of connective tissue and smooth muscle).<sup>44</sup>

Li et al. described an increased density of microvessels and small nerves in the distal third of the anterior vaginal wall; Song et al. found higher density of terminal nerve branches at the second distal one-fifth of the anterior vaginal wall.<sup>37,33</sup> Two other studies found no differences in the innervation of the anterior vaginal wall, but with one showing increased innervation along the urethra (which the authors did not consider evidence of the existence of the G-spot).<sup>41,42</sup>

D’Amati evaluated the presence of type 5-phosphodiesterase (PDE5) in the anterior vaginal wall and found it expressed in smooth muscle of vessels, which formed a pseudocavernous tissue in the vaginal wall and endothelium. The authors concluded that, similarly to males, the system nitric oxide synthase-PDE5 may also contribute to sexual female arousal.<sup>29</sup>

Hoag et al. found no macroscopic structure in the putative location of the G-spot, namely “spongy” tissue.<sup>40</sup>

Ostrzenski described the G-spot as having a diagonal orientation, measuring  $8 \pm 5$  mm of longest dimension, making a diagonal angle with the urethra and often sided (more often to the left side), 4.5–5.5 cm from the urethral meatus.<sup>9,43,38</sup> (Table 2)

**Table 2.** Studies on clinical, anatomical, imaging, surgical, or histological evidence of the existence of the G-spot

Author	Country	Year	Type of study	N =	Age range	Results
Perry J et al. <sup>24</sup>	USA	1981	<ul style="list-style-type: none"> <li>Vaginal and uterine myograph to evaluate EMG levels</li> <li>Women recruited by sex therapists, educators and counselors</li> <li>Questionnaire (Likert scale, Sexual Arousal Inventory)</li> <li>Digital examination of the pubococcygeus muscle</li> <li>Two finger palpation of the whole anterior wall (G-spot), sometimes with abdominal pressure</li> </ul>	47	21-63 (mean 34)	<ul style="list-style-type: none"> <li>G-spot identified in all subjects in the anterior wall (most at 12 o'clock)</li> <li>1 cm deep in the vaginal wall</li> <li>Swelling upon stimulation (can take more than 1 minute)</li> </ul>
Goldberg D et al. <sup>27</sup>	USA	1983	<ul style="list-style-type: none"> <li>Self-recruited women (news-papers, talk shows)</li> </ul>	11 (6 self-identified as ejaculators)	24-61	<ul style="list-style-type: none"> <li>G-spot identified in 4/11 (36%) women</li> <li>No differences between ejaculators and nonejaculators</li> </ul>
Alzate H <sup>25</sup>	Colombia	1985	<ul style="list-style-type: none"> <li>Digital exploration of both vaginal walls</li> <li>Healthy women (82% sex workers)</li> </ul>	27	(mean 24.1±4.2)	<ul style="list-style-type: none"> <li>89% had an orgasmic response in response to vaginal stimulation (not a specific area) – findings do not support the existence of a G-spot, but rather the existence of clitoral and vaginal orgasms</li> <li>85% reported it in the posterior wall (mostly in the lower half) and 74% in the anterior one</li> <li>Of the ones who had erotic sensitivity in the anterior wall, 100% referred it in the upper half and 60% in the lower half of the vagina)</li> <li>No perception of “swollen structures)</li> <li>Some of the women who reached orgasm with stimulation of the posterior wall also did it with ano-rectal stimulation</li> <li>No evidence of ejaculation</li> <li>(Bias: mostly sex workers)</li> </ul>
Hoch Z(25)	Israel	1986	<ul style="list-style-type: none"> <li>Women with coital anorgasmia but orgasmic with external genitalia stimulation</li> <li>Israeli Minnesota Multiphasic Psychological Inventory, Bem Sex-Role Inventory, Sim-Fam game, anthropometric measures of sexual dimorphism, detailed sociological questionnaire, Lief and Ebert Sexual Performance Evaluation Questionnaire</li> <li>Sexual examination of the vagina (finger) – performed by the investigator and partner</li> </ul>	56	19-64	<ul style="list-style-type: none"> <li>No evidence of a G-spot</li> <li>96% with pleasure upon exploration of the entire anterior vaginal wall, including the deeper situated urinary bladder, periurethral tissues and Halban's fascia</li> <li>6% with exploration of the cervix, 3% in the posterior wall of the vagina and 2% in the lateral vaginal walls (4–8 h)</li> <li>64% orgasmic with stimulation of the anterior vaginal wall</li> <li>(Bias: anorgasmic women)</li> </ul>
Zaviaci M et al. <sup>23</sup>	Czechoslovakia	1988	<ul style="list-style-type: none"> <li>Mostly women referred for infertility</li> <li>Questionnaire</li> <li>Digital stimulation (1-2 fingers) (in 2 “hard-to-induce expulsions” cases a vibrator was used)</li> </ul>	27	20-40	<ul style="list-style-type: none"> <li>G-spot identified in all the participants, in the anterior vaginal wall</li> <li>Tumescence of the spot upon stimulation</li> <li>37% ejaculated</li> </ul>
Whipple B et al. <sup>28</sup>	USA	2002	<ul style="list-style-type: none"> <li>Women with midthoracic complete spinal cord injury</li> <li>Brain PET-MRI during cervical and vaginal self-stimulation</li> </ul>	2 women with SCI and one for control		<ul style="list-style-type: none"> <li>All had activation of the region of the nucleus of the solitary tract, via vagus nerve (by-pass of the spinal cord)</li> </ul>
D'Amati G et al. <sup>29</sup>	Italy	2002	<ul style="list-style-type: none"> <li>Immunohistochemistry (antibodies against 5 phosphodiesterase [PDE5]) performed on vaginal tissue obtained at autopsy and on scraped cells</li> </ul>	14 vaginal specimens obtained at autopsy 5 vaginal samples from healthy donors	18-40 (mean 30.0±3.6) 24-38 years	<ul style="list-style-type: none"> <li>PDE5 immunoreactivity was mostly localized in the smooth muscle of vessels, forming a pseudocavernous tissue in the vaginal wall and endothelium</li> </ul>

(continued)

Table 2. Continued

Author	Country	Year	Type of study	N =	Age range	Results
			from the anterior vaginal epithelium of healthy female donors			<ul style="list-style-type: none"> <li>• It was also present in exfoliated vaginal cells and Skene glands</li> <li>• cavernous or pseudocavernous tissue within the vaginal wall was observed in 12/14 (86%)</li> <li>• Skene glands identified in 9/14 (64%)</li> </ul>
Shafik A et al. <sup>30</sup>	Egypt	2003	<ul style="list-style-type: none"> <li>• Healthy volunteers</li> <li>• Electrovaginogram and manometry</li> </ul>	24	26-52 (mean 28.6±8.2)	<ul style="list-style-type: none"> <li>• Electrical waves detected in the proximal vagina (pacemaker)</li> <li>• Intensity responsive to vaginal pressure</li> </ul>
Foldes P et al. <sup>31</sup>	France	2007	<ul style="list-style-type: none"> <li>• Healthy volunteers</li> <li>• Vaginal ultrasound with doppler</li> <li>• Placement of a hypercogenic stick in the supposed area of the clitoris</li> </ul>	Not referred	38-48	<ul style="list-style-type: none"> <li>• Identification of a hypercogenic area between the clitoris and the vagina and two lateral hypoecic areas (cavernous veins)</li> <li>• Descending movement of these structures with the contraction of the <i>levator ani</i>.</li> </ul>
Gravina G et al. <sup>32</sup>	Italy	2009	<ul style="list-style-type: none"> <li>• Healthy women with and without vaginal orgasm</li> <li>• Urodynamic study</li> <li>• Ultrasound for evaluation of the urethrovaginal space thickness</li> </ul>	20 (9 with vaginal orgasms, 11 without vaginal orgasm)	29-36	<ul style="list-style-type: none"> <li>• The urethrovaginal space and distal, middle, and proximal urethrovaginal segments were thinner in women without vaginal orgasm</li> <li>• Correlation between thickness and vaginal orgasm (higher for the distal segment)</li> <li>• Excellent correlation between observers</li> </ul>
Thabet S(16)	Egypt	2009	<ul style="list-style-type: none"> <li>• Evaluation of women with cystocele, prior to surgery</li> <li>• Evaluation of the circumcision and anterior vaginal surgery in the function of the G-spot</li> <li>• Preoperative "sexual examination" to map the G-spot</li> <li>• Pre and post-operative sexual function assessment (Kasr El Aini Sexual Assessment Sheet)</li> <li>• Histological examination of surgical specimens</li> </ul>	175 (125 circumcised and 50 noncircumcised)	25-35	<ul style="list-style-type: none"> <li>• Functional evidence of the presence of the G-spot in 82,3%</li> <li>• Anatomical evidence in 65.9%</li> <li>• Histological evidence (epithelial, glandular and erectile tissue) in 47,4%</li> <li>• Ejaculation in 12.6%</li> <li>• Sex scores higher in women with histological evidence of the G-spot</li> <li>• Significant drop in sex scores after anterior vaginal wall surgery</li> <li>• Circumcision rarely affects the G-spot function</li> </ul>
Song Y et al. <sup>33</sup>	Korea	2009	<ul style="list-style-type: none"> <li>• Microdissection and protein gene product (PGP) 9.5 immunohistochemistry of the vagina of fresh cadavers</li> </ul>	7	50-81 (mean 66.63±12.11)	<ul style="list-style-type: none"> <li>• Terminal nerve branches in the vaginal wall were most dense at the second 1/5 partition from the inferior anterior wall</li> <li>• Lower density at the fourth partition and scarcity in the fifth 1/5 partitions from the bottom</li> <li>• The mucosa and the vaginal muscle were thicker in the distal third of the vagina</li> <li>• Small and large nerve fibers more common in the distal third of the vagina</li> <li>• The second 1/5 partition of the distal anterior wall had significantly richer innervation</li> <li>• The authors assume the 1/5 partition of the distal anterior wall as the G-spot</li> </ul>
Foldes P et al. <sup>34</sup>	France	2009	<ul style="list-style-type: none"> <li>• Dynamic sonographic of the clitoris (flat and vaginal probe)</li> <li>• Healthy, heterosexual, sexually active women reporting vaginal orgasm and having a G-spot</li> <li>• No contraception</li> <li>• Observation between day 4-12 of the cycle</li> </ul>	5	34	<ul style="list-style-type: none"> <li>• During perineal contraction and finger penetration, the coronal planes demonstrated a close relationship between the root of the clitoris and the anterior vaginal wall (descending movement of 2,5-5 mm)</li> <li>• The G-spot may be the root of the clitoris</li> </ul>

(continued)

Table 2. Continued

Author	Country	Year	Type of study	N =	Age range	Results
			<ul style="list-style-type: none"> <li>No sexual dysfunction</li> <li>Evaluation of the size of the clitoris and of its movements during voluntary perineal contractions and finger pressure in the most pleasurable area of the vagina (without stimulation)</li> </ul>			<ul style="list-style-type: none"> <li>No glandular structures visualized along the urethra</li> </ul>
Battaglia C et al. <sup>35</sup>	Italy and Spain	2010	<ul style="list-style-type: none"> <li>2D and 3D translabial ultrasound evaluation of the clitoris and urethrovaginal space, without sexual stimulation</li> <li>Women on a stable heterosexual relationship, sexually active, and without sexual dysfunction</li> <li>Eumenorrheic women with and without vaginal orgasm</li> </ul>	39 women: <ul style="list-style-type: none"> <li>19 with vaginal orgasm</li> <li>20 without vaginal orgasm</li> </ul>	24-30	<ul style="list-style-type: none"> <li>The 3D reconstruction showed the presence of a gland-like structure ("female prostate") in the urethrovaginal space, with small vessels feeding it</li> <li>Length and volume of structures contained in the urethrovaginal space were significantly higher in women who experienced vaginal orgasms</li> <li>Volume of these structures correlated with time since intercourse, levels of testosterone and androstenedione</li> <li>The authors do not assume this area as the G-spot, but rather as a system of glands and ducts</li> </ul>
Battaglia C et al. <sup>35</sup>	Italy and Spain	2010	<ul style="list-style-type: none"> <li>2D and 3D translabial ultrasound evaluation of the clitoris and urethrovaginal space, without sexual stimulation</li> <li>Women on a stable heterosexual relationship, sexually active, and without sexual dysfunction</li> <li>Lean women with PCOS vs. eumenorrheic women</li> <li>No hormonal therapies in the previous 6 months</li> </ul>	48 women: <ul style="list-style-type: none"> <li>23 with PCOS</li> <li>25 eumenorrheic</li> </ul>	18-35	<ul style="list-style-type: none"> <li>The 3D reconstruction showed the presence of a gland-like structure ("female prostate") in the urethrovaginal space, with small vessels feeding it</li> <li>This structure was larger in women with PCOS</li> <li>Size of the "female prostate" correlated with time since intercourse, length of the urethrovaginal space and levels of testosterone</li> <li>Middle portion of the urethra significantly thicker in women with PCOS</li> <li>No differences in terms of vascularization or clitoral body volume</li> <li>No differences in terms of vaginal orgasm (56 vs. 52%)</li> <li>The authors do not assume this area as the G-spot, but rather as a system of glands and ducts</li> </ul>
Buisson O et al. <sup>13</sup>	France	2010	<ul style="list-style-type: none"> <li>Ultrasound of the anterior vaginal wall during coitus</li> <li>Coronal section on the top of the vulva during the penetration using a flat probe</li> <li>Woman not using contraception</li> <li>Exam performed during the late follicular phase</li> </ul>	1 couple	Not referred	<ul style="list-style-type: none"> <li>The penis inflated the vagina and stretched the root of the clitoris, thus becoming in a very close relationship with the anterior vaginal wall (clito-urethrovaginal complex), moving and pressuring it against the pubic symphysis.</li> <li>This could explain the pleasurable sensitivity of this anterior vaginal area.</li> <li>The authors assume the clito-urethrovaginal complex is the G-spot)</li> </ul>
Ostrzenski A. <sup>9</sup>	USA	2012	<ul style="list-style-type: none"> <li>Stratum-by-stratum vaginal wall dissection on a fresh cadaver</li> </ul>	1	83	<ul style="list-style-type: none"> <li>The G-spot has a distinguishable anatomic structure that is located on the dorsal perineal membrane, 16.5 mm from the upper part of the urethral meatus</li> <li>The lower pole and the upper pole were located 3 and 15 mm next to the lateral border of the urethra</li> <li>Appearance of a well-delineated sac with walls that resembled fibroconnective tissues and resembled erectile tissues.</li> </ul>

(continued)



Table 2. Continued

Author	Country	Year	Type of study	N =	Age range	Results
						<ul style="list-style-type: none"> <li>• Upon opening the sac blue grape-like anatomic compositions of the G-spot emerged with dimensions of 8.1 × 3.6–1.5 × 0.4 mm (stretchable to 33 mm).</li> <li>• The G-spot structure had three distinct areas</li> <li>• From the distal tail, a rope-like structure (vessel) emerged, which was seen for approximately 1.6 mm and then disappeared into the surrounding tissue.</li> <li>• Limited mobility together with the dorsal perineal membrane on which the G-spot was situated</li> </ul>
Thabet S <sup>15</sup>	Egypt	2013	<ul style="list-style-type: none"> <li>• Prospective, randomized</li> <li>• Clinical examination, operative findings and histopathological examination of surgical specimens (n=350)</li> <li>• Kasr El Aini Sexual Questionnaire Sheet</li> </ul>	1500 women: <ul style="list-style-type: none"> <li>• 500 had vulvar/vaginal surgery</li> <li>• 1000 outpatients</li> <li>• (39 with vaginal agenesis)</li> </ul>	20-35	<ul style="list-style-type: none"> <li>• 52.7% with local response upon examination ("irritation" and protrusion)</li> <li>• Localized structure in 58% and diffuse in the rest</li> <li>• Ejaculation in 100% of those with a localized G-spot and in 24.5% if diffuse</li> <li>• 100% histologically proved ("erectile tissue")</li> <li>• 100% of the cases with connection with the hymen</li> <li>• Pictures in the paper localize the "G-spot bodies" in the vulvar, below the urethral meatus.</li> </ul>
Buisson O et al. <sup>12</sup>	France, Italy	2013	<ul style="list-style-type: none"> <li>• Ultrasound evaluation of the "clitorourethrovaginal complex" using a vaginal (sagittal plane) and a linear probe</li> <li>• Functional sonography of the stimulated clitoris (manual self-stimulation of the external clitoris or during vaginal penetration with a wet tampon)</li> <li>• Heterosexual women, sexually active, in a stable relationship</li> </ul>	3	27-33	<ul style="list-style-type: none"> <li>• The sagittal scans obtained during external stimulation and vaginal penetration demonstrated that the root of the clitoris is not involved with external clitoral stimulation (vaginal wet tampon immobile when external clitoral stimulation was performed)</li> <li>• During vaginal stimulation, the whole CUV complex and the clitoral roots in particular are involved</li> <li>• The color signal indicating flow speed in the veins</li> <li>• mirrored the anatomical changes.</li> </ul>
Li T et al. <sup>37</sup>	China	2014	<ul style="list-style-type: none"> <li>• Biopsies of the distal- and proximal-third of the anterior vagina</li> <li>• Women with stress urinary incontinence or pelvic organ prolapse</li> <li>• H&amp;E, neural marker protein gene product 9.5 and smooth muscle actin</li> </ul>	32	41-77	<ul style="list-style-type: none"> <li>• Increased density of small nerves and microvessels in the distal-third of the anterior vaginal wall</li> <li>• Small nerve fibers detected in the lamina propria and muscle layers of distal- and proximal-third areas.</li> <li>• Nerve bundles less abundant in the muscle layer and rarely appeared in the lamina propria.</li> <li>• No small vessels in the lamina propria and few in the muscle layer</li> </ul>
Ostrzinski A et al. <sup>38</sup>	USA, Poland	2014	<ul style="list-style-type: none"> <li>• Fresh human cadavers</li> <li>• Anterior vaginal wall dissections and G-spot</li> <li>• All specimens stained with haematoxylin and eosin</li> <li>• Two random cases selected at random for immunohistochemical staining</li> </ul>	8	<ul style="list-style-type: none"> <li>• 37-68</li> </ul>	<ul style="list-style-type: none"> <li>• The G-spot was identified in all women and was in a diagonal plane</li> <li>• The G-spot complex was located within the distal anterior vaginal wall (average 4.5 cm from the urethral meatus) and has 3 fused parts</li> <li>• In seven (87.5%) it was on the left side and in one (12.5%) on right side</li> <li>• Angle with the urethra ranging between 18-35°</li> <li>• The G-spot was intimately fused with vessels</li> </ul>

(continued)

Table 2. Continued

Author	Country	Year	Type of study	N =	Age range	Results
						<ul style="list-style-type: none"> <li>• Sac thickness &lt;2 mm</li> <li>• A vein-like vascular structure with a few smaller feeding arteries was identified</li> <li>• A band-like structure protruded from the tail of the G-spot.</li> <li>• z.bull; The size of the G-spot was variable (mean 7 mm) and could expand on average of 5 times</li> <li>• Histologically, the G-spot was determined as a neurovascular complex structure within a fibroadipose tissue bed.</li> <li>• The neural component contained abundant peripheral nerve bundles and a nerve ganglion.</li> <li>• The vascular component comprised large vein-like vessels and smaller feeding arteries and resembled arteriovenous malformations.</li> <li>• The vascular component did not resemble erectile tissue</li> <li>• Circular and longitudinal muscles covered the G-complex.</li> </ul>
Ostrzenski A(34)	USA	2014	<ul style="list-style-type: none"> <li>• Prospective case series on fresh female cadavers</li> <li>• Anterior vaginal wall stratum-by-stratum macro-dissections</li> <li>• The G-spot tissues were stained with hematoxylin and eosin for histology</li> </ul>	11	27-83	<ul style="list-style-type: none"> <li>• The cylindrical G-spot complex was identified in all subjects on the distal vagina</li> <li>• More often on the left side<sup>8</sup> of the margin of the urethra</li> <li>• Localized laterally to the urethra (distance between tail and urethra 3.1 to 5.7 mm)</li> <li>• 5.5 cm from the urethral meatus</li> <li>• Mean size of 8±5 mm, expanding an average 5 times when released from the sac</li> <li>• Diagonal orientation plane with an angle of 18-35°</li> <li>• Sac wall with a thickness of 1.6±0.4 mm and merges with the anterior vaginal wall</li> <li>• At the lower pole of the G-spot, a tiny band-like structure was identified, grossly resembling a vascular structure</li> <li>• Blood vessels with ability to expand</li> <li>• The G-spot complex expansion elevated the anterior vaginal walls in all subjects (vascular distention)</li> <li>• The autonomic parasympathetic nervous system was the dominant division at the time of female subject sudden death.</li> <li>• No secretory glands, ducts, cavernous, spongiosum or erectile tissues were identified.</li> </ul>
Maratos YK et al. <sup>39</sup>	France, Germany	2016	<ul style="list-style-type: none"> <li>• MRI study</li> <li>• Observational, retrospective, single center</li> <li>• Consecutive women</li> <li>• MRI performed for clinical reasons</li> <li>• With or without vaginal gel opacification</li> <li>• Assumed Ostrzenski's previous papers as reference of the anatomical description of the G-spot</li> </ul>	21	17-72	<ul style="list-style-type: none"> <li>• "G-spot complex" present in the anterior vaginal wall in 62% (13/21)</li> <li>• 10/10 cases in which gel opacification used</li> <li>• 3/11 cases in the group in which gel was not used</li> </ul>
Australia, Canada	2017		<ul style="list-style-type: none"> <li>• Dissection of the anterior vaginal wall in cadavers</li> </ul>	13 (8 fixed and 5 fresh)	32-97	<ul style="list-style-type: none"> <li>• No macroscopic structure other than the urethra and vaginal wall</li> </ul>

(continued)

**Table 2.** Continued

Author	Country	Year	Type of study	N =	Age range	Results
Mazloomdoost D et al. <sup>41</sup>	USA	2017	<ul style="list-style-type: none"> <li>• Section for macroscopic inspection (4.5x magnification) and histologic examination</li> </ul>	4	67-97	<ul style="list-style-type: none"> <li>• lining in the location of the putative G-spot</li> <li>• No erectile or “spongy” tissue in the anterior vaginal wall</li> <li>• In 1 specimen, a small amount of vascular tissue was noted in the subepithelial space</li> <li>• Systematic high density of veins in the lateral to the urethral and vaginal walls (number and thickness variable)</li> </ul>
Aydın S et al. <sup>42</sup>	Turkey	2020	<ul style="list-style-type: none"> <li>• En bloc removal of 4 female fresh-frozen cadaveric pelvises</li> <li>• 18 to 25 serial sections obtained from each</li> </ul>	17	33-70	<ul style="list-style-type: none"> <li>• Epithelial, lamina propria, and muscular layer surrounded the urethral lumen in all specimens</li> <li>• Innervation and vasculature concentrated in the lamina propria</li> <li>• No differences in nerve distribution along the anterior vaginal wall</li> <li>• Differences in the innervation and vascularization along the urethra</li> <li>• No evidence of any structure corresponding to a possible G-spot described</li> </ul>
			<ul style="list-style-type: none"> <li>• Specimens were taken from women with anterior vaginal wall prolapse undergoing colporrhaphy anterior repair</li> <li>• 15 mm proximal to the external urethral orifice</li> <li>• Immunohistochemically stained: actin, smooth muscle Ab-1 and S100 Protein Ab-1</li> <li>• Microvessels and nerves in the lamina propria and muscularis were counted in five consecutive high-power fields of a light microscope</li> <li>• Comparisons with proximal, distal, right and left paravaginal microvessel and nerve fiber density</li> </ul>	<ul style="list-style-type: none"> <li>• 4 premenopausal</li> <li>• 13 postmenopausal</li> </ul>		<ul style="list-style-type: none"> <li>• Vaginal nerve fibers in the lamina propria and muscularis have a fairly even distribution in the anterior vaginal wall</li> <li>• Vaginal small vessel vascularization and microvascularization also evenly distributed</li> </ul>

EMG = electromyography; MRI = magnetic resonance imaging; PET-MRI = positron emission tomography–magnetic resonance imaging; PCOS = polycystic ovarian syndrome.

## DISCUSSION

### Main Findings

We were unable to identify agreement regarding the existence of the G-spot, on its location, size or nature. Therefore, we must conclude that its existence remains to be scientifically proven.

### Interpretation

The discussion on the existence of the G-spot has been polarized in 2 extremes and rich in letters and rebuttals to each paper published, as well as reviews aimed at supporting one view or the other. Ostrzenski (for) and Puppo (against) have spearheaded these discussions.<sup>45,46</sup> Reviewing the literature it easily becomes apparent that few groups have published original investigation regarding this topic.

The surveys on self-awareness on this topic clearly show that the majority of women believe in its existence and that they consider themselves to have this special erotogenic area. However, this belief may be biased by the current assumption that it does

exist. Also, can one assume that the source of pleasure and/or orgasm is independent of clitoral stimulation? Also, the psycho-relational aspects of orgasm — at least as important as any genital stimulus — cannot be ignored.<sup>8</sup> Some studies concluded that women with higher education levels and better sexual function were more likely to report having a G-spot, which can be the consequence of a higher exposition to the concept.

In the 1980’s several clinical studies were conducted, searching for the G-spot. In some studies, it was systematically identified while in others it could not be found. Combining all studies, it was identified in more than half of the women. When identified, the report of a swelling in the anterior vaginal wall was often reported. Replication of these studies, currently, would raise ethical concerns (investigator sexually stimulating the subjects). In case of G-spot existence, it would be expected that some (but not all) women would be inhibited and less likely to have an excitatory sexual response in an experimental environment, in accordance with the Bancroft’s Dual Control Model of Sexual Response.<sup>47</sup> Some of the criticisms to this kind of studies include

that both subjects and investigators are aware of the objectives of the experiment, and the fact that genital stimulation can lead to arousal and orgasm.<sup>11,26</sup> We could not find any study on the male perception of the question.

The imaging studies dominated since the beginning of the XXI century. The available data derives from a few investigation groups and premise that the G-spot exists.<sup>39</sup> Ultrasound, due to its dynamic and real time evaluation seems to be a good tool to explore this question. The data derived from these studies is conflicted, specially on the nature of a possible G-spot: some found glandular structures (without assuming it to be the G-spot), other reported it as vascular, others just found a thickening in the area or no structures in the area apart from the ones already consecrated in the anatomy books. The studies that described a thickening, associated it with capacity of vaginal orgasm and with androgen levels.<sup>32,35,36</sup> However, in the literature there is a lack of support to an association between androgen levels and sexual function.<sup>48,49</sup> The use of ultrasound brought to light the finding that during vaginal penetration there is a descendent movement of the anterior vaginal wall, increasing the contact between this region and the penis/fingers/object. From this, derived a new concept: the clitourethrovaginal complex. This concept could harmonize the role of the anterior vaginal wall and clitoris with orgasms.<sup>40</sup> Earlier, in 2003, Levin already supported that the concept of G-spot should shift to that of “anterior wall erogenous complex,” encompassing the urethra-clitoral-G area-Halban's fascia.<sup>50</sup> This complex can be seen more as a functional than anatomical entity; the stimulation of this area can be the key for vaginally activated orgasms (of which the clitoris would still be among its triggers, by stimulation through the anterior vaginal wall. Hoag et al. described a merging between the anterior aspect of the distal vaginal wall and the clitoris. In that point, individualization of the clitoris, urethra and vagina was hard to achieve.<sup>40</sup> Ostrzenski criticized the dissection technique used by this group and assumed that it could explain the different findings from his own work.<sup>51</sup> For some authors, the concept of clitourethrovaginal complex became a synonymous of G-spot. Puppo et al. disagreed from this concept, as they believe the penis cannot come in contact with the clitoris or the venous plexus of Kobelt during vaginal intercourse.<sup>52</sup>

Ostrzenski announced to the scientific community the finding of the G-spot during the dissection of a cadaver of an 83-year woman in 2012.<sup>9</sup> He described a structure with 3 sections, with a capsule of fibroconnective tissue, containing what seemed to be erectile tissue. Later, in larger series he systematically described it again, and included histological evaluation (considered it a neurovascular structure). Interestingly, he found that usually this structure was not in the midline.<sup>43,38</sup> The descriptions of Ostrzenski have been largely criticized in the literature, by both scholars pro and against the existence of such an anatomical/functional structure. Hoag et al., in a well conducted and documented study, could not replicate these findings. Also, previous

exhaustive anatomical studies about the clitoris never described the existence of the G-spot.<sup>40</sup>

The studies on innervation of the vaginal wall again were contradictory: while some found increased innervation in a specific area of the anterior vaginal wall and others a pacemaker effect, responsive to pressure, others found an even distribution in the anterior vaginal wall.<sup>30,33,42</sup> The findings of D'Amati, while encouraging for a role of PDE5 in female sexual arousal, did not compare the immunoreactivity of the anterior vaginal area with other areas.<sup>29</sup>

In a rat model, it was shown that vaginal innervation density is higher in the distal half of the vagina and that surgical menopause led to a decrease in both proximal and distal vaginal innervation, that could be reversed with estrogen treatment.<sup>53</sup> Nevertheless, possible interspecies differences do not allow generalizing the conclusions to humans.

Based on the premise that heritability can be shown in genuine anatomical traits, Burri et al. (2010) questioned 902 pairs of twins about their own perception of the existence of a G-spot to conclude that it is secondary pseudo-phenomenon to life experiences, without a genetic background.<sup>20</sup>

Two years later, Ostrzensky wrote that “the G-spot gene has been identified and been incorporated into the Affymetrix Gene-Chip microarrays of probes to match specified genes.”<sup>46</sup> However, upon consulting the supporting reference, it is clear that there was a misinterpretation: the “G-spots” referred in that paper are probes that contain 4 or more guanines.<sup>54</sup> In conclusion, up to now, there is no evidence from a hereditary or genetic point of view supporting the existence of the G-spot.

Assuming the existence of a suburethral erogenous structure, it is fair to assume that anterior vaginal wall and urinary incontinence surgeries (specially midurethral slings) may interfere with it. The available evidence does not support that these procedures deteriorate sexual function.<sup>55</sup> Kuhn et al. evaluated 18 women with pain after placement of a sling for urinary incontinence and showed that its removal did not improve orgasmic function.<sup>56</sup> One study that placed the G-spot in the vestibule, related vaginal surgery with worsening of sexual function.<sup>16</sup> If that was the case, vestibulectomies (used to treat some women with localized provoked vulvodynia) would have an ominous impact, which is not the case.<sup>57</sup>

Based on the finding of unequal nerve distribution in the vagina, Song et al. recommend that when possible approach of the posterior wall may be preferable.<sup>33</sup> There are reports of patients refusing prolapse surgery due to fear that the G-spot could be affected.<sup>55</sup>

Part of the success of the concept of G-spot is due to the flourishing market around it, ranging from specially designed dildos to stimulate it, to more or less complex surgeries aiming at increasing its size or function.<sup>20,40</sup>

Treatment for female sexual arousal disorder, by applying the vasodilator alprostadil (prostaglandin E1) to the clitoris and the

putative area of the G-spot, have shown moderate success.<sup>58</sup> A recent study on 52 women showed improvement of sexual function following sessions of platelet-rich plasma administration at the G-spot (no control group).<sup>59</sup>

Besides augmentation (with fat or hyaluronic acid), which some classify as a form of genital mutilation<sup>60</sup>, more complex procedures, such as the “g-spotplasty” have been proposed, without clear evidence of efficacy.<sup>61–63</sup>

The available data does not support the efficacy of these procedures and those are not recommended, namely by the ACOG, SOGC, and ISSVD.<sup>64–66</sup> For the SOGC “augmentation of the G-spot” is a mere marketing term, as the available studies are anecdotal.<sup>65</sup> In a survey among medical doctors and students, 71.0% indicated that there is never or rarely an indication for those procedures (56.0% if only plastic surgeons were considered).<sup>67</sup>

The separation between a clitoral and a vaginal (G-spot) orgasm seems to be reminiscence of the Freudian concepts – the pressure is the same: those who can only achieve it through “direct” clitoral stimulation are considered to be in a more immature sexual stage.<sup>68</sup> There is no evidence that vaginal and clitoral orgasms are different; pure vaginal stimulation during coitus probably does not happen. The concept of vaginal *vs.* clitoral orgasm strengthens the male role (penetration) in detriment of the independence gained by localizing the center of female sexual pleasure at the clitoris.<sup>8,46</sup>

In a 2019 review, Ostrzenski notes that “a physiological response cannot exist without an anatomical basis,”<sup>46</sup> however the fact that the perceived stimulation of the anterior vaginal wall elicits orgasm does not imply the existence of a particular anatomical structure.

This review shows the need for further studies on this topic, including reassessing women’s opinions and their partner’s, stimulation studies, and larger anatomical studies, including women of different ages and ethnic groups. While out of the scope of this review, animal studies are scarce, which is surprising given how much can be learned from phylogenetics.<sup>69</sup> The answer to this question may lie, for instance, in the reflex ovulator species, to whom female orgasm is part of survival.

While the G-spot is a field of debate, other erotogenic spots have already been added to the list, despite the lack of evidence: A or T (anterior fornix of the vagina), U (above and lateral to the urethral opening)<sup>15</sup> and more recently C (clitourethrovaginal complex).<sup>11</sup>

### Strengths and Limitations

This was a comprehensive review of the literature, including several perspectives of the question. However, the level of evidence available is low to very low, weakening the possible conclusions of the review. Since only G-spot/Gräfenberg spot was

searched, it is possible that relevant anatomical studies, addressing the anterior vaginal wall were missed.

## CONCLUSIONS

Female sexuality, including orgasm, is much more complex than a mere formula including hormones, psychological aspects, culture, religion, anatomy, and previous experience.

Most studies published so far about the G-spot favor its existence, but there is substantial disagreement even between these. This disagreement starts with its location: most authors describe it as a suburethral structure, but some place its location in the vulva.

Unanswered questions remain: does it exist? If so, where is it located, what size is it, what is its histological nature, what is its role in female sexuality, is it associated with female ejaculation?

Pressure put on women on the need to have a G-spot – those who cannot find it may feel “inadequate or abnormal.”<sup>70</sup> On the other hand, if it indeed exists, neglecting it may be equivalent to denying women the way to pleasurable experiences.<sup>17</sup>

The clitoris is still an unexplored continent, but the G-spot may just be another Atlantis.

**Corresponding Author:** Pedro Vieira-Baptista, MD, Hospital Lusíadas Porto, Av. da Boavista 171, 4050-115 Porto, Portugal; E-mail: [pedrovieirabaptista@gmail.com](mailto:pedrovieirabaptista@gmail.com)

*Conflict of Interest:* The authors report no conflicts of interest.

*Funding:* None.

## STATEMENT OF AUTHORSHIP

Pedro Vieira-Baptista: conceptualization, methodology, analysis, writing – original Draft; Joana Lima-Silva: conceptualization, methodology, writing – Review & Editing; Mario Preti: conceptualization, methodology, writing – Review & Editing; Joana Xavier: writing – Review & Editing; Pedro Vendeira: conceptualization, analysis, writing – Review & Editing; Colleen K. Stockdale: conceptualization, methodology, writing – Review & Editing.

## REFERENCES

1. Korda JB, Goldstein SW, Sommer F. The history of female ejaculation. *J Sex Med* 2010;7:1965–1975.
2. Syed R. [Knowledge of the “Gräfenberg zone” and female ejaculation in ancient Indian sexual science. A medical history contribution]. *Sudhoffs Arch* 1999;83:171–190.
3. Gräfenberg E. The role of urethra in female orgasm. *Int J Sexolog* 1950;III:145–148.
4. Addiego F, Belzer E, Comolli J, et al. Female ejaculation: A case study. *J Sex Res* 1981;17:13–21.

5. Ladas A, Whipple B, Perry J. The G spot: And other discoveries about human sexuality. New York: Holt, Rinehart, and Winston; 1982.
6. Belzer E. A review of female ejaculation and the Grafenberg Spot. *Women Health* 1984;9:5–16.
7. Semm K, Giese KP. [Ernst Gräfenberg, the life and medical accomplishments of the gynaecologist from Kiel. In honor of his 100th birthday on September 26, 1981 (author's transl)]. *Geburtshilfe Frauenheilkd* 1981;41:444–448.
8. Jannini EA, Whipple B, Kingsberg SA, et al. Who's afraid of the G-spot? *J Sex Med* 2010;7(1 Pt 1):25–34.
9. Ostrzenski A. G-spot anatomy: A new discovery. *J Sex Med* 2012;9:1355–1359.
10. Kanhai RC. Sensate vagina pedicled-spot for male-to-female transsexuals: The experience in the first 50 patients. *Aesthetic Plast Surg* 2016;40:284–287.
11. Jannini EA, Buisson O, Rubio-Casillas A. Beyond the G-spot: Clitourethrovaginal complex anatomy in female orgasm. *Nat Rev Urol* 2014;11:531–538.
12. Buisson O, Jannini EA. Pilot echographic study of the differences in clitoral involvement following clitoral or vaginal sexual stimulation. *J Sex Med* 2013;10:2734–2740.
13. Buisson O, Foldes P, Jannini E, et al. Coitus as revealed by ultrasound in one volunteer couple. *J Sex Med* 2010;7:2750–2754.
14. Shamseer L, Moher D, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: Elaboration and explanation. *Bmj* 2015;350:g7647.
15. Thabet SM. New findings and concepts about the G-spot in normal and absent vagina: Precautions possibly needed for preservation of the G-spot and sexuality during surgery. *J Obstet Gynaecol Res* 2013;39:1339–1346.
16. Thabet SM. Reality of the G-spot and its relation to female circumcision and vaginal surgery. *J Obstet Gynaecol Res* 2009;35:967–973.
17. Davidson Sr. JK, Darling CA, Conway-Welch C. The role of the Grafenberg Spot and female ejaculation in the female orgasmic response: An empirical analysis. *J Sex Marital Ther* 1989;15:102–120.
18. Wang X, Norris JL, Liu Y, et al. Risk behaviors for reproductive tract infection in women who have sex with women in Beijing, China. *PLoS One* 2012;7:e40114.
19. Darling CA, Davidson Sr. JK, Conway-Welch C. Female ejaculation: Perceived origins, the Grafenberg spot/area, and sexual responsiveness. *Arch Sex Behav* 1990;19:29–47.
20. Burri AV, Cherkas L, Spector TD. Genetic and environmental influences on self-reported G-spots in women: A twin study. *J Sex Med* 2010;7:1842–1852.
21. Ellibeş Kaya A, Çalışkan E. Women self-reported G-spot existence and relation with sexual function and genital perception. *Turk J Obstet Gynecol* 2018;15:182–187.
22. Kratochvíl S. [Sexual stimulation and the female orgasm]. *Cesk Psychiatr* 1993;89:191–199.
23. Zaviacíc M, Zaviacícová A, Holomán IK, et al. Female urethral expulsions evoked by local digital stimulation of the G-spot: Differences in the response patterns. *J Sex Res* 1988;24:311–318.
24. Perry J, Whipple B. Pelvic muscle strength of female ejaculators: Evidence in support of a new theory of orgasm. *J Sex Res* 1981;17:22–39.
25. Alzate H. Vaginal eroticism: A replication study. *Arch Sex Behav* 1985;14:529–537.
26. Hoch Z. Vaginal erotic sensitivity by sexological examination. *Acta Obstet Gynecol Scand* 1986;65:767–773.
27. Goldberg DC, Whipple B, Fishkin RE, et al. The Grafenberg spot and female ejaculation: A review of initial hypotheses. *J Sex Marital Ther* 1983;9:27–37.
28. Whipple B, Komisaruk BR. Brain (PET) responses to vaginal-cervical self-stimulation in women with complete spinal cord injury: Preliminary findings. *J Sex Marital Ther* 2002;28:79–86.
29. D'Amati G, di Gioia CR, Bologna M, et al. Type 5 phosphodiesterase expression in the human vagina. *Urology* 2002;60:191–195.
30. Shafik A, El Sibai O, Shafik AA, et al. The electrovaginogram: study of the vaginal electric activity and its role in the sexual act and disorders. *Arch Gynecol Obstet* 2004;269:282–286.
31. Foldes P, Buisson O. [Clitoris and G spot: An intimate affair]. *Gynecol Obstet Fertil* 2007;35:3–5.
32. Gravina GL, Brandetti F, Martini P, et al. Measurement of the thickness of the urethrovaginal space in women with or without vaginal orgasm. *J Sex Med* 2008;5:610–618.
33. Song YB, Hwang K, Kim DJ, et al. Innervation of vagina: Microdissection and immunohistochemical study. *J Sex Marital Ther* 2009;35:144–153.
34. Foldes P, Buisson O. The clitoral complex: A dynamic sonographic study. *J Sex Med* 2009;6:1223–1231.
35. Battaglia C, Nappi RE, Mancini F, et al. 3-D volumetric and vascular analysis of the urethrovaginal space in young women with or without vaginal orgasm. *J Sex Med* 2010;7(4 Pt 1):1445–1453.
36. Battaglia C, Nappi RE, Mancini F, et al. PCOS and urethrovaginal space: 3-D volumetric and vascular analysis. *J Sex Med* 2010;7:2755–2764.
37. Li T, Liao Q, Zhang H, et al. Anatomic distribution of nerves and microvascular density in the human anterior vaginal wall: Prospective study. *PLoS One* 2014;9:e110239.
38. Ostrzenski A, Krajewski P, Ganjei-Azar P, et al. Verification of the anatomy and newly discovered histology of the G-spot complex. *BJOG* 2014;121:1333–1339.
39. Maratos YK, Gombergh R, Cornier E, et al. The G-spot: an observational MRI pilot study. *Bjog* 2016;123:1542–1549.
40. Hoag N, Keast JR, O'Connell HE. The "G-Spot" is not a structure evident on macroscopic anatomic dissection of the vaginal wall. *J Sex Med* 2017;14:1524–1532.

41. Mazloomdoost D, Westermann LB, Mutema G, et al. Histologic anatomy of the anterior vagina and urethra. *Female Pelvic Med Reconstr Surg* 2017;23:329–335.
42. Aydın S, Sönmez FC, Karasu AFG, et al. Search for the G spot: Microvessel and nerve mapping of the paraurethral anterior vaginal wall. *Int Urogynecol J* 2020;31:2565–2572.
43. Ostrzenski A. Anatomic documentation of the G-spot complex role in the genesis of anterior vaginal wall ballooning. *Eur J Obstet Gynecol Reprod Biol* 2014;180:186–191.
44. Velazquez E, Barreto J, Cañete-Portillo S, et al. Penis and Distal Urethra. In: Milley S, editor. *Histology for pathologists*. 5th ed. Philadelphia: Wolters Kluwer; 2019.
45. Puppo V, Gruenwald I. Does the G-spot exist? A review of the current literature. *Int Urogynecol J* 2012;23:1665–1669.
46. Ostrzenski A. G-Spot anatomy and its clinical significance: A systematic review. *Clin Anat* 2019;32:1094–1101.
47. Bancroft J, Graham C, Janssen E, et al. The dual control model: Current status and future directions. *J Sex Res* 2009;46:121–142.
48. Zheng J, Islam RM, Skiba MA, et al. Associations between androgens and sexual function in premenopausal women: A cross-sectional study. *Lancet Diabetes Endocrinol* 2020;8:693–702.
49. Noroozadeh M, Tehrani FR, Mobarakabadi SS, et al. Sexual function and hormonal profiles in women with and without polycystic ovary syndrome: A population-based study. *Int J Impot Res* 2017;29:1–6.
50. Levin R. The G-spot—Reality or illusion? *Sexual and Relationship Therapy* 2003;18:117–119.
51. Ostrzenski A. The G-Spot anatomical structure spotted. *J Sex Med* 2018;15:803.
52. Puppo V, Puppo G. Anatomy of sex: Revision of the new anatomical terms used for the clitoris and the female orgasm by sexologists. *Clin Anat* 2015;28:293–304.
53. Li T, Ma Y, Zhang H, et al. Estrogen replacement regulates vaginal innervations in ovariectomized adult virgin rats: A histological study. *Biomed Res Int* 2017;2017:7456853.
54. Upton GJ, Langdon WB, Harrison AP. G-spots cause incorrect expression measurement in Affymetrix microarrays. *BMC Genomics* 2008;9:613.
55. Dwyer PL. Skene's gland revisited: Function, dysfunction and the G spot. *Int Urogynecol J* 2012;23:135–137.
56. Kuhn A, Burkhard F, Eggemann C, et al. Sexual function after suburethral sling removal for dyspareunia. *Surg Endosc* 2009;23:765–768.
57. David A, Bornstein J. Evaluation of long-term surgical success and satisfaction of patients after vestibulectomy. *J Low Genit Tract Dis* 2020;24:399–404.
58. Liao Q, Zhang M, Geng L, et al. Efficacy and safety of alprostadil cream for the treatment of female sexual arousal disorder: A double-blind, placebo-controlled study in chinese population. *J Sex Med* 2008;5:1923–1931.
59. Sukgen G, Ellibeş Kaya A, Karagün E, et al. Platelet-rich plasma administration to the lower anterior vaginal wall to improve female sexuality satisfaction. *Turk J Obstet Gynecol* 2019;16:228–234.
60. Puppo V. [G-spot does not exist: G-spot amplification (i.e. G-spot Augmentation, G-Spotplasty) is a female genital mutilation type IV]. *Ann Chir Plast Esthet* 2015;60:84–86.
61. Ostrzenski A. G-Spotplasty: A new surgical plastic intervention—the preliminary study. *Aesthetic Plast Surg* 2018;42:1126–1132.
62. Herold C, Motamedi M, Hartmann U, et al. G-spot augmentation with autologous fat transplantation. *J Turk Ger Gynecol Assoc* 2015;16:187–188.
63. Bachelet JT, Mojallal A, Boucher F. [Female genital surgery, G-spot amplification techniques—State of the science]. *Ann Chir Plast Esthet* 2014;59:344–347.
64. ACOG Committee Opinion No. 378. Vaginal “rejuvenation” and cosmetic vaginal procedures. *Obstet Gynecol* 2007;110:737–738.
65. Shaw D, Lefebvre G, Bouchard C, et al. Chirurgie esthétique génitale chez la femme. *J Obstet Gynaecol Can* 2016;38: S370–S375.
66. Vieira-Baptista P, Almeida G, Bogliatto F, et al. International society for the study of vulvovaginal disease recommendations regarding female cosmetic genital surgery. *J Low Genit Tract Dis* 2018;22:415–434.
67. Vieira-Baptista P, Lima-Silva J, Fonseca-Moutinho J, et al. Survey on aesthetic vulvovaginal procedures: What do portuguese doctors and medical students think? *Rev Bras Ginecol Obstet* 2017;39:415–423.
68. Pfaus JG, Quintana GR, Mac Cionnaith C, et al. The whole versus the sum of some of the parts: Toward resolving the apparent controversy of clitoral versus vaginal orgasms. *Socioaffect Neurosci Psychol* 2016;6:32578.
69. Pavličev M, Wagner G. The evolutionary origin of female orgasm. *J Exp Zool B Mol Dev Evol* 2016;326:326–337.
70. Whipple B, Perry JD. The G-spot: A modern gynecologic myth. *Am J Obstet Gynecol* 2002;187:519.