



## Role of Magnetic Resonance Imaging in the Management of Patients With Multiple Myeloma: A Consensus Statement

Meletios A. Dimopoulos, Jens Hillengass, Saad Usmani, Elena Zamagni, Suzanne Lentzsch, Faith E. Davies, Noopur Raje, Orhan Sezer, Sonja Zweegman, Jatin Shah, Ashraf Badros, Kazuyuki Shimizu, Philippe Moreau, Chor-Sang Chim, Juan José Lahuerta, Jian Hou, Artur Jurczynski, Hartmut Goldschmidt, Pieter Sonneveld, Antonio Palumbo, Heinz Ludwig, Michele Cavo, Bart Barlogie, Kenneth Anderson, G. David Roodman, S. Vincent Rajkumar, Brian G.M. Durie, and Evangelos Terpos

See accompanying editorial on page 537

Author affiliations appear at the end of this article.

Published online ahead of print at [www.jco.org](http://www.jco.org) on January 20, 2015.

Written on behalf of the International Myeloma Working Group.

Terms in [blue](#) are defined in the glossary, found at the end of this article and online at [www.jco.org](http://www.jco.org).

Authors' disclosures of potential conflicts of interest are found in the article online at [www.jco.org](http://www.jco.org). Author contributions are found at the end of this article.

Corresponding author: Meletios A. Dimopoulos, MD, Department of Clinical Therapeutics, National and Kapodistrian University of Athens School of Medicine, Alexandra General Hospital, 80 Vas. Sofias Ave, 11528, Athens, Greece; e-mail: [mdimop@med.uoa.gr](mailto:mdimop@med.uoa.gr).

© 2015 by American Society of Clinical Oncology

0732-183X/15/3306w-657w/\$20.00

DOI: 10.1200/JCO.2014.57.9961

### ABSTRACT

#### Purpose

The aim of International Myeloma Working Group was to develop practical recommendations for the use of magnetic resonance imaging (MRI) in multiple myeloma (MM).

#### Methods

An interdisciplinary panel of clinical experts on MM and myeloma bone disease developed recommendations for the value of MRI based on data published through March 2014.

#### Recommendations

MRI has high sensitivity for the early detection of marrow infiltration by myeloma cells compared with other radiographic methods. Thus, MRI detects bone involvement in patients with myeloma much earlier than the myeloma-related bone destruction, with no radiation exposure. It is the gold standard for the imaging of axial skeleton, for the evaluation of painful lesions, and for distinguishing benign versus malignant osteoporotic vertebral fractures. MRI has the ability to detect spinal cord or nerve compression and presence of soft tissue masses, and it is recommended for the workup of solitary bone plasmacytoma. Regarding smoldering or asymptomatic myeloma, all patients should undergo whole-body MRI (WB-MRI; or spine and pelvic MRI if WB-MRI is not available), and if they have > one focal lesion of a diameter > 5 mm, they should be considered to have symptomatic disease that requires therapy. In cases of equivocal small lesions, a second MRI should be performed after 3 to 6 months, and if there is progression on MRI, the patient should be treated as having symptomatic myeloma. MRI at diagnosis of symptomatic patients and after treatment (mainly after autologous stem-cell transplantation) provides prognostic information; however, to date, this does not change treatment selection.

*J Clin Oncol* 33:657-664. © 2015 by American Society of Clinical Oncology

### INTRODUCTION

Bone disease, characterized by the presence of osteolytic lesions, bone fractures, or osteoporosis, is a significant cause of morbidity and mortality in multiple myeloma (MM). Therefore, the guidelines of the International Myeloma Working Group (IMWG) suggest that the presence of even asymptomatic bone disease on conventional radiography is a criterion of symptomatic MM that requires treatment.<sup>1</sup>

In 2009, the IMWG indicated that whole-body (WB) x-ray (WBXR) remains the gold standard for the evaluation of MM-related bone disease.<sup>2</sup> However, the detection limit of WBXR is low; to detect an

osteolytic lesion by WBXR, a proportion of at least 30% to 50% of the trabecular bone has to be resorbed.<sup>3</sup> Moreover, WBXR is not a suitable technique for the diagnosis of myeloma-related osteoporosis, has low visualization of the spine and pelvis, and cannot accurately depict the cause of painful lesions in patients with MM. In previous recommendations, the IMWG supported the implementation of magnetic resonance imaging (MRI) in the absence of osteolytic lesions on WBXR.<sup>2</sup> However, the IMWG did not suggest the use of MRI for the definition of symptomatic myeloma. Thus, to date, a patient with focal lesions on MRI but with no lytic lesions on WBXR and with no other CRAB (hypercalcemia, renal failure, anemia, and bone disease)

**Table 1.** Levels of Evidence and Grades of Recommendations

Rating	Description
Level	
I	Evidence obtained from meta-analysis of multiple well-designed, controlled studies; randomized trials with low false-positive and low false-negative errors (high power)
II	Evidence obtained from at least one well-designed experimental study; randomized trials with high false-positive and/or false-negative errors (low power)
III	Evidence obtained from well-designed, quasiexperimental studies (eg, nonrandomized, controlled single-group, pre-post, cohort, time, or matched case-control studies)
IV	Evidence from well-designed, nonexperimental studies (eg, comparative and correlational descriptive and case studies)
V	Evidence from case reports and clinical examples
Grade	
A	Evidence of type I or consistent findings from multiple studies of types II, III, or IV
B	Evidence of type II, III, or IV; findings are generally consistent
C	Evidence of type II, III, or IV; findings are inconsistent
D	Little or no systematic empiric evidence

criteria is considered to have smoldering or asymptomatic myeloma (SMM), and follow-up with no treatment is recommended. Several novel data stress the value of MRI in this setting, and we suggest that the current treatment practice be changed for these patients. Our aim was to produce useful recommendations for the use of MRI in everyday clinical practice for the management of patients with myeloma and introduce novel MRI criteria for the definition of SMM.

## METHODS

An interdisciplinary panel of experts on myeloma bone disease and MRI performance in patients with myeloma developed the recommendations based on evidence of published clinical or observational studies, meta-analyses, and systematic reviews through March 2014. Expert consensus was used to propose recommendations in the absence of sufficiently published data. Levels of evidence and grades of recommendations were used according to established criteria (Table 1). The statement was drafted and circulated among all panel members, followed by subsequent rounds of revision until consensus was achieved.

## MRI TECHNIQUES FOR MYELOMA

Several MRI techniques have been developed for the assessment of bone marrow involvement in MM: T1 weighted, T2 weighted with fat suppression, short time inversion recovery, and gadolinium T1 weighted with fat suppression.<sup>4</sup> Myeloma lesions typically show a low signal intensity on T1-weighted images, a high signal intensity on T2-weighted and short time inversion recovery images, and often enhancement on gadolinium-enhanced images.<sup>5,6</sup> Table 2 summarizes the minimum technical requirements of these techniques for the management of patients with myeloma.

Limitations of MRI are the prolonged acquisition time, high cost, exclusion of patients with metal devices in their bodies, difficulties in cases of claustrophobic patients, and limited field of view. To override these restrictions, a WB-MRI methodology, which does not usually require contrast infusion, was developed. The time of WB-MRI is

**Table 2.** Minimum Technical Requirements for Use of MRI in Patients With Myeloma

Type	Requirement
Coil	Target volume adapted surface-reception coil system
Examination volume	Whole spine and pelvic MRI (if WB-MRI is not available)
Orientation	Sagittal axial for spine; axial coronal for pelvis
Parameter	
Slice thickness	4 mm (definitely not > 5 mm)
Matrix	512 pixels
Weighting	T1, T2
Sequences	Sagittal T1 TSE, sagittal or coronal TIRM (STIR), axial T1 and T2 TSE, chemical shift, and T1 post contrast
Abbreviations: MRI, magnetic resonance imaging; STIR, short time inversion recovery; TIRM, turbo inversion recovery magnitude; TSE, turbo spin echo; WB, whole body.	

approximately 45 minutes. Although of interest, this newer technique is not yet widely employed.

The MRI methods mentioned use MRI exquisite contrast and spatial resolution for the depiction of WB anatomy and specific tissue composition in detail. A newer MRI sequence is diffusion-weighted imaging (DWI). This functional technique demonstrates alterations in intra- and extracellular water content from disruption of the transmembrane water flux that are visible before identified changes on the morphologic routine sequences.<sup>7-9</sup> One disadvantage of DWI is that the apparent diffusion coefficient (ADC) is not exclusively influenced by diffusion but also by perfusion. However, improved sequences are under development to differentiate both influences.<sup>10</sup> DWI can be used to detect regions with bone marrow infiltration for both diagnosis and monitoring of treatment response.<sup>11,12</sup> In patients with MM, the ADC was reproducible<sup>13</sup> and correlated with bone marrow cellularity and microvessel density (MVD).<sup>14</sup>

The dynamic contrast-enhanced MRI (DCE-MRI) is another MRI technique in which the distribution of a contrast agent inside and outside the blood vessels is assessed by computer-based analysis of repeated images over time. The analysis provides data for blood volume and vessel permeability for the assessment of microcirculation.<sup>15,16</sup> More importantly, in patients with MM, DCE-MRI measurements have been correlated with marrow angiogenesis and MVD<sup>17</sup> as well as with angiogenic response to therapy.<sup>18</sup> Regarding DCE-MRI sampling rate and model, there are two pharmacokinetic models (proposed by Brix and Tofts) that have been applied in the literature. However, a comparison of these models demonstrated that the Brix model is slightly more robust.<sup>19</sup> Because DCE-MRI has not been established in clinical routine, no definite sequence can be recommended.

Positron emission tomography (PET) in combination with MRI is a novel and promising new methodology in which PET detects active focal lesions, while MRI shows the location of the lesions and provides information on myeloma cell infiltration of the bone marrow. Especially in patients who reach a complete remission (CR), this technique might be able to localize residual sites of disease activity and therefore may help to guide treatment in the future.<sup>20</sup>

## MRI PATTERNS OF MARROW INVOLVEMENT

Five MRI patterns of marrow involvement in myeloma have been recognized: normal appearance of bone marrow, focal involvement (positive focal lesion is considered lesion of diameter  $\geq 5$  mm), homogeneous diffuse infiltration, combined diffuse and focal infiltration, and variegated or salt-and-pepper pattern with inhomogeneous bone marrow with interposition of fat islands.<sup>21,22</sup> Low tumor burden is usually associated with a normal MRI pattern, but a high tumor burden is usually suspected when there is diffuse hypointense change on T1-weighted images, diffuse hyperintensity on T2-weighted images, and enhancement with gadolinium injection.<sup>23</sup> In several studies, the percentage of symptomatic patients with each of the abnormal MRI bone marrow patterns has ranged from 18% to 50% for focal pattern, 25% to 43% for diffuse pattern, and 1% to 5% for variegated pattern.<sup>16</sup> The Durie-Salmon PLUS system uses the number of focal lesions (from focal or combined focal and diffuse patterns) for the staging of a patient with myeloma rather than the diffuse or salt-and-pepper pattern.<sup>24</sup>

## MRI IN SYMPTOMATIC MYELOMA

### **MRI Versus Conventional Radiography and Other Imaging Techniques for Detection of Bone Involvement**

MRI is more sensitive compared with WBXR for the detection of bone involvement in MM. In the largest series of patients published to date, MRI was compared with WBXR in 611 patients who received tandem autologous stem-cell transplantation (ASCT). MRI and WBXR detected focal and osteolytic lesions in 74% and 56% of the imaged anatomic sites, respectively. Furthermore, 52% of 267 patients with normal WBXR had focal lesions on MRI. More precisely, MRI detected more focal lesions compared with lytic lesions in WBXR in the spine (78% v 16%;  $P < .001$ ), pelvis (64% v 28%;  $P < .001$ ), and sternum (24% v 3%;  $P < .001$ ). WBXR had better performance than MRI in the ribs (10% v 43%;  $P < .001$ ) and long bones (37% v 48%;  $P = .006$ ) and equal results in the skull and shoulders.<sup>25</sup> Similar results had been previously reported in smaller studies, where MRI was superior to WBXR for the detection of focal versus osteolytic lesions in the pelvis (75% v 46% of patients) and spine (76% v 42%), especially in the lumbar spine.<sup>26-30</sup> A recent meta-analysis confirmed the superiority of MRI over WBXR regarding the detection of focal lesions and showed that MRI especially outscores WBXR in the axial skeleton but not in the ribs.<sup>31</sup>

Although it is clear that MRI can detect bone marrow focal lesions long before the development of osteolytic lesions on WBXR, other imaging techniques such as PET combined with computed tomography (CT), CT, or WB-CT detect more osteolytic lesions compared with WBXR.<sup>31</sup> Do we have any evidence that MRI is superior to the other techniques in depicting bone involvement in myeloma? In a study with 41 patients with newly diagnosed MM, WB-MRI was found superior to WB-CT in detecting lesions in the skeleton.<sup>32</sup> In a prospective study, Zamagni et al<sup>33</sup> compared MRI of the spine and pelvis with WBXR versus PET-CT in 46 patients with MM at diagnosis. Although PET-CT was superior to WBXR in detecting lytic lesions in 46% of patients (19% had negative WBXR), it failed to reveal abnormal findings in 30% of patients who had abnormal MRI in the

same areas, mainly of diffuse pattern. In that study, the combination of spine and pelvic MRI with PET-CT detected both medullary and extramedullary active myeloma sites in almost all patients (92%). Nevertheless, the Arkansas group was not able to confirm any superiority of MRI over PET-CT in the detection of more focal lesions in a large number of patients ( $n = 303$ ) within the Total Therapy 3 protocols.<sup>34</sup> Still, in 188 patients who had  $\geq 1$  focal lesion on MRI, MRI was superior to PET-CT regarding the detection of higher number of focal lesions ( $P = .032$ ). Furthermore, in this study, the presence of diffuse marrow pattern was not taken into consideration as an abnormal MRI finding.<sup>34</sup> Compared with MIBI (sestamibitechnetium-99m) scan, WB-MRI detected more lesions in the vertebrae and long bones, produced similar results in the skull, and was inferior in the ribs.<sup>35</sup> One important question on this point is the value of WB-MRI, which is not available everywhere, over MRI of the spine and pelvis. In 100 patients with MM or monoclonal gammopathy of undetermined significance (MGUS) who underwent WB-MRI, 10% presented with focal lesions merely in the extra-axial skeleton. These lesions would have been ignored if only MRI of the spine and pelvis had been performed.<sup>36</sup>

Other advantages of MRI over WBXR and CT include the discrimination of myeloma from normal marrow<sup>4,37</sup>; this finding can help in the differential diagnosis between myeloma and benign cause of a vertebral fracture. This is of extreme importance in cases of patients with a vertebral fracture and no other CRAB criteria or lytic lesions. MRI can also accurately illustrate the spinal cord and/or nerve root compression for surgical intervention or radiation therapy.<sup>2,4</sup> Furthermore, the presence of soft tissue extension of MM and presence of extramedullary plasmacytomas, which develop in approximately 10% to 20% of patients during the course of their disease, can be precisely visualized by WB-MRI.<sup>38-40</sup> MRI can also help in the better evaluation of avascular necrosis of the femoral head<sup>41</sup> and presence of soft tissue amyloid deposits.<sup>42</sup> Moreover, tumor load can be assessed and monitored by MRI, even in patients with nonsecretory or oligosecretory MM.<sup>43</sup>

### **Consensus Statement**

MRI is the imaging gold-standard method for the detection of bone marrow involvement in MM (grade A). We stress that MRI detects bone marrow involvement and not bone destruction. MRI of the spine and pelvis can detect approximately 90% of focal lesions in MM, and thus, it can be used in cases where WB-MRI is not available (grade B). MRI is the procedure of choice to evaluate a painful lesion in patients with myeloma, mainly in the axial skeleton, and detect spinal cord compression (grade A). MRI is particularly useful in the evaluation of collapsed vertebrae, especially when myeloma is not active, where the possibility of osteoporotic fracture is high (grade B).

### **Prognostic Value of MRI**

The prognostic significance of MRI findings in symptomatic myeloma has been evaluated. The largest study in the literature included 611 patients who received tandem ASCT-based protocols. Focal lesions detected by spinal MRI and not seen on WBXR independently correlated with overall survival (OS). Resolution of the focal lesions on MRI after treatment occurred in 60% of the patients who had superior survival. At disease progression after CR, MRI revealed new focal lesions in 26% of patients, enlargement of previous focal lesions in 28%, and both features in 15%.<sup>25</sup> In a more recent analysis

by the same group involving 429 patients, patients who had > seven focal lesions on MRI (n = 147) had a 73% probability of 3-year OS versus 86% for those who had zero to seven focal lesions (n = 235) and 81% for those who had diffuse pattern of marrow infiltration (n = 47;  $P = .04$ ). PET-CT and WBXR also produced similar results in the univariable analysis. In the multivariable analysis, from the imaging variables, only the presence of > two osteolytic lesions on WBXR at diagnosis and presence of > three focal lesions on PET-CT 7 days after ASCT had independent prognostic value for inferior OS ( $P = .01$  and  $.03$ , respectively). However, we have to mention the high percentage of patients (232 [54%] of 429) who had no detectable osteolytic lesions by WBXR and the absence of evaluation of diffuse MRI pattern in this study.<sup>44</sup>

The MRI pattern of marrow infiltration has also been reported to have prognostic significance in newly diagnosed patients with symptomatic disease.<sup>23,45,46</sup> In the conventional chemotherapy era, Moulopoulos et al<sup>23</sup> reported that the median OS of patients with newly diagnosed MM was 24 months if they had diffuse MRI pattern versus 51, 52, and 56 months for those with focal, variegated, and normal patterns, respectively ( $P = .001$ ). This is possibly because diffuse MRI marrow pattern correlates with increased angiogenesis and advanced disease features.<sup>47,48</sup> The same group also reported the prognostic value of MRI patterns in 228 patients with symptomatic MM who received first-line regimens based on novel agents. Patients with diffuse pattern had inferior survival compared with patients with other MRI patterns; moreover, the combination of diffuse MRI pattern, International Staging System stage III, and high-risk cytogenetics could identify a group of patients with poor survival (ie, median, 21 months; probability of 3-year OS, 35%).<sup>46</sup> Another study in 126 patients with newly diagnosed symptomatic myeloma who underwent ASCT showed that the diffuse and variegated MRI patterns had an independent predictive value for disease progression (hazard ratio [HR], 1.922;  $P = .008$ ).<sup>48</sup> Finally, in patients with progressive or relapsed MM, an increased DCE-MRI signal indicated shorter progression-free survival, possibly because of its association with higher MVD.<sup>15</sup>

### Consensus Statement

The number of MRI focal lesions (> seven; grade A) and presence of diffuse pattern (grade B) correlate with inferior survival. Prospective clinical studies are needed to define if these patients have to be treated in a different or more aggressive way.

### MRI and Response to Antimyeloma Therapy

An interesting finding is that a change in MRI pattern correlates with response to therapy. Moulopoulos et al<sup>49</sup> first reported in the era of conventional chemotherapy that CR is characterized by complete resolution of the preceding marrow abnormality, whereas partial response is characterized by changeover of diffuse pattern to variegated or focal pattern. In a retrospective study that was conducted in the era of novel agents, response to treatment was compared with changes in infiltration pattern on WB-MRI before and after ASCT (n = 100). There was a strong correlation between response to antimyeloma therapies and changes in both diffuse ( $P = .004$ ) and focal ( $P = .01$ ) MRI patterns. Furthermore, the number of focal lesions on second MRI was of prognostic significance for OS ( $P = .001$ ).<sup>50</sup> Another study in 33 patients who underwent ASCT showed that WB-MRI data demonstrated progressive disease in 10 patients (30%) and response to high-dose therapy in 23 (70%). Eight (80%) of the 10 patients with progressive disease revealed intramedullary lesions, and two patients

(20%) had intra- and extramedullary lesions. WB-MRI had a sensitivity of 64%, specificity of 86%, positive predictive value of 70%, negative predictive value of 83%, and accuracy of 79% for detection of remission.<sup>51</sup> This study supports that one of the disadvantages of MRI is that it often provides false-positive results because of persistent nonviable lesions. Thus, PET-CT might be more suitable than MRI for determination of remission status.<sup>52</sup> Indeed, in a large study of 191 patients, PET-CT revealed faster change of imaging findings than MRI in patients who responded to therapy.<sup>53</sup> It seems that PET-CT normalization after treatment can offer more information compared with MRI for the better definition of CR.<sup>54</sup>

To improve the results of MRI for the most accurate detection of remission, DW-MRI has been recently used. In a first preliminary report, ADC values in active myeloma were significantly higher than marrow in remission.<sup>55</sup> Furthermore, the mean ADC increased in 95% of responding patients and decreased in all nonresponders (n = 5;  $P = .002$ ). An increase of ADC by 3.3% was associated with response, having a sensitivity of 90% and specificity of 100%. Furthermore, there was a negative correlation between changes of ADC and changes of biochemical markers of response ( $r = -0.614$ ;  $P = .001$ ).<sup>56</sup> Large prospective clinical studies are definitely justified by these results.

### Consensus Statement

MRI might help in the better definition of CR (grade D; panel consensus). Nevertheless, the high number of false-positive results suggests that its combination with methods that reveal active lesions (ie, PET-MRI) or another imaging method, such as PET-CT, might be of more value in this setting. Thus, the systematic performance of MRI for the follow-up of patients, before or after different therapies, in the absence of clinical indications is not recommended. Novel clinical studies have to include MRI for the response evaluation in an effort to clarify the role of MRI in this important field of myeloma therapy.

### VALUE OF MRI IN DEFINITION OF SMM

The presence of lytic lesions by WBXR is included in the definition of symptomatic myeloma, based on studies showing that patients with  $\geq$  one lytic lesion on WBXR have a median time to progression (TTP) of 10 months.<sup>57</sup> However, in patients with no osteolytic lesions on WBXR, MRI reveals abnormal marrow appearance in 20% to 50%<sup>22,23,58-60</sup>; these patients are at higher risk for progression. Moulopoulos et al<sup>58</sup> reported that patients with SMM and abnormal MRI studies required therapy after a median of 16 months versus 43 months for those with normal MRI ( $P < .01$ ). Hillengass et al<sup>59</sup> evaluated WB-MRI in 149 patients with SMM. Focal lesions were detected in 42 patients (28%), while > one focal lesion was present in 23 patients (15%) who had high risk of progression (HR, 4.05;  $P < .001$ ). The median TTP was 13 months, and the progression rate at 2 years was 70%. On multivariable analysis, presence of > one focal lesion remained a significant predictor of progression after adjusting for other risk factors, including bone marrow plasmacytosis, serum and urine M protein levels, and suppression of uninvolved immunoglobulins. In the same study, diffuse marrow infiltration on MRI was also associated with increased risk for progression (HR, 3.5;  $P < .001$ ). Kastiris et al<sup>60</sup> also showed in a study of 98 patients with SMM that abnormal marrow pattern on MRI of the spine, which was present in

21% of patients, was associated with high risk of progression, with a median TTP to symptomatic myeloma of 15 months ( $P = .001$ ).

The identification of patients with SMM who are at high risk for progression is of great importance, because these patients may benefit from immediate therapy. A recent randomized study from the Spanish Myeloma Study Group compared the combination of lenalidomide plus low-dose dexamethasone (Rd) versus observation in patients with high-risk SMM (MRI was not used for defining high-risk SMM). TTP was significantly longer with Rd compared with observation (median, not reached  $\nu$  21 months, respectively;  $P < .001$ ). More importantly, Rd offered OS advantage (probability of 3-year survival, 94%  $\nu$  80%, respectively;  $P = .03$ ).<sup>61</sup>

An important issue is whether patients who have  $\geq$  two small focal lesions ( $< 5$  mm) should be considered as having symptomatic myeloma and how to manage them. Recently, the Heidelberg group analyzed data of 63 patients with SMM who had  $\geq$  two WB-MRIs performed for follow-up before progression to symptomatic disease. The definition of radiologic progression according to MRI findings included one of the following: development of a new focal lesion, increase of the diameter of an existing focal lesion, or detection of novel or progressive diffuse MRI pattern. The second MRI was performed 3 to 6 months after the performance of the first MRI. Evaluation of response according to IMWG criteria was also performed. Progressive disease according to MRI was observed in approximately 50% of patients, whereas 40% of patients developed symptomatic MM based on the CRAB criteria. In the multivariable analysis, progressive disease according to MRI was an independent prognostic factor for progression. Patients with stable MRI findings had no higher risk of progression, even when focal lesions were present on the initial MRI.<sup>62</sup> Prospective clinical trials should be conducted to confirm these findings.

### Consensus Statement

We recommend that patients with  $>$  one unequivocal focal lesion (diameter of  $> 5$  mm) should be considered to have symptomatic myeloma that requires therapy (grade B). Patients with equivocal focal lesions should repeat the MRI after 3 to 6 months, and in cases of MRI progression, patients should be considered as symptomatic patients who need therapy (grade C; panel consensus). The biopsy of such lesions should be encouraged. Regarding diffuse MRI marrow pattern, we need additional studies before its incorporation into the definition of symptomatic myeloma.

### MRI FINDINGS IN MGUS

MGUS by definition is characterized by the absence of osteolytic lesions. However, patients with MGUS have higher incidences of osteoporosis and vertebral fractures compared with the normal population.<sup>63,64</sup> In a small study that included 37 patients with MGUS or SMM, MRI abnormalities were detected in 20%. These patients had a higher TTP to symptomatic myeloma compared with patients with a normal MRI who did not experience progression after a median follow-up of 30 months.<sup>65</sup> A prospective study involving 331 patients with MGUS or SMM revealed that the detection of multiple ( $>$  one) focal lesions by MRI conferred an increased risk of progression.<sup>66</sup> In another large study, which included only patients with MGUS ( $n = 137$ ) who underwent WB-MRI at diagnosis, a focal infiltration pattern was detected in 23%. Independent prognostic factors for progression

to symptomatic myeloma included the presence and number of focal lesions and value of M-protein.<sup>67</sup>

### Consensus Statement

WB-MRI identifies patients with MGUS with focal lesions that possibly reflect infiltration by monoclonal plasma cells in the bone marrow. These patients seem to have increased risk of progression to myeloma. To date, MRI is not recommended as part of the routine workup for patients with MGUS unless there are clinical features that increase suspicion.

### MRI AND SOLITARY BONE PLASMACYTOMA

The diagnosis of solitary plasmacytoma of the bone (SBP) includes the presence of a solitary bone lesion, with a confirmed infiltration by plasma cells on biopsy of the lesion, absence of clonal plasma cells on the trephine bone marrow biopsy, and no CRAB criteria met. Although definitive radiotherapy usually eradicates the local disease, a majority of patients will develop MM because of the growth of previously occult lesions not detected by WBXR.<sup>39</sup> Mouloupoulos et al<sup>38</sup> reported that spinal MRI revealed additional focal lesions in four of 12 patients with SBP. After treatment with radiotherapy to the painful lesion, three patients developed systemic disease within 18 months from diagnosis. Furthermore, Liebross et al<sup>68</sup> observed that among patients with SBP with spinal disease, seven of eight staged by WBXR

**Table 3.** Recommendations for Use of MRI in Myeloma, MGUS, and SBP

Recommendation	Grade
<b>Symptomatic myeloma</b>	
MRI is imaging gold-standard method for detection of bone marrow involvement in MM	A
MRI of spine and pelvis can be used in cases where WB-MRI is not available	B
MRI is procedure of choice to evaluate painful lesion and spinal cord compression	A
MRI can differentiate malignant from nonmalignant vertebral fractures	B
MRI provides accurate visualization of soft tissue extension and extramedullary plasmacytoma	A
Focal pattern provides prognostic information	A
Diffuse pattern also correlates with inferior prognosis	B
MRI might help in better definition of CR	D
MRI is not recommended for treatment follow-up	
<b>Asymptomatic myeloma</b>	
Patients with $>$ one unequivocal focal lesion (diameter $> 5$ mm) should be considered to have symptomatic myeloma that requires therapy	B
Patients with equivocal focal lesions should repeat MRI after 3 to 6 months; in cases of MRI progression, patients should be considered to be symptomatic and to require therapy	C
<b>MGUS and SBP</b>	
MRI is not recommended as part of routine workup of patients with MGUS unless there are clinical features that increase suspicion	
MRI should be part of staging procedures in patients with SBP	A

Abbreviations: CR, complete response; MGUS, monoclonal gammopathy of undetermined significance; MM, multiple myeloma; MRI, magnetic resonance imaging; SBP, solitary bone plasmacytoma; WB, whole body.

alone developed MM, compared with only one of seven patients who also underwent spinal MRI.

### Consensus Statement

MRI should be part of the staging procedures in patients with SBP to better assess the extent of the local tumor and reveal occult lesions elsewhere (grade A).

## DISCUSSION

In summary, MRI describes the pattern of myelomatous infiltration of the bone marrow and is the procedure of choice for the evaluation of painful lesions in patients with myeloma for the detection of spinal cord compression and differentiation of malignant from nonmalignant vertebral fractures. MRI provides significant prognostic information in patients with symptomatic disease and may be found useful in the better definition of CR. More importantly, MRI can reclassify patients with SMM to symptomatic MM, and patients with > one unequivocal focal lesion (diameter of > 5 mm) should be considered as having symptomatic myeloma that requires therapy. Finally, MRI is part of the staging workup of SBP. A summary of the recommendations is listed in Table 3.

## REFERENCES

1. International Myeloma Working Group: Criteria for the classification of monoclonal gammopathies, multiple myeloma and related disorders: A report of the International Myeloma Working Group. *Br J Haematol* 121:749-757, 2003
2. Dimopoulos M, Terpos E, Comenzo RL, et al: International Myeloma Working Group consensus statement and guidelines regarding the current role of imaging techniques in the diagnosis and monitoring of multiple myeloma. *Leukemia* 23:1545-1556, 2009
3. Durie BG, Salmon SE: A clinical staging system for multiple myeloma: Correlation of measured myeloma cell mass with presenting clinical features, response to treatment, and survival. *Cancer* 36:842-854, 1975
4. Mouloupoulos LA, Dimopoulos MA: Magnetic resonance imaging of the bone marrow in hematologic malignancies. *Blood* 90:2127-2147, 1997
5. Libshitz HI, Malthouse SR, Cunningham D, et al: Multiple myeloma: Appearance at MR imaging. *Radiology* 182:833-837, 1992
6. Weininger M, Lauterbach B, Knop S, et al: Whole-body MRI of multiple myeloma: Comparison of different MRI sequences in assessment of different growth patterns. *Eur J Radiol* 69:339-345, 2008
7. Attariwala R, Picker W: Whole body MRI: Improved lesion detection and characterization with diffusion weighted techniques. *J Magn Reson Imaging* 38:253-268, 2013
8. Müller MF, Edelman RR: Echo planar imaging of the abdomen. *Top Magn Reson Imaging* 7:112-119, 1995
9. Wang Y: Description of parallel imaging in MRI using multiple coils. *Magn Reson Med* 44:495-499, 2000
10. Lemke A, Stieltjes B, Schad LR, et al: Toward an optimal distribution of B values for intravoxel incoherent motion imaging. *Magn Reson Imaging* 29:766-776, 2011

11. Xu X, Ma L, Zhang JS, et al: Feasibility of whole body diffusion weighted imaging in detecting bone metastasis on 3.0T MR scanner. *Chin Med Sci J* 23:151-157, 2008
12. Padhani AR, Koh DM: Diffusion MR imaging for monitoring of treatment response. *Magn Reson Imaging Clin N Am* 19:181-209, 2011
13. Messiou C, Collins DJ, Morgan VA, et al: Optimising diffusion weighted MRI for imaging metastatic and myeloma bone disease and assessing reproducibility. *Eur Radiol* 21:1713-1718, 2011
14. Hillengass J, Bäuerle T, Bartl R, et al: Diffusion-weighted imaging for non-invasive and quantitative monitoring of bone marrow infiltration in patients with monoclonal plasma cell disease: A comparative study with histology. *Br J Haematol* 153:721-728, 2011
15. Hillengass J, Wasser K, Delorme S, et al: Lumbar bone marrow microcirculation measurements from dynamic contrast-enhanced magnetic resonance imaging is a predictor of event-free survival in progressive multiple myeloma. *Clin Cancer Res* 13:475-481, 2007
16. Hillengass J, Landgren O: Challenges and opportunities of novel imaging techniques in monoclonal plasma cell disorders: Imaging "early myeloma." *Leuk Lymphoma* 54:1355-1363, 2013
17. Huang SY, Chen BB, Lu HY, et al: Correlation among DCE-MRI measurements of bone marrow angiogenesis, microvessel density, and extramedullary disease in patients with multiple myeloma. *Am J Hematol* 87:837-839, 2012
18. Zechmann CM, Traine L, Meissner T, et al: Parametric histogram analysis of dynamic contrast-enhanced MRI in multiple myeloma: A technique to evaluate angiogenic response to therapy? *Acad Radiol* 19:100-108, 2012
19. Zwick S, Brix G, Tofts PS, et al: Simulation-based comparison of two approaches frequently used for dynamic contrast-enhanced MRI. *Eur Radiol* 20:432-442, 2010
20. Fraioli F, Punwani S: Clinical and research applications of simultaneous positron emission tomography and MRI. *Br J Radiol* 87:20130464, 2014

## AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Disclosures provided by the authors are available with this article at [www.jco.org](http://www.jco.org).

## AUTHOR CONTRIBUTIONS

**Conception and design:** Meletios A. Dimopoulos, Evangelos Terpos

**Administrative support:** Evangelos Terpos

**Provision of study materials or patients:** Jens Hillengass

**Collection and assembly of data:** Meletios A. Dimopoulos, Jens Hillengass, Hartmut Goldschmidt, Pieter Sonneveld, Evangelos Terpos

**Data analysis and interpretation:** Meletios A. Dimopoulos, Saad Usmani, Elena Zamagni, Suzanne Lentzsch, Faith E. Davies, Noopur Raje, Orhan Sezer, Sonja Zweegman, Jatin Shah, Ashraf Badros, Kazuyuki Shimizu, Philippe Moreau, Chor-Sang Chim, Juan José Lahuerta, Jian Hou, Artur Jurczynszyn, Hartmut Goldschmidt, Pieter Sonneveld, Antonio Palumbo, Heinz Ludwig, Michele Cavo, Bart Barlogie, Kenneth Anderson, G. David Roodman, S. Vincent Rajkumar, Brian G.M. Durie, Evangelos Terpos

**Manuscript writing:** All authors

**Final approval of manuscript:** All authors

21. Baur-Melnyk A, Buhmann S, Dürr HR, et al: Role of MRI for the diagnosis and prognosis of multiple myeloma. *Eur J Radiol* 55:56-63, 2005
22. Mouloupoulos LA, Varma DG, Dimopoulos MA, et al: Multiple myeloma: Spinal MR imaging in patients with untreated newly diagnosed disease. *Radiology* 185:833-840, 1992
23. Mouloupoulos LA, Gika D, Anagnostopoulos A, et al: Prognostic significance of magnetic resonance imaging of bone marrow in previously untreated patients with multiple myeloma. *Ann Oncol* 16:1824-1828, 2005
24. Durie BG: The role of anatomic and functional staging in myeloma: Description of Durie/Salmon plus staging system. *Eur J Cancer* 42:1539-1543, 2006
25. Walker R, Barlogie B, Haessler J, et al: Magnetic resonance imaging in multiple myeloma: Diagnostic and clinical implications. *J Clin Oncol* 25:1121-1128, 2007
26. Ludwig H, Frühwald F, Tscholakoff D, et al: Magnetic resonance imaging of the spine in multiple myeloma. *Lancet* 2:364-366, 1987
27. Ghanem N, Lohrmann C, Engelhardt M, et al: Whole-body MRI in the detection of bone marrow infiltration in patients with plasma cell neoplasms in comparison to the radiological skeletal survey. *Eur Radiol* 16:1005-1014, 2006
28. Lecouvet FE, Malghem J, Michaux L, et al: Skeletal survey in advanced multiple myeloma: Radiographic versus MR imaging survey. *Br J Haematol* 106:35-39, 1999
29. Tertti R, Alanen A, Remes K: The value of magnetic resonance imaging in screening myeloma lesions of the lumbar spine. *Br J Haematol* 91:658-660, 1995
30. Narquin S, Ingrand P, Azais I, et al: Comparison of whole-body diffusion MRI and conventional radiological assessment in the staging of myeloma. *Diagn Interv Imaging* 94:629-636, 2013
31. Regelink JC, Minnema MC, Terpos E, et al: Comparison of modern and conventional imaging techniques in establishing multiple myeloma-related

bone disease: A systematic review. *Br J Haematol* 162:50-61, 2013

32. Baur-Melnyk A, Buhmann S, Becker C, et al: Whole-body MRI versus whole-body MDCT for staging of multiple myeloma. *AJR Am J Roentgenol* 190:1097-1104, 2008

33. Zamagni E, Nanni C, Patriarca F, et al: A prospective comparison of 18F-fluorodeoxyglucose positron emission tomography-computed tomography, magnetic resonance imaging and whole-body planar radiographs in the assessment of bone disease in newly diagnosed multiple myeloma. *Haematologica* 92:50-55, 2007

34. Waheed S, Mitchell A, Usmani S, et al: Standard and novel imaging methods for multiple myeloma: Correlates with prognostic laboratory variables including gene expression profiling data. *Haematologica* 98:71-78, 2013

35. Khalafallah AA, Snarski A, Heng R, et al: Assessment of whole body MRI and sestamibi technetium-99m bone marrow scan in prediction of multiple myeloma disease progression and outcome: A prospective comparative study. *BMJ Open* 3:pii:e002025, 2013

36. Bäuerle T, Hillengass J, Fechtner K, et al: Multiple myeloma and monoclonal gammopathy of undetermined significance: Importance of whole-body versus spinal MR imaging. *Radiology* 252:477-485, 2009

37. Baur A, Stähler A, Brüning R, et al: Diffusion-weighted MR imaging of bone marrow: Differentiation of benign versus pathologic compression fractures. *Radiology* 207:349-356, 1998

38. Mouloupoulos LA, Dimopoulos MA, Weber D, et al: Magnetic resonance imaging in the staging of solitary plasmacytoma of bone. *J Clin Oncol* 11:1311-1315, 1993

39. Dimopoulos MA, Mouloupoulos LA, Maniatis A, et al: Solitary plasmacytoma of bone and asymptomatic multiple myeloma. *Blood* 96:2037-2044, 2000

40. Varettoni M, Corso A, Pica G, et al: Incidence, presenting features and outcome of extramedullary disease in multiple myeloma: A longitudinal study on 1003 consecutive patients. *Ann Oncol* 21:325-330, 2010

41. Lafforgue P, Dahan E, Chagnaud C, et al: Early-stage avascular necrosis of the femoral head: MR imaging for prognosis in 31 cases with at least 2 years of follow-up. *Radiology* 187:199-204, 1993

42. Syed IS, Glockner JF, Feng D, et al: Role of cardiac magnetic resonance imaging in the detection of cardiac amyloidosis. *JACC Cardiovasc Imaging* 3:155-164, 2010

43. Carlson K, Aström G, Nyman R, et al: MR imaging of multiple myeloma in tumour mass measurement at diagnosis and during treatment. *Acta Radiol* 36:9-14, 1995

44. Usmani SZ, Mitchell A, Waheed S, et al: Prognostic implications of serial 18-fluoro-deoxyglucose emission tomography in multiple myeloma treated with total therapy. *Blood* 121:1819-1823, 2013

45. Lecouvet FE, Vande Berg BC, Michaux L, et al: Stage III multiple myeloma: Clinical and prognostic value of spinal bone marrow MR imaging. *Radiology* 209:653-660, 1998

46. Mouloupoulos LA, Dimopoulos MA, Kastritis E, et al: Diffuse pattern of bone marrow involvement on magnetic resonance imaging is associated with high risk cytogenetics and poor outcome in newly diagnosed, symptomatic patients with multiple myeloma: A single center experience on 228 patients. *Am J Hematol* 87:861-864, 2012

47. Mouloupoulos LA, Dimopoulos MA, Christoulas D, et al: Diffuse MRI marrow pattern correlates with increased angiogenesis, advanced disease features and poor prognosis in newly diagnosed myeloma treated with novel agents. *Leukemia* 24:1206-1212, 2010

48. Song MK, Chung JS, Lee JJ, et al: Magnetic resonance imaging pattern of bone marrow involvement as a new predictive parameter of disease progression in newly diagnosed patients with multiple myeloma eligible for autologous stem cell transplantation. *Br J Haematol* 165:777-785, 2014

49. Mouloupoulos LA, Dimopoulos MA, Alexanian R, et al: Multiple myeloma: MR patterns of response to treatment. *Radiology* 193:441-446, 1994

50. Hillengass J, Ayyaz S, Kilk K, et al: Changes in magnetic resonance imaging before and after autologous stem cell transplantation correlate with response and survival in multiple myeloma. *Haematologica* 97:1757-1760, 2012

51. Bannas P, Hentschel HB, Bley TA, et al: Diagnostic performance of whole-body MRI for the detection of persistent or relapsing disease in multiple myeloma after stem cell transplantation. *Eur Radiol* 22:2007-2012, 2012

52. Derlin T, Peldschus K, Münster S, et al: Comparative diagnostic performance of <sup>18</sup>F-FDG PET/CT versus whole-body MRI for determination of remission status in multiple myeloma after stem cell transplantation. *Eur Radiol* 23:570-578, 2013

53. Spinnato P, Bazzocchi A, Brioli A, et al: Contrast enhanced MRI and <sup>18</sup>F-FDG PET-CT in the assessment of multiple myeloma: A comparison of results in different phases of the disease. *Eur J Radiol* 81:4013-4018, 2012

54. Bartel TB, Haessler J, Brown TL, et al: F18-fluorodeoxyglucose positron emission tomography in the context of other imaging techniques and prognostic factors in multiple myeloma. *Blood* 114:2068-2076, 2009

55. Messiou C, Giles S, Collins DJ, et al: Assessing response of myeloma bone disease with diffusion-weighted MRI. *Br J Radiol* 85:e1198-e1203, 2012

56. Giles SL, Messiou C, Collins DJ, et al: Whole-body diffusion-weighted MR imaging for assessment of treatment response in myeloma. *Radiology* 271:785-794, 2014

57. Dimopoulos MA, Mouloupoulos A, Smith T, et al: Risk of disease progression in asymptomatic multiple myeloma. *Am J Med* 94:57-61, 1993

58. Mouloupoulos LA, Dimopoulos MA, Smith TL, et al: Prognostic significance of magnetic resonance imaging in patients with asymptomatic multiple myeloma. *J Clin Oncol* 13:251-256, 1995

59. Hillengass J, Fechtner K, Weber MA, et al: Prognostic significance of focal lesions in whole-body magnetic resonance imaging in patients with asymptomatic multiple myeloma. *J Clin Oncol* 28:1606-1610, 2010

60. Kastritis E, Terpos E, Mouloupoulos L, et al: Extensive bone marrow infiltration and abnormal free light chain ratio identifies patients with asymptomatic myeloma at high risk for progression to symptomatic disease. *Leukemia* 27:947-953, 2013

61. Mateos MV, Hernández MT, Giraldo P, et al: Lenalidomide plus dexamethasone for high-risk smoldering multiple myeloma. *N Engl J Med* 369:438-447, 2013

62. Merz M, Hielscher T, Wagner B, et al: Predictive value of longitudinal whole-body magnetic resonance imaging in patients with smoldering multiple myeloma. *Leukemia* 28:1902-1908, 2014

63. Pepe J, Petrucci MT, Nofroni I, et al: Lumbar bone mineral density as the major factor determining increased prevalence of vertebral fractures in monoclonal gammopathy of undetermined significance. *Br J Haematol* 134:485-490, 2006

64. van de Donk NW, Palumbo A, Johnsen HE, et al: The clinical relevance and management of monoclonal gammopathy of undetermined significance and related disorders: Recommendations from the European Myeloma Network. *Haematologica* 99:984-996, 2014

65. Vande Berg BC, Michaux L, Lecouvet FE, et al: Nonmyelomatous monoclonal gammopathy: Correlation of bone marrow MR images with laboratory findings and spontaneous clinical outcome. *Radiology* 202:247-251, 1997

66. Dhodapkar MV, Sexton R, Waheed S, et al: Clinical, genomic, and imaging predictors of myeloma progression from asymptomatic monoclonal gammopathies (SWOG S0120). *Blood* 123:78-85, 2014

67. Hillengass J, Weber MA, Kilk K, et al: Prognostic significance of whole-body MRI in patients with monoclonal gammopathy of undetermined significance. *Leukemia* 28:174-178, 2014

68. Liebross RH, Ha CS, Cox JD, et al: Solitary bone plasmacytoma: Outcome and prognostic factors following radiotherapy. *Int J Radiat Oncol Biol Phys* 41:1063-1067, 1998

### Affiliations

Meletios A. Dimopoulos and Evangelos Terpos, National and Kapodistrian University of Athens School of Medicine, Athens, Greece; Jens Hillengass and Hartmut Goldschmidt, University Hospital Heidelberg, Heidelberg, Germany; Saad Usmani, Carolinas Healthcare System, Charlotte, NC; Elena Zamagni and Michele Cavo, Bologna University School of Medicine, Bologna; Antonio Palumbo, S. Giovanni Battista Hospital, University of Turin, Turin, Italy; Suzanne Lentzsch, Columbia University College of Physicians and Surgeons, New York, NY; Faith E. Davies, Institute of Cancer Research, Sutton, United Kingdom; Noopur Rajee, Massachusetts General Hospital Cancer Center; Kenneth Anderson, Dana-Farber Brigham and Women's Cancer Center, Harvard Medical School, Boston, MA; Orhan Sezer, Memorial Sisli Hospital, Istanbul, Turkey; Sonja Zweegman, VU University Medical Center, Amsterdam; Pieter Sonneveld, Erasmus University Medical Center, Rotterdam, the Netherlands; Jatin Shah, MD Anderson Cancer Center, Houston, TX; Ashraf Badros, University of Maryland Medical Center, Baltimore, MD; Kazuyuki Shimizu, Tokai Central Hospital, Kakamigahara, Japan; Philippe Moreau, University Hospital Hôtel-Dieu,


Nantes, France; Chor-Sang Chim, Queen Mary Hospital, University of Hong Kong, Hong Kong, Special Administrative Region; Jian Hou, Changzheng Hospital, Second Military Medical University, Shanghai, People's Republic of China; Juan José Lahuerta, Hospital Universitario 12 de Octubre, Madrid, Spain; Artur Jurczyszyn, University Hospital, Krakow, Poland; Heinz Ludwig, Wilhelminenspital, Vienna, Austria; Bart Barlogie, University of Arkansas for Medical Sciences, Little Rock, AR; G. David Roodman, Indiana University School of Medicine, Indianapolis, IN; S. Vincent Rajkumar, Mayo Clinic, Rochester, MN; and Brian G.M. Durie, Samuel Oschin Comprehensive Cancer Institute, Los Angeles, CA.

## GLOSSARY TERMS

**dynamic contrast-enhanced magnetic resonance imaging (DCE-MRI):** a magnetic resonance imaging acquisition strategy involving multiple scans over a set volume during injection of a magnetic resonance contrast agent.

**microvessel density (MVD):** a quantification technique used to assess the number of vessels in a particular tumor specimen using immunohistochemical stains for endothelial markers. High MVD has been found to be associated with poor prognosis in patients with solid tumors.

## Not an ASCO Member? Subscribe to *Journal of Oncology Practice*



**Journal of Oncology Practice (JOP)** is ASCO's bimonthly forum for providing its subscribers with information, news, and tools to enhance practice efficiency and promote a high standard of quality for patient care.

Every issue of *JOP* includes important features on cancer policy issues and their practical effect on cancer care, methods for enhancing the quality of patient care, and tools for improving practice management.

Whether practitioners are in an office or hospital setting, a community or academic environment, *JOP* provides practical information and advice that oncologists and other oncology professionals can apply immediately to their practices. Key features include:

- Articles for all members of the practice—physicians, nurses, and administrators
- Timely and relevant information to help practices succeed
- Focus on improving practice efficiency and quality of care
- Coverage of legal, financial, technology, and personnel issues

Subscribe today at [jop.ascopubs.org](http://jop.ascopubs.org).



American Society of Clinical Oncology



## AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

### Role of Magnetic Resonance Imaging in the Management of Patients With Multiple Myeloma: A Consensus Statement

The following represents disclosure information provided by authors of this manuscript. All relationships are considered compensated. Relationships are self-held unless noted. I = Immediate Family Member, Inst = My Institution. Relationships may not relate to the subject matter of this manuscript. For more information about ASCO's conflict of interest policy, please refer to [www.asco.org/rwc](http://www.asco.org/rwc) or [jco.ascopubs.org/site/ifc](http://jco.ascopubs.org/site/ifc).

#### Meletios A. Dimopoulos

**Consulting or Advisory Role:** Ortho Biotech (Inst), Celgene (Inst), Onyx Pharmaceuticals (Inst)

#### Jens Hillengass

**Honoraria:** Celgene, Janssen-Cilag, Medtronic

**Travel, Accommodations, Expenses:** Celgene, Janssen-Cilag, Medtronic

#### Saad Usmani

**Honoraria:** Celgene, Onyx Pharmaceuticals, sanofi-aventis, Millennium Pharmaceuticals

**Consulting or Advisory Role:** Celgene, Onyx Pharmaceuticals, sanofi-aventis, Millennium Pharmaceuticals

**Speakers' Bureau:** Celgene, Onyx Pharmaceuticals, Millennium Pharmaceuticals

**Research Funding:** Celgene, Onyx Pharmaceuticals, Janssen-Cilag, Array BioPharma, Pharmaclics

#### Elena Zamagni

No relationship to disclose

#### Suzanne Lentzsch

**Consulting or Advisory Role:** Celgene, Bristol-Myers Squibb, Millennium Pharmaceuticals, Janssen

**Speakers' Bureau:** Axiom

**Research Funding:** Celgene

#### Faith E. Davies

**Honoraria:** Celgene, Janssen-Cilag

**Consulting or Advisory Role:** Millennium Pharmaceuticals, Onyx Pharmaceuticals, Celgene, Janssen-Cilag

#### Noopur Raje

**Consulting or Advisory Role:** Celgene, Millennium Pharmaceuticals, Amgen, Novartis

**Research Funding:** Eli Lilly, Acetylon Pharmaceuticals

#### Orhan Sezer

**Honoraria:** Celgene, Janssen-Cilag

**Consulting or Advisory Role:** Celgene, Janssen-Cilag

#### Sonja Zweegman

**Consulting or Advisory Role:** Celgene (Inst), Janssen-Cilag (Inst), Millennium Pharmaceuticals (Inst)

**Research Funding:** Celgene (Inst), Janssen-Cilag (Inst), Millennium Pharmaceuticals (Inst)

#### Jatin Shah

**Consulting or Advisory Role:** Celgene, Novartis, Millennium Pharmaceuticals, Onyx Pharmaceuticals, sanofi-aventis, Array BioPharma

**Research Funding:** Celgene, Millennium Pharmaceuticals, Novartis, Onyx Pharmaceuticals, Array BioPharma

#### Ashraf Badros

No relationship to disclose

#### Kazuyuki Shimizu

**Consulting or Advisory Role:** Daiichi Sankyo

**Expert Testimony:** Amgen

#### Philippe Moreau

No relationship to disclose

#### Chor-Sang Chim

**Honoraria:** Janssen Oncology, Novartis

**Consulting or Advisory Role:** Novartis

**Speakers' Bureau:** Janssen Oncology

**Travel, Accommodations, Expenses:** Janssen Oncology, Novartis

#### Juan José Lahuerta

**Consulting or Advisory Role:** Celgene, Janssen-Cilag

**Travel, Accommodations, Expenses:** Celgene

#### Jian Hou

No relationship to disclose

#### Artur Jurczynszyn

**Travel, Accommodations, Expenses:** Celgene, Teva

#### Hartmut Goldschmidt

**Honoraria:** Janssen Pharmaceuticals, Celgene, Novartis, Chugai Pharmaceutical, Onyx Pharmaceuticals, Millennium Pharmaceuticals

**Consulting or Advisory Role:** Janssen Pharmaceuticals (Inst), Celgene (Inst), Novartis (Inst), Onyx Pharmaceuticals (Inst), Millennium Pharmaceuticals (Inst)

**Research Funding:** Janssen Pharmaceuticals (Inst), Celgene (Inst), Novartis (Inst), Chugai Pharmaceutical (Inst)

#### Pieter Sonneveld

**Consulting or Advisory Role:** Celgene (Inst), Onyx Pharmaceuticals (Inst), Janssen-Cilag (Inst), Millennium Pharmaceuticals (Inst), Novartis

**Research Funding:** Celgene (Inst), Onyx Pharmaceuticals (Inst), Janssen-Cilag (Inst)

#### Antonio Palumbo

**Honoraria:** Amgen, Bristol-Myers Squibb, Array BioPharma, Genmab, Celgene, Janssen-Cilag, Millennium Pharmaceuticals, BiPar/sanofi-aventis, Onyx Pharmaceuticals

**Consulting or Advisory Role:** Bristol-Myers Squibb, Amgen, Genmab, Celgene, Janssen-Cilag, Millennium Pharmaceuticals, Onyx Pharmaceuticals

#### Heinz Ludwig

**Consulting or Advisory Role:** Celgene (Inst), Janssen-Cilag (Inst), Onyx Pharmaceuticals (Inst), Bristol-Myers Squibb (Inst)

**Speakers' Bureau:** Celgene, Janssen-Cilag

**Research Funding:** Takeda Pharmaceuticals (Inst)

**Travel, Accommodations, Expenses:** Sandoz

#### Michele Cavo

**Honoraria:** Janssen-Cilag, Celgene, Millennium Pharmaceuticals, Onyx Pharmaceuticals, Bristol-Myers Squibb

**Consulting or Advisory Role:** Janssen-Cilag, Celgene, Millennium Pharmaceuticals

**Speakers' Bureau:** Janssen-Cilag, Celgene

#### Bart Barlogie

**Consulting or Advisory Role:** Celgene, Millennium Pharmaceuticals

**Research Funding:** Celgene

**Patents, Royalties, Other Intellectual Property:** Myeloma Health

**Travel, Accommodations, Expenses:** Celgene, Multiple Myeloma Research Foundation, Dana-Farber Cancer Institute

#### Kenneth Anderson

No relationship to disclose

**G. David Roodman**

**Consulting or Advisory Role:** Amgen

**Research Funding:** Eli Lilly (Inst)

**S. Vincent Rajkumar**

No relationship to disclose

**Brian G.M. Durie**

No relationship to disclose

**Evangelos Terpos**

**Honoraria:** Celgene, Medtronic, Onyx Pharmaceuticals, Novartis, Janssen-Cilag, Amgen

**Consulting or Advisory Role:** Roche (Inst), Teva (Inst), Acetylon Pharmaceuticals (Inst), Amgen (Inst), Celgene (Inst)

**Research Funding:** Amgen (Inst), Janssen-Cilag (Inst)

**Travel, Accommodations, Expenses:** Janssen-Cilag, Genesis

### Appendix

*Members of the International Myeloma Working Group:* Niels Abildgaard, Syddansk Universitet, Odense, Denmark; Rafat Abonour, Indiana University School of Medicine, Indianapolis, IN; Melissa Alsina, H. Lee Moffitt Cancer Center and Research Institute, Tampa, FL; Kenneth C. Anderson, Dana-Farber Cancer Institute, Boston, MA; Michel Attal, Purpan Hospital, Toulouse, France; Hervé Avet-Loiseau, University of Toulouse, Toulouse, France; Ashraf Badros, University of Maryland, Baltimore, MD; Bart Barlogie, Myeloma Institute for Research and Therapy, University of Arkansas for Medical Sciences, Little Rock, AR; Régis Bataille, Institute de Biologie, Nantes, France; Meral Bektaş, Ankara University, Ankara, Turkey; Andrew Belch, University of Alberta, Edmonton, Alberta, Canada; Dina Ben-Yehuda, Hadassah University Hospital, Hadassah, Israel; Bill Bensinger, Fred Hutchinson Cancer Center, Seattle, WA; P. Leif Bergsagel, Mayo Clinic Scottsdale, Scottsdale, AZ; Joan Bladé, Hospital Clinica, Barcelona, Spain; Mario Boccadoro, University of Torino, Torino, Italy; Jo Caers, Centre Hospitalier Universitaire de Liège, Liège, Belgium; Michele Cavo, Università di Bologna, Bologna, Italy; Asher Chanan-Khan, Mayo Clinic, Jacksonville, FL; Wen Ming Chen, Beijing Chaoyang Hospital, Beijing, People's Republic of China; Marta Chesi, Mayo Clinic Scottsdale, Scottsdale, AZ; J. Anthony Child, Leeds General Hospital, Leeds, United Kingdom; Chor Sang Chim, Queen Mary Hospital, Hong Kong, Special Administrative Region, People's Republic of China; Wee-Joo Chng, National University Health System, Singapore, Singapore; Ray Comenzo, Tufts Medical School, Boston, MA; John Crowley, Cancer Research and Biostatistics, Seattle, WA; William Dalton, H. Lee Moffitt, Tampa, FL; Faith E. Davies, Royal Marsden Hospital, London, United Kingdom; Javier de la Rubia, Hospital Universitario La Fe, Valencia, Spain; Cármino de Souza, Universidade de Campinas, Campinas, Brazil; Michel Delforge, University Hospital Gasthuisberg, Leuven, Belgium; Meletios A. Dimopoulos, University of Athens School of Medicine, Athens, Greece; Angela Dispenzieri, Mayo Clinic, Rochester, MN; Johannes Drach, University of Vienna, Vienna, Austria; Matthew Drake, Mayo Clinic, Rochester, MN; Juan Du, Changzhen Hospital, Shanghai, People's Republic of China; Brian G.M. Durie, Cedars-Sinai Samuel Oschin Cancer Center, Los Angeles, CA; Hermann Einsele, Universitätsklinik Würzburg, Würzburg, Germany; Thierry Facon, Centre Hospitalier Régional Universitaire de Lille, Lille, France; Dorotea Fantl, Sociedad Argentina de Hematología, Buenos Aires, Argentina; Jean-Paul Fermand, Hôpital Saint-Louis, Paris, France; Carlos Fernández de Larrea, Hospital Clínic de Barcelona, Barcelona, Spain; Rafael Fonseca, Mayo Clinic Scottsdale, Scottsdale, AZ; Gösta Gahrton, Karolinska Institute for Medicine, Huddinge, Sweden; Ramón García-Sanz, University Hospital of Salamanca, Salamanca, Spain; Laurent Garderet, Hôpital Saint Antoine, Paris, France; Christina Gasparetto, Duke University Medical Center, Durham, NC; Morie Gertz, Mayo Clinic, Rochester, MN; Irene Ghobrial, Dana-Farber Cancer Institute, Boston, MA; John Gibson, Royal Prince Alfred Hospital, Sydney, New South Wales, Australia; Peter Gimsing, University of Copenhagen, Copenhagen, Denmark; Sergio Giralt, Memorial Sloan-Kettering Cancer Center, New York, NY; Hartmut Goldschmidt, University Hospital Heidelberg, Heidelberg, Germany; Jingli Gu, The First Hospital, Sun Yat-Sen University, Guangdong, People's Republic of China; Roman Hajek, University Hospital Ostrava and University of Ostrava School of Medicine, Ostrava, Czech Republic; Izhar Hardan, Tel Aviv University, Tel Aviv, Israel; Parameswaran Hari, Medical College of Wisconsin, Milwaukee, WI; Hiroyuki Hata, Kumamoto University Hospital, Kumamoto, Japan; Yutaka Hattori, Keio University School of Medicine, Tokyo, Japan; Tom Heffner, Emory University, Atlanta, GA; Jens Hillengass, University of Heidelberg, Heidelberg, Germany; Joy Ho, Royal Prince Alfred Hospital, Sydney, New South Wales, Australia; Antje Hoering, Cancer Research and Biostatistics, Seattle, WA; Jian Hou, Shanghai Chang Zheng Hospital, Shanghai, People's Republic of China; Jeffrey Huang, National Taiwan University Hospital, Taipei, Taiwan; Vania Hungria, Clinica San Germano, Sao Paulo, Brazil; Shinsuke Ida, Nagoya City University Medical School, Nagoya, Japan; Peter Jacobs, Constantiaberg Medi-Clinic, Plumstead, South Africa; Sundar Jagannath, Mt. Sinai Cancer Institute, New York, NY; Andrzej J. Jakubowiak, University of Chicago, Chicago, IL; Hans Johnsen, Aalborg Hospital Science and Innovation Center, Aalborg, Denmark; Douglas Joshua, Royal Prince Alfred Hospital, Sydney, New South Wales, Australia; Artur Jurczyszyn, University Hospital, Krakow, Poland; Efstathios Kastritis, University of Athens, Athens, Greece; Jonathan Kaufman, Emory Clinic, Atlanta, GA; Michio Kawano, Yamaguchi University, Ube, Japan; Neha Korde, National Institutes of Health, Bethesda, MD; Eva Kovacs, Cancer Immunology Research-Life, Birsfelden, Switzerland; Amrita Krishnan, City of Hope, Duarte, CA; Sigurdur Kristinsson, Karolinska University Hospital and Karolinska Institutet, Stockholm, Sweden; Nicolaus Kröger, University Hospital Hamburg, Hamburg, Germany; Shaji Kumar, Mayo Clinic, Rochester, MN; Robert A. Kyle, Mayo Clinic, Rochester, MN; Chara Kyriacou, Northwick Park Hospital, London, United Kingdom; Martha Lacy, Mayo Clinic, Rochester, MN; Juan José Lahuerta, Grupo Español de Mieloma, Hospital Universitario 12 de Octubre, Madrid, Spain; Ola Landgren, Memorial Sloan-Kettering Cancer Center, New York, NY; Alessandra LaRocca, Divisione Universitaria di Ematologia, Torino, Italy; Jacob Laubach, Dana-Farber Cancer Institute, Boston, MA; Fernando Leal da Costa, Instituto Portugues De Oncologia, Lisbon, Portugal; Jae-Hoon Lee, Gachon University Gil Hospital, Incheon, Korea; Merav Leiba, Sheba Medical Center, Tel Hashomer, Israel; Xavier LeLeu, Hôpital Huriez, Centre Hospitalier Régional Universitaire, Lille, France; Suzanne Lentzsch, Columbia University, New York, NY; Nelson Leung, Mayo Clinic, Rochester, MN; Henk Lokhorst, University Medical Center Utrecht, Utrecht, the Netherlands; Sagar Lonial, Emory University Medical School, Atlanta, GA; Jin Lu, People's Hospital, Beijing University, Beijing, People's Republic of China; Heinz Ludwig, Wilhelminenspital der Stat Wien, Vienna, Austria; Anuj Mahindra, University of California at San Francisco, San Francisco, CA; Angelo Maiolino, Rua fonte da Saudade, Rio de Janeiro, Brazil; María-Marivi Mateos, University Hospital of Salamanca–Instituto de Investigación Biomédica de Salamanca (IBSAL), Instituto de Biología Molecular y Celular del Cancer (IBMCC; Universidad de Salamanca [USAL] –Centro Superior de Investigaciones Científicas [CSIC]), Salamanca, Spain; Amitabha Mazumder, New York University Comprehensive Cancer Center, New York, NY; Philip McCarthy, Roswell Park Cancer Center, Buffalo, NY; Jayesh Mehta, Northwestern

University, Chicago, IL; Ulf-Henrik Mellqvist, Sahlgrenska University Hospital, Gothenburg, Sweden; Giampaolo Merlini, University of Pavia, Pavia, Italy; Joseph Mikhael, Mayo Clinic Scottsdale, Scottsdale, AZ; Philippe Moreau, University Hospital, Nantes, France; Gareth Morgan, Royal Marsden Hospital, London, United Kingdom; Nikhil Munshi, Dana-Farber Cancer Institute, Boston, MA; Hareth Nahi, Karolinska University Hospital, Stockholm, Sweden; Weerasak Nawarawong, Chiang Mai University, Chiang Mai, Thailand; Ruben Niesvizky, Weill Cornell Medical College, New York, NY; Amara Nouel, Hospital Rutz y Paez, Bolivar, Venezuela; Yana Novis, Hospital Sírío Libanês, Bela Vista, Brazil; Enrique Ocio, University Hospital of Salamanca-IBSAL, IBMCC (USAL-CSIC), Salamanca, Spain; Alberto Orfao, University Hospital of Salamanca-IBSAL, IBMCC (USAL-CSIC), Salamanca, Spain; Robert Orłowski, MD Anderson Cancer Center, Houston, TX; Bruno Paiva, Clínica Universitaria de Navarra, Centro de Investigación Médica Aplicada (CIMA), Pamplona, Spain; Antonio Palumbo, University of Torino, Torino, Italy; Santiago Pavlovsky, Fundaleu, Buenos Aires, Argentina; Linda Pilarski, University of Alberta, Edmonton, Alberta, Canada; Raymond Powles, Parkside Cancer Centre, London, United Kingdom; Noopur Raje, Massachusetts General Hospital, Boston, MA; S. Vincent Rajkumar, Mayo Clinic, Rochester, MN; Donna Reece, Princess Margaret Hospital, Toronto, Ontario, Canada; Anthony Reiman, Saint John Regional Hospital, Saint John, New Brunswick, Canada; Paul G. Richardson, Dana-Farber Cancer Institute, Boston, MA; Angelina Rodríguez Morales, Bonco Metro Político de Sangre, Caracas, Venezuela; Kenneth R. Romeril, Wellington Hospital, Wellington, New Zealand; David Roodman, Indiana University, Indianapolis, IN; Laura Rosiñol, Hospital Clinic, Barcelona, Spain; Murielle Rousseau, University of Toulouse, Toulouse, France; Stephen Russell, Mayo Clinic, Rochester, MN; Jesús San Miguel, Clínica Universitaria de Navarra, CIMA, Pamplona, Spain; Rik Schots, Universitair Ziekenhuis Brussel, Brussels, Belgium; Sabina Sevcikova, Masaryk University, Brno, Czech Republic; Orhan Sezer, Memorial Sisli Hastanesi, Istanbul, Turkey; Jatin J. Shah, MD Anderson Cancer Institute, Houston, TX; Kazuyuki Shimizu, Tokai Central Hospital, Kakamigahara, Japan; Chaim Shustik, McGill University, Montreal, Quebec, Canada; David Siegel, Hackensack University Medical Center, Hackensack, NJ; Seema Singhal, Northwestern University, Chicago, IL; Pieter Sonneveld, Erasmus University Medical Center, Rotterdam, the Netherlands; Andrew Spencer, Alfred Hospital, Melbourne, Victoria, Australia; Edward Stadtmauer, University of Pennsylvania, Philadelphia, PA; Keith Stewart, Mayo Clinic Scottsdale, Scottsdale, AZ; Daryl Tan, Singapore General Hospital, Singapore, Singapore; Evangelos Terpos, University of Athens School of Medicine, Athens, Greece; Patrizia Tosi, Italian Cooperative Group, Istituto di Ematologia Seragnoli, Bologna, Italy; Guido Tricot, University of Iowa Hospital and Clinics, Iowa City, IA; Ingemar Turesson, Skane University Hospital, Malmo, Sweden; Saad Usmani, Levine Cancer Institute/Carolinas Healthcare System, Charlotte, NC; Ben Van Camp, Vrije Universiteit Brussel, Brussels, Belgium; Brian Van Ness, University of Minnesota, Minneapolis, MN; Ivan Van Riet, Vrije Universiteit Brussel (VUB), Brussels, Belgium; Isabelle Vande Broek, VUB, Brussels, Belgium; Karin Vanderkerken, VUB, Brussels, Belgium; Robert Vescio, Cedars-Sinai Cancer Center, Los Angeles, CA; David Vesole, Hackensack University Medical Center, Hackensack, NJ; Peter Voorhees, University of North Carolina, Chapel Hill, NC; Anders Waage, University Hospital, Trondheim, Norway; Michael Wang, MD Anderson Cancer Center, Houston, TX; Donna Weber, MD Anderson Cancer Center, Houston, TX; Jan Westin, Sahlgrenska University Hospital, Gothenburg, Sweden; Keith Wheatley, University of Birmingham, Birmingham, United Kingdom; Elena Zamagni, University of Bologna, Bologna, Italy; Jeffrey Zonder, Karmanos Cancer Institute, Detroit, MI; and Sonja Zweegman, VU University Medical Center, Amsterdam, the Netherlands.