












ORIGINAL RESEARCH

Effect of Maternal Prepregnancy/Early-Pregnancy Body Mass Index and Pregnancy Smoking and Alcohol on Congenital Heart Diseases: A Parental Negative Control Study

Kurt Taylor , BSc; Ahmed Elhakeem , PhD; Johanna Lucia Thorbjørnsrud Nader , PhD; Tiffany C Yang , PhD; Elena Isaevska, PhD; Lorenzo Richiardi , PhD; Tanja Vrijkotte , PhD; Angela Pinot de Moira , PhD; Deirdre M. Murray, MB, MAO, BCH, PhD; Daragh Finn, MB, MD; Dan Mason , PhD; John Wright, FRCP; Sam Oddie, MD; Nel Roeleveld, PhD; Jennifer R. Harris , PhD; Anne-Marie Nybo Andersen , PhD; Massimo Caputo, MD; Deborah A. Lawlor , PhD

BACKGROUND: Congenital heart diseases (CHDs) are the most common congenital anomaly. The causes of CHDs are largely unknown. Higher prenatal body mass index (BMI), smoking, and alcohol consumption are associated with increased risk of CHDs. Whether these are causal is unclear.

METHODS AND RESULTS: Seven European birth cohorts, including 232 390 offspring (2469 CHD cases [1.1%]), were included. We applied negative exposure paternal control analyses to explore the intrauterine effects of maternal BMI, smoking, and alcohol consumption during pregnancy, on offspring CHDs and CHD severity. We used logistic regression, adjusting for confounders and the other parent's exposure and combined estimates using a fixed-effects meta-analysis. In adjusted analyses, maternal overweight (odds ratio [OR], 1.15 [95% CI, 1.01–1.31]) and obesity (OR, 1.12 [95% CI, 0.93–1.36]), compared with normal weight, were associated with higher odds of CHD, but there was no clear evidence of a linear increase in odds across the whole BMI distribution. Associations of paternal overweight, obesity, and mean BMI were similar to the maternal associations. Maternal pregnancy smoking was associated with higher odds of CHD (OR, 1.11 [95% CI, 0.97–1.25]) but paternal smoking was not (OR, 0.96 [95% CI, 0.85–1.07]). The positive association with maternal smoking appeared to be driven by nonsevere CHD cases (OR, 1.22 [95% CI, 1.04–1.44]). Associations with maternal moderate/heavy pregnancy alcohol consumption were imprecisely estimated (OR, 1.16 [95% CI, 0.52–2.58]) and similar to those for paternal consumption.

CONCLUSIONS: We found evidence of an intrauterine effect for maternal smoking on offspring CHDs, but no evidence for higher maternal BMI or alcohol consumption. Our findings provide further support for the importance of smoking cessation during pregnancy.

Key Words: congenital heart disease ■ negative control ■ risk factors

Congenital heart diseases (CHDs) are the most common congenital anomaly (CA), affecting 6 to 8 per 1000 live births and 10% of stillbirths, and are the leading cause of death from CAs.¹ Many patients with CHD present with sequela from surgical intervention and late complications related to the

Correspondence to: Kurt Taylor, BSc, Oakfield House, Medical Research Council Integrative Epidemiology Unit Bristol, Population Health Science Institute, University of Bristol, Bristol BS8 2BN, United Kingdom. E-mail: kurt.taylor@bristol.ac.uk

Preprint posted on MedRxiv, November 4, 2020. doi: <https://doi.org/10.1101/2020.09.29.20203786>.

Supplementary Material for this article is available at <https://www.ahajournals.org/doi/suppl/10.1161/JAHA.120.020051>

For Sources of Funding and Disclosures, see page 15.

© 2021 The Authors. Published on behalf of the American Heart Association, Inc., by Wiley. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

JAHA is available at: www.ahajournals.org/journal/jaha

CLINICAL PERSPECTIVE

What Is New?

- Previous studies showing associations of higher maternal body mass index, smoking, and alcohol consumption in pregnancy were not able to establish causality.
- Using parental negative control analyses, our study provides stronger evidence that maternal pregnancy smoking may increase offspring congenital heart disease risk via intrauterine mechanisms, whereas it does not suggest maternal overweight or obesity increases risk.

What Are the Clinical Implications?

- Emphasizing the potential adverse effect of smoking on congenital heart diseases might help in supporting women of reproductive age not to start smoking and women who are smoking at the start of pregnancy to be encouraged to quit.
- Understanding the mechanisms through which maternal smoking influences congenital heart disease risk could identify novel targets for prevention beyond smoking cessation.

Nonstandard Abbreviations and Acronyms

ABCD	The Amsterdam Born Children and Their Development Study
ALSPAC	Avon Longitudinal Study of Parents and Children
BASELINE	Cork Scope Baseline Study
BiB	Born in Bradford
CA	congenital anomaly
DNBC	Danish National Birth Cohort
MoBa	Norwegian Mother, Father and Child Cohort Study
NINFEA	Nascita e Infanzia: gli Effetti dell'Ambiente (Birth and Childhood: Effects of the Environment)

anomaly, resulting in health problems that persist into adulthood.^{2,3} The causes of CHDs are largely unknown, but intrauterine mechanisms may play a role in their underlying pathophysiological characteristics.⁴ Identifying modifiable risk factors for CHDs is important for improving causative understanding and developing preventive interventions.

Several modifiable maternal characteristics have been found to be associated with increased risk of CHDs, including maternal prepregnancy/early-pregnancy body mass index (BMI),⁵⁻⁷ smoking,⁸ and

alcohol⁹ consumption in pregnancy. Whether these are causal is unclear. A recent systematic review and meta-analysis of the association of BMI with CHDs found that risk of CHDs was higher in those whose mothers were overweight or obese at the start of pregnancy, compared with those who were normal weight. Results for underweight mothers were not reported,⁵ but a large cohort study consisting of >2 000 000 singletons found no clear association for maternal underweight status and CHDs.⁶ These results from conventional multivariable approaches may be explained by residual confounding because of incomplete identification or adjustment for confounders. Maternal active smoking⁸ and maternal exposure to alcohol⁹ were both associated with offspring CHDs in recent meta-analyses. However, 68% and 69% of the studies within the meta-analyses (for maternal smoking and alcohol, respectively) did not adjust for confounders. Therefore, those studies showing associations for smoking and alcohol cannot determine whether these reflect the magnitude of a causal effect or are biased by confounding.

Negative control studies are widely used in laboratory science and in recent years have become increasingly used to explore causal effects in epidemiology.¹⁰ The idea behind negative control studies is that either the exposure or the outcome in the real experiment is substituted for a negative control exposure (or outcome) that is not a plausible risk factor but would have similar sources of bias or confounding as in the main experiment. In epidemiology, this approach has been primarily used for determining the extent to which hypothesized intrauterine and early life exposures might be associated with outcomes as a result of residual confounding.^{10,11} Negative parental exposure control studies are used for this purpose. This involves comparing the confounder-adjusted associations of maternal pregnancy exposures with the offspring outcome of interest to similarly adjusted associations of the same characteristics (negative controls) in the father. The assumptions of this approach are that: (1) measured and unmeasured confounders influence the exposures in the same direction and with a similar magnitude in mothers and fathers and (2) there is no plausible reason why the exposure in the father would affect the offspring outcome (or at a minimum the paternal association would be much weaker than in the mother). In the present study, we are assuming that paternal BMI, smoking, and alcohol cannot causally influence offspring CHDs through intrauterine mechanisms. Under these assumptions, if there is a causal intrauterine effect of any of the maternal pregnancy exposures, we would expect to see a maternal-specific association, with no (or a much weaker) association with the equivalent paternal exposure. Similar associations in mothers and fathers would suggest that

these are largely driven by residual confounding. It is plausible that passive smoking from fathers could influence offspring outcomes via intrauterine exposure; however, we would expect a much weaker association for fathers. As proof of concept, maternal smoking relates strongly to lower birth weight (a known causal intrauterine effect), whereas paternal smoking has a much weaker association; and when the 2 are mutually adjusted, the maternal association remains strong, whereas the weak paternal association attenuates to the null.^{10,12}

We aimed to explore the causal intrauterine effects of maternal pregnancy BMI, smoking, and alcohol on CHDs using data from the Horizon 2020 LifeCycle project.¹³ As well as the negative parental control study providing scope to explore residual confounding, the use of a large existing collaboration of birth cohorts adds benefit to this study in comparison to previous studies. First, both offspring with and without CHDs are from the same underlying populations and have been selected for inclusion and assessed in identical ways. Second, most studies of risk factors for CHDs are case-control studies, and these dominate meta-analysis results. These have advantages in that they have large numbers of CHD cases and hence greater statistical power than most cohorts, but they are prone to selection bias as response rates in controls are commonly low, and in some studies controls are selected from hospitals or clinics and do not reflect exposure status in the population from which the cases came.¹⁴ Furthermore, case-control studies are susceptible to information bias because of differential recall and reporting of the exposure between cases and controls.¹⁴ Third, we have harmonized data on all exposures, confounders, and outcomes. Fourth, we have large numbers, with 232 390 participants in total and 2469 CHD cases. Last, the ethos of the LifeCycle collaboration is that all studies contribute to each research question unless they do not have data on either exposure or outcome, meaning publication bias is minimized.

METHODS

Requests to access the data used in this study may be sent individually to the included cohorts. We have included details on how researchers can access each cohort at the end of the article under “data access.” Materials supporting the findings of this study are available from the corresponding author on reasonable request.

Inclusion Criteria and Participating Cohorts

This study was part of the Horizon2020 LifeCycle Project. LifeCycle is a collaboration of largely European

birth cohorts that aims to determine the impact of early-life stressors on risk of developing adverse cardiovascular/metabolic, respiratory, cognitive, and mental health outcomes (<http://lifecycle-project.eu>).¹³ A LifeCycle cohort was eligible for inclusion if it had information on CHD in the offspring ascertained by any method and data on at least one of the following: (1) mother’s prepregnancy/early-pregnancy BMI, (2) maternal smoking during pregnancy, (3) maternal alcohol consumption during pregnancy, or (4) the same exposures (1–3) measured in the father at a similar time to their pregnant partners. Eligible LifeCycle cohorts could be from any geographical area and with participants from any ethnic background. In total, 7 cohorts were eligible, and all participated: ABCD (The Amsterdam Born Children and Their Development Study),¹⁵ ALSPAC (Avon Longitudinal Study of Parents and Children),^{16,17} BASELINE (Cork Scope Baseline Study),¹⁸ BiB (Born in Bradford) study,¹⁹ DNBC (Danish National Birth Cohort) study,²⁰ MoBa (Norwegian Mother, Father and Child Cohort Study),^{21,22} and NINFEA (Nascita e Infanzia: gli Effetti dell’Ambiente [Birth and Childhood: Effects of the Environment]) study.^{23,24} Individual cohort descriptions can be found in Data S1. We excluded multiple births from the study population because they differ from single births for CA outcomes.^{25,26} Some previous studies have excluded infants with any known chromosomal or genetic defects on the assumption that modifiable risk factors are unlikely to contribute in the presence of known causes. We have not made these exclusions in our main analyses because it is plausible that CHD in children with these complex syndromes is also influenced by the modifiable exposures we explore herein. Furthermore, from a public health and clinical perspective, we believe it is useful to know effects for all CHD cases. In additional analyses, we explore whether their removal alters our main results.

BMI, Smoking, and Alcohol Measurements

We used harmonized LifeCycle data for exposure and confounder data, with the exclusion of paternal alcohol consumption, which had not been harmonized by LifeCycle when we started this project.²⁷ ABCD and BASELINE were not part of the core LifeCycle cohorts and therefore not part of phase 1 harmonized data that we used herein. We harmonized the data for these cohorts to resemble the harmonized LifeCycle variables. Cohort-specific information on methods of data collection can be found in Table S1.

LifeCycle-harmonized maternal BMI used measured or self-reported prepregnancy/early-pregnancy weight and height. Prepregnancy weight was prioritized, and if not available, the earliest pregnancy measures were used. Paternal BMI was similarly reported

(by the father or their pregnant partner) or measured, and we prioritized the timing to be prepregnancy or as early as possible in their partners pregnancy. BMI was used as a continuous variable for the main analyses. In cohorts that had >100 CHD cases, we also categorized BMI as underweight (BMI <18.5 kg/m²), normal weight (BMI 18.5–<25 kg/m²), overweight (BMI 25–<30 kg/m²), and obese (BMI ≥30 kg/m²). ALSPAC, BiB study, DNBC study, and MoBa contributed to these analyses.

We used 2 LifeCycle smoking variables for maternal and paternal smoking at the time of pregnancy: (1) smoking in the first trimester (yes/no) where this was available, otherwise any smoking during pregnancy (yes/no); and (2) categorized into nonsmokers, light smokers (<10 cigarettes smoked per day), and heavy smokers (≥10 cigarettes per day) throughout the entire pregnancy. Paternal smoking was categorized as “any smoking (yes/no)” at the time of their partners pregnancy.

We used 2 LifeCycle variables for maternal alcohol consumption: (1) binary (yes/no), which like smoking prioritized the first trimester if available but was otherwise any alcohol intake during pregnancy; and (2) categorized into nondrinkers (none), light drinkers (>0 and <3 units per week), and moderate/heavy drinkers (≥3 units per week) during pregnancy. Two studies (ALSPAC and MoBa) had data on paternal alcohol consumption in pregnancy and thus were able to harmonize variables relating to paternal alcohol for this project. We generated one variable, categorized as: nondrinkers, light drinkers (>0 and <7 units per week), or moderate/heavy drinkers (≥7 units per week) (Data S2).

The rationale for prioritizing maternal pregnancy smoking and alcohol during the first trimester is because fetal cardiac development starts early in pregnancy and much of the development occurs in the first trimester.²⁸ A total of 47% and 96% of mothers had measures specifically in the first trimester for smoking and alcohol, respectively.

CHD Outcomes

Information on CHDs was retrieved from a variety of sources, depending on the cohort. ALSPAC, BiB study, DNBC study, and NINFEA study had *International Classification of Diseases, Tenth Revision (ICD-10)*, coded data. BASELINE had individual CHD diagnoses assigned by a cardiologist based on echocardiography. For ABCD and MoBa, we had a nonspecific CHD diagnosis (yes/no). Data in ABCD, BASELINE, DNBC study, and NINFEA study were restricted to liveborn infants, whereas ALSPAC, BiB study, and MoBa included stillbirths.

In the ABCD cohort, data on CHDs in liveborn children were obtained from 3 different sources: (1) the

infant questionnaire, which was filled out by the mother at an average infant age of 12.9 weeks; (2) the questionnaire filled out by the mother at an average child age of 5.1 years; and (3) clinical data of the Youth Health Care Registration. In the ALSPAC cohort, cases were obtained from a range of data sources, including health record linkage and questionnaire data up until age 25 years following European Surveillance of Congenital Anomalies guidelines.²⁹ In BASELINE, at 2 months, mothers were asked of any medical problems and/or referrals. If a baby had been referred to a specialist, he/she was checked by a cardiologist to see if he/she had results from an echocardiogram with exact diagnoses reported. Further diagnoses up until age 12 years were identified through records from the echocardiogram. In the BiB study cohort, there were 2 separate sources to identify CAs. Both sources were used in this study: (1) CAs up to 5 years of age, identified in primary care records by Bishop et al,³⁰ following European Surveillance of Congenital Anomalies guidelines. *ICD-10* codes were mapped to clinical term-V3 codes before extraction from primary care records. (2) Data extracted from the Yorkshire and Humber CA register database. Data were *ICD-10* coded. All of these were confirmed postnatally. In the DNBC study, all diagnoses of CAs (according to European Surveillance of Congenital Anomalies guide 1.4, sections 3.2 and 3.3) up until the age of 15 years were extracted from the Danish National Patient Register, which is linked to the cohort data.^{31,32} Diagnoses were *ICD-10* coded. These data were restricted to children born alive. In MoBa, information on whether a child had a CHD or not was obtained through linkage to the Medical Birth Registry of Norway. All maternity units in Norway must notify births to the Medical Birth Registry of Norway. In the NINFEA study cohort, CHDs were reported in the second questionnaire, compiled 6 months after birth. Mothers compiled a checklist that included prespecified anomalies. If the child died or had any surgery performed in the first 6 months, the cause of death and type of surgery were also checked to see if any CA was reported. Data were coded using *ICD-10* codes by an experienced pediatrician and were reassessed by an independent physician. Further details of the sources of data for CHDs in each cohort are provided in Data S3.

In all studies, our main outcome was any CHD. Where data allowed (ie, when we had full *ICD-10* codes), any CHD was defined according to European Surveillance of Congenital Anomalies, which excludes isolated patent ductus arteriosus and peripheral pulmonary artery stenosis in preterm births (gestational age, <37 weeks) (Table S2). We also categorized cases into severe CHD (heterotaxia, conotruncal defect, atrioventricular septal defect, anomalous pulmonary venous return, left ventricle outflow tract obstruction, right ventricle outflow

tract obstruction, or other complex defects) and the remainder as nonsevere CHD (patent ductus arteriosus [in full-term infants], valvular pulmonary stenosis, ventricular septal defect, atrial septal defects, unspecified septal defects, isolated valve defects, other specified heart defects, or unspecified heart defects)^{33,34} (Table S2).

Confounders

Analyses were adjusted for several confounders based on their known or plausible influence on ≥ 1 of the maternal pregnancy exposures and on CHD: maternal age (all exposures), parity (all exposures), ethnicity (all exposures), socioeconomic position (all exposures), smoking (for BMI and alcohol analyses), and alcohol use (for BMI and smoking analyses). In the paternal negative control analyses, confounders were similar: fathers' age (all exposures), number of children (all exposures), ethnicity (all exposures), socioeconomic position (all exposures), smoking (for BMI and alcohol), and alcohol use (for BMI and smoking). We also adjusted for offspring sex in all adjusted analyses. We used educational attainment for both parents' measures of socioeconomic position. Full details of our selection and harmonization of confounders are provided in Data S4.

Statistical Analysis

Analyses were conducted in either R (version 3.6.1) or Stata (version 16). An analysis plan was written and published in October 2019, with any subsequent changes and their rationale documented in the publication.³⁵ All associations between exposures and CHDs were performed within participating studies using logistic regression (binary for main analyses and multinomial for CHD severity analyses). In the 2 largest cohorts (DNBC study and MoBa), we assessed deviation from linearity in our models in the BMI analyses by running our main confounder-adjusted model with BMI split into fifths. We ran regression models with these fifths as 4 indicator variables (nonlinear) and compared this model with one in which the fifths were treated as a continuous (score) variable. We used a likelihood ratio comparison to compare these 2 models. All analyses were run (1) unadjusted; (2) adjusted for maternal/paternal age, socioeconomic position, parity, ethnicity, smoking and/or alcohol (depending on exposure), and offspring sex; and (3) adjusted for all confounders (as in 2) as well as the other parent's exposure. In the adjusted models, studies were asked to adjust for as many of the confounders as possible. All analyses were performed with maximal numbers (ie, numbers included in each model will vary because of missing data on exposure/outcome or confounders). In a

sensitivity analysis, we repeated our main analyses using complete-case data to assess whether missing data were influencing the results.

For the main negative control analyses (ie, where we directly compared maternal with paternal exposure-CHD associations), we used multivariable logistic regression in which both maternal and paternal exposures were adjusted for the other parent's exposure (model 3 above). This produces a maternal association that adjusts for maternal confounders as well as the paternal exposure, and similarly a paternal association adjusting for paternal confounders and the maternal exposure. The rationale for mutually adjusting for the other parent's exposure is that parental BMI, smoking, and alcohol may relate to each other through assortative mating and/or convergence of behaviors that occurs over time in couples.³⁶ Causal structural graphs together with simulated data show failure to undertake this mutual adjustment will bias the negative control analysis results.³⁷ Also, paternal exposures may have some intrauterine impact (eg, via passive smoking or paternal support for the mother to reduce alcohol and have a normal BMI during her preconceptional period or in pregnancy).³⁸ Mutual adjustment for maternal and paternal confounders was necessary for ensuring both parental results were fully adjusted. Comparisons between maternal and paternal associations from this model were assessed by visually comparing the 2 results. In addition, statistical evidence of any differences was obtained by calculating differences in log odds of CHD between the fathers' and mothers' associations and report of the corresponding P value ($P_{\text{difference}}$), under the null hypothesis that there is no difference between the maternal and paternal estimate.

Analyses were conducted separately in each study and then meta-analysed using the *meta* package in R.³⁹ All the data used in the present study originated from European birth cohorts, with broadly similar methods, and therefore, we assumed that they were each estimating an association from the same underlying populations and used a fixed-effects meta-analysis. To explore this assumption, differences between studies were assessed using I^2 and Cochrane Q P values for heterogeneity.⁴⁰

Additional Analyses

We repeated the main and subgroup (by CHD severity) analyses after excluding infants with any known chromosomal/genetic or maternal drug defects. Methods of data collection and definition of these variables can be found in Table S3. We also repeated analyses in mothers only including those with smoking data in the first trimester. Folic acid supplementation has been shown to lower risk of birth defects

and adverse pregnancy outcomes.^{41,42} We repeated the adjusted maternal analyses with additional adjustment for first-trimester folic acid supplementation (yes/no).

RESULTS

Participant Characteristics

Figures S1 through S7 show flowcharts designating the assignment of participants into analysis groups for each cohort. In total, 7 cohorts, including 232 390 offspring with 2469 CHD cases (1.1%), were included. The prevalence of CHD was close to 1% in most cohorts, with the lowest being in ABCD (0.4%) and the highest in DNBC study (1.4%) (Table). The Table shows the distributions of maternal and paternal characteristics for each cohort. Mean maternal age across the cohorts was broadly similar (all late 20s to early 30s). Mean BMI was also similar across the cohorts, but proportions in different categories varied, with the lowest prevalence of prepregnancy/early-pregnancy obesity seen in NINFEA study (5%) and the highest in BiB study (21%). There was also variation in maternal smoking and alcohol consumption across the cohorts, with notably high levels of both smoking (25% and 26%) and alcohol (55% and 45%) in ALSPAC and DNBC study, respectively. Fathers were generally older than mothers and more likely to smoke and drink alcohol, with the overall patterns of between-study differences being similar to those for the mothers. There were differing levels of missing data in each cohort (summarized in Table S4 and also illustrated in cohort-specific flow charts [Figures S1 through S7]). To check whether missing data influenced any of our results, we report complete-case analysis results for our main analyses in the Supplementary Material. Overall, complete-case results from meta-analyses were comparable (Tables S5 through S8). Below, we present our main results separated by exposure. We include supplementary results for BMI (Figures S8 through S20, Tables S9 and S10), smoking (Figures S21 through S27), and alcohol (Figures S28 through S32 and Table S11) analyses in the Supplementary Material.

BMI and CHDs

In confounder and other parent BMI-adjusted analyses, there was no difference in the odds of offspring CHD per 1-kg/m² difference in maternal BMI (odds ratio [OR], 1.00; 95% CI, 0.99–1.02) or paternal mean BMI (OR, 1.01; 95% CI, 0.99–1.03) ($P_{\text{difference}}=0.43$), with both being close to the null (Figure 1A). Unadjusted and confounder-only adjusted results did not differ notably from those presented in Figure 1 (Figure S8). The odds of CHD did not clearly increase linearly in mothers or fathers in DNBC

study or MoBa (Figures S9 and S10). Analyses of continuously measured BMI with CHD cases separated into nonsevere and severe showed similar null associations for both mothers and fathers (Figure S11).

In analyses of BMI categories, there were increased odds of offspring CHD in overweight and obese mothers and fathers compared with those of a normal BMI, with similar magnitudes of association in both parents ($P_{\text{difference}} \text{ overweight}=0.65$ and $P_{\text{difference}} \text{ obese}=0.83$) (Figure 1B). Underweight mothers had an increased odds of offspring CHD, whereas underweight fathers had a decreased odds of offspring CHD. Because of small numbers of underweight parents, particularly fathers, however, results were imprecise, with wide CIs, and there was no statistical evidence for between parental differences for underweight ($P_{\text{difference}} \text{ underweight}=0.27$). Individual study results for BMI categories are shown in Figures S15 through S17. Positive parental associations of overweight and obesity were also observed for both nonsevere (Figure 1C) and severe (Figure 1D) CHDs, with similar magnitudes of association in mothers and fathers. Individual study results for BMI categories and CHD severity are shown in Figures S18 through S20.

Smoking and CHDs

In confounder and other parental smoking-adjusted analyses, maternal smoking in pregnancy was associated with increased odds of CHD (OR, 1.11; 95% CI, 0.97–1.25), whereas paternal smoking at the time of their partners pregnancy did not increase odds of offspring CHD (OR, 0.96; 95% CI, 0.85–1.07) ($P_{\text{difference}}=0.09$) (Figure 2A). When removing offspring with a chromosomal/genetic defect, there was stronger statistical evidence of a difference between maternal and paternal smoking ($P_{\text{difference}}=0.02$) (Figure 2B). Results for unadjusted analyses were consistent with the confounder and mutual parent smoking-adjusted result, whereas confounder-only analyses were slightly attenuated for maternal smoking (Figure S21). Maternal smoking results were similar when analyses were restricted to studies with confirmed first-trimester smoking (Figure S22). A positive association between maternal smoking and offspring CHD was also seen with nonsevere CHDs (OR, 1.22; 95% CI, 1.04–1.44), although not with severe CHDs (OR, 0.90; 95% CI, 0.69–1.17) (Figure 2C and 2D and Figure S23). When we analyzed maternal smoking frequency categories (ie, none, light, and heavy smoking), the results did not support an effect of heaviness over and above what we saw with any smoking (Figure S24). The maternal and paternal associations for these categories were statistically consistent ($P_{\text{difference}}=0.25$ and $P_{\text{difference}}=0.38$ for light and heavy smoking, respectively).

Table. Characteristics of the Participating Cohorts

Characteristics	Category	ABCD (N=8131)	ALSPAC (N=13 049)	BASELINE (N=1436)	BIB Study (N=12 799)	DNBC Study (N=89 107)	MoBa (N=101 975)	NINFEA Study (N=5893)
Country		The Netherlands	United Kingdom	Ireland	United Kingdom	Denmark	Norway	Italy
Recruitment period		2003–2004	1991–1992	2008–2011	2007–2011	1996–2002	1999–2008	2005–2016
Offspring								
CHD	Any	34 (0.4)	103 (0.8)	10 (0.7)	145 (1.1)	1264 (1.4)	879 (0.9)	34 (0.6)
CHD severity in those with CHD	Nonsevere	...	73/103 (70.9)	...	93/145 (64.1)	896/1264 (70.9)	...	27/34 (79.4)
	Severe	...	30/103 (29.1)	...	52/145 (35.9)	368/1264 (29.1)	...	7/34 (20.6)
Chromosomal/genetic defects*		26 (0.3)	58 (0.4)	...	198 (1.5)	698 (0.8)	169 (0.2)	7 (0.1)
Maternal								
Age, y		30.7 (5.3)	28.9 (4.8)	30.7 (4.4)	26.0 (5.7)	29.9 (4.3)	30.2 (4.6)	33.1 (4.3)
BMI, kg/m ²		23.1 (4.1)	22.6 (4.4)	24.4 (4.1)	26.0 (5.7)	23.6 (4.3)	24.0 (4.3)	22.5 (3.8)
BMI categories	Underweight (<18.5 kg/m ²)	360 (4.9)	1271 (11.6)	23 (1.6)	444 (4.4)	3861 (4.5)	3077 (3.2)	501 (8.5)
	Normal (18.5–<25 kg/m ²)	5270 (71.8)	7426 (67.7)	914 (63.6)	4586 (45.4)	57 894 (67.8)	63 706 (65.4)	4156 (70.5)
	Overweight (25–<30 kg/m ²)	1245 (17.0)	1537 (14.0)	345 (24.0)	2952 (29.2)	16 578 (19.4)	21 280 (21.8)	826 (14.0)
	Obese (≥30 kg/m ²)	467 (6.4)	736 (6.7)	154 (10.7)	2127 (21.0)	7017 (8.2)	9337 (9.6)	286 (4.9)
Pregnancy smoking	Yes†	769 (9.5)	3147 (24.7)†	357 (24.9)†	1788 (16.4)	22 514 (26.0)†	9650 (9.6)	472 (8.1)†
	Light	...	1684 (15.7)	...	1362 (12.5)	15 777 (17.9)	7856 (7.7)	438 (7.5)
	Heavy	...	1096 (10.2)	...	426 (3.9)	7431 (8.5)	1587 (1.6)	30 (0.5)
Pregnancy alcohol	Yes†	1686 (20.8)	6894 (54.6)†	527 (36.7)	...	38 733 (44.7)†	22 799 (27.7)†	1508 (25.8)†
	Light	...	3044 (46.8)	46 774 (52.9)	10 461 (12.4)	1416 (24.4)
	Moderat/heavy	...	871 (13.4)	3717 (4.2)	509 (0.6)	230 (3.9)
Parity	Nulliparous	4500 (55.3)	5645 (45.0)	1436 (100)	4912 (39.8)	42 203 (47.4)	46 988 (46.9)	4070 (72.4)
Education	Low	4035 (49.6)	2374 (20.0)	...	5717 (56.9)	22 225 (27.6)	2735 (2.9)	278 (4.8)
	Medium	2225 (27.4)	7985 (67.1)	208 (14.6)	1563 (15.6)	17 756 (22.0)	31 430 (33.1)	1892 (32.4)
	High	1871 (23.0)	1538 (12.9)	1219 (85.4)	2769 (27.6)	40 675 (50.4)	60 847 (64.0)	3677 (62.9)
Folic acid supplementation	Yes	5677 (70.7)	1070 (8.5)	56 998 (69.0)	74 466 (74.3)	4741 (82.5)
Paternal								
Age, y		35.1 (5.8)	30.9 (5.8)	32.2 (4.8)	30.4 (6.6)	32.2 (5.2)	32.7 (5.4)	36.2 (5.2)
BMI, kg/m ²		25.0 (3.5)	25.2 (3.3)	26.8 (3.6)	26.8 (4.7)	25.2 (3.2)	25.8 (3.3)	24.8 (3.2)
BMI categories	Underweight (<18.5 kg/m ²)	28 (0.8)	41 (0.5)	2 (0.2)	53 (1.9)	271 (0.4)	242 (0.2)	43 (0.75)

(Continued)

Table . Continued

Characteristics	Category	ABCD (N=8131)	ALSPAC (N=13 049)	BASELINE (N=1436)	BIB Study (N=12 799)	DNBC Study (N=89 107)	MoBa (N=101 975)	NINFEA Study (N=5893)
	Normal (18.5–<25 kg/m ²)	1966 (54.8)	4308 (53.3)	345 (30.9)	953 (35.0)	33 502 (53.5)	42 952 (44.4)	3332 (58.4)
	Overweight (25–<30 kg/m ²)	1372 (38.2)	3111 (38.5)	594 (53.3)	1137 (41.7)	24 529 (39.2)	43 888 (45.3)	1977 (34.6)
	Obese (≥30 kg/m ²)	223 (6.2)	616 (7.6)	174 (15.6)	582 (21.4)	4335 (6.9)	9759 (10.1)	355 (6.2)
Smoking	Yes	...	3459 (37.9)	277 (24.9)	1021 (32.0)	26 242 (30.9)	27 803 (27.3)	...
	None	...	449 (5.5)	2963 (4.1)	...
Alcohol	Light drinking	...	4251 (51.8)	59 577 (82.3)	...
	Moderate/heavy drinking	...	3505 (42.7)	9882 (13.6)	...
Education	Low	190 (8.5)	2959 (25.9)	...	4299 (52.9)	17 069 (21.8)	4245 (4.4)	956 (16.6)
	Medium	398 (17.9)	6391 (55.9)	...	1115 (13.7)	28 230 (36.0)	43 576 (45.1)	2464 (42.8)
	High	1670 (73.9)	2079 (18.2)	...	2709 (33.3)	33 118 (42.2)	48 782 (50.5)	2335 (40.6)

Data are given as mean (SD) or number (percentage). Study numbers are based on singletons with data on at least one outcome and one exposure. Light smoking, <10 cigarettes per day; heavy smoking, ≥10 cigarettes per day; maternal light drinking, >0 and <3 units per week during pregnancy; maternal moderate/heavy drinking, ≥3 units per week during pregnancy; paternal light drinking, >0 and <7 units per week; paternal moderate/heavy drinking, ≥7 units per week. ... Indicates data were not available; ABCD, The Amsterdam Born Children and Their Development Study; ALSPAC, Avon Longitudinal Study of Parents and Children; BASELINE, Cork Scope Baseline Study; BIB, Born in Bradford; BMI, body mass index; CHD, congenital heart disease; DNBC, Danish National Birth Cohort; MoBa, Norwegian Mother, Father and Child Cohort Study; and NINFEA, Nascita e Infanzia: gli Effetti dell'Ambiente (Birth and Childhood: Effects of the Environment).

*Chromosomal/genetic/teratogenic anomalies with a cause thought to be already known (see Table S3 for classifications).

[†]Denotes that the study had data specifically during the first trimester. Numbers in the moderate/heavy columns for smoking and alcohol do not add up to the number of any smoking/alcohol because some studies used trimester-specific data for the binary data, whereas moderate/heavy is an assessment of the exposure throughout pregnancy.




Figure 1. Associations between maternal and paternal prepregnancy/early-pregnancy body mass index (BMI) and offspring congenital heart disease (CHD).

A, Odds ratios (ORs) of CHD for a 1-unit (1-kg/m²) difference in maternal BMI (top) and paternal BMI (bottom) in each study and pooled across studies. **B**, The pooled (across ALSPAC [Avon Longitudinal Study of Parents and Children], BiB [Born in Bradford] study, DNBC [Danish National Birth Cohort] study, and MoBa [Norwegian Mother, Father and Child Cohort Study]) results for maternal (top) and paternal (bottom) BMI categories. Results are ORs of CHD in comparison to normal BMI. **C** and **D**, ORs of nonsevere CHD and severe CHD, respectively, for BMI categories in comparison to normal BMI (pooled across ALSPAC, BiB study, DNBC study, and MoBa). All results are adjusted for confounders (depending on cohort: maternal and paternal age, education, ethnicity, smoking, alcohol, maternal parity, and offspring sex) as well as the other parent's BMI. The study-specific results for BMI categories are shown in Figures S15 through S20. In **D**, there were too few cases with paternal BMI data to report results. These analyses are from the LifeCycle project, a consortium that brings birth cohorts together and harmonizes individual-level data for their use in research.¹³ All LifeCycle studies with eligible data were included in this study. More information on each can be found as follows: ABCD (The Amsterdam Born Children and Their Development Study),¹⁵ ALSPAC,^{16,17} BASELINE (Cork Scope Baseline Study),¹⁸ BiB study,¹⁹ DNBC study,²⁰ MoBa,^{21,22} and NINFEA (Nascita e Infanzia: gli Effetti dell'Ambiente [Birth and Childhood: Effects of the Environment]) study.^{23,24}

Downloaded from <http://ahajournals.org> by on September 3, 2021




Figure 2. Associations in each study and pooled across studies for maternal and paternal pregnancy smoking and offspring congenital heart disease (CHD).

Maternal first-trimester smoking was prioritized and used where possible. **A**, Odds ratios (ORs) of any CHD for maternal smoking during pregnancy (top) and paternal smoking (bottom). **B**, ORs of any CHD after removing those with a chromosomal/genetic defect from the study population. **C** and **D**, ORs of nonsevere CHD and severe CHD, respectively. All results are adjusted for confounders (depending on cohort: maternal and paternal age, education, ethnicity, alcohol, maternal parity, and offspring sex) as well as the other parent's smoking. These analyses are from the LifeCycle project, a consortium that brings birth cohorts together and harmonizes individual-level data for their use in research.¹³ All LifeCycle studies with eligible data were included in this study. More information on each can be found as follows: ABCD (The Amsterdam Born Children and Their Development Study),¹⁵ ALSPAC (Avon Longitudinal Study of Parents and Children),^{16,17} BASELINE (Cork Scope Baseline Study),¹⁸ BiB (Born in Bradford) study,¹⁹ DNBC (Danish National Birth Cohort) study,²⁰ MoBa (Norwegian Mother, Father and Child Cohort Study),^{21,22} and NINFEA (Nascita e Infanzia: gli Effetti dell'Ambiente [Birth and Childhood: Effects of the Environment]) study.^{23,24}

Alcohol and CHDs

Because of lack of relevant paternal data, we were unable to undertake negative control analyses for any first-trimester alcohol consumption. Maternal-only associations for that exposure are presented herein, followed by the negative control analyses for levels of alcohol intake at any time in pregnancy. With adjustment for all confounders, any maternal first-trimester

alcohol consumption was not associated with odds of offspring CHD in meta-analyses from 5 cohorts (OR, 1.03; 95% CI, 0.94–1.13) (Figure S28). There was a small increase in risk when restricting these analyses to nonsevere CHD (OR, 1.07; 95% CI, 0.93–1.22), although CIs included the null. Associations for severe CHD were null (OR, 0.91; 95% CI, 0.73–1.12) (Figure S29).

In confounder and other parental alcohol-adjusted analyses, there was weak evidence of an association between maternal light alcohol intake and CHDs (OR, 1.15; 95% CI, 0.90–1.48), which appeared stronger than that seen for paternal alcohol (OR, 1.01; 95% CI, 0.63–1.62), although with no strong statistical support for a difference ($P_{\text{difference}}=0.63$). Associations for moderate/heavy intake were consistent for maternal and paternal alcohol ($P_{\text{difference}}=0.90$), with point estimates showing weak positive associations, but with wide CIs that included the null (Figure 3A and 3B). We did not test associations between levels of alcohol intake and CHD severity because of small numbers. Because of the small number of cohorts having paternal alcohol data, we also show confounder-adjusted models

(without mutual paternal adjustment) for maternal alcohol intake (Figure 3C). The point estimate for maternal light drinking was close to the null and that for heavy drinking suggested it resulted in increased risk of offspring CHD. However, both of these estimates had wide CIs because of relatively few women reporting drinking (particularly heavily) during pregnancy. Results in unadjusted analyses were unchanged (Figure S30).

Between-Study Heterogeneity and Additional Analyses

We have included heterogeneity statistics (I^2 and $P_{\text{heterogeneity}}$) in all figures. Analyses of continuously




Figure 3. Associations in each study and pooled across studies for maternal and paternal pregnancy alcohol intake and any offspring congenital heart disease (CHD).

A, Confounder and other parent's alcohol adjusted odds ratios (ORs) of any CHD for maternal light drinking during pregnancy (top) and paternal light drinking (bottom). **B**, Confounder and other parent's alcohol-adjusted ORs of any CHD for maternal moderate/heavy drinking during pregnancy (top) and paternal moderate/heavy drinking (bottom). **C**, Confounder-adjusted ORs of any CHD for maternal light drinking during pregnancy (top) and maternal moderate/heavy drinking (bottom). Confounders (depending on cohort): maternal and paternal age, education, ethnicity, smoking, maternal parity, and offspring sex (and other parental alcohol intake in **A** and **B**). Definitions for maternal/paternal alcohol intake are described in the Methods section. These analyses are from the LifeCycle project, a consortium that brings birth cohorts together and harmonizes individual-level data for their use in research.¹³ All LifeCycle studies with eligible data were included in this study. More information on each can be found as follows: ABCD (The Amsterdam Born Children and Their Development Study),¹⁵ ALSPAC (Avon Longitudinal Study of Parents and Children),^{16,17} BASELINE (Cork Scope Baseline Study),¹⁸ BiB (Born in Bradford) study,¹⁹ DNBC (Danish National Birth Cohort) study,²⁰ MoBa (Norwegian Mother, Father and Child Cohort Study),^{21,22} and NINFEA (Nascita e Infanzia: gli Effetti dell'Ambiente [Birth and Childhood: Effects of the Environment]) study.^{23,24}

measured BMI and severe CHDs in additional analyses (Figure S14) and BMI analyzed as categories with severe CHDs (Figures S19 and S20) were the only results where we found any statistical evidence of heterogeneity. Across the remaining analyses for all exposures, there was no strong evidence of between-study heterogeneity. Removal of those with any known genetic/chromosomal defects from the study population did not notably alter any main or severity subgroup analyses for BMI and alcohol consumption. However, for smoking, removal of offspring with a chromosomal/genetic defect increased the magnitude of the association for maternal smoking and CHDs (OR, 1.15; 95% CI, 1.01–1.32), and slightly decreased that for paternal smoking (OR, 0.93; 95% CI, 0.83–1.05) ($P_{\text{difference}}=0.02$) (Figure 2B). Furthermore, the positive association between maternal smoking and nonsevere CHDs was slightly stronger when removing those with chromosomal/genetic defects from the study population (OR, 1.25; 95% CI, 1.05–1.49) (Figure S26). All maternal results were materially unchanged after additional adjustment for folic acid supplementation (Figures S12, S27, and S32).

DISCUSSION

In this large multicohort study, we found evidence that maternal pregnancy smoking may increase offspring CHD risk via intrauterine mechanisms and that this may be driven by a specific effect on nonsevere CHDs. We did not find robust evidence to suggest a causal intrauterine effect of higher maternal prepregnancy/early-pregnancy mean BMI or overweight or obesity on offspring CHD risk. Nor did we find evidence of an intrauterine effect of alcohol consumption on offspring CHD risk, although we acknowledge that for alcohol, we had less data and limited statistical power. To our knowledge, this is the first study to use a parental negative control method to explore whether maternal exposures have a causal intrauterine effect on offspring CHDs or whether associations are explained by residual confounding, which would generate a similar association for parental exposures.

We found increased odds of offspring CHD in mothers who were overweight and obese. This is consistent with the most recent systematic review and meta-analysis, which included 2 416 546 participants (57 172 with offspring CHD) from 19 studies and reported increased risk of any offspring CHD in women who were overweight or obese during pregnancy.⁵ However, adjustment for confounders was poor, with 10 of the 19 included studies not providing information on confounder adjustment or not adjusting for any confounders. With more stringent confounder adjustment and the findings from a negative control study, our results suggest that the increased risk of offspring CHD

in overweight and obese mothers is largely the result of residual confounding. We also found that mothers who were underweight at the start of pregnancy were at increased risk of having offspring with CHD, whereas underweight in fathers appeared to be protective of offspring CHD. There were 9537 underweight mothers (4.4%) but only 680 underweight fathers (0.4%) in our study population, making the paternal analyses imprecise and our negative control analyses lacking in power to reliably identify parental differences. The recent systematic review mentioned above did not report on associations of underweight with CHDs because too few studies looked at this.

A large Swedish linkage study of >2 million singleton live born infants (born between 1992 and 2012, with 28 628 CHD cases) has explored associations with maternal underweight, as well as overweight and 3 grades of obesity.⁷ It is difficult to directly compare the results from that study with ours as we only present results for any CHD (and CHD stratified by severity), whereas they only present associations of maternal BMI with specific subtypes of CHDs. The fact that we lack statistical power in our study to explore associations with specific subtypes is a limitation. However, magnitudes of associations of BMI categories and nonsevere CHDs in our study appear to be broadly consistent with several nonsevere defects in the large Swedish study, including atrial septal defects and isolated valve defects. In their study, risks of offspring CHD were similar in underweight compared with normal weight women for all types of CHD (analyzed individually), except for mitral to tricuspid valve defects (14 cases), pulmonary valve defects (24 cases), and right ventricular defects (5 cases), where there was some evidence of increased prevalence with underweight. However, these estimates were based on small numbers and hence imprecise, with CIs including the null. Although our findings suggest maternal underweight might increase offspring risk of CHDs, we lacked power to rule out residual confounding in our negative control analyses, and as noted above the large Swedish study had limited power to determine precise effects in relation to maternal underweight for specific types of CHD where point estimates suggested potentially important magnitudes of increased risk. Other studies that we are aware of have not explored associations of maternal underweight. Thus, any possible effect of maternal underweight on CHD risk remains unclear. As the prevalence of CHD in some low- and middle-income countries is high,⁴³ and these countries currently experience the double burden of undernutrition and overnutrition, we would argue that further exploration of any possible impact of maternal underweight is warranted.

Consistent with our findings, a recent meta-analysis of >8 million participants (137 575 CHD cases) from 125 studies reported positive associations between

maternal pregnancy smoking and offspring CHDs.⁸ There was substantial heterogeneity ($I^2=89%$) in their pooled results, and only 68% of the included studies report adjustment for confounders. The authors also report positive associations between maternal passive smoking and paternal active smoking with offspring CHDs, both of which (somewhat unexpectedly) had stronger magnitudes of association than results from maternal active smoking. Our results, including the negative control study, add to the previous research findings by providing more robust evidence that these associations for maternal smoking are unlikely to be explained by residual confounding and are potentially causal. Other research has shown that pregnancy smoking is a risk factor for orofacial clefts.⁴⁴ The prevalence of CHD is around 1% in the general population, as shown in our study, yet in those with orofacial clefts, CHD prevalence rates of up to 20% have been reported.⁴⁵ Both the heart and the palate develop during early pregnancy, around weeks 5 to 9. Therefore, it is plausible that smoking in early pregnancy could disturb common biological pathways in these conditions. We found that the associations for maternal smoking were possibly largely driven by an effect in nonsevere CHDs, with the association strengthening when those with chromosomal or genetic defects were removed. Previous research has reported positive associations between maternal smoking and septal defects, in particular for atrial septal defects,^{46–48} which are defined as nonsevere according to the classification system used in our study. However, caution is needed in interpreting results by subgroups based on severity. First, one of the largest studies (MoBa) did not have information on case severity and so the severity subgroup analyses are based on different participants and have lower statistical power than in the main analyses. Second, even had all studies been included in the severity analyses, by definition, subgroup analyses have limited power in comparison to main analyses. Third, and more important, caution is required with any subgroup analyses as it is common for multiple characteristics to differ between subgroups in addition to the subgroup defining feature (herein, CHD severity).

In confounder-adjusted analyses, maternal alcohol consumption in the first trimester of pregnancy was not associated with offspring CHD. There was some evidence that maternal moderate or heavy alcohol consumption any time in pregnancy was associated with increased risk of offspring CHD. Although associations between mothers and fathers light, moderate, and heavy alcohol consumption, compared with none, were statistically consistent, only 2 cohorts (80 627 participants, 703 with offspring CHD) had alcohol information on fathers around the time of their partners pregnancy. Associations for fathers in particular were imprecise, with wide CIs. Two recent meta-analyses found consistent modest increases in risk of offspring CHD in

mothers reporting alcohol consumption in pregnancy (OR, 1.11 [95% CI, 0.96–1.29]⁴⁹ and OR, 1.16 [95% CI, 1.05–1.27]⁹). Although the first of these concluded “no association,” it can be seen that the results for the 2 are consistent, and the larger sample size of the second has increased precision. Of note, the second of these studies also explored paternal consumption and found increased risk of offspring CHD related to fathers’ alcohol consumption (OR, 1.44 [95% CI, 1.19–1.74]).⁹ Although the OR for fathers’ consumption suggests a stronger effect, the CIs are wide, and the result is statistically consistent with that for mothers’ alcohol consumption. As in our study, there were fewer studies with data on paternal alcohol consumption around the time of their partners pregnancy. Taken together with our findings, these suggest that positive associations of maternal alcohol consumption with offspring CHD may be attributable to residual confounding rather than a causal intrauterine effect.

The key strengths of this study are its large sample size, the use of a negative paternal exposures control study, and the pooling of results from several cohort studies that are less prone to selection bias that can occur in case-control studies and are not selected on the basis of publication, but on being part of an existing collaboration. The latter reduces the risk of publication bias as studies were included if they had data and not on the basis of (published) results. This also allowed us to explore replication across studies, and the consistency of findings between studies in our main analyses adds confidence to our conclusions.

The use of harmonized data from LifeCycle is a strength that limits between-study heterogeneity. However, harmonizing data across several studies, as we have done in LifeCycle, can mean that some variables lose detail. Herein, that is particularly relevant for the exposure and confounding variables. For example, we were not able to explore pack weeks of smoking across the entire pregnancy. Simplified confounder measurements, such as Western versus non-Western for ethnicity, could result in residual confounding if more specific ethnic groups have strong influences on exposure and outcome. Furthermore, there were other confounders that we considered, including type 1/pre-existing diabetes mellitus and physical activity, but had too few numbers (diabetes mellitus) across all cohorts or too few studies with data (physical activity) to include. However, we aimed to address any form of residual confounding in our paternal negative control analyses. Under the assumption that adjusted for but poorly measured (eg, ethnicity) or unadjusted for (eg, physical activity) confounders influence paternal exposures in the same direction and to the same extent as in mothers, observing parental consistency of association implies that the maternal association is influenced by residual confounding.

We were not able to fully harmonize outcome data, with the key differences between studies being the extent to which they only included cases that were diagnosed antenatally or at birth or whether they included cases later in life. MoBa (N=101 975 participants and N=879 cases) only had cases diagnosed antenatally or around the time of birth, with the remaining cohorts having diagnoses beyond antenatal care, ranging from 6 months to 25 years. Many previous studies have only included cases diagnosed at birth or early infancy. They, and the cohorts included herein that only have these early life cases, may be biased by outcome misclassification (ie, the offspring who would have been diagnosed later in life are treated as not having CHD). This is an important point for consideration because although most CHDs are identified in utero or at birth, many are diagnosed after discharge from hospital during childhood or even adulthood.⁵⁰ Therefore, it is reassuring that our main results are largely consistent across studies. In confounder and other parent-adjusted smoking analyses, the weakest association was found in the MoBa cohort. It is likely that we missed some nonsevere cases in MoBa, which were diagnosed later in life. Given that we demonstrate the smoking results were largely driven by nonsevere CHDs, this could have biased MoBa (and therefore meta-analysis) results toward the null.

The negative control analyses assume that factors that would confound the maternal exposure-offspring CHD associations would have a similar magnitude and direction of confounding for the equivalent paternal associations, irrespective of whether the confounders are measured or if measured how accurately and precisely they are measured. This is likely to be true for paternal negative control exposure studies, as used herein.^{10,11} Both maternal and paternal BMI, smoking, and alcohol consumption could have preconceptional effects via influences on gametes, including epigenetic changes. Any such effects would plausibly differ between mothers and fathers, and for the mother would be in addition to potential intrauterine effects, such that we may still expect stronger maternal associations. Furthermore, there is little conclusive evidence of effects of factors, such as smoking, on gametes that do not render them infertile but are sufficient to influence embryo development and hence CHDs, as such studies are difficult in humans. Heart development occurs in utero (specifically in early pregnancy), and we would expect passive paternal smoke inhalation to expose the fetus to a lower level of exposure than active maternal smoking. As proof of concept, paternal smoking does not associate with offspring birth weight or fetal growth parameters (assessed by repeated ultrasound), in contrast to maternal smoking, which has marked effects.¹² It is possible

that potential differences in misreporting smoking and alcohol consumption between mothers and fathers could produce spurious parental differences. Pregnant women are likely to underreport whether they smoke or drink alcohol and the amount they smoke or drink, because of the social stigma of these, particularly in recent decades. As the report of alcohol and smoking in the LifeCycle cohorts was collected early in pregnancy, it is likely to be random in relation to an offspring CHD as the vast majority would not have been diagnosed. Hence, this underreporting would be expected to attenuate any true effect of smoking/alcohol on CHD toward the null. This misclassification is less likely in fathers. Thus, the specific positive association of maternal smoking on CHDs and its difference to the paternal association may be underestimated.

Finally, only 47% of mothers with smoking data in our study had this specifically during the first trimester. Paternal smoking was defined as smoking around the time of pregnancy, with no specific trimester measurements. However, although amount smoked may change across pregnancy, it is highly likely that any smoking in later trimesters is a strong proxy for smoking in the first trimester. More important, we have shown that our results using only maternal first-trimester smoking are consistent with our main results. Similarly, paternal smoking at any time during pregnancy is likely to be a good proxy for smoking in early pregnancy. However, we acknowledge it would be useful to have more detailed data on both parents across all trimesters to explore whether association magnitudes vary by trimester.

In summary, we found evidence to support a causal intrauterine effect of maternal smoking on any CHD, particularly with nonsevere CHDs, but did not find robust evidence for a causal effect of maternal BMI or alcohol on offspring CHD risk. Although everyone should be encouraged not to smoke, and all clinical guidelines advocate not starting smoking, and if women do smoke, to quit before becoming pregnant, there are still high rates of smoking in some groups, particularly those from deprived backgrounds. In the studies included in this article, in 2 contemporary cohorts, BASELINE (Ireland), with births occurring between 2008 and 2011, and the BiB cohort (United Kingdom), with births occurring between 2007 and 2011, smoking prevalence rates were 25% and 16%, respectively. The prevalence in the BiB cohort masks the high rate in White British women (33%) who are from socioeconomically deprived backgrounds, as >50% of births in that cohort are to Pakistani women who have low rates of smoking (3%).¹⁹ It is possible that emphasizing the potential adverse effect on CHDs in specific groups might help in supporting women of reproductive age not to start smoking and women who are smoking at the start of pregnancy to

be encouraged to quit. Furthermore, understanding the specific mechanisms that link maternal smoking to increased offspring CHD risk could identify targets for interventions for its prevention.

ARTICLE INFORMATION

Received November 4, 2020; accepted March 29, 2021.

Affiliations

Population Health Science, Bristol Medical School, Bristol, United Kingdom (K.T., A.E., D.A.L.); Medical Research Council Integrative Epidemiology Unit at the University of Bristol, United Kingdom (K.T., A.E., D.A.L.); Division of Health Data and Digitalisation, Department of Genetics and Bioinformatics, Norwegian Institute of Public Health, Oslo, Norway (J.L.T.N.); Bradford Institute for Health Research, Bradford Teaching Hospitals National Health Service Foundation Trust, Bradford, United Kingdom (T.C.Y., D.M., J.W.); Cancer Epidemiology Unit, Department of Medical Sciences, University of Turin and CPO Piemonte, Turin, Italy (E.I., L.R.); Department of Public and Occupational Health, Amsterdam Public Health Research Institute, Amsterdam University Medical Center, University of Amsterdam, the Netherlands (T.V.); Section for Epidemiology, Department of Public Health, University of Copenhagen, Denmark (A.P.d., A.N.A.); The Irish Centre for Fetal and Neonatal Translational Research (D.M.M., D.F.); and Department of Paediatrics and Child Health (D.M.M., D.F.), University College Cork, Cork, Ireland; Centre for Reviews and Dissemination, University of York, Heslington, York, United Kingdom (S.O.); Department for Health Evidence, Radboud Institute for Health Sciences, Radboud University Medical Center, Nijmegen, the Netherlands (N.R.); Division of Health Data and Digitalisation (J.R.H.); and Centre for Fertility and Health (J.R.H.), Norwegian Institute of Public Health, Oslo, Norway; Translational Science, Bristol Medical School, Bristol, United Kingdom (M.C.); and Bristol National Institute for Health Research Biomedical Research Center, Bristol, United Kingdom (M.C., D.A.L.).

Acknowledgments

We are grateful to all participating hospitals, obstetric clinics, and general practitioners for their assistance in implementing ABCD (The Amsterdam Born Children and Their Development Study) and thank all of the women who participated for their cooperation. We are grateful to all the families who took part in the ALSPAC (Avon Longitudinal Study of Parents and Children), the midwives for their help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists, and nurses. The authors thank the families for their continued support and the BASELINE (Cork Scope Baseline Study) Birth Cohort Study research team. BiB (Born in Bradford) study is only possible because of the enthusiasm and commitment of the children and parents. We are grateful to all the participants, practitioners, and researchers who have made BiB study happen. The DNBC (Danish National Birth Cohort) study was established with a significant grant from the Danish National Research Foundation. Additional support was obtained from the Danish Regional Committees, the Pharmacy Foundation, the Egmont Foundation, the March of Dimes Birth Defects Foundation, the Health Foundation, and other minor grants. The DNBC Biobank has been supported by the Novo Nordisk Foundation and the Lundbeck Foundation. We are grateful to all the participating families in Norway who take part in the ongoing MoBa (Norwegian Mother, Father and Child Cohort Study). The authors thank all families participating in the NINFEA (Nascita e Infanzia: gli Effetti dell'Ambiente [Birth and Childhood: Effects of the Environment]) study cohort. Data access: The Horizon2020 LifeCycle Project is a cross-cohort collaboration that brings together data from multiple birth cohorts from across Europe and Australia. A major product of this collaboration has been the establishment of a findable, accessible, interoperable, and reusable data resource known as the European Union Child Cohort Network (<https://lifecycle-project.eu/for-scientists/variable-catalogue/>). The ABCD study welcomes collaboration and the interest of colleagues. A set of guidelines for researchers interested in using data for research purposes is available at <http://www.abcd-study.nl>. The ALSPAC data management plan (<http://www.bristol.ac.uk/alspac/researchers/data-access/documents/alspac-data-management-plan.pdf>) describes in detail the policy on data sharing, which is through a system of managed open access. Researchers interested in using BASELINE data can contact the study (baseline@ucc.ie), and further

information can be found (<http://www.baselinestudy.net/> or <http://www.birthcohorts.net/>). Scientists are encouraged to make use of the BiB study data, which are available through a system of managed open access. Before you contact BiB study, please make sure you have read the Guidance for Collaborators. The DNBC study operates an open access policy. More information can be found on the study website (<https://www.dnbc.dk/access-to-dnbc-data>). MoBa data are used by researchers and research groups at both the Norwegian Institute of Public Health and other research institutions nationally and internationally. The research must adhere to the aims of MoBa and the participants' given consent. All use of data and biological material from MoBa is subject to Norwegian legislation. More information can be found on the study website (<https://www.fhi.no/en/studies/moba/for-forskere-artikler/research-and-data-access/>). Researchers interested in using NINFEA study data can contact info@progettoninfea.it, and more information can be found at the study website (<https://www.progettoninfea.it/>).

Sources of Funding

The LifeCycle project received funding from the 452 European Union's Horizon 2020 Research and Innovation Programme 453 (Grant Agreement No. 733206 LifeCycle). This study is also supported by the British Heart Foundation (AA/18/7/34219), Bristol National Institute for Health Research (NIHR) Biomedical Research Centre grant, European Research Council (Advanced Grant; 669545), and US National Institutes of Health (R01 DK10324). K. Taylor is supported by a British Heart Foundation Doctoral Training Program (FS/17/60/33474). K. Taylor, Dr Elhakeem, and Prof Lawlor work in a unit that is supported by the University of Bristol and UK Medical Research Council (MC_UU_00011/6). Prof Lawlor is supported by a British Heart Foundation Chair in Cardiovascular Science and Clinical Epidemiology (CH/F/20/90003) and a NIHR Senior Investigator (NF-0616-10102). Prof Caputo is supported by the British Heart Foundation Chair in Congenital Heart Disease (CH/1/32804). Dr Pinot de Moira is funded by a Lundbeck Foundation Fellowship (R264-2017-3099). Work by Dr Harris is partly supported by the Research Council of Norway through its Centers of Excellence Funding Scheme, project No. 262700. Details of funding for each individual LifeCycle cohort that has contributed to this study are provided below. ABCD (The Amsterdam Born Children and Their Development Study) was supported by the Netherlands Organization for Health Research and Development (grant 2100.0076) and the Electromagnetic Fields and Health Research Program (grants 85600004 and 85800001). Core funding for ALSPAC (Avon Longitudinal Study of Parents and Children) is provided by the UK Medical Research Council and Wellcome (217065/Z/19/) and the University of Bristol. Many grants have supported different data collections, including for some of the data used in this publication, and a comprehensive list of grant funding is available on the ALSPAC website (<http://www.bristol.ac.uk/alspac/external/documents/grant-acknowledgements.pdf>). The BASELINE (Cork Scope Baseline Study) Birth Cohort Study was funded by the National Children's Research Centre (Project grant: Jan 2009). BiB (Born in Bradford) study has received core support from Wellcome Trust (WT101597MA), a joint grant from the UK Medical Research Council and UK Economic and Social Science Research Council (MR/N024397/1), the British Heart Foundation (CS/16/4/32482), and the NIHR Applied Research Collaboration Yorkshire and Humber (NIHR200166) and Clinical Research Network. The views expressed in this publication are those of the author(s) and not necessarily those of the National Institute for Health Research or the Department of Health and Social Care. The DNBC (Danish National Birth Cohort) study was supported by the Danish Epidemiology Science Centre, the Lundbeck Foundation (grant 195/04), the Egmont Foundation, the March of Dimes Birth Defect Foundation, the Augustinus Foundation, and the Medical Research Council (grant SSVF 0646). The MoBa (Norwegian Mother, Father and Child Cohort Study) is supported by the Norwegian Ministry of Health and Care Services and the Ministry of Education and Research. The NINFEA (Nascita e Infanzia: gli Effetti dell'Ambiente [Birth and Childhood: Effects of the Environment]) study was partially funded by the Compagnia San Paolo Foundation.

Disclosures

Prof Lawlor reports support from Roche Diagnostics and Medtronic Ltd for research unrelated to that presented herein. The remaining authors have no disclosures to report.

Supplementary Material

Data S1–S4
Tables S1–S11
Figures S1–S32
References S1–59

REFERENCES

- Wang H, Naghavi M, Allen C, Barber RM, Bhutta ZA, Carter A, Casey DC, Charlson FJ, Chen AZ, Coates MM, et al. Global, regional, and national life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death, 1980–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet*. 2016;388:1459–1544. DOI: 10.1016/S0140-6736(16)31012-1.
- McCracken C, Spector LG, Menk JS, Knight JH, Vinocur JM, Thomas AS, Oster ME, St Louis JD, Moller JH, Kochilas L. Mortality following pediatric congenital heart surgery: an analysis of the causes of death derived from the national death index. *J Am Heart Assoc*. 2018;7:e010624. DOI: 10.1161/JAHA.118.010624.
- van der Bom T, Zomer AC, Zwiderman AH, Meijboom FJ, Bouma BJ, Mulder BJM. The changing epidemiology of congenital heart disease. *Nat Rev Cardiol*. 2011;8:50–60. DOI: 10.1038/nrcardio.2010.166.
- Jenkins KJ, Correa A, Feinstein JA, Botto L, Britt AE, Daniels SR, Elixson M, Warnes CA, Webb CL. Noninherited risk factors and congenital cardiovascular defects: current knowledge: a scientific statement from the American Heart Association Council on Cardiovascular Disease in the Young: endorsed by the American Academy of Pediatrics. *Circulation*. 2007;115:2995–3014. DOI: 10.1161/CIRCULATIONAHA.106.183216.
- Liu X, Ding G, Yang W, Feng X, Li Y, Liu H, Zhang Q, Ji L, Li D. Maternal body mass index and risk of congenital heart defects in infants: a dose-response meta-analysis. *Biomed Res Int*. 2019;2019:1315796. DOI: 10.1155/2019/1315796.
- Persson M, Cnattingius S, Villamor E, Söderling J, Pasternak B, Stephansson O, Neovius M. Risk of major congenital malformations in relation to maternal overweight and obesity severity: cohort study of 1.2 million singletons. *BMJ*. 2017;357:j2563. DOI: 10.1136/bmj.j2563.
- Persson M, Razaz N, Edstedt Bonamy A-K, Villamor E, Cnattingius S. Maternal overweight and obesity and risk of congenital heart defects. *J Am Coll Cardiol*. 2019;73:44–53. DOI: 10.1016/j.jacc.2018.10.050.
- Zhao L, Chen L, Yang T, Wang L, Wang T, Zhang S, Chen L, Ye Z, Zheng Z, Qin J. Parental smoking and the risk of congenital heart defects in offspring: an updated meta-analysis of observational studies. *Eur J Prev Cardiol*. 2019;27:1284–1293. DOI: 10.1177/2047487319831367.
- Zhang S, Wang L, Yang T, Chen L, Zhao L, Wang T, Chen L, Ye Z, Zheng Z, Qin J. Parental alcohol consumption and the risk of congenital heart diseases in offspring: an updated systematic review and meta-analysis. *Eur J Prev Cardiol*. 2020;27:410–421. DOI: 10.1177/2047487319874530.
- Davey SD. Negative control exposures in epidemiologic studies. *Epidemiology*. 2012;23:350–351. DOI: 10.1097/EDE.0b013e318245912c.
- Sanderson E, Macdonald-Wallis C, Davey SG. Negative control exposure studies in the presence of measurement error: implications for attempted effect estimate calibration. *Int J Epidemiol*. 2018;47:587–596. DOI: 10.1093/ije/dyx213.
- Brand JS, Gaillard R, West J, McEachan RRC, Wright J, Voerman E, Felix JF, Tilling K, Lawlor DA. Associations of maternal quitting, reducing, and continuing smoking during pregnancy with longitudinal fetal growth: findings from Mendelian randomization and parental negative control studies. *PLoS Med*. 2019;16:e1002972. DOI: 10.1371/journal.pmed.1002972.
- Jaddoe VVW, Felix JF, Andersen A-M, Charles M-A, Chatzi L, Corpeleijn E, Donner N, Elhakeem A, Eriksson JG, Foong R, et al. The LifeCycle Project-EU Child Cohort Network: a federated analysis infrastructure and harmonized data of more than 250,000 children and parents. *Eur J Epidemiol*. 2020;35:709–724. DOI: 10.1007/s10654-020-00662-z.
- Sutton-Tyrrell K. Assessing bias in case-control studies: proper selection of cases and controls. *Stroke*. 1991;22:938–942. DOI: 10.1161/01.STR.22.7.938.
- van Eijsden M, Vrijkotte TG, Gemke RJ, van der Wal MF. Cohort profile: the Amsterdam Born Children and their Development (ABCD) Study. *Int J Epidemiol*. 2011;40:1176–1186. DOI: 10.1093/ije/dyq128.
- Boyd A, Golding J, Macleod J, Lawlor DA, Fraser A, Henderson J, Molloy L, Ness A, Ring S, Davey SG. Cohort profile: the “Children of the 90s”—the index offspring of the Avon Longitudinal Study of Parents and Children. *Int J Epidemiol*. 2013;42:111–127. DOI: 10.1093/ije/dys064.
- Fraser A, Macdonald-Wallis C, Tilling K, Boyd A, Golding J, Davey Smith G, Henderson J, Macleod J, Molloy L, Ness A, et al. Cohort profile: the Avon Longitudinal Study of Parents and Children: ALSPAC mothers cohort. *Int J Epidemiol*. 2013;42:97–110. DOI: 10.1093/ije/dys066.
- O'Donovan SM, Murray DM, Hourihane JO, Kenny LC, Irvine AD, Kiely M. Cohort profile: the Cork BASELINE Birth Cohort Study: babies after SCOPE: evaluating the longitudinal impact on neurological and nutritional endpoints. *Int J Epidemiol*. 2015;44:764–775. DOI: 10.1093/ije/dyu157.
- Wright J, Small N, Raynor P, Tuffnell D, Bhopal R, Cameron N, Fairley L, Lawlor DA, Parslow R, Petherick ES, et al. Cohort profile: the Born in Bradford multi-ethnic family cohort study. *Int J Epidemiol*. 2013;42:978–991. DOI: 10.1093/ije/dys112.
- Olsen J, Melbye M, Olsen SF, Sørensen TIA, Aaby P, Nybo Andersen A-M, Taxbøl D, Hansen KD, Juhl M, Schow TB, et al. The Danish National Birth Cohort—its background, structure and aim. *Scand J Public Health*. 2001;29:300–307. DOI: 10.1177/14034948010290040201.
- Magnus P, Irgens LM, Haug K, Nystad W, Skjaerven R, Stoltenberg C; MoBa Study Group. Cohort profile: the Norwegian Mother and Child Cohort Study (MoBa). *Int J Epidemiol*. 2006;35:1146–1150. DOI: 10.1093/ije/dyl170.
- Magnus P, Birke C, Vejrup K, Haugan A, Alsaker E, Daltveit AK, Handal M, Haugen M, Høiseith G, Knudsen GP, et al. Cohort profile update: the Norwegian Mother and Child Cohort Study (MoBa). *Int J Epidemiol*. 2016;45:382–388. DOI: 10.1093/ije/dyw029.
- Firestone R, Cheng S, Pearce N, Douwes J, Merletti F, Pizzi C, Pivetta E, Rusconi F, Richiardi L. Internet-based birth-cohort studies: is this the future for epidemiology? *JMIR Res Protoc*. 2015;4:e71. DOI: 10.2196/resprot.3873.
- Richiardi L, Baussano I, Vizzini L, Douwes J, Pearce N, Merletti F; NINFEA cohort. Feasibility of recruiting a birth cohort through the Internet: the experience of the NINFEA cohort. *Eur J Epidemiol*. 2007;22:831–837. DOI: 10.1007/s10654-007-9194-2.
- Best KE, Rankin J. Increased risk of congenital heart disease in twins in the North of England between 1998 and 2010. *Heart*. 2015;101:1807–1812. DOI: 10.1136/heartjnl-2015-307826.
- Glinianaia SV, Rankin J, Wright C. Congenital anomalies in twins: a register-based study. *Hum Reprod*. 2008;23:1306–1311. DOI: 10.1093/humrep/den104.
- Pinot de Moira A, Haakma S, Strandberg-Larsen K, van Enckevort E, Kooijman M, Cadman T, Cardol M, Corpeleijn E, Crozier SR, Duijts L, et al. The EU Child Cohort Network's core data: establishing a set of findable, accessible, interoperable and re-usable (FAIR) variables. *Eur J Epidemiol*. 2020. Jul 23 [epub ahead of print]. DOI: 10.1007/s10654-021-00733-9.
- Gittenberger-de Groot AC, Bartelings MM, Deruiter MC, Poelmann RE. Basics of cardiac development for the understanding of congenital heart malformations. *Pediatr Res*. 2005;57:169–176. DOI: 10.1203/01.PDR.0000148710.69159.61.
- Taylor K, Thomas R, Mumme M, Golding J, Boyd A, Northstone K, Caputo M, A Lawlor D. Ascertaining and classifying cases of congenital anomalies in the ALSPAC birth cohort. *Wellcome Open Res*. 2020;5:231. DOI: 10.12688/wellcomeopenres.16339.1.
- Bishop C, Small N, Mason D, Corry P, Wright J, Parslow RC, Bittles AH, Sheridan E. Improving case ascertainment of congenital anomalies: findings from a prospective birth cohort with detailed primary care record linkage. *BMJ Paediatr Open*. 2017;1:e000171. DOI: 10.1136/bmjpo-2017-000171.
- Lynge E, Sandegaard JL, Rebolj M. The Danish National Patient Register. *Scand J Public Health*. 2011;39:30–33. DOI: 10.1177/1403494811401482.
- Schmidt M, Schmidt SAJ, Sandegaard JL, Ehrenstein V, Pedersen L, Sørensen HT. The Danish National Patient Registry: a review of content, data quality, and research potential. *Clin Epidemiol*. 2015;7:449. DOI: 10.2147/CLEP.S91125.
- Øyen N, Olsen SF, Basit S, Leirgul E, Strøm M, Carstensen L, Granstrøm C, Tell GS, Magnus P, Vollset SE, et al. Association between maternal folic acid supplementation and congenital heart defects in offspring in birth cohorts from Denmark and Norway. *J Am Heart Assoc*. 2019;8:e011615. DOI: 10.1161/JAHA.118.011615.
- Dale MTG, Magnus P, Leirgul E, Holmstrøm H, Gjessing HK, Brodwall K, Haugen M, Stoltenberg C, Øyen N. Intake of sucrose-sweetened soft beverages during pregnancy and risk of congenital heart defects (CHD) in offspring: a Norwegian pregnancy cohort study. *Eur J Epidemiol*. 2019;34:383–396. DOI: 10.1007/s10654-019-00480-y.
- Taylor K, Elhakeem A. LifeCycle Project: the effect of maternal pre-/early-pregnancy BMI and pregnancy smoking and alcohol on congenital

- heart diseases: a parental negative control study. medRxiv. Preprint. November 4, 2020.
36. Di Castelnuovo A, Quacquarello G, Donati MB, de Gaetano G, Iacoviello L. Spousal concordance for major coronary risk factors: a systematic review and meta-analysis. *Am J Epidemiol*. 2008;169:1–8. DOI: 10.1093/aje/kwn234.
 37. Madley-Dowd P, Rai D, Zammit S, Heron J. Simulations and directed acyclic graphs explained why assortative mating biases the prenatal negative control design. *J Clin Epidemiol*. 2020;118:9–17. DOI: 10.1016/j.jclinepi.2019.10.008.
 38. Sharp GC, Lawlor DA. Paternal impact on the life course development of obesity and type 2 diabetes in the offspring. *Diabetologia*. 2019;62:1802–1810. DOI: 10.1007/s00125-019-4919-9.
 39. Balduzzi S, Rücker G, Schwarzer G. How to perform a meta-analysis with R: a practical tutorial. *Evid Based Ment Health*. 2019;22:153–160. DOI: 10.1136/ebmental-2019-300117.
 40. Higgins JPT, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ*. 2003;327:557–560. DOI: 10.1136/bmj.327.7414.557.
 41. Xu A, Cao X, Lu Y, Li H, Zhu Q, Chen X, Jiang H, Li X. A meta-analysis of the relationship between maternal folic acid supplementation and the risk of congenital heart defects. *Int Heart J*. 2016;57:725–728. DOI: 10.1536/ihj.16-054.
 42. He Y, Pan A, Hu FB, Ma X. Folic acid supplementation, birth defects, and adverse pregnancy outcomes in Chinese women: a population-based mega-cohort study. *Lancet*. 2016;388:S91. DOI: 10.1016/S0140-6736(16)32018-9.
 43. Hoffman JIE. The global burden of congenital heart disease: review article. *Cardiovasc J Afr*. 2013;24:141–145. DOI: 10.5830/CVJA-2013-028.
 44. Leite M, Albieri V, Kjaer SK, Jensen A. Maternal smoking in pregnancy and risk for congenital malformations: results of a Danish register-based cohort study. *Acta Obstet Gynecol Scand*. 2014;93:825–834. DOI: 10.1111/aogs.12433.
 45. Genisca AE, Frias JL, Broussard CS, Honein MA, Lammer EJ, Moore CA, Shaw GM, Murray JC, Yang W, Rasmussen SA, et al. Orofacial clefts in the National Birth Defects Prevention Study, 1997–2004. *Am J Med Genet A*. 2009;149A:1149–1158. DOI: 10.1002/ajmg.a.32854.
 46. Zhang D, Cui H, Zhang L, Huang Y, Zhu J, Li X. Is maternal smoking during pregnancy associated with an increased risk of congenital heart defects among offspring? A systematic review and meta-analysis of observational studies. *J Matern Fetal Neonatal Med*. 2017;30:645–657. DOI: 10.1080/14767058.2016.1183640.
 47. Malik S, Cleves MA, Honein MA, Romitti PA, Botto LD, Yang S, Hobbs CA. Maternal smoking and congenital heart defects. *Pediatrics*. 2008;121:e810–e816. DOI: 10.1542/peds.2007-1519.
 48. Källén K. Maternal smoking and congenital heart defects. *Eur J Epidemiol*. 1999;15:731–737.
 49. Wen Z, Yu DI, Zhang W, Fan C, Hu L, Feng YU, Yang L, Wu Z, Chen R, Yin K-J, et al. Association between alcohol consumption during pregnancy and risks of congenital heart defects in offspring: meta-analysis of epidemiological observational studies. *Ital J Pediatr*. 2016;42:12. DOI: 10.1186/s13052-016-0222-2.
 50. Massin MM, Dessy H. Delayed recognition of congenital heart disease. *Postgrad Med J*. 2006;82:468–470. DOI: 10.1136/pgmj.2005.044495.
 51. Northstone K, Lewcock M, Groom A, Boyd A, Macleod J, Timpson N, Wells N. The Avon Longitudinal Study of Parents and Children (ALSPAC): an update on the enrolled sample of index children in 2019. *Wellcome Open Res*. 2019;4:51. DOI: 10.12688/wellcomeopenres.15132.1.
 52. Shire K, Andrews E, Barber S, Bruce A, Corkett J, Hill LJB, Kelly B, McEachan R, Mon-Williams M, Tracey L, et al. Starting School: a large-scale start of school assessment within the “Born in Bradford” longitudinal cohort. *Wellcome Open Res*. 2020;5:47. DOI: 10.12688/wellcomeopenres.15610.1.
 53. Popovic M, Pizzi C, Rusconi F, Galassi C, Gagliardi L, Marco LD, Migliore E, Merletti F, Richiardi L. Infant weight trajectories and early childhood wheezing: the NINFEA birth cohort study. *Thorax*. 2016;71:1091–1096. DOI: 10.1136/thoraxjnl-2015-208208.
 54. Nederlof M, de Walle HEK, van Poppel MNM, Vrijkotte TGM, Gademan MGJ. Deviant early pregnancy maternal triglyceride levels and increased risk of congenital anomalies: a prospective community-based cohort study. *BJOG*. 2015;122:1176–1183. DOI: 10.1111/1471-0528.13393.
 55. Freathy RM, Kazeem GR, Morris RW, Johnson PCD, Paternoster L, Ebrahim S, Hattersley AT, Hill A, Hingorani AD, Holst C, et al. Genetic variation at CHRNA5-CHRNA3-CHRNA4 interacts with smoking status to influence body mass index. *Int J Epidemiol*. 2011;40:1617–1628. DOI: 10.1093/ije/dyr077.
 56. Åsvold BO, Bjørngaard JH, Carlslake D, Gabrielsen ME, Skorpen F, Smith GD, Romundstad PR. Causal associations of tobacco smoking with cardiovascular risk factors: a Mendelian randomization analysis of the HUNT study in Norway. *Int J Epidemiol*. 2014;43:1458–1470. DOI: 10.1093/ije/dyu113.
 57. Morris RW, Taylor AE, Fluharty ME, Bjørngaard JH, Åsvold BO, Gabrielsen ME, Campbell A, Marioni R, Kumari M, Korhonen T, et al. Heavier smoking may lead to a relative increase in waist circumference: evidence for a causal relationship from a Mendelian randomisation meta-analysis: the CARTA consortium. *BMJ Open*. 2015;5:e008808. DOI: 10.1136/bmjopen-2015-008808.
 58. Lawlor DA, Nordestgaard BG, Benn M, Zuccolo L, Tybjaerg-Hansen A, Davey SG. Exploring causal associations between alcohol and coronary heart disease risk factors: findings from a Mendelian randomization study in the Copenhagen General Population Study. *Eur Heart J*. 2013;34:2519–2528. DOI: 10.1093/eurheartj/ehd081.
 59. Pastorino S, Bishop T, Crozier SR, Granström C, Kordas K, Küpers LK, O'Brien EC, Polanska K, Sauder KA, Zafarmand MH, et al. Associations between maternal physical activity in early and late pregnancy and offspring birth size: remote federated individual level meta-analysis from eight cohort studies. *BJOG*. 2019;126:459–470. DOI: 10.1111/1471-0528.15476.

SUPPLEMENTARY MATERIAL

The effect of maternal pre-/early-pregnancy BMI and pregnancy smoking and alcohol on congenital heart diseases: a parental negative control study

Contents and navigation menu

Table of Contents

Supplementary Methods	3
<i>Data S1. Cohort descriptions</i>	3
<i>Table S1. Study-specific methods for data collection</i>	6
<i>Data S2. Paternal alcohol consumption</i>	9
<i>Data S3. Definition of congenital heart disease (CHD) and other congenital anomalies (CAs)</i>	10
<i>Data S4. Confounder data</i>	14
Supplementary Results – Tables S4-S11	16
<i>Missing data</i>	16
<i>Sensitivity analysis: complete-case analyses</i>	17
<i>Supplementary Tables S9-S11</i>	23
Supplementary Results – Figures S1-S32	25
<i>Participant flow charts for each cohort</i>	25
<i>BMI analyses – Supplementary Figures</i>	32
<i>Smoking analyses – Supplementary Figures</i>	47
<i>Alcohol analyses - Supplementary Figures</i>	55

Supplementary Methods

Data S1. Cohort descriptions

The Amsterdam Born Children and their Development Study (ABCD)

The following text was adapted from the ABCD cohort profile where full study details are described (<https://doi.org/10.1093/ije/dyq128>)¹⁵:

Between January 2003 and March 2004, all pregnant women living in Amsterdam were asked to participate in the ABCD study during their first prenatal visit to an obstetric care provider (general practitioner, midwife or gynaecologist). Altogether, 12 373 women were approached—by estimate, ≥99% of the target population. According to Dutch law, all pregnant women, including illegal immigrants and asylum-seekers, are entitled to receive prenatal care, which is free of charge if costs are a problem. For all of the women approached, the care provider completed a registration form which included personal data such as name, address and date of birth. Based on this information, a questionnaire covering socio-demographic characteristics, obstetric history, lifestyles and psychosocial conditions was sent to the pregnant women within 2 weeks, to be filled out at home and returned to the Public Health Service by prepaid mail. A reminder was sent 2 weeks later. The questionnaire included an informed consent sheet the women could use to grant permission for follow-up of their infants at the age of 3 months and every 5 years thereafter, and for the perusal of their medical files. Approval for the ABCD study was obtained from the Central Committee on Research involving Human Subjects in the Netherlands, the Medical Ethical Committees of the participating hospitals, and from the Registration Committee of the Municipality of Amsterdam. Written informed consent was obtained from all participating mothers.

Of the 12 373 women approached, 8266 women filled out the pregnancy questionnaire (response rate: 67%). Of this group, 7050 women granted permission for follow-up (85%) and 7043 women granted permission for perusal of her and her child's medical files (85%). To enhance participation among foreign-born women, two supportive measures were taken: (i) a Turkish, Arabic or English translation was provided to women born in Turkey, Morocco or other non-Dutch-speaking countries and (ii) the possibility of completing the questionnaire orally was offered to women who were illiterate or had reading difficulties.

The Avon Longitudinal Study of Parents and Children (ALSPAC)

ALSPAC is a prospective birth cohort study which was devised to investigate the environmental and genetic factors of health and development. Detailed information about the methods and procedures of ALSPAC is available elsewhere^{16,17,51}. 14,541 pregnant women with an expected delivery date of April 1991 and December 1992, residing in the former region of Avon, UK were eligible to take part. Additional enrolment provided a baseline sample of 14,901 participants⁵¹. The study website contains details of all the data that is available through a fully searchable data dictionary. Ethical approval for the study was obtained from the ALSPAC Ethics and Law Committee and the Local Research Ethics Committees (<http://www.bristol.ac.uk/alspac/researchers/research-ethics/>). Informed consent for the use of data collected via questionnaires and clinics was obtained from participants following the recommendations of the ALSPAC Ethics and Law Committee at the time.

The Cork SCOPE BASELINE Birth Cohort Study (BASELINE)

The following text was adapted from the BASELINE cohort profile where full study details are described: <https://doi.org/10.1093/ije/dyu157>¹⁸.

The study is based in Cork, Ireland. The SCOPE Ireland pregnancy cohort formed the basis of recruitment of infants to BASELINE (n = 1537). In 2007, the amalgamation of all three Cork maternity units into one centre, Cork University Maternity Hospital (CUMH), provided a unique opportunity to conduct research in pregnancy in Cork. CUMH, which is co-located with the Cork University Hospital, is the third largest maternity hospital in Ireland, with 8563 deliveries in 2012. As recruitment was regionally based, the generalizability of the data may be limited. In 2008, all primiparous women in Cork were invited to take part in the Screening for Pregnancy Endpoints (SCOPE) pregnancy cohort. The SCOPE cohort is an international collaboration of research groups interested in the study of major adverse outcomes in late pregnancy, particularly but not exclusively, pre-eclampsia, fetal growth restriction and spontaneous preterm birth⁸ and as a consequence strict exclusion criteria were applied.⁹ Detailed maternal, fetal and paternal information was obtained antenatally, as well as blood samples at 15 and 20 weeks' gestation, see Table 1. All women who participated in the SCOPE study were informed about the birth cohort, and if consent was obtained infants were registered to the Cork BASELINE birth cohort.

The Born in Bradford Cohort (BiB)

The Born in Bradford study is a population-based prospective birth cohort including 12,453 women who experienced 13,776 pregnancies between 2007 and 2011. The study is unique in that it has almost an equal split between White European and South Asian women, all residing in Bradford, UK. Bradford is a city in the North of England with high levels of socioeconomic deprivation, and the cohort was started due to a high prevalence of poor child health in the city⁵². Full details of the study methodology were reported previously¹⁹. The study website provides more information, including protocols, questionnaires and information on how researchers can access data and a full list of all available data (<https://borninbradford.nhs.uk/research/documents-data/>). Mothers, and their partners, recruited into the study provided detailed interview questionnaire data, measurements, and biological samples. They also consented to the linkage of theirs and their child's data.

The Danish National Birth Cohort (DNBC)

The DNBC is a nationwide cohort of pregnant women, recruited from 1996 through 2002 consisting of 100,415 pregnancies²⁰. Informed consent was obtained from participants upon enrolment, and the study was approved by the Danish Data Protection Agency through the joint notification of the Faculty of Health and Medical Sciences at the University of Copenhagen (Sund-2017-09), according to Danish regulations. Information on lifestyle and environmental factors potentially associated with offspring health was collected through 4 prenatal and postnatal telephone interviews at target ages gestational weeks 12 and 30 and child ages 6 and 18 months. The parent-child dyads were then invited for follow-up at 7, 11, and 18 years.

The Norwegian Mother, Father and Child Cohort Study (MoBa)

MoBa is a nationwide, pregnancy cohort comprising family triads (mother-father-offspring) who are followed longitudinally. All pregnant women in Norway who were able to read Norwegian were

eligible for participation. The first child was born in October 1999 and the last in July 2009. Invitations were sent to women in 277 702 pregnancies, the participation rate was 41%. The cohort includes more than 114 000 children, 95 000 mothers and 75 000 fathers^{21,22}. Extensive longitudinal data were collected using nine questionnaires: three during pregnancy, and then follow-up questionnaires when the children were 6 months, 18 months, 36 months, 5 years, 7 years and 8 years of age. In addition, a single questionnaire was administered to fathers during gestational weeks 15-18. Data collected include general background and health information, including diet and lifestyle, a semi-quantitative food frequency questionnaire, information on birth and pregnancy outcomes, and on several aspects of child nutrition and development, as well as the physical and mental health of both mother and child. MoBa is linked to the Medical Birth Registry of Norway, which provides standardized information about the health of the mother during pregnancy, other essential medical information related to the pregnancy and birth, and standard post-natal measures of the child. The establishment of MoBa and initial data collection was based on a license from the Norwegian Data Protection Agency and approval from The Regional Committees for Medical and Health Research Ethics. The MoBa cohort is now based on regulations related to the Norwegian Health Registry Act.

NINFEA study

The NINFEA study is internet-based birth cohort established in 2005 in Italy (<http://www.progettoninfea.it>)^{23,24,53}. The cohort consists of children born to mothers who have access to the internet and enough knowledge of Italian to complete online questionnaires. The recruitment was conducted actively, through obstetrics clinics, and passively, via internet and the media. A baseline questionnaire on general health and exposures before and during pregnancy is completed by mothers at enrolment, which may occur at any time during pregnancy. During the period 2005-2016 around 7500 mothers were recruited. Further follow-up information was obtained with repeated questionnaires completed 6 and 18 months after delivery and when children turn 4, 7, 10 and 13 years. The response rates for each questionnaire are available at <https://www.progettoninfea.it/attachments/70>. The study was approved by the ethical committee of the San Giovanni Battista Hospital and CTO/CRF/Maria Adelaide Hospital (Turin, Italy) (approval N.0048362 and following amendments).

Table S1. Study-specific methods for data collection.

Measurement	Study-specific details
<i>BMI data</i>	
Maternal BMI	<p>ABCD: Women filled out a questionnaire containing questions on sociodemographic characteristics, medical history, lifestyle and dietary habits (16 weeks of gestation; IQR 12–20 weeks). BMI was based on pre-pregnancy height and weight as reported in the pregnancy questionnaire.</p> <p>ALSPAC: In the 2nd pregnancy questionnaire (12 weeks' gestation) women were asked to report their pre-pregnancy weight and height. No definition of pre-pregnancy was provided in the question. Subsequently for the majority of women all weight measurements from any time of pregnancy have been extracted from obstetric records (height was not routinely measured antenatally in the UK when these women were pregnant). First antenatal clinic measurements of weight correlated strongly with the women's self-report (Pearson correlation = 0.93).</p> <p>Baseline: At 15 weeks' gestation sociodemographic and anthropometric measurements, including objectively measured weight and height, were collected.</p> <p>BiB: Weight and height (unshod and in light clothing and following a standard protocol) were measured at the recruitment assessment. As women were recruited at the oral glucose tolerance test (26-28 weeks of gestation for the majority) this would not provide an accurate measure of pre-/early-pregnancy weight and would include fetal and amniotic weight and pregnancy related weight gain. All measurements of weight from all antenatal clinics were extracted from the obstetric records and pre-/early-pregnancy BMI was calculated using weight from the first antenatal clinic (median 12 weeks' gestation) and height at recruitment (26-28 weeks' gestation).</p> <p>DNBC: Self-reported information on pre-pregnancy weight and height from the first pregnancy interview at around 16 weeks' gestation.</p> <p>MoBa: Pre-pregnancy weight and height were self-reported during the first interview at week 17 in pregnancy.</p> <p>NINFEA: Pre-pregnancy weight and height were self-reported in the baseline questionnaire (completed at any time during pregnancy).</p>
Paternal BMI	<p>ABCD: Paternal weight was maternally reported in questionnaire when child was aged 5-6 years (the closest timepoint available to pregnancy). Paternal height was maternally reported in the pregnancy questionnaire at around 16 weeks' gestation.</p> <p>ALSPAC: Paternal weight and height were self-reported from the first partner questionnaire completed around 18 weeks' gestation.</p> <p>Baseline: Paternal weight and height were measured around the time of pregnancy.</p> <p>BiB: Paternal weight and height were self-reported from the first partner questionnaire mostly completed at recruitment (26–28 weeks' gestation).</p>

	<p>DNBC: Paternal weight and height were reported by the mother during the first pregnancy interview conducted at around 16 weeks' gestation.</p> <p>MoBa: Paternal weight and height were maternally reported by questionnaire at around 18 weeks' gestation.</p> <p>NINFEA: Paternal weight and height were maternally reported in the baseline questionnaire (completed at any time during pregnancy).</p>
<i>Smoking data</i>	
Maternal smoking	<p>ABCD: Asked number of cigarettes per day during pregnancy in first questionnaire (16 weeks of gestation; IQR 12–20 weeks). Binary variable used any smoking during pregnancy.</p> <p>ALSPAC: Asked number of cigarettes per day during pregnancy in questionnaire at around 18 weeks' gestation. Binary variable used any smoking during the first trimester.</p> <p>Baseline: Reported in early pregnancy questionnaire around 14 weeks gestation. Binary variable used any smoking during the first trimester. Baseline smoking data only used to adjust for BMI analyses.</p> <p>BiB: Asked number of cigarettes per day during pregnancy in first questionnaire (26-28 weeks' gestation). Binary variable used any smoking during pregnancy.</p> <p>DNBC: Maternal smoking in the first trimester was ascertained from a computer-assisted telephone interview conducted at approximately 16 weeks' gestation. Binary variable used any smoking during the first trimester. Smoking heaviness was based on the average number of cigarettes smoked per day reported in interviews 1 and 2.</p> <p>MoBa: Smoking habits were assessed from questionnaires sent by mail at 13-17 and 30 weeks. Binary variable used any smoking during pregnancy.</p> <p>NINFEA: Smoking habits in the first two trimesters were assessed in the baseline questionnaire (completed any time during pregnancy). Binary variable used any smoking during the first trimester.</p>
Paternal smoking	<p>ABCD: NA</p> <p>ALSPAC: Asked about smoking habits within the partner questionnaire during pregnancy at around 18 weeks' gestation.</p> <p>Baseline: Maternally reported in pregnancy questionnaire around 14 weeks' gestation.</p> <p>BiB: Asked about smoking habits within partner questionnaire during pregnancy (26-28 weeks' gestation).</p> <p>DNBC: Maternally reported at 16 weeks' gestation.</p> <p>MoBa: Self-reported within first partner questionnaire around 15 weeks' gestation.</p> <p>NINFEA: NA</p>
<i>Alcohol data</i>	
Maternal alcohol	<p>ABCD: Mothers asked how many glasses of alcohol they drunk during first period of pregnancy (16 weeks of gestation; IQR 12–20 weeks). Binary variable used any alcohol intake during pregnancy.</p> <p>ALSPAC: Self-reported from pregnancy questionnaire at around 18 weeks' gestation. Binary variable used any alcohol intake during the first trimester.</p>

	<p>Baseline: Reported in early pregnancy questionnaire around 14 weeks gestation. Binary variable used any alcohol intake during the first trimester. Baseline alcohol data only used to adjust for BMI analyses.</p> <p>BiB: NA</p> <p>DNBC: Self-reported at 16 weeks' gestation. Binary variable used alcohol intake during the first trimester. Drinking heaviness was based on the average number of units drank per week reported in interviews 1 and 2.</p> <p>MoBa: Assessed via questionnaire around 17 weeks' gestation. Binary variable used any alcohol intake during the first trimester.</p> <p>NINFEA: Drinking habits in the first trimester were assessed in the baseline questionnaire (completed at any time during pregnancy). Binary variable used any alcohol intake during the first trimester.</p>
Paternal alcohol	<p>ALSPAC: Self-reported within first partner questionnaire at around 18 weeks' gestation.</p> <p>MoBa: Self-reported within first partner questionnaire at around 15 weeks' gestation.</p>

Data S2. Paternal alcohol consumption**ALSPAC**

We used data from the partners questionnaire which was filled in by partners at around 18 weeks' gestation. We used data from questions B18 and B19 from the PB questionnaire (<http://www.bristol.ac.uk/alspac/researchers/our-data/>).

B18b. How often have you drunk alcoholic drinks during the last 3 months: 1) Never, 2) less than once a week, 3) at least once a week, 4) 1-2 glasses every day, 5) 3-9 glasses every day, 6) at least 10 glasses every day.

B19b. How many days in the past month did you drink the equivalent of 2 pints of beer, 4 glasses of wine or 4 pub measures of spirit? 1) Every day, 2) more than 10 days, 3) 5-10 days, 4) 3-4 days, 5) 1-2 days, 6) none.

We coded paternal alcohol consumption as follows: non-drinkers = If answered 1 to B18b; light drinkers = answered 5 to B19b; mod/heavy drinkers = answered 1,2,3 or 4 to B19b.

MoBa

Question FF244. How often do you drink alcohol now that your partner is pregnant? Response options: 1) Approximately 6-7 times per week, 2) Approximately 4-5 times per week, 3) Approximately 2-3 times per week, 4) Approximately once per week, 5) Approximately 1-3 times per month, 6) Less than once per month, 7) Never.

Using data from FF244, we coded paternal alcohol consumption as follows: non-drinkers = Answered number 7; light drinkers = Answered 4, 5 or 6; mod/heavy drinkers = Answered 1, 2 or 3

Data S3. Definition of congenital heart disease (CHD) and other congenital anomalies (CAs)

Here we describe ascertainment of CA cases for each cohort. International Classification of Diseases (ICD; version 10) codes were used to define CA cases when possible (see Table S2 above for classifications). However, in some cohorts these data were not available. The following cohorts were used to define CA cases with ICD codes: ALSPAC, BiB, DNBC, NINFEA.

ABCD

The ABCD cohort has previously published research involving CAs⁵⁴. The same methods for data extraction were used for the present study. Data on CAs were obtained from three different sources: the infant questionnaire, which was filled out by the mother at an average infant age of 12.9 weeks (IQR 12.4–13.4 weeks); the questionnaire filled out by the mother at an average infant age of 5.07 years (IQR 5.04–5.13 years), and clinical data of the Youth Health Care Registration (health and development registration of all children in the Netherlands, which is mandatory under the law on medical treatment agreement). The questionnaires were screened by a researcher, and in the case of missing or unclear answers the mothers were contacted. Subsequently, the questionnaires were scanned and transferred to a database by a certified company (Scan serv, Nootdorp, the Netherlands). Missing data in the questionnaires could be supplemented by data from the Youth Health Care Registration, and in the case of any discrepancy the data from the Youth Health Care Registration prevailed. CA data in ABCD was restricted to live-born children.

CAs were categorized as follows: 0 = no defect 1 = congenital malformations of the nervous system 2 = congenital malformations of eye, ear, face, throat 3 = congenital malformations of the cardiovascular system 4 = congenital malformations of the respiratory tract 5 = split lip and/or palate 6 = congenital malformations of the digestive tract 7 = congenital malformations of the kidneys, urinary tract, genitalia 8 = congenital malformations of the musculoskeletal system 9 = neoplasms 10 = other congenital malformations 11 = chromosomal defect 12 = monogenic defect 13 = microdeletions and uniparental disomy 14 = other syndromes 15 = complex cardiovascular defects 16 = multiple defects of the extremities 17 = other multiple defects within an organ system 18 = multiple defects (in multiple organ systems) 21 = minor defect 22 = unclear/uncertain diagnosis 23 = "don't know which defect" 24 = "not applicable" 25 = missing information.

We coded CHD cases if they were "Yes" for category 3. We coded chromosomal/genetic aberrations if "Yes" for any of the following categories: 11, 12, 13, 14.

ALSPAC

Case ascertainment of CAs in the ALSPAC cohort has been described in detail in a recently published data note²⁹. Data were combined from multiple sources: NHS records (primary care, paediatric cardiology database, data on fetal deaths and local child health services), midwifery and birth records and maternal self-report via child-based questionnaires. Each source was coded using ICD-10 codes. By combining sources, there would be a greater possibility of capturing all of possible cases within the cohort. The majority of cases of CAs were identified by primary care records (79% for any CA and 68% for any CHD). We included diagnoses made at any age (from birth up until age 25/26). There were no restrictions in cases of CAs in ALSPAC, we included all cases whether live-born or not. However, it is possible that some

cases that were terminated earlier in pregnancy were missed due to them never having an NHS number and thus not being identified through record linkage.

BASELINE

At 2 months, mothers were asked of any medical problems and/or referrals. If a baby had been referred to a specialist, it was checked to see if they had results from an echocardiogram. Echocardiograms were checked by a cardiologist. Exact CHD diagnoses were reported based on the echo. At 6 months, there was one additional baby that had cardiac surgery and added as a case. If a baby had been diagnosed after 6 months, they would have been identified through records on the Echo. Therefore, in BASELINE we obtained all CHDs up until ~age 12.

BiB

In the BiB cohort, there were two separate sources to identify CAs. Both sources were used in this study: (i) CAs up to 5 years of age, identified in GP records by Bishop et al³⁰ following EUROCAT guidelines. ICD-10 codes were mapped to clinical term (CT)-V3 codes prior to extraction from GP records. (ii) Data extracted from the Yorkshire and Humber CAs register database. Data were ICD-10 coded. All of these were confirmed postnatally. BiB includes data on the birth outcome of each child (live birth, miscarriage, still birth). Therefore, diagnoses were not necessarily restricted to live born children. However, there is the possibility that some women would have terminated the pregnancy after the 12- or 20-week scans which would lead to an under-representation of congenital anomaly cases.

DNBC

In the DNBC, all diagnoses of congenital anomalies (according to EUROCAT guide 1.4 section 3.2 and 3.3) up until the age of 15 years were extracted from the Danish National Patient Register (DNPR) which is linked to the cohort data^{31,32}. Diagnoses were ICD-coded. These data were restricted to children born alive.

MoBa

Information on whether a child had a CHD or not was obtained through linkage to the Medical Birth Registry of Norway (MBRN). All maternity units in Norway must notify births to the MBRN. The notification form includes the name and personal identity number of the child and parents, as well as information about maternal health before and during pregnancy, and any complications during pregnancy or at birth, including the presence of any heart defects. The MBRN contains information on all births and pregnancies ended after the 12th week of gestation, including stillbirths and abortions after the 12th week, including on heart defects. Heart defects are registered in the MBRN through notifications from clinical staff identifying these defects at delivery or any hospital in patient treatments occurring immediately after birth until the child is discharged. The medical notification is made at discharge, which can be several months after birth. Details of the notified heart defects, such as specific diagnosis or treatment are not provided. Whilst most of the heart defects would have been diagnosed at birth it is possible that some children were admitted to hospital after delivery for non-specific reasons of for diagnoses that at the time were not considered to be related to a heart defect. Therefore, MOBA contribute only to analyses of any CHD and we considered diagnosis to have been made between birth and 6 months (few would remain in hospital after this length).

NINFEA

Congenital anomalies in the NINFEA cohort were reported in the second questionnaire compiled 6 months after birth. Mothers compiled a checklist that included pre-specified anomalies (namely cryptorchidism (also assessed 18 months after birth), congenital hip dysplasia, cleft palate, spina bifida and pyloric stenosis) and anomalies divided by major systems (namely cardiovascular, gastrointestinal, genitourinary, musculoskeletal, respiratory and nervous system, and genetic/chromosomal or metabolic/endocrine disease). If the mother reported an anomaly from a specific system, the exact name of the anomaly was asked. If the child died or had any surgery performed in the first 6 months, the cause of death and type of surgery were also checked to see if any congenital anomaly was reported. All congenital anomalies were coded using ICD-10 codes by an experienced pediatrician and were reassessed by an independent physician. NINFEA included live-born infants only.

Studies with ICD coded data

Table S2 shows how cases of CHD were defined in the studies with ICD codes (ALSPAC, BiB, DNBC, NINFEA).

Table S2. Subcategories of CHD.

Category	CHDs included/excl	ICD codes
All	Any CHD as defined by EUROCAT* Patent ductus arteriosus (PDA) with gestational age (GA) < 37 weeks not considered a CHD case. Peripheral pulmonary artery stenosis with GA < 37weeks not considered as a CHD case.	Q20-Q25, Q260, Q262-Q269**
Severe	Heterotaxia, Conotruncal defect, Atrioventricular septal defect, Anomalous pulmonary venous return, Left ventricle outflow tract obstruction, Right ventricle outflow tract obstruction, Other complex defects	Q240, Q241, Q206, Q200, Q251, Q252, Q253, Q254, Q203, Q213, Q201, Q214, Q212, Q26, Q262, Q264, Q268, Q269, Q234, Q251, Q230, Q231, Q221, Q224, Q225, Q255, Q204
Non-severe	PDA (in full term infants), valvular pulmonary stenosis, ventricular septal defect (VSD), atrial septum defects (ASD), unspecified septal defects, isolated valve defects, other specified heart defects, unspecified heart defects	Non-severe cases that are All=1 and Severe=0.
* Definitions taken from here: https://eu-rd-platform.jrc.ec.europa.eu/sites/default/files/EUROCAT-Guide-1.4-Section-3.3.pdf		
**Q250 and Q256 not a case if isolated and GA<37weeks		

Additional analysis - excluding infants with any known chromosomal/genetic/teratogenic defects

ABCD, ALSPAC, BiB, DNBC, MoBa and NINFEA were able to contribute to this additional analysis. In ALSPAC, BiB, DNBC and NINFEA, we used the ICD codes in Table S3 to exclude cases. In ABCD, there were specific categories (described above) which corresponded to chromosomal and genetic anomalies (11 = chromosomal defect 12 = monogenic defect 13 = microdeletions and uniparental disomy 14 = other syndromes). In MoBa, we used questionnaire data which was maternally reported at 6 months after birth: “Is your child suspected of having a syndrome?” and “Is your child suspected of having a chromosomal defect?”.

Table S3. Subcategories of congenital anomalies with a ‘known cause’ used in additional analyses.

Category	ICD-10 Codes
Teratogenic/genetic syndromes, microdeletions and chromosomal abnormalities (additional analysis).	D821, P350-P352, P371, Q619, Q751, Q754, Q771-Q772, Q780, Q796, Q85, Q861-Q869, Q87, Q90-Q92, Q930-Q939, Q95-Q99

Data S4. Confounder data

By definition a confounder has to cause (or be a plausible cause) of exposure and outcome. The maximum number of confounders used in fully adjusted models are listed below. Confounder and other parent exposure adjusted models are the same as fully adjusted but with additional adjustment for the other parent's exposure and additional adjustment for maternal parity in paternal models.

Exposure = BMI: age, education, parity (maternal), ethnicity, smoking, alcohol, offspring sex.

Exposure = Smoking: age, education, parity (maternal), ethnicity, alcohol, offspring sex.

Exposure = Alcohol: age, education, parity (maternal), ethnicity, smoking, offspring sex.

There is evidence that smoking and alcohol influence BMI⁵⁵⁻⁵⁸. We therefore treated those as confounders for the association of maternal/paternal BMI with CHD. Smoking and alcohol are associated with each other in most populations but whether one causes the other is unclear. It is possible that most of their association is due to socioeconomic and cultural factors. Despite being unclear about whether they could be confounders of each other's effect on CHD (e.g. alcohol a confounder for smoking and vice versa) in the final confounder adjusted model we included alcohol as a confounder for smoking and vice versa.

We used maternal/paternal age at birth in complete years. We used educational attainment for both parents' measures of socioeconomic position (SEP). In the harmonized LifeCycle data education has been defined according to the international classification (High: Short cycle tertiary, Bachelor, Masters, Doctoral or equivalent (ISCED-2011: 5-8, ISCED-97: 5-6) Medium: Upper secondary, Post-secondary non-tertiary (ISCED-2011: 3-4, ISCED-97: 3-4) Low: No education; early childhood; pre-primary; primary; lower secondary or second stage of basic education). Mothers' parity was based on previous born children (previous stillbirths included, abortions excluded) (coded as 0, 1, 2, 3, ≥ 4). For ethnicity we used the best estimate of the mother's/father's ethnic background based on the cohort's discretion (Western, Non-western, Mixed). Offspring sex was a binary variable (male/female). In additional analyses, we adjusted for folic acid supplementation in fully adjusted maternal models. This was a yes/no variable defined as intake of folic acids (folate, vitamin B9) during the period from conception to early pregnancy (12 weeks).

In NINFEA, due to the smaller sample size, maternal parity and maternal/paternal education were categorized as binary variables (parity: nulliparous and multiparous, education: low and medium combined together).

In ALSPAC, BASELINE, DNBC, MoBa and NINFEA we did not adjust for ethnicity in any analyses. 98% of women were of Western origin in ALSPAC. >98.5% of women in BASELINE were of Western origin. Ethnicity in the DNBC is said to be of >99% White European origin with a recent paper reporting their DNBC population to be 100% of White origin⁵⁹. There were no data available on ethnicity in MoBa, however, it is believed that 99-100% are of Western origin. Ethnicity data were not available in NINFEA, although, the large majority of mothers (>98%) were born in Europe. Data on paternal country of birth was available for approximately half of the cohort and >98% of them were born in Europe.

In BiB only ~28% of mothers had harmonized data on alcohol intake during pregnancy, therefore this was not included in any models within BiB analyses as an exposure and also as a confounder in BMI and smoking models.

ABCD and BASELINE did not have harmonized LifeCycle data available. We describe methods for data harmonization here:

We used available ABCD data and tried to harmonize it as best as possible to match the LifeCycle data. BMI, sex, age, parity and folic acid supplementation were identical variables to the harmonized LifeCycle ones. Paternal height was self-reported by the mother and paternal weight was from 11 months after pregnancy (the closest timepoint available). We used any pregnancy smoking or drinking (yes/no) for the smoking and alcohol variables as there was no trimester specific exposure data. ABCD did not contribute to paternal alcohol or smoking analyses as there were no data for these exposures around the time of pregnancy. Maternal education was defined as: high (Short cycle tertiary, Bachelor, Masters, Doctoral or equivalent (9 or more years)), medium (Upper secondary, Post-secondary non-tertiary (6-9 years)) or low (No education; early childhood; pre-primary; primary; lower secondary or second stage of basic education (<6 years)). Paternal education was from the 11-year questionnaire and split into 3 groups as this was the only data available. For ethnicity, we defined Western and non-western as appropriate from physiological ethnicity of grandmother's birth country for maternal ethnicity. Paternal ethnicity was reported by the mother and recoded to Western/Non-Western/Mixed.

All women were experiencing their first pregnancy in BASELINE; therefore we did not adjust for parity in any analyses. BMI, sex, age and smoking were coded the same as the harmonized LifeCycle data. Education in BASELINE was binary defined as medium or high. This was left unchanged and used as a measure of SEP as in other analyses.

In the analysis plan, we originally stated that we would treat type-1 diabetes (T1D) as a confounder. The rationale for this was that diabetes is a known teratogen for CHDs and could also influence pregnancy lifestyle factors through changes in behaviours. However, after exploring the data, the prevalence of T1D was low in those cohorts with data (0.2% in ALSPAC, 0.1% in BiB and 0.2% in DNBC for maternal T1D) and the other cohorts did not have data on specific diabetes diagnoses. For cohorts with T1D data, the number of CHD cases in those with a diagnosis was either zero or less than 10, making adjustment not meaningful or impossible through complete separation in the logistic model.

Supplementary Results – Tables S4-S11

Missing data

Table S4. Summary of missing data in each cohort.

	ABCD N = 8,131	ALSPAC N = 13,049	BASELINE N = 1,436	BiB N = 12,799	DNBC N = 89,107	MoBa N = 101,975	NINFEA N = 5,893
Country	Netherlands	UK	RoI	UK	Denmark	Norway	Italy
Recruitment period	2003-2004	1991-1992	2008-2011	2007-2011	1996-2002	1999-2008	2005-2016
Maternal (n missing (%))							
Age, years	0	2062 (15.8)	0	0	0	181 (0.2)	1 (0.0)
BMI, kg/m ²	789 (9.7)	2079 (15.9)	0	2690 (21.0)	3757 (4.2)	4575 (4.5)	124 (2.1)
Preg smoking yes/no	14 (0.2)	333 (2.6)	0	1912 (14.9)	2367 (2.7)	933 (0.9)	92 (1.6)
Preg smoking heaviness	-	2350 (18.0)	-	1912 (14.9)	1184 (1.3)	390 (0.4)	72 (1.2)
Preg alcohol yes/no	6 (0.1)	427 (3.3)	43 (3.0)	-	2399 (2.7)	19617 (19.2)	50 (0.9)
Preg alcohol heaviness	-	6548 (50.2)	-	-	758 (0.9)	17539 (17.2)	79 (1.3)
Parity	0	502 (3.8)	0	470 (3.7)	0	1805 (1.8)	272 (4.6)
Education	83 (1.0)	1152 (8.8)	9 (0.6)	2750 (21.5)	8451 (9.5)	6963 (6.8)	46 (0.8)
Ethnicity	14 (0.2)	-	0	1906 (14.9)	-	-	-
Folic acid supp	98 (1.2)	424 (3.2)	-	-	6510 (7.3)	1805 (1.8)	148 (2.5)
Paternal (n missing (%))							
Age, years	4378 (53.8)	5488 (42.1)	321 (22.4)	9439 (73.7)	1371 (1.5)	521 (0.5)	2506 (42.5)
BMI, kg/m ²	4542 (55.9)	4973 (38.1)	321 (22.4)	10074 (78.7)	26470 (29.7)	5134 (5.0)	186 (3.2)
Smoking	-	3915 (30.0)	323 (22.5)	9612 (75.1)	4181 (4.7)	171 (0.2)	-
Alcohol	-	4844 (37.1)	-	-	-	29553 (28.9)	-
Education	5873 (72.2)	1620 (12.4)	0	4676 (36.5)	10690 (12.0)	5372 (5.3)	138 (2.3)
Ethnicity	197 (2.4)	-	321 (22.4)	9625 (75.2)	-	-	-
Offspring sex	203 (2.5)	0	0	0	0	196 (0.2)	1 (0.0)

Sensitivity analysis: complete-case analyses**Table S5.** Comparison between maximal numbers from **main analyses presented in the manuscript (black, top rows)** and **complete case models (red, bottom rows)**. Results are odds ratios (95% CIs) of any offspring CHD per unit difference in BMI.

Model	ABCD	ALSPAC	BASELINE	BiB	DNBC	MoBa	NINFEA	Meta-analysis results
Maternal BMI unadjusted	1.02 (0.94, 1.09) N = 7,342	1.05 (1.00, 1.09) N = 10,970	1.07 (0.92, 1.20) N = 1,436	1.01 (0.97, 1.04) N = 10,109	1.02 (1.01, 1.03) N = 85,350	0.99 (0.98, 1.01) N = 97,400	0.93 (0.83, 1.03) N = 5,769	1.01 (1.00, 1.02) N = 218,376
	1.07 (0.95, 1.16) N = 3,415	1.01 (0.93, 1.08) N = 6,452	1.06 (0.87, 1.23) N = 1,078	0.99 (0.89, 1.10) N = 1,753	1.02 (1.00, 1.03) N = 55,564	0.99 (0.97, 1.01) N = 73,637	0.93 (0.83, 1.04) N = 5,393	1.01 (0.99, 1.02) N = 147,292
Maternal BMI confounder adjusted	1.04 (0.95, 1.11) N = 7,103	1.05 (0.99, 1.10) N = 9,179	1.08 (0.93, 1.21) N = 1,386	1.02 (0.98, 1.05) N = 7,279	1.02 (1.00, 1.03) N = 78,180	0.99 (0.97, 1.01) N = 75,448	0.94 (0.84, 1.05) N = 5,476	1.01 (1.00, 1.02) N = 184,051
	1.05 (0.93, 1.15) N = 3,415	1.01 (0.94, 1.08) N = 6,452	1.06 (0.87, 1.23) N = 1,078	0.98 (0.87, 1.09) N = 1,753	1.01 (1.00, 1.03) N = 55,564	0.99 (0.97, 1.01) N = 73,637	0.95 (0.85, 1.06) N = 5,393	1.01 (0.99, 1.02) N = 147,292
Maternal BMI confounder and other parent BMI adjusted	1.05 (0.93, 1.15) N = 3,415	1.02 (0.94, 1.10) N = 6,452	1.05 (0.85, 1.23) N = 1,078	0.99 (0.88, 1.09) N = 1,753	1.01 (1.00, 1.03) N = 55,564	0.99 (0.97, 1.01) N = 73,637	0.94 (0.84, 1.06) N = 5,393	1.00 (0.99, 1.02) N = 147,292
	1.05 (0.93, 1.15) N = 3,415	1.02 (0.94, 1.10) N = 6,452	1.05 (0.85, 1.23) N = 1,078	0.99 (0.88, 1.09) N = 1,753	1.01 (1.00, 1.03) N = 55,564	0.99 (0.97, 1.01) N = 73,637	0.94 (0.84, 1.06) N = 5,393	1.00 (0.99, 1.02) N = 147,292
Paternal BMI unadjusted	0.99 (0.84, 1.08) N = 3,589	0.99 (0.91, 1.06) N = 8,076	1.07 (0.86, 1.21) N = 1,115	1.03 (0.94, 1.12) N = 2,706	1.02 (1.00, 1.04) N = 62,637	0.99 (0.97, 1.01) N = 96,841	1.02 (0.92, 1.13) N = 5,707	1.01 (0.99, 1.02) N = 180,690
	1.04 (0.88, 1.11) N = 1,732	0.97 (0.86, 1.07) N = 5,044	1.07 (0.86, 1.21) N = 1,113	1.01 (0.89, 1.13) N = 1,572	1.02 (1.00, 1.04) N = 53,922	0.99 (0.96, 1.01) N = 67,071	0.96 (0.81, 1.13) N = 3,166	1.01 (0.99, 1.03) N = 133,620
Paternal BMI confounder adjusted	1.03 (0.84, 1.10) N = 1,800	0.96 (0.86, 1.06) N = 5,550	1.06 (0.86, 1.21) N = 1,113	1.04 (0.93, 1.14) N = 2,085	1.02 (1.00, 1.05) N = 54,710	1.00 (0.97, 1.02) N = 68,623	1.03 (0.89, 1.19) N = 3,294	1.01 (1.00, 1.03) N = 137,175
	1.03 (0.84, 1.10) N = 1,732	0.97 (0.86, 1.08) N = 5,044	1.06 (0.86, 1.21) N = 1,113	1.04 (0.92, 1.16) N = 1,572	1.02 (1.00, 1.04) N = 53,922	1.00 (0.97, 1.02) N = 67,071	0.96 (0.81, 1.14) N = 3,166	1.01 (1.00, 1.03) N = 133,620
Paternal BMI confounder and other parent BMI adjusted	1.03 (0.85, 1.10) N = 1,732	0.97 (0.86, 1.08) N = 5,044	1.05 (0.84, 1.21) N = 1,113	1.04 (0.92, 1.15) N = 1,572	1.02 (1.00, 1.04) N = 53,922	1.00 (0.97, 1.02) N = 67,071	0.99 (0.83, 1.18) N = 3,166	1.01 (0.99, 1.03) N = 133,620
	1.03 (0.85, 1.11) N = 1,732	0.97 (0.86, 1.08) N = 5,044	1.05 (0.84, 1.21) N = 1,113	1.04 (0.92, 1.15) N = 1,572	1.02 (1.00, 1.04) N = 53,922	1.00 (0.97, 1.02) N = 67,071	(0.99, 0.83, 1.18) N = 3,166	1.01 (0.99, 1.03) N = 133,620

Covariates used for each study in fully adjusted models (mutually adjusted models the same as fully adjusted but with additional adjustment for the other parent's BMI and parity in paternal models);
 ABCD: Maternal: offspring sex, age, education, parity, ethnicity, smoking, alcohol. Paternal: offspring sex, age, education, ethnicity.
 ALSPAC: Maternal: offspring sex, age, education, parity, smoking, alcohol. Paternal: offspring sex, age, education, smoking, alcohol.
 BASELINE: Maternal: offspring sex, age, education, smoking, alcohol. Paternal: offspring sex, age, smoking.
 BiB: Maternal: offspring sex, age, education, parity, ethnicity, smoking. Paternal: offspring sex, age, education, ethnicity, smoking.
 DNBC: Maternal: offspring sex, age, education, parity, smoking, alcohol. Paternal: offspring sex, age, education, smoking.
 MoBa: Maternal: offspring sex, age, education, parity, smoking, alcohol. Paternal: offspring sex, age, education, smoking, alcohol.
 NINFEA: Maternal: offspring sex, age, education, parity, smoking, alcohol. Paternal: offspring sex, age, education.

Table S6. Comparison between **maximal numbers (black, top rows)** and **complete case models (red, bottom rows)**. Results are odds ratios (95% CIs) of any offspring CHD for a BMI category in comparison to normal BMI. Categories: underweight (BMI <18.5 kg/m²), normal weight (BMI 18.5 to <25 kg/m²), overweight (BMI 25 to <30 kg/m²) and obese (BMI ≥30 kg/m²).

Exposure	ALSPAC	BiB	DNBC	MoBa	Meta-analysis results
Maternal underweight unadjusted	0.69 (0.26, 1.48) N = 10,970	0.67 (0.17, 0.89) N = 10,109	1.36 (1.05, 1.73) N = 85,350	1.03 (0.70, 1.52) N = 97,400	1.19 (0.97, 1.46) N = 203,829
	0.63 (0.15, 1.78) N = 6,452	NA	1.35 (0.95, 1.86) N = 55,564	1.06 (0.66, 1.71) N = 73,637	1.21 (0.92, 1.57) N = 135,653
Maternal underweight confounder adjusted	0.63 (0.19, 1.57) N = 9,179	0.64 (0.10, 2.11) N = 7,360	1.33 (1.01, 1.71) N = 79,288	1.06 (0.66, 1.71) N = 75,448	1.20 (0.96, 1.50) N = 171,275
	0.68 (0.16, 1.93) N = 6,452	NA	1.34 (0.94, 1.84) N = 55,564	1.08 (0.67, 1.74) N = 73,637	1.21 (0.93, 1.58) N = 135,653
Maternal underweight confounder and other parent BMI adjusted	0.65 (0.15, 1.84) N = 6,452	NA	1.35 (0.95, 1.86) N = 55,564	1.07 (0.67, 1.73) N = 73,637	1.21 (0.93, 1.58) N = 135,653
	0.65 (0.15, 1.84) N = 6,452	NA	1.35 (0.95, 1.86) N = 55,564	1.07 (0.67, 1.73) N = 73,637	1.21 (0.93, 1.58) N = 135,653
Maternal overweight unadjusted	1.23 (0.64, 2.20) N = 10,970	1.35 (0.87, 2.08) N = 10,109	1.24 (1.07, 1.42) N = 85,350	1.01 (0.85, 1.20) 1.02 N = 97,400	1.15 (1.04, 1.28) N = 203,829
	0.71 (0.21, 1.82) N = 6,452	1.46 (0.41, 5.29) N = 1,753	1.28 (1.07, 1.53) N = 55,564	1.04 (0.86, 1.27) N = 73,637	1.16 (1.02, 1.32) N = 137,406
Maternal overweight confounder adjusted	0.85 (0.35, 1.80) N = 9,179	1.34 (0.80, 2.22) N = 7,360	1.23 (1.06, 1.42) N = 79,288	1.06 (0.87, 1.29) N = 75,448	1.17 (1.04, 1.31) N = 171,275
	0.72 (0.21, 1.87) N = 6,452	1.45 (0.39, 5.37) N = 1,753	1.26 (1.05, 1.51) N = 55,564	1.04 (0.85, 1.27) N = 73,637	1.15 (1.01, 1.31) N = 137,406
Maternal overweight confounder and other parent BMI adjusted	0.77 (0.23, 1.99) N = 6,452	1.46 (0.39, 5.42) N = 1,753	1.24 (1.04, 1.49) N = 55,564	1.05 (0.86, 1.29) N = 73,637	1.15 (1.01, 1.31) N = 137,406
	0.77 (0.23, 1.99) N = 6,452	1.46 (0.39, 5.42) N = 1,753	1.24 (1.04, 1.49) N = 55,564	1.05 (0.86, 1.29) N = 73,637	1.15 (1.01, 1.31) N = 137,406
Maternal obesity unadjusted	1.99 (0.95, 3.78) N = 10,970	1.05 (0.62, 1.74) N = 10,109	1.30 (1.06, 1.57) N = 85,350	1.07 (0.85, 1.35) N = 97,400	1.21 (1.05, 1.39) N = 203,829
	1.56 (0.46, 4.00) N = 6,452	0.84 (0.12, 3.93) N = 1,753	1.16 (0.88, 1.51) N = 55,564	1.10 (0.83, 1.44) N = 73,637	1.14 (0.94, 1.37) N = 137,406
Maternal obesity confounder adjusted	2.16 (0.93, 4.43) N = 9,179	1.20 (0.66, 2.11) N = 7,360	1.21 (0.97, 1.49) N = 79,288	1.09 (0.83, 1.43) N = 75,448	1.19 (1.02, 1.40) N = 171,275
	1.72 (0.50, 4.49) N = 6,452	0.67 (0.10, 3.33) N = 1,753	1.14 (0.86, 1.48) N = 55,564	1.09 (0.83, 1.44) N = 73,637	1.12 (0.93, 1.36) N = 137,406
Maternal obesity confounder and other parent BMI adjusted	1.88 (0.55, 4.93) N = 6,452	0.70 (0.09, 3.44) N = 1,753	1.10 (0.83, 1.43) N = 55,564	1.12 (0.85, 1.49) N = 73,637	1.12 (0.93, 1.36) N = 137,406
	1.88 (0.55, 4.93) N = 6,452	0.70 (0.09, 3.44) N = 1,753	1.10 (0.83, 1.43) N = 55,564	1.12 (0.85, 1.49) N = 73,637	1.12 (0.93, 1.36) N = 137,406
Paternal underweight unadjusted	NA	NA	0.59 (0.10, 1.84) N = 62,637	1.97 (0.73, 5.31) N = 96,841	1.31 (0.58, 2.95) N = 159,478

Taylor et al Supplementary Material

	NA	NA	0.38 (0.02, 1.71) N = 53,922	0.81 (0.11, 5.80) N = 67,071	0.56 (0.14, 2.24) N = 120,993
<i>Paternal underweight confounder adjusted</i>	NA	NA	0.36 (0.02, 1.63) N = 54,710	0.82 (0.11, 5.87) N = 68,623	0.54 (0.13, 2.19) N = 123,333
	NA	NA	0.37 (0.02, 1.67) N = 53,922	0.85 (0.12, 6.09) N = 67,071	0.56 (0.14, 2.26) N = 120,993
<i>Paternal underweight confounder and other parent BMI adjusted</i>	NA	NA	0.36 (0.02, 1.65) N = 53,922	0.85 (0.12, 6.08) N = 67,071	0.55 (0.14, 2.24) N = 120,993
	NA	NA	0.36 (0.02, 1.64) N = 53,922	0.85 (0.12, 6.08) N = 67,071	0.55 (0.14, 2.24) N = 120,993
<i>Paternal overweight unadjusted</i>	0.90 (0.53, 1.49) N = 8,076	0.60 (0.18, 1.88) N = 2,725	1.10 (0.95, 1.27) N = 62,637	1.02 (0.88, 1.18) N = 96,841	1.05 (0.95, 1.16) N = 159,478
	0.73 (0.32, 1.54) N = 5,044	0.53 (0.11, 2.17) N = 1,572	1.18 (1.01, 1.38) N = 53,922	1.03 (0.86, 1.23) N = 67,071	1.10 (0.98, 1.23) N = 127,609
<i>Paternal overweight confounder adjusted</i>	1.07 (0.37, 3.20) N = 5,550	0.67 (0.17, 2.39) N = 2,085	1.20 (0.95, 1.53) N = 54,710	1.08 (0.90, 1.28) N = 68,623	1.11 (0.97, 1.28) N = 130,968
	1.11 (0.33, 3.78) N = 5,044	0.66 (0.13, 2.76) N = 1,572	1.22 (0.97, 1.56) N = 53,922	1.05 (0.88, 1.25) N = 67,071	1.10 (0.96, 1.27) N = 127,609
<i>Paternal overweight confounder and other parent BMI adjusted</i>	1.10 (0.33, 3.73) N = 5,044	0.67 (0.13, 2.82) N = 1,572	1.22 (0.96, 1.56) N = 53,922	1.05 (0.88, 1.26) N = 67,071	1.10 (0.96, 1.27) N = 127,609
	1.10 (0.33, 3.73) N = 5,044	0.67 (0.13, 2.82) N = 1,572	1.22 (0.96, 1.56) N = 53,922	1.05 (0.88, 1.26) N = 67,071	1.10 (0.96, 1.27) N = 127,609
<i>Paternal obesity unadjusted</i>	1.33 (0.54, 2.81) N = 8,076	1.65 (0.56, 4.83) N = 2,725	1.31 (1.00, 1.67) N = 62,637	1.00 (0.79, 1.37) N = 96,841	1.15 (0.97, 1.37) N = 159,478
	1.12 (0.26, 3.31) N = 5,044	1.40 (0.34, 5.31) N = 1,572	1.35 (1.01, 1.76) N = 53,922	0.95 (0.71, 1.25) N = 67,071	1.15 (0.95, 1.40) N = 127,609
<i>Paternal obesity confounder adjusted</i>	2.03 (0.19, 18.64) N = 5,550	1.79 (0.50, 6.16) N = 2,085	1.48 (0.89, 2.48) N = 54,710	1.02 (0.76, 1.37) N = 68,623	1.15 (0.90, 1.47) N = 130,968
	2.96 (0.24, 33.50) N = 5,044	1.93 (0.46, 7.70) N = 1,572	1.47 (0.88, 2.49) N = 53,922	1.02 (0.76, 1.37) N = 67,071	1.15 (0.89, 1.48) N = 127,609
<i>Paternal obesity confounder and other parent BMI adjusted</i>	2.99 (0.25, 33.86) N = 5,044	1.96 (0.47, 7.78) N = 1,572	1.46 (0.87, 2.46) N = 53,922	1.03 (0.76, 1.39) N = 67,071	1.16 (0.90, 1.50) N = 127,609
	2.99 (0.25, 33.86) N = 5,044	1.96 (0.47, 7.78) N = 1,572	1.46 (0.87, 2.46) N = 53,922	1.03 (0.76, 1.39) N = 67,071	1.16 (0.90, 1.50) N = 127,609
Covariates used for each study in fully adjusted models (mutually adjusted models the same as fully adjusted but with additional adjustment for the other parent's BMI and parity in paternal models); ABCD: Maternal: offspring sex, age, education, parity, ethnicity, smoking, alcohol. Paternal: offspring sex, age, education, ethnicity. ALSPAC: Maternal: offspring sex, age, education, parity, smoking, alcohol. Paternal: offspring sex, age, education, smoking, alcohol. BASELINE: Maternal: offspring sex, age, education, smoking, alcohol. Paternal: offspring sex, age, smoking. BiB: Maternal: offspring sex, age, education, parity, ethnicity, smoking. Paternal: offspring sex, age, education, ethnicity, smoking. DNBC: Maternal: offspring sex, age, education, parity, smoking, alcohol. Paternal: offspring sex, age, education, smoking. MoBa: Maternal: offspring sex, age, education, parity, smoking, alcohol. Paternal: offspring sex, age, education, smoking, alcohol. NINFEA: Maternal: offspring sex, age, education, parity, smoking, alcohol. Paternal: offspring sex, age, education.					

Table S7. Comparison between maximal numbers (black, top rows) and complete case models (red, bottom rows). Results are odds ratios (95% CIs) of any offspring CHD for smoking during pregnancy.

Model	ABCD	ALSPAC	BiB	DNBC	MoBa	NINFEA	Meta-analysis results
<i>Maternal smoking unadjusted</i>	2.06 (0.77, 4.65) N = 8,117	1.23 (0.78, 1.87) N = 12,716	0.89 (0.53, 1.42) N = 10,887	1.11 (0.98, 1.26) N = 86,740	1.07 (0.86, 1.34) N = 101,042	0.77 (0.18, 3.21) N = 5,801	1.11 (1.00, 1.23) N = 225,303
	2.04 (0.76, 4.62) N = 7,824	1.40 (0.71, 2.56) N = 7,626	1.62 (0.52, 4.20) N = 2,624	1.10 (0.96, 1.26) N = 78,229	1.03 (0.78, 1.37) N = 77,266	0.79 (0.19, 3.29) N = 5,527	1.11 (0.99, 1.25) N = 179,096
<i>Maternal smoking confounder adjusted</i>	2.02 (0.73, (4.77) N = 7,824	1.22 (0.69, 2.06) N = 10,217	0.93 (0.50, 1.60) N = 9,646	1.05 (0.91, 1.20) N = 80,571	1.02 (0.77, 1.36) N = 77,311	0.92 (0.22, 3.96) N = 5,527	1.06 (0.94, 1.18) N = 191,096
	2.02 (0.73, (4.77) N = 7,824	1.31 (0.65, 2.46) N = 7,626	2.09 (0.64, 5.84) N = 2,624	1.07 (0.93, 1.23) N = 78,229	1.02 (0.77, 1.37) N = 77,266	0.92 (0.22, 3.96) N = 5,527	1.09 (0.97, 1.23) N = 179,096
<i>Maternal smoking confounder and other parent smoking adjusted</i>	-	1.27 (0.61, 2.50) N = 7,626	1.77 (0.51, 5.36) N = 2,624	1.11 (0.96, 1.28) N = 79,000	1.05 (0.78, 1.41) N = 77,266	-	1.11 (0.97, 1.25) N = 166,516
	-	1.27 (0.61, 2.50) N = 7,626	1.77 (0.51, 5.36) N = 2,624	1.13 (0.98, 1.30) N = 78,229	1.05 (0.78, 1.41) N = 77,266	-	1.12 (0.99, 1.28) N = 165,745
<i>Paternal smoking unadjusted</i>	-	1.29 (0.79, 2.10) N = 9,134	1.20 (0.50, 2.66) N = 3,187	0.95 (0.84, 1.08) N = 84,926	0.96 (0.82, 1.11) N = 101,804	-	0.97 (0.88, 1.06) N = 198,421
	-	1.28 (0.68, 2.35) N = 6,182	1.46 (0.54, 3.73) N = 2,373	0.96 (0.84, 1.10) N = 77,477	1.00 (0.83, 1.20) N = 70,018	-	0.99 (0.89, 1.10) N = 156,050
<i>Paternal smoking confounder adjusted</i>	-	1.17 (0.61, 2.19) N = 6,308	1.43 (0.51, 3.76) N = 2,424	0.95 (0.83, 1.08) N = 77,526	1.05 (0.87, 1.26) N = 70,766	-	0.99 (0.89, 1.10) N = 157,024
	-	1.23 (0.64, 2.30) N = 6,182	1.51 (0.53, 4.06) N = 2,373	0.95 (0.83, 1.08) N = 77,477	1.05 (0.87, 1.27) N = 70,018	-	0.99 (0.89, 1.10) N = 156,050
<i>Paternal smoking confounder and other parent BMI adjusted</i>	-	1.14 (0.56, 2.23) N = 6,182	1.18 (0.38, 3.41) N = 2,373	0.90 (0.79, 1.04) N = 77,499	1.04 (0.85, 1.26) N = 70,018	-	0.96 (0.85, 1.07) N = 156,072
	-	1.14 (0.64, 2.30) N = 6,182	1.18 (0.38, 3.41) N = 2,373	0.91 (0.79, 1.04) N = 77,477	1.04 (0.85, 1.26) N = 70,018	-	0.96 (0.86, 1.07) N = 156,050

Covariates used for each study in fully adjusted models (mutually adjusted models the same as fully adjusted but with additional adjustment for the other parent's smoking);
 ABCD: Maternal: offspring sex, age, education, parity, ethnicity, alcohol.
 ALSPAC: Maternal: offspring sex, age, education, parity, alcohol. Paternal: offspring sex, age, education, alcohol.
 BiB: Maternal: offspring sex, age, education, parity, ethnicity. Paternal: offspring sex, age, education, ethnicity.
 DNBC: Maternal: offspring sex, age, education, parity, alcohol. Paternal: offspring sex, age, education.
 MoBa: Maternal: offspring sex, age, education, parity, alcohol. Paternal: offspring sex, age, education, alcohol.
 NINFEA: Maternal: offspring sex, age, education, parity, alcohol.

Table S8. Comparison between **maximal numbers (black, top rows)** and **complete case models (red, bottom rows)**. Results are odds ratios (95% CIs) of any offspring CHD for alcohol intake during pregnancy in comparison to non-drinkers.

Model	ABCD	ALSPAC	DNBC	MoBa	NINFEA	Meta-analysis results
Maternal alcohol (yes/no) unadjusted	1.38 (0.61, 2.85) N = 8,125	1.20 (0.81, 1.80) N = 12,622	1.00 (0.89, 1.12) N = 86,708	1.04 (0.88, 1.23) N = 82,358	1.20 (0.57, 2.51) N = 5,843	1.03 (0.94, 1.12) N = 195,656
	1.36 (0.60, 2.81) N = 7,824	1.18 (0.56, 2.55) N = 4,585	1.00 (0.89, 1.13) N = 79,648	1.06 (0.86, 1.31) N = 51,006	1.19 (0.57, 2.49) N = 5,527	1.03 (0.93, 1.14) N = 148,590
Maternal alcohol (yes/no) confounder adjusted	1.17 (0.50, 2.56) N = 7,824	1.24 (0.78, 2.01) N = 10,217	1.01 (0.90, 1.14) N = 80,571	1.03 (0.86, 1.23) N = 77,311	1.18 (0.56, 2.49) N = 5,527	1.03 (0.94, 1.13) N = 181,450
	1.17 (0.50, 2.56) N = 7,824	1.20 (0.56, 2.63) N = 4,585	1.01 (0.89, 1.14) N = 79,648	1.06 (0.85, 1.31) N = 51,066	1.18 (0.56, 2.49) N = 5,527	1.03 (0.93, 1.14) N = 148,590
Maternal light drinking unadjusted	-	0.93 (0.52, 1.67) N = 6,501	0.92 (0.82, 1.03) N = 88,349	1.10 (0.88, 1.36) N = 84,436	-	0.96 (0.87, 1.06) N = 179,286
	-	1.27 (0.58, 2.93) N = 4,585	0.93 (0.82, 1.05) N = 79,648	1.24 (0.94, 1.63) N = 51,006	-	0.98 (0.88, 1.09) N = 135,239
Maternal light drinking confounder adjusted	-	0.92 (0.48, 1.78) N = 5,797	0.95 (0.85, 1.08) N = 80,214	1.13 (0.90, 1.41) N = 79,695	-	0.99 (0.89, 1.10) N = 165,706
	-	1.35 (0.61, 3.14) N = 4,585	0.94 (0.83, 1.06) N = 79,648	1.22 (0.92, 1.61) N = 51,006	-	0.99 (0.88, 1.10) N = 135,239
Maternal light drinking confounder and other parent alcohol adjusted	-	1.40 (0.62, 3.27) N = 4,585	-	1.13 (0.87, 1.47) N = 59,571	-	1.15 (0.90, 1.48) N = 64,156
	-	1.40 (0.62, 3.27) N = 4,585	-	1.19 (0.90, 1.57) N = 51,006	-	1.21 (0.93, 1.57) N = 55,591
Maternal mod/heavy drinking unadjusted	-	0.67 (0.22, 1.65) N = 6,501	1.14 (0.87, 1.48) N = 88,349	1.85 (0.92, 3.73) N = 84,436	-	1.17 (0.92, 1.49) N = 179,286
	-	0.92 (0.21, 3.01) N = 4,585	1.19 (0.89, 1.56) N = 79,648	1.77 (0.66, 4.78) N = 51,006	-	1.21 (0.93, 1.58) N = 135,239
Maternal mod/heavy drinking confounder adjusted	-	0.64 (0.18, 1.75) N = 5,797	1.21 (0.90, 1.58) N = 80,214	1.47 (0.65, 3.32) N = 79,695	-	1.19 (0.92, 1.54) N = 165,706
	-	0.89 (0.20, 2.98) N = 4,585	1.19 (0.89, 1.57) N = 79,648	1.73 (0.64, 4.69) N = 51,006	-	1.21 (0.93, 1.58) N = 135,239
Maternal mod/heavy drinking confounder and other parent alcohol adjusted	-	0.94 (2.06, 3.19) N = 4,585	-	1.31 (0.48, 3.56) N = 59,571	-	1.16 (0.52, 2.58) N = 64,156
	-	0.94 (2.06, 3.19) N = 4,585	-	1.57 (0.58, 4.27) N = 51,006	-	1.30 (0.59, 2.89) N = 55,591
Paternal light drinking unadjusted	-	0.90 (0.36, 3.02) N = 8,205	-	0.90 (0.61, 1.32) N = 72,422	-	0.90 (0.63, 1.29) N = 80,627
	-	1.90 (0.39, 34.09) N = 5,228	-	1.01 (0.62, 1.65) N = 58,847	-	1.05 (0.65, 1.68) N = 64,075
Paternal light drinking confounder adjusted	-	2.11 (0.44, 37.99) N = 5,346	-	0.86 (0.58, 1.28) N = 70,766	-	0.89 (0.60, 1.31) N = 76,112
	-	2.04 (0.42, 36.80) N = 5,228	-	0.97 (0.60, 1.58) N = 58,847	-	1.01 (0.63, 1.63) N = 64,075

Taylor et al Supplementary Material

<i>Paternal light drinking confounder and other parent alcohol adjusted</i>		1.77 (0.36, 32.20) N = 5,316	-	0.97 (0.63, 1.62) N = 58,847		1.01 (0.63, 1.62) N = 64,163
	-	1.74 (0.35, 31.60) N = 5,228	-	0.97 (0.60, 1.59) N = 58,847	-	1.01 (0.62, 1.62) N = 64,075
<i>Paternal mod/heavy drinking unadjusted</i>	-	0.86 (0.34, 2.93) N = 8,205	-	1.11 (0.73, 1.70) N = 72,422	-	1.08 (0.73, 1.59) N = 80,627
	-	1.83 (0.37, 33.05) N = 5,228	-	1.28 (0.76, 2.17) N = 58,847	-	1.31 (0.79, 2.18) N = 64,075
<i>Paternal mod/heavy drinking confounder adjusted</i>	-	2.00 (0.40, 36.05) N = 5,346	-	1.07 (0.69, 1.66) N = 70,766	-	1.10 (0.72, 1.69) N = 76,112
	-	1.94 (0.39, 35.05) N = 5,228	-	1.20 (0.71, 2.04) N = 58,847	-	1.24 (0.74, 2.07) N = 64,075
<i>Paternal mod/heavy drinking confounder and other parent alcohol adjusted</i>	-	1.72 (0.34, 31.20) N = 5,316	-	1.21 (0.71, 2.05) N = 58,847	-	1.23 (0.74, 2.06) N = 64,163
	-	1.70 (0.34, 30.83) N = 5,228	-	1.21 (0.71, 2.05) N = 58,847	-	1.23 (0.74, 2.06) N = 64,075
Covariates used for each study in fully adjusted models (mutually adjusted models the same as fully adjusted but with additional adjustment for the other parent's alcohol intake); ABCD: Maternal: offspring sex, age, education, parity, ethnicity, smoking. ALSPAC: Maternal: offspring sex, age, education, parity, smoking. Paternal: offspring sex, age, education, smoking. DNBC: Maternal: offspring sex, age, education, parity, smoking. MoBa: Maternal: offspring sex, age, education, parity, smoking. Paternal: offspring sex, age, education, smoking. NINFEA: Maternal: offspring sex, age, education, parity, ethnicity, smoking.						

Supplementary Tables S9-S11**Table S9.** Meta-analysis results from 4 cohorts (ALSPAC, BiB, DNBC, MoBa) for associations between BMI categories and CHDs with and without removing chromosomal/genetic defects from the study population. Results reported as odds ratios for CHD for parental underweight, overweight or obesity in comparison to parental normal weight.

<i>Model</i>	<i>Main analysis Outcome = CHD</i>	<i>Additional analysis Outcome = CHD with chromo/gen defects removed from study population</i>
Confounder adjusted	M-Underweight: 1.20 (0.96, 1.50) P-Underweight: 0.54 (0.13, 2.19)	M- Underweight: 1.16 (0.90, 1.48) P-Underweight: 0.67 (0.16, 2.70)
	M-Overweight: 1.17 (1.04, 1.31) P-Overweight: 1.11 (0.97, 1.28)	M-Overweight: 1.20 (1.06, 1.35) P-Overweight: 1.09 (0.94, 1.27)
	M-Obesity: 1.19 (1.02, 1.40) P-Obesity: 1.15 (0.90, 1.47)	M-Obesity: 1.21 (1.02, 1.44) P-Obesity: 1.19 (0.91, 1.58)
Confounder and other parent BMI adjusted	M-Underweight: 1.21 (0.93, 1.58) P-Underweight: 0.55 (0.14, 2.24)	M-Underweight: 1.22 (0.90, 1.63) P-Underweight: 0.67 (0.17, 2.72)
	M-Overweight: 1.15 (1.01, 1.31) P-Overweight: 1.10 (0.96, 1.27)	M-Overweight: 1.20 (1.04, 1.38) P-Overweight: 1.08 (0.93, 1.27)
	M-Obesity: 1.12 (0.93, 1.36) P-Obesity: 1.16 (0.90, 1.50)	M-Obesity: 1.15 (0.93, 1.42) P-Obesity: 1.20 (0.90, 1.59)
M = maternal P = paternal ^ICD codes used to remove these cases from the population can be found in Table S3.		

Table S10. Meta-analysis results from 3 cohorts (ALSPAC, BiB and DNBC) for associations between BMI categories and CHD severity with and without removing chromosomal/genetic defects from the study population. Results reported as odds ratios for CHD for parental underweight, overweight or obesity in comparison to parental normal weight.

<i>Model</i>	<i>Outcome = Non-severe CHD</i>	<i>Outcome = Non-severe CHD (excluding chromo/gen defects)^</i>	<i>Outcome = Severe CHD</i>	<i>Outcome = Severe CHD (excluding chromo/gen defects)^</i>
Confounder adjusted	M-Underweight: 1.24 (0.91, 1.68) P-Underweight: 0.49 (0.07, 3.56)	M- Underweight: 1.32 (0.95, 1.83) P-Underweight: 0.55 (0.07, 4.10)	M-Underweight: 1.25 (0.79, 1.97) P-Underweight: *	M- Underweight: 1.27 (0.75, 2.16) P-Underweight: *
	M-Overweight: 1.24 (1.05, 1.47) P-Overweight: 1.16 (0.89, 1.51)	M-Overweight: 1.27 (1.06, 1.52) P-Overweight: 1.09 (0.82, 1.45)	M-Overweight: 1.19 (0.93, 1.53) P-Overweight: 1.19 (0.77, 1.84)	M-Overweight: 1.29 (0.98, 1.70) P-Overweight: 1.09 (0.66, 1.79)
	M-Obesity: 1.36 (1.08, 1.71) P-Obesity: 1.49 (0.86, 2.59)	M-Obesity: 1.36 (1.06, 1.74) P-Obesity: 1.51 (0.83, 2.74)	M-Obesity: 1.07 (0.73, 1.56) P-Obesity: 1.65 (0.71, 3.87)	M-Obesity: 1.12 (0.74, 1.71) P-Obesity: 1.58 (0.60, 4.19)
Confounder and other parent BMI adjusted	M-Underweight: 1.34 (0.92, 1.93) P-Underweight: 0.48 (0.07, 3.54)	M-Underweight: 1.41 (0.95, 2.09) P-Underweight: 0.56 (0.08, 4.10)	M-Underweight: 1.16 (0.61, 2.22) P-Underweight: *	M-Underweight: 1.42 (0.69, 2.94) P-Underweight: *
	M-Overweight: 1.29 (1.05, 1.58) P-Overweight: 1.22 (0.93, 1.59)	M-Overweight: 1.33 (1.07, 1.66) P-Overweight: 1.13 (0.84, 1.51)	M-Overweight: 1.11 (0.78, 1.57) P-Overweight: 1.13 (0.72, 1.77)	M-Overweight: 1.39 (0.95, 2.02) P-Overweight: 0.97 (0.58, 1.62)
	M-Obesity: 1.14 (0.84, 1.55) P-Obesity: 1.61 (0.91, 2.84)	M-Obesity: 1.19 (0.85, 1.66) P-Obesity: 1.57 (0.84, 2.91)	M-Obesity: 1.17 (0.71, 1.93) P-Obesity: 1.36 (0.56, 3.32)	M-Obesity: 1.31 (0.74, 2.32) P-Obesity: 1.17 (0.42, 3.28)
M = maternal P = paternal ^ICD codes used to remove these cases from the population can be found in Table S3. * = not enough data to compute results.				

Table S11. Meta-analysis results for associations between alcohol intake and CHDs after removing chromosomal/genetic defects from the study population. Results reported as odds ratios and 95% confidence intervals for CHD in comparison to non-drinkers.

<i>Model</i>	<i>Main analysis Outcome = CHD</i>	<i>Additional analysis Outcome = CHD with chromo/gen removed from study population</i>
Confounder adjusted	M – y/n: 1.03 (0.93, 1.13)	M – y/n: 1.04 (0.94, 1.15)
	M – light: 0.99 (0.89, 1.10) P – light: 0.89 (0.60, 1.31)	M – light: 0.95 (0.85, 1.07) P – light: 1.13 (0.69, 1.87)
	M – mod/heavy: 1.19 (0.92, 1.54) P – mod/heavy: 1.10 (0.72, 1.69)	M – mod/heavy: 1.24 (0.94, 1.63) P – mod/heavy: 1.36 (0.79, 2.34)
Confounder and other parent BMI adjusted	M – light: 1.15 (0.90, 1.48) P – light: 1.01 (0.63, 1.62)	M – light: 1.17 (0.88, 1.55) P – light: 1.21 (0.68, 2.16)
	M – mod/heavy: 1.16 (0.52, 2.58) P – mod/heavy: 1.23 (0.74, 2.06)	M – mod/heavy: 1.20 (0.52, 3.17) P – mod/heavy: 1.52 (0.82, 2.80)
<p>M = maternal P = paternal y/n = alcohol as a binary variable, yes or no. Estimates from yes/no analyses derived from 5 cohorts (ABCD, ALSPAC, DNBC, MoBa, NINFEA). Estimates from maternal light and mod/heavy drinking analyses derived from ALSPAC, DNBC and MoBa in fully adjusted results, but only ALSPAC and MoBa in paternal and mutually adjusted results.</p>		

Supplementary Results – Figures S1-S32

Participant flow charts for each cohort

LifeCycle CHD analysis in the ABCD cohort




Figure S1. Study flow chart illustrating participant selection in the ABCD cohort.

LifeCycle CHD analysis in the ALSPAC cohort




Figure S2. Study flow chart illustrating participant selection in the ALSPAC cohort.

LifeCycle CHD analysis in the Baseline cohort




Figure S3. Study flow chart illustrating participant selection in the BASELINE cohort. We included 1436 participants in our study (Stream 1). Adapted from: <https://doi.org/10.1093/ije/dyu157>

LifeCycle CHD analysis in the BiB cohort




Figure S4. Study flow chart illustrating participant selection in the BiB cohort.

LifeCycle CHD analysis in the DNBC




Figure S5. Study flow chart illustrating participant selection in the DNBC cohort.

LifeCycle CHD analysis in MoBa




Figure S6. Study flow chart illustrating participant selection in the MoBa cohort. MBR = Medical birth registry.

LifeCycle CHD analysis in the NINFEA cohort




Figure S7. Study flow chart illustrating participant selection in the NINFEA cohort.

BMI analyses – Supplementary Figures

- **Figure S8.** Main analysis associations between parental BMI as a continuous measurement in kg/m² (maternal top, paternal bottom) and offspring congenital heart disease.
- **Figure S9.** Confounder adjusted associations between maternal BMI split into fifths and offspring CHDs in the DNBC (A) and MoBa (B).
- **Figure S10.** Confounder adjusted associations between paternal BMI split into fifths and offspring CHDs in the DNBC (A) and MoBa (B).
- **Figure S11.** Linear associations between parental BMI and offspring CHD severity.
- **Figure S12.** Linear associations between maternal BMI and offspring congenital heart disease with additional adjustment of folic acid supplementation.
- **Figure S13.** Linear associations between parental BMI and offspring congenital heart disease with chromosomal/genetic defects removed from the study population.
- **Figure S14.** Linear associations between parental BMI and offspring CHD severity with cases of chromosomal/genetic defects removed from the study population.
- **Figure S15.** Meta-analysis results for unadjusted BMI categories using World Health Organization cut-offs with normal BMI as the reference.
- **Figure S16.** Meta-analysis results for confounder adjusted BMI categories using World Health Organization cut-offs with normal BMI as the reference.
- **Figure S17.** Meta-analysis results for confounder and other parent BMI adjusted BMI categories using World Health Organization cut-offs with normal BMI as the reference.
- **Figure S18.** Meta-analysis results for unadjusted BMI categories using World Health Organization cut-offs with normal BMI as the reference.
- **Figure S19.** Meta-analysis results for confounder adjusted BMI categories using World Health Organization cut-offs with normal BMI as the reference.
- **Figure S20.** Meta-analysis results for confounder and other parent BMI adjusted BMI categories using World Health Organization cut-offs with normal BMI as the reference.




Figure S8. Main analysis associations between parental BMI as a continuous measurement in kg/m² (maternal top, paternal bottom) and offspring congenital heart disease. Panel A results are unadjusted, panel B results are fully adjusted for all confounders and panel C results are adjusted for all confounders as well as other parent’s BMI. Confounders: **ABCD**: parental age, education, parity, ethnicity, smoking, alcohol, offspring sex; **ALSPAC**: parental age, education, parity, smoking, alcohol, offspring sex; **BASELINE**: parental age, education, smoking, alcohol, offspring sex; **BiB**: parental age, education, parity, ethnicity, smoking, offspring sex; **DNBC**: parental age, education, parity, smoking, alcohol, offspring sex; **MoBa**: parental age, education, parity, smoking, alcohol, offspring sex; **NINFEA**: parental age, education, parity, smoking, alcohol, offspring sex.

Figures S9 and S10 show the odds ratios of CHD by fifths of the BMI distribution for mothers and fathers respectively in DNBC and MoBa. Whilst there was statistical evidence for a linear trend in DNBC mothers (p -value for per fifth increase = 0.05) the graph shows this was driven by increased risk only in the highest fifth, with the 2nd, 3rd and 4th fifth (compared to the first) consistent with the null. In MoBa mothers there was no clear pattern with some evidence that the 4th compared to the 1st fifth was associated with lower risk with the 3 other categories being consistent with the null (p -value for linear trend in MoBa = 0.22). Whilst the p -values for the likelihood ratio comparing the linear model with the category model (0.03 and 0.09, for DNBC and MoBa mothers, respectively) provide statistical support for the category model in each, this is based on just one of the fifths. Results for the fathers are broadly consistent with those for the mothers, and overall, these results are consistent with no association of maternal or paternal mean BMI with offspring CHD risk.




Figure S9. Confounder adjusted associations between maternal BMI split into fifths and offspring CHDs in the DNBC (A) and MoBa (B). Results are odds ratios and 95% CIs for maternal BMI quintile and offspring CHD in comparison to BMI quintile 1.






Figure S10. Confounder adjusted associations between paternal BMI split into fifths and offspring CHDs in the DNBC (A) and MoBa (B). Results are odds ratios and 95% CIs for paternal BMI quintile and offspring CHD in comparison to BMI quintile 1.

A: Non-severe CHD confounder adjusted

B: Severe CHD confounder adjusted



C: Non-severe CHD confounder and other parent BMI adjusted

D: Severe CHD confounder and other parent BMI adjusted





Figure S11. Linear associations (top (A&B): confounder adjusted, bottom (C&D): confounder and other parent BMI adjusted) between parental BMI and offspring non-severe congenital heart disease (left) and severe congenital heart disease (right). Definitions for CHD subtypes can be found in Table S2.




Figure S12. Linear associations between maternal BMI and offspring congenital heart disease. Results are fully adjusted for all confounders (top) and all confounders plus additional adjustment for folic acid supplementation during weeks 0-12 of pregnancy (bottom).

Mean BMI results with chromosomal/genetic defects removed from study population

A: Additional analysis confounder adjusted

B: Additional analysis confounder and other parent BMI adjusted






Figure S13. Additional analysis: linear associations between parental BMI and offspring congenital heart disease with chromosomal/genetic defects removed from the study population. **A** is adjusted for all confounders, and **B** is adjusted for all confounders and the other parent's BMI. The rationale here is to see if estimates differ when we remove offspring from the population with an anomaly associated with a pre-specified cause such as a genetic, chromosomal or teratogenic aberration. ICD codes used to remove these cases from the population can be found in Table S3. For comparison the pooled associations from main analyses (without removal of genetic/chromo disorders) were: 1.01 (1.00, 1.02) & 1.01 (0.99, 1.02) for maternal (top graphs, left and right respectively) and 1.01 (1.00, 1.03) & 1.01 (0.99, 1.03) for paternal (bottom graphs left and right respectively).

Mean BMI CHD severity results with chromosomal/genetic defects removed

A: Non-severe CHD confounder adjusted

B: Severe CHD confounder adjusted



C: Non-severe CHD confounder and other parent BMI adjusted

D: Severe CHD confounder and other parent BMI adjusted





Figure S14. Linear associations (top (A&B): confounder adjusted, bottom (C&D): confounder and other parent BMI adjusted) between parental BMI and offspring non-severe congenital heart disease (left) and severe congenital heart disease (right) with cases of chromosomal/genetic defects removed from the study population. Definitions for CHD subtypes can be found in Table S2.


BMI analyses using World Health Organization (WHO) categories

In this section, we present meta-analysis results from the WHO BMI analyses. All Odds ratios should be interpreted as an increase/decrease odds of CHD for a maternal/paternal BMI category in comparison to normal weight. The BMI categories are: underweight (BMI <18.5 kg/m²), normal weight (BMI 18.5 to <25 kg/m², reference range), overweight (BMI 25 to <30 kg/m²) and obesity (BMI ≥30 kg/m²).


We present fully adjusted (adjusted for all confounders) and mutually adjusted (adjusted for all confounders plus other parents' exposure) models. ALSPAC, BiB, DNBC and MoBa contributed to these analyses. Covariates adjusted for by each study are:

- ALSPAC - parental age, education, parity, smoking, alcohol, offspring sex
- BiB - parental age, education, parity, ethnicity, smoking, offspring sex
- DNBC – parental age, education, parity, smoking, alcohol, offspring sex
- MoBa - parental age, education, parity, smoking, alcohol, offspring sex

A: Underweight



B: Overweight



C: Obesity






Figure S15. Meta-analysis results for unadjusted BMI categories using World Health Organization cut-offs with normal BMI as the reference. Outcome = any CHD in the offspring.

A: Underweight



B: Overweight



C: Obesity






Figure S16. Meta-analysis results for confounder adjusted BMI categories using World Health Organization cut-offs with normal BMI as the reference. Outcome = any CHD in the offspring.

A: Underweight



B: Overweight



C: Obesity






Figure S17. Meta-analysis results for confounder and other parent BMI adjusted BMI categories using World Health Organization cut-offs with normal BMI as the reference. Outcome = any CHD in the offspring.


A: Underweight and non-severe CHDs




Underweight and severe CHDs




B: Overweight and non-severe CHDs



Overweight and severe CHDs



C: Obesity and non-severe CHDs



Obesity and severe CHDs






Figure S18. Meta-analysis results for unadjusted BMI categories using World Health Organization cut-offs with normal BMI as the reference. Outcome = non-severe CHDs (left) and severe CHDs (right). Ns represent total numbers included in the non-severe/severe analyses presented.


A: Underweight and non-severe CHDs




Underweight and severe CHDs




B: Overweight and non-severe CHDs



Overweight and severe CHDs



C: Obesity and non-severe CHDs



Obesity and severe CHDs






Figure S19. Meta-analysis results for confounder adjusted BMI categories using World Health Organization cut-offs with normal BMI as the reference. Outcome = non-severe CHDs (left) and severe CHDs (right). Ns represent total numbers included in the non-severe/severe analyses presented.



A: Underweight and non-severe CHDs

Underweight and severe CHDs



B: Overweight and non-severe CHDs

Overweight and severe CHDs



C: Obesity and non-severe CHDs

Obesity and severe CHDs






Figure S20. Meta-analysis results for confounder and other parent BMI adjusted BMI categories using World Health Organization cut-offs with normal BMI as the reference. Outcome = non-severe CHDs (left) and severe CHDs (right). Ns represent total numbers included in the non-severe/severe analyses presented.


Smoking analyses – Supplementary Figures

- **Figure S21.** Main analysis associations between parental smoking (maternal top, paternal bottom) and offspring congenital heart disease.
- **Figure S22.** Showing the smoking results in those cohorts that had confirmed data on maternal first trimester smoking.
- **Figure S23.** Unadjusted and confounder adjusted results for the smoking and CHD severity analyses presented in the main manuscript Figure 2.
- **Figure S24.** Associations between parental smoking heaviness and offspring congenital heart disease.
- **Figure S25.** Associations between parental smoking and offspring congenital heart disease with chromosomal/genetic defects removed from the study population.
- **Figure S26.** Smoking and CHD severity results with chromosomal/genetic defects removed from the study population.
- **Figure S27.** Associations between maternal smoking and offspring congenital heart disease with additional adjustment for folic acid supplementation.

A: Unadjusted



B: Confounder adjusted



C: Confounder and other parent smoking adjusted






Figure S21. Main analysis associations between parental smoking (maternal top, paternal bottom) and offspring congenital heart disease. Panel A results are unadjusted, B results are fully adjusted for all confounders and C results are adjusted for all confounders as well as other parent’s smoking. Confounders: **ABCD**: parental age, education, parity, ethnicity, alcohol, offspring sex; **ALSPAC**: parental age, education, parity, alcohol, offspring sex; **BiB**: parental age, education, parity, ethnicity, offspring sex; **DNBC**: parental age, education, parity, alcohol, offspring sex; **MoBa**: parental age, education, parity, alcohol, offspring sex; **NINFEA**: parental age, education, parity, alcohol, offspring sex.

Results including cohorts with confirmed first trimester maternal smoking (yes/no)

A: Unadjusted



B: Confounder adjusted



C: Confounder and other parent smoking adjusted





Figure S22. Showing the smoking results in those cohorts that had confirmed data on maternal first trimester smoking. Panel A results are unadjusted, B results are fully adjusted for all confounders and C results are adjusted for all confounders as well as other parent’s smoking. Confounders: **ALSPAC:** parental age, education, parity, alcohol, offspring sex; **DNBC:** parental age, education, parity, alcohol, offspring sex; **NINFEA:** parental age, education, parity, alcohol, offspring sex.

Taylor et al Supplementary Material

A: Non-severe CHD unadjusted

B: Severe CHD unadjusted



C: Non-severe CHD confounder adjusted

D: Severe CHD confounder adjusted






Figure S23. Unadjusted (A & B) and confounder adjusted (C & D) results for the smoking and CHD severity analyses presented in the main manuscript Figure 2.

Taylor et al Supplementary Material


A: (Maternal) Unadjusted




B: (Maternal) Confounder adjusted




C: (Maternal) Confounder and other parent smoking adjusted



D: (Paternal) Unadjusted



E: (Paternal) Confounder adjusted



F: (Paternal) Confounder and other parent smoking adjusted






Figure S24. Associations between parental smoking heaviness (**top (A, B & C): maternal, bottom (D, E & F): paternal**) and offspring congenital heart disease. Results are unadjusted (left), adjusted for all confounders (middle) as well as all confounders and other parents smoking (right). Smoking categorized as none (non-smoker), light (< 10 cigarettes smoked per day during pregnancy) and heavy (≥ 10 cigarettes per day). Results presented as odds ratios and 95% confidence intervals for offspring CHD in comparison to non-smokers.

Adjusted smoking results with chromosomal/genetic defects removed from study population


A: Yes/no confounder adjusted



B: Yes/no confounder and other parent smk adjusted



C: Heaviness confounder adjusted



D: Heaviness confounder and other parent smk adjusted






Figure S25. Additional analysis: Associations between parental smoking and offspring congenital heart disease with chromosomal/genetic defects removed from the study population. The plots on the top half (A & B) are for smoking yes/no analyses and the plots on the bottom half (C & D) are for smoking heaviness analyses. The rationale here is to see if estimates differ when we remove offspring from the population with an anomaly associated with a pre-specified cause such as a genetic, chromosomal or teratogenic aberration. ICD codes used to remove these cases from the population can be found in Table S3.

Smoking and CHD severity results with chromosomal/genetic defects removed


A: Non-severe CHD unadjusted




B: Severe CHD unadjusted




C: Non-severe CHD confounder adjusted



D: Severe CHD confounder adjusted



E: Non-severe CHD confounder and other parent smk adjusted



F: Severe CHD confounder and other parent smk adjusted






Figure S26. Smoking and CHD severity results with chromosomal/genetic defects removed from the study population.

A: Smk yes/no FA supp



B: Smk categories: confounder adjusted



C: Smk categories & additional adjustment for FA supp




Figure S27. Associations between maternal smoking and offspring congenital heart disease. Results are fully adjusted for all confounders and all confounders plus additional adjustment for folic acid supplementation during weeks 0-12 of pregnancy. Panel A shows the results for the yes/no smoking analyses and panels B and C show results for the smoking heaviness analyses.

Alcohol analyses - Supplementary Figures

- **Figure S28.** Associations between maternal alcohol consumption in the first trimester and offspring congenital heart disease.
- **Figure S29.** Confounder adjusted associations between maternal alcohol consumption during the first trimester and offspring non-severe congenital heart disease and severe congenital heart disease.
- **Figure S30.** Associations between parental alcohol intake and offspring congenital heart disease (all models).
- **Figure S31.** Confounder adjusted associations between maternal alcohol consumption during the first trimester and offspring non-severe congenital heart disease and severe congenital heart disease with chromosomal/genetic defects removed from the study population.
- **Figure S32.** Associations between maternal drinking during the first trimester and offspring congenital heart disease with additional adjustment for folic acid supplementation.



A: Unadjusted**B: Confounder adjusted**

Figure S28. Associations between maternal alcohol consumption in the first trimester and offspring congenital heart disease. *ABCD did not have trimester-specific data, therefore analyses presented for ABCD are any alcohol consumption during pregnancy. Results are adjusted for all confounders. Confounders: **ABCD**: parental age, education, parity, ethnicity, smoking, offspring sex; **ALSPAC**: parental age, education, parity, smoking, offspring sex; **DNBC**: parental age, education, parity, smoking, offspring sex; **MoBa**: parental age, education, parity, smoking, offspring sex; **NINFEA**: parental age, education, parity, smoking, offspring sex.




A: Non-severe CHD**B: Severe CHD**


Figure S29. Confounder adjusted associations between maternal alcohol consumption during the first trimester and offspring non-severe congenital heart disease **(A)** and severe congenital heart disease **(B)**. Definitions for CHD subtypes can be found in Table S2.

Taylor et al Supplementary Material


A: (Maternal) Unadjusted




B: (Maternal) Confounder adjusted




C: (Maternal) Confounder and other parent smoking adjusted



D: (Paternal) Unadjusted



E: (Paternal) Confounder adjusted



F: (Paternal) Confounder and other parent smoking adjusted




Figure S30. Associations (top (A, B & C): maternal, bottom (D, E & F): paternal) between parental alcohol intake and offspring congenital heart disease. Results are unadjusted (left), adjusted for all confounders (middle) as well as all confounders and other parents smoking (right). Maternal alcohol intake categorized as none (non-drinker), light (< 3 units per week during pregnancy) and moderate/heavy (≥ 3 units per week during pregnancy). Paternal alcohol intake categorized as none (non-drinker), light (< 7 units per week during pregnancy) and moderate/heavy (≥ 7 units per week during pregnancy). Results presented as odds ratios and 95% confidence intervals for offspring CHD in comparison to non-drinkers.



A: Non-severe CHD**B: Severe CHD**

Figure S31. Confounder adjusted associations between maternal alcohol consumption during the first trimester and offspring non-severe congenital heart disease **(A)** and severe congenital heart disease **(B)** with chromosomal/genetic defects removed from the study population. Definitions for CHD subtypes can be found in Table S2.




Figure S32. Associations between maternal drinking during the first trimester and offspring congenital heart disease. Results are adjusted for all confounders (top) and all confounders plus additional adjustment for folic acid supplementation during weeks 0-12 of pregnancy (bottom). Results are for first trimester drinking or any drinking during pregnancy where trimester data were not available (denoted by *).

Full reference list can be found within the main manuscript.