

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Plasticity, exudation and microbiome-association of the root system of Pellitory-of-the-wall plants grown in environments impaired in iron availability**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1808979> since 2021-10-04T14:51:51Z

*Published version:*

DOI:10.1016/j.plaphy.2021.09.040

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Original Article

# **Plasticity, exudation and microbiome-association of the root system of Pellitory-of-the-wall plants grown in environments impaired in iron availability**

Liliana Tato<sup>1</sup>, Vincenzo Lattanzio<sup>2</sup>, Enrico Ercole<sup>3</sup>, Marta Dell'Orto<sup>1</sup>, Agostino Sorgonà<sup>5</sup>, Vito Linsalata<sup>4</sup>, Alessandra Salvioli<sup>3</sup>, Mara Novero<sup>3</sup>, Stefania Astolfi<sup>6</sup>, Maria Rosa Abenavoli<sup>5</sup>, Irene Murgia<sup>7</sup>, Graziano Zocchi<sup>1</sup> and Gianpiero Vigani<sup>3</sup>

<sup>1</sup>Dipartimento di Scienze Agrarie e Ambientali, Produzioni, Territorio, Agroenergia, Università degli Studi di Milano Italy

<sup>2</sup>Dipartimento di Scienze Agrarie, degli Alimenti e dell'Ambiente, Università degli Studi di Foggia, Italy

<sup>3</sup>Dipartimento di Scienze della Vita e Biologia dei Sistemi, Università degli Studi di Torino, Italy

<sup>4</sup>C.N.R. Istituto di Scienze delle Produzioni Alimentari, Bari, Italy

<sup>5</sup>Dipartimento Agraria, Università "Mediterranea" di Reggio Calabria Feo di Vito, 89124 Reggio Calabria

<sup>6</sup>DAFNE, Università della Tuscia, Viterbo, Italy

<sup>7</sup>Dipartimento di Bioscienze, Università degli Studi di Milano, Milano, Italy

*corresponding author:*

Dr Gianpiero Vigani

Plant Physiology Unit,

Centro dell'Innovazione- Dip. Scienze della Vita e Biologia dei Sistemi

Università degli studi di Torino, Torino, Italy

Email: gianpiero.vigani@unito.it

Tel +39 0116706360

<https://www.sciencedirect.com/science/article/pii/S0981942821005052?dgcid=coauthor>

<https://doi.org/10.1016/j.plaphy.2021.09.040>

## **Abstract**

The investigation of the adaptive strategies of wild plant species to extreme environments is a challenging issue, which favors the identification of new traits for plant resilience. We investigated different traits which characterize the root-soil interaction of *Parietaria judaica*, a wild plant species commonly known as “Pellitory-of-the-wall”. *P. judaica* adopts the acidification-reduction strategy (Strategy I) for iron (Fe) acquisition from soil, and it can complete its life cycle in highly calcareous environments without any symptoms of chlorosis. In a field-to-lab approach, the microbiome associated with *P. judaica* roots was analyzed in spontaneous plants harvested from an urban environment consisting of an extremely calcareous habitat. Also, the phenolics and carboxylates content and root plasticity and exudation were analyzed in *P. judaica* plants grown under three different controlled conditions mimicking the effect of calcareous environments on Fe availability: results show that *P. judaica* differentially modulates root plasticity under different Fe availability-impaired conditions and that it induces, to a high extent, the exudation of caffeoylquinic acid derivatives under calcareous conditions, positively impacting Fe solubility.

**Keywords:** calcareous soil, iron deficiency, microbiome, *Parietaria judaica*, pellitory-of-the-wall, phenols, rhizosphere, root architecture, urban habitat

## Introduction

Variations in soil pH are among the abiotic factors affecting most plant growth, since they influence the bioavailability of essential nutrients and toxic elements for plants. The presence of carbonates in soils causes a decrease in the solubility of iron (Fe) and other micronutrients (Lindsay and Schwab, 1982; Schenkeveld and Kraemer 2018). Calcareous soils, representing 30% of the Earth's land surface, are characterized by high pH values and may contain high  $\text{HCO}_3^-$  ions in the soil solution. Iron deficiency-induced chlorosis, otherwise known as lime chlorosis, is a major nutritional disorder observed in crops growing in calcareous soils;—on one hand, alkaline pHs dramatically reduce Fe solubility in soil and on the other hand, the presence of  $\text{HCO}_3^-$  may interfere with the physiological processes of Fe uptake (Chen and Barak, 1982; Römheld, 1987; Kim and Guerinot 2007; Díaz et al., 2012). Plant roots grown under calcareous conditions may exhibit a higher Fe content than those grown under non-calcareous conditions, indicating that low Fe availability in soil is not the unique factor causing lime-induced chlorosis; indeed, the composition of the soil itself can also influence Fe uptake from the apoplast (Mengel, 1994).

Iron is an essential micronutrient for the energy-yielding electron transfer reactions of respiration and photosynthesis and other major metabolic processes in plants (Kobayashi and Nishizawa, 2012; Vignani and Murgia, 2018); the elucidation of plant adaptive growth strategies in calcareous habitats represents a valuable approach for the identification of tolerance traits under soil constraints. In this context, the characterization of *Arabidopsis thaliana* demes locally adapted in their native habitat to soils with high carbonate has recently been reported (Teres et al., 2019).

The understanding of plant resilience to different biotic and abiotic stresses and to various natural habitats and climate changes is an emerging and urgent goal, as outlined in the Plant Science Decadal Vision 2020-2030 (Henkhaus et al., 2020). In particular, Decadal Vision's proposed actions include, among others, the selection of ecologically diverse plant lineages for an in-depth analysis of their morphology, anatomy, ecology in their natural environments, as well as the exploration and characterization of as-yet-undiscovered plant-associated biota.

In the context of Fe nutrition, the issues raised by Plant Science Decadal Vision ~~described above~~, suggest that the adapting processes of non-crop plants to calcareous habitats, can potentially uncover traits ~~already~~ lost in domesticated crops. Moreover, they also support the importance of studying the root microbiota associated with non-crop plants (or wild crop relatives). Interestingly, urban calcareous habitats can also constitute a potential source of information of the resilience of plants in inhospitable soils.

In most cases, crop domestication led to modifications of the composition of root microbiota, with adverse effects on the association with beneficial plant microbes (Pérez -Jaramillo et al., 2016). Root

exudation of metabolic compounds plays a crucial role in influencing the recruitment of functional microbiota. In general, the variety of biotic and abiotic stresses, such as Fe deficiency, encountered by plants during *in vitro* and *in vivo* growth conditions impacts the biosynthesis of secondary metabolites (Cheynier et al., 2013; Isah, 2019). Several works have reported an increased number of phenolic compounds in plants and root exudates as a response to different environmental stresses (Cesco et al., 2010; Caretto et al., 2015), and as an adjustment of secondary metabolism occurring in response to Fe mobilization (Jin et al., 2007). Some species, such as *P. judaica*, can secrete ortho-dihydroxy phenolic compounds showing reducing and metal chelating properties that enhance Fe availability in the rhizosphere. Indeed, plant metabolites released to rhizosphere can have diverse effects on soil microbial communities by changing soil properties or nutrient availability in the root vicinity, directly attracting microbes or being toxic for others (Vives-Peris et al., 2020).

In some cases, specific plant root exudates initiate a molecular dialogue mediated by both partners' exometabolite production, resulting in the establishment of beneficial plant-microbe associations (Sasse et al., 2018). Badri and co-workers found a strong positive correlation between phenolics and Plant Growth Promoting Bacteria (PGPB) species, including *Rhizobium*, *Bacillus*, *Sphingomonas*, *Streptomyces* and *Frankia* (Badri et al., 2013).

A similar mechanism of recruitment of PGPB has been demonstrated for coumarins in Fe-limiting soils: besides their established role in Fe mobilization in the rhizosphere (Rajniak et al., 2018), coumarins are also involved in shaping root-associated microbiomes (Voges et al., 2019). Furthermore, Harbort et al. (2020), by using synthetic microbiota and *A. thaliana* plants deficient in the exudation of secondary metabolites, demonstrated that coumarins are important drivers for the assembly of the rhizospheric bacterial community under Fe deprivation. All these recent works on the tripartite interaction between roots' exudates, soils and microorganisms support the emerging view that metabolites exuded under peculiar environmental conditions can recruit microbiota components able to alleviate the specific stress experienced by the plant.

The present work aims to unravel root-related processes of *Parietaria judaica*'s response to Fe deficiency, particularly its root plasticity and metabolite exudation under Fe deficiency. *P. judaica* (L. 1753) is a wild perennial dicotyledonous plant capable to grow in acidic and alkaline soils (Tato et al., 2020); it represents the most widespread plant species to be found in highly calcareous and hostile environments such as wall cracks exposed to the sun, where it displays phenotypic changes, though without any chlorotic symptoms (Dell'Orto et al., 2003; Donnini et al., 2012; Tato et al., 2020). In this work, we adopted a field-to-lab approach to first investigate the microbiome associated with *P. judaica* roots in plants growing spontaneously in an urban environment and harvested from an extremely calcareous habitat. We then analyzed phenolics content and root plasticity and exudation

of *P. judaica* plants grown under various controlled laboratory conditions mimicking the effects of calcareous environments—conditions; in particular, we applied conditions that allowed us to discriminate the effects of low Fe availability from those caused by the presence of carbonate and alkaline conditions (Tato et al., 2020).

## **Material and Methods**

### ***P. judaica* growth conditions**

*P. judaica* plants were sampled in an urban area of Milan (45°28'36.8"N, 9°13'39.2"E; 45°28'37.0"N, 9°14'00.1"E; 45°28'35.2"N, 9°14'03.7"E): to sample the whole root apparatus, the walls and the substrates where plants were growing were broken if necessary.

Cuttings of *P. judaica* were allowed to radicate in aerated half-strength nutrient solution for 10 days (Tato et al., 2020). Rooted plants were then transferred into 10 L plastic tanks (40 plants/tank) under four different conditions: +Fe (control, complete nutrient solution adjusted to pH 6.2 with NaOH), -Fe (complete nutrient solution without Fe, adjusted to pH 6.2), Bic (complete nutrient solution supplemented with 5 mM CaCO<sub>3</sub> and 15 mM NaHCO<sub>3</sub>, pH 8.3) and Tric (complete nutrient solution, buffered with Tricine at pH 8.3); the pH was adjusted with NaOH if required. The nutrient solution was changed every two days. Plant size at the harvesting time has been previously reported in Tato et al (2013)

Treatments were carried out for 7 days in a growth chamber under 16/8 h light/dark regime with cool-white light 200  $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ , 27/21°C temperature range, 65-75% relative humidity.

### **Sampling of *P. judaica* in urban sites and DNA extraction of their root-associated microbes**

From each of the three urban sampling sites considered, three replicates of bulk soil (i.e. the soil portion not interested by the root presence, according to Bulgarelli et al., 2012) and root samples from *P. judaica* were collected. Large soil aggregates were removed from the roots by shaking them as described in Bulgarelli et al., 2012, in order to leave only the rhizospheric soil attached to the roots. Bulk soil and root samples with attached soil were then stored at -20°C until DNA extraction. To obtain the samples that represent the root-associated microbial diversity, the root-adhering soil plus the root itself were processed together. This condition accounts both for the externally associated microbes and for the root endophytes (the rhizosphere + the endorhiza, called rhizosphere for the sake of brevity hereinafter).

The rhizosphere samples were first lyophilized for 24 h and then ground to a fine powder with liquid nitrogen. DNA was obtained as described in Edwards et al., (2015) extracting from a 300-mg bulk soil samples and 50 mg rhizospheric samples (dry weight), by using the MoBio PowerSoil DNA

Isolation Kit (Qiagen Inc., Hilden, Germany) and DNeasy Plant Mini Kit (Qiagen Inc., Hilden, Germany), respectively, according to the manufacturer's instructions.

Extracted DNA was quantified using NanoDrop spectrophotometry (NanoDrop, Wilmington, DE, USA) and normalized prior to library construction for the high-throughput sequencing.

### **Molecular, bioinformatic and statistical analyses of the microbes associated with *P. judaica* roots**

The extracted genomic DNA was used to amplify the V3–V4 region of the prokaryotic 16S rRNA gene, using the modified primer pair pro341f/pro805r (Takahashi et al., 2014) with the standard Illumina overhang. Fungal ITS2 rDNA cistron was amplified using the modified primers fITS7 (Ihrmark et al., 2012) and ITS4ngs (Tedersoo et al., 2014), with the standard Illumina overhang adapters. Purified PCR products were combined in equimolar amounts, and the corresponding metabarcoding libraries were sequenced on the Illumina MiSeq platform (Illumina, San Diego, CA, USA) with paired-end 2 x 300 bp sequencing mode at the IGA Technology Services (Udine, Italy). Raw data were processed and analyzed following the pipelines of QIIME2 version 2019.7.0 (Bolyen et al., 2019; Caporaso et al., 2010). The high-quality reads were clustered into operational taxonomic units (OTUs) at a 97% identity level and chimeric sequences were filtered using UCHIME (Edgar, 2011), as implemented in the QIIME2 pipeline.

Taxonomy assignment of both prokaryotic and fungal OTUs was performed using ~~respectively~~ the SILVA database (version 132, release date 13.12.2017; Quast et al., 2012; Yilmaz et al., 2013) and the ITS UNITE database (UNITE QIIME release for Fungi, Version 18.11.2018. <https://doi.org/10.15156/BIO/786334>) respectively using *sklearn* algorithm as implemented in QIIME2 (Pedregosa, 2011).

All statistical analyses were conducted in R v3.6.1 (R Development Core Team, 2016). Rarefaction species richness curves were generated using the R package *vegan* (Oksanen et al., 2017). Alpha diversity indexes were calculated for Prokaryotic and Fungal OTU tables using the *vegan* package (Oksanen et al., 2017). The differences between soil and root samples were tested by using Tukey's Honest Significant Differences test, with the R package *TukeyC* (Faria et al., 2016).

To correct for difference in sequencing depth, a subsampling at even sequencing depth from each sample (9,935 for Prokaryotic and 2,632 for Fungal samples) was performed before the downstream analysis using the R package *phyloseq* (McMurdie and Holmes, 2013), generating also the taxonomical composition of the whole microbial community. The significance of Bray-Curtis dissimilarity between the soil and root samples were tested by permutational multivariate analysis of variance (PerMANOVA) using the *adonis* function in the R package *vegan* (Oksanen et al., 2017)

with 9999 permutations. The multivariate homogeneity of group dispersions was first assessed by means of the *betadisper* and *permutest* (with 9999 permutations) functions in the R package *vegan* (Oksanen et al., 2013). The differences in the composition of Prokaryotic and Fungal communities in *P. judaica* soil and root samples were visualized by means of a Non-metric Multidimensional Scaling (NMDS) ordination carried out using *metaMDS* function in the R package *vegan* (Oksanen et al., 2013). The R package *gunifrac* (Chen et al., 2012) was used to test differences in the microbial composition of soil and root samples. Co-occurrences in the Prokaryotic and Fungal communities were assessed by performing network analysis using the Spearman rank correlations between OTUs ( $\rho > 0.7$  and  $p < 0.001$ ). All networks were visualized with the Fruchterman-Reingold layout with 9999 permutations in the R package *igraph* (Csardi and Nepusz, 2006). Descriptive and topological network properties, as well as network modules (substructures of nodes with a higher density of edges within the group than outside it) were calculated as described in Hartman et al., (2018).

### **Morphological analysis of arbuscular mycorrhizal (AM) root colonization**

Roots of *P. judaica* plants sampled in urban sites (described above, n=8) were carefully washed with tap water, stained overnight in a solution of methyl blue (0.1% w/v) in 80% lactic acid (v/v) and clarified in lactic acid in order to remove the excess of staining solution. 80 segments, 1 cm long each, were obtained from each root apparatus, placed on glass slides and observed under a light microscope.

### **HPLC analysis of phenolic compounds**

For qualitative and quantitative determination of phenolic compounds, plant tissues (1 g FW) of *P. judaica* were first homogenized for 3 min with 30 mL hot MeOH-EtOH (1:1) and then refluxed under nitrogen for 30 min (2×). After centrifugation and pooling of the extracts, the combined solutions were first concentrated under vacuum, depigmented with petroleum ether (bp 40-70 °C), filtered through 0.45 µm Millipore Millex-HN, and then used for the determination of total phenolic content and HPLC-DAD determination of phenolic compounds (Lattanzio et al., 2009). Identification of phenolics was made by using retention times (tR) and spectral data of different peaks compared with standard compounds (Extrasynthèse, Genay, France). In addition, HPLC-MS/MS analyses of main peaks identified in *P. judaica* and standard compounds were performed in order to obtain useful data for structure characterization.

Metabolic compounds released by root (root exudates fraction) were collected according to Tato et al. (2020) from plants grown hydroponically. The collected materials were acidified to pH 3.5-4.0 with HCl to maintain the structural stability of phenolic compounds, freeze-dried, suspended in 3 mL methanol and filtered through 0.45 µm Millipore Millex-HN; the filtered solution was analyzed for



total phenolic content and HPLC determination of phenolic compounds. HPLC analyses were performed with a Hewlett Packard Series 1100 liquid chromatograph equipped with a binary gradient pump G1312A, a G1315A spectrophotometric photodiode array detector was set at 325 nm, and G1316A Column with the thermostat set at 45°C. The Hewlett Packard Chem Station (Rev. A. 06.03) software was used for spectra and data processing. An analytical Phenomenex (Torrance, CA, USA) Luna C18 (5) column (4.6 x 250 mm) was used throughout this work. The solvent system consisted of (A) MetOH and (B) acetic acid-water (5/95, v/v). The elution profile was as reported by Lattanzio and Van Sumere (1987). The flow rate was 1 ml min<sup>-1</sup>. Samples (25 µl) were applied to the column using a 25 µl loop valve. UV absorption spectra were acquired in the 235-450 nm range.

HPLC-MS/MS analyses were performed on a QTrap MS/MS system, (~~from~~ Applied Biosystems, Foster City, CA, USA), equipped with an ESI interface and a 1100 series micro-LC system comprising a binary pump and a microautosampler (Agilent Technologies, Waldbronn, Germany). The ESI interface was used in positive ion mode, with the following settings: temperature (TEM) 350 °C; curtain gas, nitrogen, 30 psi; nebuliser gas, air, 10 psi; heater gas, air, 30 psi; ion spray voltage + 4500 V. Full scan chromatograms were acquired in the mass range 100 –800 amu, MS/MS chromatograms were acquired at collision energy of 20 V. LC conditions were as for the HPLC-DAD analysis.

### **Root morphology and biomass allocation**

After 7 d treatments (+Fe, -Fe, Bic, Tric), three independent biological samples from each treatment were collected randomly, and their shoots and roots were harvested. Shoots were dried at 70°C for 48 h, and their dry weight (WS, g) was measured. The root system was stained with 0.1% (w/v) toluidine blue O for 5 min and then divided into two root orders: ‘shoot-borne’ or adventitious roots (AR), and their 1<sup>st</sup>- order lateral roots (LR) as defined by Atkinson et al., (2014). Each root was scanned at 300 dpi resolution (WinRhizo STD 1600, Instruments Régent Inc., Canada) to determine length (LA, cm), volume (VA, cm<sup>3</sup>) and surface area (SA, cm<sup>2</sup>) of the adventitious roots and the total length (LI, cm), total volume (VI, cm<sup>3</sup>) and total surface area (SI, cm<sup>2</sup>) of the 1st-order laterals by the WinRhizo Pro v. 4.0 software package (Instruments Régent Inc.). Length (LT), surface area (ST) and volume (VT) of the whole root system were calculated as sum of the two root types. Then, dry weights of the adventitious roots (WA, g) and total dry weight of 1st-order lateral roots (WI, g) were measured after drying in the oven at 70°C for 48 h. Total root dry weight (WT, g) was the sum of the WA and WI. Plant dry weight (WP, g) was obtained by the sum of WT and WS. Based on the measurements above, the following parameters were calculated for the whole root system:

root length ratio RLR= LT/WP (cm g<sup>-1</sup>) (1)

root mass ratio  $RMR = WT/WP$  ( $g\ g^{-1}$ ) (2)

fineness  $F = LT/VT$  ( $cm\ cm^{-3}$ ) (3)

tissue density  $TD = WT/VT$  ( $g\ cm^{-3}$ ) (4)

where the RLR expresses the root order's potential for the acquisition of below-ground resources, the RMR indicates the relative biomass allocated to the root and the F and TD represent the structural root parameters. The relationship among the above parameters is the following:  $RLR = RMR \times F/TD$  (Ryser and Lambers, 1995).

The adventitious (NA) number and the 1st-order laterals (NI) were directly counted from the images. The average length of the adventitious [ $aLA=LA/NA$ ] and the 1st-order laterals [ $aLI = LI/NI$ ] (cm) were also calculated. The 'branching zone' length (BZ) that extends rootwards from the shoot base to the youngest emerged LR and the 'lateral formation zone' (LFZ) that spreads from below the youngest emerged LR to the 2 to 6 mm from the root apex, were also measured, as described in Dubrovsky and Forde (2012).  $NI/BZ$  calculated the root branching density (BD, number of laterals in cm of branching zone).

Two-way ANOVA tested the effects of the different treatments on the root parameters. Tukey's post hoc test comparison was applied to test the effects of each treatment at  $P<0.05$ . To correct for allometric effects (Coleman et al., 1994), the ln-transformed plant dry weight (ln WP) was used as a covariate to analyse the root morphology and biomass allocation, when significant correlations between lnWP and these root traits were found. A multivariate statistical PCA approach (principal components analysis) and a cluster analysis were performed by using SPSS software. To unveil the impact of the root morphology pattern on plant growth, Pearson's test was used to test the correlation between the PC factor scores and plant DW. Statistical analysis was conducted using the Systat v. 8.0 software package (SPSS Inc., Evanston, IL, USA).

### **Estimation of total ortho-dihydroxy phenolic compounds (Arnow's reagent)**

One mL of extract sample was placed in a test tube and 1 mL 0.5 N HCl was added. The tube was well mixed and then 1 mL Arnow's reagent (10 g Na nitrite and 10 g Na molybdate in a final volume of 100 mL distilled water) was added (which resulted in a yellow color), mixed, and 1 mL 1 N NaOH was added (solution turned into red color). The solution was then brought to a final volume of 5 mL with distilled water and absorbance was measured at 500 nm. The concentration was calculated and expressed as  $mg\ g^{-1}\ FW$ . Chlorogenic acid was used for the standard curve prepared in a range of 0-0.15  $mg\ mL^{-1}$ .

### **Iron reduction by phenolics compounds**

The phenolics concentration in root exudates and the caffeic and citric acids ability to reduce Fe(III)-EDTA was measured spectrophotometrically by using BPDS (Chaney et al., 1972). Root exudates (100 µg), prepared as described above, were incubated for 120 min in 1 mL 100 mM Fe(III)-EDTA, 100 mM BPDS solution in the dark, 26 °C, under shaking. A solution containing caffeic acid (50 mM) and/or citric acid (50 mM) 100 mM Fe(III)-EDTA and 100 µM BPDS in the dark at 26 °C under shaking, was also prepared, according to Hu et al. (2005). The absorbance at 535 nm was measured as in Donnini et al. (2009).

### **Soil incubation with caffeic and citric acids**

Solutions of caffeic and citric acids were adjusted to pH 5.5 using diluted NaOH and were then added to soil at rates 50 µmol g<sup>-1</sup> soil according to Hu et al., (2005). The soil was watered to field capacity, and then incubated at 20°C for 30 min. Soluble fractions of soils were collected according to Mimmo et al. (2008). Extracts were filtered through 0.2 µm filters and then analysed by an Agilent 7100 Capillary Electrophoresis System (Agilent Technologies, Santa Clara, CA, US). Phosphate anions were determined by capillary electrophoresis, using a bare fused silica capillary with extended light path BF3 (i.d. = 50 µm, I = 72 cm, L = 80.5 cm). Sample injection was at 50 mbar for 4 s with -30 kV voltage and detection at 350/80 nm wavelength. Compounds were identified using pure standards and anions contents were expressed as µg g<sup>-1</sup> FW.

### **Miscellaneous**

Iron and P content were determined by ICP-MS on oven-dried tissue samples (n=3) mineralized in HNO<sub>3</sub>. and carboxylic acids contents in roots were determined according to Tato et al. (2020). Apoplastic Fe was determined according to Tato et al., (2020). Briefly, roots from 5 plants per treatment were transferred to a beaker with 0.5 mM CaSO<sub>4</sub> under vigorous aeration. After 10–15 min, roots were transferred to 40 mL tubes with 21 mL of 10 mM 2-(N-morpholino)ethanesulfonic acid (MES), 0.5 mM Ca(NO<sub>3</sub>)<sub>2</sub>, 1.5 mM 2,2' bipyridyl (pH 5.5) at 25 °C. Tubes were covered with a cotton plug and N<sub>2</sub> was bubbled through the solution. After 5 min, 1 mL of 250 mM Na<sub>2</sub>S<sub>2</sub>O<sub>4</sub> was added. The A<sub>520</sub> of the solution (A<sub>520</sub> of 1mM Fe[bipyridyl]<sub>3</sub> = 8.650) was determined on 2 mL aliquots. The aliquots employed for the determinations were returned to the tube and left for 1 h in the dark. The determination was carried out every 1 h 30 min until a constant value was obtained. Lignin visualization was performed according to Donnini et al. (2011). After being fixed at 4 °C overnight in 100 mM Na-phosphate buffer (pH 7.00) containing 4% paraformaldehyde (w/v), root segments were dehydrated through an ethanol–tertiary butanol series and embedded in paraffin

(Paraplast Plus, Sigma). Serial sections (5  $\mu\text{m}$ ) were cut with a microtome, mounted on silanized slides, deparaffinized in xylene and rehydrated through an ethanol series. Sections were then stained with the safranin/fast green method (Johansen, 1940), mounted with cover slides, observed by optical microscopy (Leica DMR) and images were acquired using a digital camera (Leica DC300F).

## Results

### The root-associated microbiome of *P. judaica* collected from urban sites

*P. judaica* plants were harvested from soil of an urban site displaying alkaline and calcareous conditions, with a pH value of  $7.9 \pm 1.3$  and an average 20% CaCO<sub>3</sub> content.

The root-associated microbiota of the collected plants (rhizosphere samples) and that of the soil not interested by the root presence were then analyzed. The rarefaction curves indicated that a satisfactory sequencing depth was obtained for each considered sample, for both fungal and bacterial amplicons (Fig. S1). The clustering of high-quality reads allowed us to obtain 349 fungal and 2327 prokaryotic operational taxonomic units (OTUs), that were further assigned taxonomically (Supplemental files 1a, 1b). The standardized datasets were used to generate Venn diagrams of the community composition and unique or shared OTUs among bulk soil and rhizosphere samples (Fig. 1). The results show a reasonable degree of microbiota diversity, especially for prokaryotes (more than 2000 OTUs overall). Also, a reduced number of OTUs unique to the rhizosphere compartment has been detected, while the highest OTU number was present in the bulk soil only. The OTUs shared between rhizosphere and bulk soil likely represent the core of microbes recruited by roots under the occurring environmental conditions.

The analysis of Alpha diversity indexes outlines the structure of the considered microbial communities (Fig. S2). All the indexes point to a higher microbial richness in the bulk soil than the rhizosphere, in agreement with the higher OTU number detected in the former compartment. A general analysis of Beta diversity on prokaryotic and fungal communities ~~comparing soil with rhizosphere samples~~ was also conducted. The Bray-Curtis index does not record a significant dissimilarity in the microbial composition of bulk soil and rhizosphere; however, the two compartments appear more diverse when phylogenetic distances weighted by relative abundances are considered in relation to fungal microbiota (weighted GUnifrac, Fig. S3). Notably, the prokaryotic taxa typically present and dominant in soils (Janssen, 2006) were all detected in the current experiment (Fig. 2), with Proteobacteria as the most represented in both compartments. However, Acidobacteria were underrepresented; this result is consistent with the high pH of the soil under investigation (around pH 7.9), as many species in this phylum are indeed acidophilic.

Bulk soil and rhizosphere compartments do not show a dramatic difference in the overall community structure, although some shifts in microbial composition can be detected already at the phylum level (Fig. 2). The rhizosphere showed an increase in the relative abundance of Proteobacteria, Actinobacteria and Firmicutes phyla in comparison to the bulk soil. In contrast, other phyla such as Bacteroidetes, Chloroflexi, Planctomycetes and Verrucomicrobia were significantly more abundant in the bulk soil compartment (Fig. 2, upper panel).

A more in-depth analysis of microbiome composition (at genus level) revealed a higher relative abundance of *Bacillus*, the *Rhizobium* group and *Streptomyces* in the rhizosphere, and for the last two genera the increase was statistically significant (Fig. 3, upper panel). However, the fungal genera *Mortierella* and *Wallemia* were more abundant in the bulk soil (Fig. 3, lower panel). This latter genus comprises a few known species for their xerotolerant and halophilic behaviour (Zajc and Gunde-Cimerman, 2018).

Arbuscular mycorrhizal (AM) fungi provide many services to plants, including the improvement of mineral nutrition, at a cost for the plant host; indeed, they take up photosynthates from the roots. Hence, the presence of AM fungi of such urban soil was also investigated; a few OTUs referring to AM fungi were retrieved in both the bulk soil and the rhizosphere compartment (data not shown). They point to *Funneliformis mosseae*, a widespread species common in diverse soils, and to another AM fungus also belonging to Glomeromycotina. Despite AM OTUs not being abundant in our dataset, arbuscule formation was observed in the same *P. judaica* roots sampled for the microbiome sequencing, suggesting that AM fungi are an active microbial component of such an urban niche.

Co-occurrences in the Prokaryotic and Fungal communities were assessed by performing network analysis and visualizing the positive, significant correlations among OTUs ( $\rho > 0.7$  and  $p < 0.001$ , Fig. S4). Similarly, meta-networks were constructed to visualize correlations between Prokaryotic and Fungal OTUs in the soil and root communities (Fig. 4). Within this network, we identified keystone OTUs (Fig. 4), defined as the top 1% node with the highest degree of interactions. These OTUs are microbial taxa that frequently co-occur with other taxa under the experimental conditions considered and are thought to be ecologically important and potentially to play a key in the structuring of the microbiota (Hartman et al., 2018, Supplemental file 2).

Also, OTUs that might act as indicator species in such networks were sought; a species is described as an “indicator” when it is characteristic of a group of samples or experimental treatments and/or is highly sensitive to the changes entailed by the treatment. Only two bacterial OTUs could be identified as indicators for the rhizosphere compartment (Supplemental file 3): the first one refers to the *Rhizobium* genus (probably *Rhizobium grahamii*, 100% sequence identity), the second points to an uncultured isolate belonging to Actinobacteria (99.75% identity). Members of Actinobacteria are widespread in soils and display tolerance to diverse extreme conditions (Ranjani et al., 2016).

Indicator species of the bulk soil comprise genera of fungi and bacteria known to be widespread in soils, including saprotrophs such as *Mortierella* as well as microbes that tolerate extreme environments, such as the fungi *Coniosporium apollinis* and *Naganishia albida*, as well as the bacteria *Microvirga*, *Brevundimonas*, *Altererythrobacter* and *Rhodospirellula* (Supplemental File 3).

The microbiome associated with the roots displayed an increase in P solubilizing microbial genera (as defined by Kalayu, 2019), mainly belonging to Rhizobiaceae and Streptomyces (Fig. 3, upper panel). Conversely, some soil generalist microbes seem to be rather excluded from the rhizosphere of *P. judaica*.

### **Root morphology of *P. judaica* grown in calcareous, alkaline or Fe-deprived media**

*P. judaica* plants sampled from the urban sites were allowed to radicate in half-strength complete nutrient solution and then transferred into one of four different media (i.e. +Fe, -Fe, Bic and Tric), to discriminate between the effects of low Fe availability due to a high pH and those of bicarbonate itself (Tato et al., 2020). Since alkaline and calcareous conditions mainly affect Fe and P availability, the leaf and root concentration of these nutrients was determined. A reduction of Fe concentration was observed in leaves of plants grown in the -Fe, Bic and Tric media, and in -Fe roots (Fig. 5). Phosphorus was slightly decreased with respect to the control only in the roots of plants grown in Bic (Fig. 5).

The morphology of the whole root systems of *P. judaica* was not significantly modified by all the treatments in comparison with the +Fe condition, with TD exception (Fig. 6A, B and C). Indeed, the TD of the whole root system was lower in the -Fe and +Fe than in Bic- and Tric-treated plants (Fig. 6C).

The root system of *P. judaica* consisted of the adventitious roots (AR), also named "shoot-borne" roots, and the lateral roots (LR), which emerged from AR, suggesting the "within-root" approach to analyze the root morphology. Differently from the whole root system, some treatments significantly modified the LI and aLI and the aLA total and average length of the LI, the aLI and the aLA in comparison to those of the +Fe plants (Fig. 6D): Bic-treated plants decreased (-49%) the aLA in comparison to the +Fe treatment and the Tric treatment reduced the LI and aLI by -83% and -71%, respectively, compared to the +Fe plants (Fig. 6D). The Fe-deficient plants exhibited similar morphology in both root types compared with the +Fe plants (Fig. 6D). As expected in plants adapted to alkaline soils (White et al., 2013), Bic-treated *P. judaica* plants exhibited an increase in the BD (+61%) determined by a reduction of the BZ, but these parameters seemed unchanged in Fe deficiency and Tric treatments (Fig. 6E).

The PCA was applied using only root parameters significantly changed by treatments as observed in the univariate ANOVA. The PC1 (49% of proportion of variability) consisted of high positive loads for the aLA and the BZ and negative loads for the TD, while the PC2 (38%) showed high positive loads for the LI and aLI (Table 1). Two-dimensional PCA score plots and subsequent hierarchical cluster analysis revealed a sharp separation among the treatments (Fig. 6A; Fig. S1). In particular, a

first cluster included the +Fe plants, a second cluster comprised the Bic-treated plants, and a third one incorporated both the Tric- and Fe deficient-treated plants (Fig. 7A; Fig. S5).

As shown by Figure 6A and Table 1, the Bic-treated plants were characterized by lower aLA and BZ associated with high TD, aLI and LI, thus exhibiting root architectures different to the +Fe plants. Conversely, the root architectures of the Fe deficient- and Tric-treated plants exhibited lower aLI and LI but intermediate values of aLA, RBZ and TD between the Bic treated and the +Fe plants (Fig. 6A). The importance of these different root architectures for the *P. judaica* fitness was tested by a Pearson correlation between the plant dry weight and the PC1 and PC2 scores. The PC1 was significantly and positively correlated with the plant growth ( $r=0.4890$ ,  $p=0.0139$ ) differently to the PC2 ( $r=0.0356$ ,  $p=0.55$ ) (Fig. 7B, C), suggesting that the high aLA, BZ and low TD but not the lateral roots (PC2) explained 58% of the *P. judaica* growth.

### **Carboxylic acids and phenolic compounds in root tissues and in exudates of *P. judaica* grown in calcareous, alkaline or Fe-deprived media**

The total phenolics and the citric and malic acids released by *P. judaica* roots were recently monitored in the four conditions, i.e. control, -Fe, Bic, Tric (Tato et al., 2020). The profile of carboxylic acids and the total phenolics of *P. judaica*, both in roots and in their exudates are shown in Fig. S6 and S7, respectively. Malic, ketoglutaric and citric acids were detected in all tested root samples, with a higher accumulation of malic and citric acid in -Fe, Bic, Tric treatments whereas ketoglutaric acid concentration increased only in -Fe and Tric-treated plant roots. Both malic and citric acid were present in all root exudates, with a higher accumulation in -Fe, Bic and Tric treatments. Notably, cis-aconitic acid was exuded only by the Tric-treated roots.

Total root phenolics were measured spectrophotometrically using Arnow's reagent, which selectively determines the concentration of ortho-dihydroxy phenolic compounds (Fig. S7A). Only the Bic treatment induced a significant increase (87%) of total phenolic concentration in roots. A non-significant increase was observed in the other two conditions, -Fe and Tric (+28% and +17% respectively).

These data are consistent with the results obtained from chromatographic HPLC-MS analyses where an increase of total phenolics in plants grown under Bic condition (+73%) and in plants grown in Tric conditions (+18%) was found (Fig. S7B). In detail, the phenolic fraction of root tissues was characterized by the presence of various positional isomers of mono- and di-caffeoylquinic acid esters. 5-*O*-Caffeoylquinic acid (chlorogenic acid) and 3,5-*O*-dicaffeoylquinic acid were the main constituents of the phenolics fraction in root extracts (Fig. 8A, left panel). Other caffeoylquinic derivatives identified in roots were: 3-*O*-caffeoylquinic acid, 4,5-*O*- and 3,4-*O*-dicaffeoylquinic



acids, and two *p*-coumaric acid glycosides. Accordingly, the phenolics content of plants harvested in the urban site revealed that the main constituents of root extracts of *P. judaica* were represented by Chlorogenic acid and 3,5-O-Dicaffeoylquinic acid that constitute together about 88% of the total phenolic compounds present in root extracts (Fig.8B).

From a qualitative viewpoint, isomerization phenomena have been observed in the phenolic fraction of stressed plants in comparison with the control. Overall, such isomerization phenomena in all stressed plant extracts cause an increase of total mono-caffeoylquinic acids while the total content of di-caffeoylquinic acids decreases (Fig.8A, left panel).

The nutritional stress conditions -Fe, Bic, Tric also affected the composition of phenolic fractions of *P. judaica* root exudates (Fig. 8A, right panel) in which 3-*O*- and 5-*O*-caffeoylquinic acids, 3,5-*O*-dicaffeoylquinic acid and the aglycone caffeic acid have been identified; the latter compound is absent in root extracts. This suggests that secretion of phenolics by *P. judaica* roots (both control and stressed roots) also leads to partial hydrolysis of caffeoylquinic esters. Although total phenolics concentration determined by Arnow's reagent method revealed an increase mainly in root exudates of -Fe plants (Fig. S7A), HPLC-MS results revealed an increase in total phenolics root exudates from -Fe, Bic and Tric treated plants (Fig. 7B). Again, the highest increase in phenolics compounds was observed in Bic exudates.

In Bic and Tric exudates, caffeic acid accounted for 67% and 44% of total phenolics content, respectively. Whereas, caffeic acid and chlorogenic acid together accounted for 76-84% of total phenolic compounds in both control and all the stressed exudates (Fig. 8A, right panel).

### **Fe mobilization properties of *P. judaica*'s root exudates in calcareous, alkaline or Fe-deprived media**

The ability of root exudates to favor Fe mobilization was tested by measuring the *in vivo* Fe reductase activities of exudates secreted from roots of *P. judaica* plants grown in -Fe, Bic and Tric media. Such Fe reductase activity was increased by all three stress conditions, and in particular, it was the highest in the Bic treatment (Fig. 9A). Since Bic-grown *P. judaica* roots exuded citric and caffeic acids, the effect of the commercially available caffeic and citric acids was assayed on the Fe(III) reduction. Caffeic acid displayed a higher Fe reduction capacity compared with citric acid (Fig. 9A). Besides Fe availability, calcareous conditions also affect phosphorus (P) availability; the urban calcareous soil where plants were collected was incubated with caffeic and citric acid to study their potential effect on PO<sub>4</sub><sup>3-</sup> solubility. Soil incubation with both caffeic acid and citric acid enhanced the concentration of PO<sub>4</sub><sup>3-</sup> in the soil soluble fraction with respect to the control, but the effect of citric acid was stronger than that of caffeic acid (Fig. S8).

Bic treatment led to a significant accumulation of Fe in the apoplast, suggesting ~~indicating~~ that the higher synthesis of phenolics compounds might be induced by the high Fe content in the intercellular spaces (Fig. 9B). Other than Fe mobilization, phenolics might be involved in other reactions, such as lignification process (Domini et al., 2011). Therefore, the lignification rate of root tissues was investigated by a staining procedure and the root cross-sections were visualised microscopically. Roots of plants grown under Bic and Tric showed high lignification signals at rhizodermis, endodermis and cortex layers (Fig. 9C). Such findings suggest that both alkaline growth conditions (Bic and Tric) induced the synthesis of lignin in the root cell walls.

## Discussion

The study of plants living in natural and extreme environments, in which different stressors naturally coexist, allows to unravel the morpho-anatomical and physiological traits enabling them to survive in these extreme environments (Bartoli et al., 2013; Bechtold, 2018). In this paper, we investigated different traits characterizing the root-soil interaction of *P. judaica*, a wild plant species commonly known as “Pellitory of the wall”: morphological plasticity, exudation, and association with the microbiome.

The analysis of the root microbial community associated to spontaneous *P. judaica* plants harvested from the urban environment revealed a good degree of microbiota diversity, detecting all the prokaryotic taxa typically present and dominant in soils. The comparison of the root-associated versus bulk soil microbiota highlighted some shifts in the microbial composition, with a higher relative abundance in the rhizosphere of some genera, including beneficial species for plants. Among these beneficial bacterial species, two OTUs are noteworthy; the first OTU refers to a bacterium from the genus *Rhizobium*, whereas the second OTU points to an uncultured actinobacterium. Besides Rhizobia's ability to form N-fixing nodules on legume roots, they can also thrive in the rhizosphere of non-leguminous plants acting as Plant Growth Promoting Rhizobacteria (PGPR, Mehboob et al., 2012), and thus providing benefits even in the absence of nodule formation. Also, besides the well-studied *Streptomyces* genus, many other Actinobacteria can associate with roots of a wide range of hosts being beneficial for the plants' health, and the interest in their application as PGPR bacteria has raised in recent years (Sathya et al., 2017).

Taken together, our analyses of the root-associated microbiome of spontaneous urban *P. judaica* plants indicate that these plants retain the competence to actively recruit beneficial soil microbes such as PGPR, phosphate solubilizers and AM fungi, possibly excluding from their rhizosphere other

components of the soil microbiota. These results are remarkable, given the limited microbial reservoir to which plant roots had access in the urban environmental niche where these plants were growing spontaneously.

It is now well acknowledged that plants play an active role in the assembly of the rhizospheric microbiota, and the outcome and magnitude of such an influence can change, depending on both fixed (e.g. plant genotype) and variable (biotic/abiotic stresses) factors. In this scenario, root exudation can indeed act as a crucial driver of microbiota recruitment (Sasse et al., 2018). Among the different fractions of *A. thaliana* root exudates, phenolics are the most effective in shaping the soil microbiome, as they significantly correlate with 31 bacterial OTUs (Badri et al., 2013). Also, a role for root-secreted coumarins in shaping the *A. thaliana* rhizospheric microbiota has been discovered (Voges et al., 2018). In particular, coumarins limit the growth of a *Pseudomonas* strain through a mechanism that involves the production of reactive oxygen species. Interestingly, we could not highlight any significant enrichment for the *Pseudomonas* genus in the *P. judaica* rhizosphere, although some other genera of PGP bacteria seemed to be actively recruited in the rhizosphere. This suggests that the negative effect demonstrated for coumarins on *Pseudomonas* growth might be extended to other phenolic compounds, thus providing indications towards the engineering of beneficial plant root microbiota.

Root morphological plasticity and exudation, which are two relevant processes driving plant-soil interaction, were investigated in *P. judaica* grown in three different controlled conditions inducing Fe-deficiency, i.e. -Fe, Bic, Tric, to discriminate between root responses to the low availability of Fe due to a high pH and that caused by bicarbonate (Tato et al., 2020). Among the morphological parameters of the whole root system, the TD of *P. judaica* was the only affected trait, mainly by the calcareous condition (Bic) (Fig. 1C). The TD is an adaptive trait positively correlated with the lignification degree and cell wall thickness (Ciamporova et al., 1998; Wahl and Ryser 2000; Hummel et al., 2007) and, in turn, is inversely related to the *Arabidopsis* adaptation to Fe deficiency (Barberon et al., 2016). As well, the root lignification degree has been interpreted as a Fe deficiency sensitivity trait in a quince rootstock (Donnini et al., 2011).

Besides the morphology of the whole root system, the 'within-root analysis' was applied, which accounted for the morphological changes of the different root types observed in the plant root system. Such a phenotyping approach could provide early information on the contributions of the different root types of *P. judaica* to the adaptation in alkaline, calcareous and Fe deficient conditions. Indeed, root types were found to respond differently to the environmental cues such as water (Romano et al., 2013; Tellah et al., 2014; Abenavoli et al., 2016), salt (Stevanato et al., 2013), and combined P/drought stress (Ho et al., 2005), N deficiency (Sorgonà et al., 2007), allelochemicals (Abenavoli et

al., 2004, 2008; Lupini et al., 2016), rot (Roman-Aviles et al., 2004) and fungal colonisation (Zadworny and Eissenstat, 2011). In the present work, lateral roots of *P. judaica* were more modified by treatments in comparison with adventitious ones. This kind of root ideotype, which is characterized by an even extended spread of roots throughout the soil, is useful for the nutrient acquisition with restricted phytoavailability in alkaline soils (White et al., 2013). Indeed, *P. judaica* exhibited an increase of the BD (+61%), maintaining the LI, in response to the Bic treatment (Fig. 5D,E).

Recently, several works pointed out the importance of the synergism and/or antagonism among the different root traits for understanding the root architecture adaptation to diverse environments (York et al., 2013; Miguel et al., 2015; Rangarajan et al., 2018), suggesting to us the multivariate rather than univariate approach for analyzing *P. judaica* root architecture. The “root multi-trait” pattern, as determined in the present work, also in agreement with the results of within-root morphology, might reflect the adaptation of *P. judaica* to low Fe availability caused by high pH (Tric-treatment) and by calcareous environment (Bic). Indeed, the root architecture of the -Fe and Tric-treated plants was characterized by the development of adventitious roots and branching zone associated with lower TD of their root axes. In particular, the low TD of the root axes observed in Fe-deficient and Tric treatments was negatively correlated with root exudation which, in turn, is a fundamental physiological trait of the Fe deficiency syndrome (Ladygina and Hedlund, 2010; Hell and Stephan, 2003; de Vries et al., 2019). This root trait in association with a high aLA and RBZ explained the higher fitness of *P. judaica* plants as shown by Pearson correlations.

Conversely to the -Fe and Tric-treated plants, the root architecture of the Bic-treated ones was characterized by higher LI but less TD and lower BZ. The Bic-induced calcareous environment affected the availability of different nutrients including P, Mn, B, and Zn (Tyler, 2003). In this study, a low P content in roots and a Fe accumulation in the root apoplast was observed in Bic-treated plants. In such conditions, the soil exploration by roots might be a strategy to survive in calcareous soils (White et al., 2013; Campestre et al., 2016; Ding et al., 2019). Indeed, *P. judaica* exposed to the calcareous condition displayed higher variability of root plasticity with respect to plants exposed to the other treatments, by reducing the branching zone, increasing the branching density and lateral spread.

Direct (-Fe) as well as induced Fe deficiency (Bic, Tric), all caused an increase in caffeoylquinic acid derivatives, especially in Bic-treated roots (both tissues and exudates). Interestingly, caffeic acid, probably arising from hydrolysis of caffeoylquinic esters, is one of the components of the root exudates. The accumulation of phenolics in plant tissues is a hallmark of plant stress: phenolic compounds may be synthesized *de novo* in plants as a response to various biotic and abiotic stresses, including nutrient deficiency (Osmond et al., 1987; Cheynier et al., 2013; Lattanzio, 2019). Several

studies have reported the increase in chlorogenic acid and/or mono- and di-caffeoylquinic acid in response to different abiotic stresses such as low-temperature (Lattanzio et al., 1987; Lattanzio, 2001; Lattanzio et al., 1994), wounding (Cantos et al., 2001), high UV-B irradiation and insect attack (Izaguirre et al., 2007).

Caffeoylquinic acid (CQA) derivatives are caffeic acid (3,4-dihydroxycinnamic acid) depsides, positional isomers of caffeic acid esters of quinic acid, which are broadly distributed in plants. The chelating activity of CQAs is attributed to their catechol ring (Kono et al., 1998). Low temperature stress induces, in artichoke tissues, an accumulation of constitutive phenolic compounds, mono- and di-caffeoylquinic acids, that protect chilled tissues from damage by free radical-induced oxidative stress (Lattanzio et al., 1994). Due to the presence of a catechol ring in its structure, chlorogenic acid promotes the reductive release of ferritin Fe as mobile  $\text{Fe}^{2+}$  that, in turn, forms colourless complexes with the excess of chlorogenic acid (Boyer et al., 1988). Hence, chlorogenic acid can act as a reductant of  $\text{Fe}^{3+}$  as well as a ligand of  $\text{Fe}^{2+}$ . In addition, this paper shows that the exudation process produces, likely by hydrolytic processes, caffeic acid, which exhibits a high Fe reduction ability. Overall, the results in the present work support the current view that secretion of phenolic compounds is a relevant component of the reduction strategy of Fe acquisition in non-graminaceous plants. In the past decades, several studies suggested that the secreted phenolics could enhance Fe availability in the rhizosphere soil, as an alternative/reinforcement of the membrane-bound reductase, through chelation and reduction of insoluble Fe. Initially, phenolics were thought to help with the solubilization and reutilization of apoplastic Fe in red clover. This feature was not considered part of the Fe uptake mechanism until coumarin derived phenolics were observed in *Arabidopsis* under high pH conditions. Other plant species such as peanut (*Arachis hypogaea* L.) and rice (*Oryza sativa*) plants secrete phenylpropanoids instead of coumarins, which also facilitate the reduction of ferric Fe (Römheld and Marschner, 1983; Ishimaru et al., 2011).

Root exudates collection has been performed by a hydroponic-only system which is useful for the characterization of specific compounds, avoiding alteration by sorption process to the soil matrix and microbial decomposition (Oburger and Jones, 2018). However, soil-specific resource availability and microbiome activity are important factors affecting plant metabolism and root exudation, therefore hydroponic-only systems are less suitable to provide useful information about the metabolites released by roots (Oburger and Jones, 2018). Nevertheless, by setting up different treatments mimicking an alkaline or calcareous condition, our approach allowed us to identify differential phenolics exudation patterns in direct (-Fe) and induced Fe deficiency (Bic and Tric) conditions. However, further analyses are required to provide further details on root exudations from *P.judaica*.

In addition, however, it has been suggested that the root Fe deficiency response also includes the dynamic use of a large Fe reservoir bound to cell wall components in the root apoplast, secretion of phenolic compounds in the apoplast, and inhibition of suberization of endodermal cells in order to allow apoplastic and transcellular radial transport of Fe (Römheld and Marschner, 1983; Jin et al., 2007, 2008; Ishimaru et al., 2011; Connorton et al., 2017). Accordingly, the increased cell walls lignification, together with the high apoplastic Fe accumulation observed in Bic-treated roots are in agreement with these findings. Recently, it has been suggested that kiwifruit plants activate two different strategies to acquire and translocate Fe from the -Fe or +Bic nutrient solution (Wang et al., 2020). Under -Fe conditions, a foraging-reusing strategy increased the mobilization of Fe (by the release of hemicellulose Fe from the cell wall and the redistribution of water-soluble Fe and apoplastic Fe in roots). However, under +Bic conditions, roots employed a resisting-inactivating strategy due to the bicarbonate-mediated inhibition of Fe translocation from root to shoot, resulting in an accumulation of water-soluble and apoplastic Fe and slowing down the release of hemicellulose Fe in the cell wall (Wang et al., 2020).

The approach employed highlighted several differences between direct (-Fe) and induced Fe deficiency (Bic and Tric) treatments. Other than the presence of bicarbonate, such differences are also attributable to the high pH. Recently it has been demonstrated that the environmental pH is an important determinant of global gene expression which tunes Fe acquisition to the prevailing edaphic conditions in *Arabidopsis* plants. Under high pH Fe deficiency responses are affected, and the production and secretion of Fe-mobilizing coumarins is induced, prioritizing the most effective strategy to mobilize Fe from otherwise inaccessible pools. Furthermore, at transcriptional level, Fe-deficient plants grown at high pH displayed increased expression of genes involved in the orchestration of defence responses to pathogens (Tsai and Schmidt 2020).

Furthermore, the exudation of phenolics into the rhizosphere influences selectively some microbial soil species that produce either siderophores or auxins that support Fe acquisition by the plant (Jin et al., 2008; Stringlis et al., 2018). The overall microbiome associated with the roots of *P. judaica* differed from that of the bulk soil, indicating that plants in urban soil carry out microbial recruitment. The coumarins mechanism of shaping root-associated microbioma is mainly due to the catechol moiety of such compounds, which can mobilize Fe and produce ROS, which play a detrimental effect on the growth of some microbial genera (Voge et al., 2019). A mechanism similar to that suggested for coumarins in the rhizosphere might also occur for *P. judaica* phenolics, as they also display both the catechol moiety in their chemical structures and Fe reducing activity.

## **Acknowledgements**

We thank Dr. Veronica Lattanzio for supporting HPLC-MS analysis.

The authors declare no conflict of interest

**Funding:** This work was supported by the local research funds of the Department of Life Science and Systems Biology, University of Turin.

## **Authorship**

All the authors have made substantial contributions to conception and design, or acquisition of data, or analysis and interpretation of data (G.V, G.Z, I.M, conception and design; LT, plant growth, exudate collection, organic acid analysis; E.E, M.N, A.S, microbiome characterization and data elaboration; V.La, V.Li phenols characterization; M.A, A.S, root architecture analysis; M.D, histological analysis; G.V., S.A., soil analysis and plant nutrient analysis); G.V and I.M. drafted the manuscript. All the authors participate in revising the manuscript.

## References

- Abenavoli, M. R., Leone, M., Sunseri, F., Bacchi, M., Sorgonà, A. (2016). Root phenotyping for drought tolerance in bean landraces from Calabria (Italy). *Journal of Agronomy and Crop Science*. 202 (1): 1-12 doi: 10.1111/jac.1212
- Abenavoli, M. R., Nicolò, A., Lupini, A., Oliva, S., Sorgonà, A. (2008) Effects of different allelochemicals on root morphology of *Arabidopsis thaliana*. *Allelopathy Journal* 22: 245-252.
- Abenavoli, MR, A. Sorgonà, S. Albano, G. Cacco. (2004). Coumarin differentially affects the morphology of different root types of maize seedlings. *Journal of Chemical Ecology* 30(9): 1871-1883.
- Atkinson, JA., Rasmussen, A., Traini, R., Voß, U., Sturrock, C., Mooney S. J., Wells Darren, M., Bennett, M. J. (2014) Branching out in roots: uncovering Form, Function, and Regulation. *Plant Physiology* 166 (2) 538-550; DOI: 10.1104/pp.114.245423
- Badri, D.V., Chaparro, J.M., Zhang, R., Shen, Q., Vivanco, J.M. (2013) Application of natural blends of phytochemicals derived from the root exudates of *Arabidopsis* to the soil reveal that phenolic-related compounds predominantly modulate the soil microbiome. *Journal of Biological Chemistry* 15;288(7):4502-12. doi: 10.1074/jbc.M112.433300.
- Barberon, M., Vermeer, J.E., De Bellis, D., Wang, P., Naseer, S, Andersen, T.G., Humbel, B.M., Nawrath, C., Takano, J., Salt, D.E., Geldner, N. (2016). Adaptation of root function by nutrient-induced plasticity of endodermal differentiation. *Cell*. 164(3):447-459.
- Bartoli, G., Bottega, S., Forino, L.M.C., Ruffini Castiglione, M., Tagliasacchi, A.M., Grilli, I., Spanò, C. (2013). Morpho-physiological plasticity contributes to tolerance of *Calluna vulgaris* in an active geothermical field. *Australina Journal of Botany*, 61: 107-18
- Bechtold, U. (2018) Plant Life in Extreme Environments: How Do You Improve Drought Tolerance? *Frontiers in Plant Science* 9:543. doi: 10.3389/fpls.2018.00543
- Bolyen, E., Rideout, J.R., Dillon, M.R., Bokulich, N.A., Abnet, C.C., Al-Ghalith, G.A., Alexander, H, ... Knight, R., and Caporaso J.G. (2019). Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nature Biotechnology* 37: 852–857. <https://doi.org/10.1038/s41587-019-0209-9>
- Boyer, R.F., Clark HM, LaRoche AP. (1988). Reduction and release of ferritin iron by plant phenolics. *Journal of Inorganic Biochemistry*. 32(3):171-81. doi: 10.1016/0162-0134(88)80025-4. PMID: 3131480.
- Bulgarelli, D., Rott, M., Schlaeppi, K., Ver Loren van Themaat, E., Ahmadinejad, N., Assenza, F., Rauf, P., Huettel, B., Reinhardt, R., Schmelzer, E. (2012). Revealing structure and assembly cues for *Arabidopsis* root-inhabiting bacterial microbiota. *Nature* 488:91–95



- Campestre MP, Antonelli C, Calzadilla PI, Maiale SJ, Rodríguez AA, Ruiz O.A. (2016). The alkaline tolerance in *Lotus japonicus* is associated with mechanisms of iron acquisition and modification of the architectural pattern of the root. *Journal of Plant Physiology* 206, 40-48,
- Cantos E., Espin J.C., Tomas-Barberan F. A. (2001). Effect of wounding on phenolic enzymes in six minimally processed lettuce cultivars upon storage. *Journal of Agricultural and Food Chemistry*, 49: 322–330
- Caporaso, J. G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F. D., Costello, E. K., et al. (2010). QIIME allows analysis of high-throughput community sequencing data. *Nature Methods* 7, 335–336.
- Caretto, S., Linsalata, V., Colella, G., Mita, G., Lattanzio, V. (2015) Carbon fluxes between primary metabolism and phenolic pathway in plant tissues under stress. *International Journal of Molecular Sciences*. 16(11):26378-26394. <https://doi.org/10.3390/ijms161125967>
- Cesco, S., Neumann, G., Tomasi, N., Pinton, R., Weisskopf, L. (2010). Release of plant-borne flavonoids into the rhizosphere and their role in plant nutrition. *Plant and Soil* 329(1-2):1-25.
- Chaney, R.L., Brown J.C. & Tiffin L.O. (1972) Obligatory reduction of ferric chelates in iron uptake by soybeans. *Plant Physiology* 50, 208–213
- Chen, J., Bittinger, K., Charlson, E.S., Hoffmann, C., Lewis, J., Wu, G.D., Collman, R.G., Bushman, F.D., Li, H. (2012). Associating microbiome composition with environmental covariates using generalised UniFrac distances. *Bioinformatics* 28(16): 2106-2113.
- Chen, Y., Barak, P. (1982). Iron Nutrition of Plants in Calcareous Soils. *Advances in Agronomy*, 35: 217–240
- Cheyrier, V., Comte, G., Davies, K.M., Lattanzio, V., Martens, S. (2013). Plant Phenolics: recent advances on their biosynthesis, genetics, and ecophysiology. *Plant Physiology and Biochemistry*, 72: 1-20.
- Ciamporova, M., Dekankova, K., Ovecka, M. (1998). Intra- and interspecific variation in root length, root turnover and the underlying parameters. In: Lambers H, Poorter H, VanVuuren MMI (eds) Variation in plant growth. *Physiological mechanisms and ecological consequences*. Backhuys Publishers, Leiden, pp 57–69.
- Coleman, J.S., McConnaughay, K.D.M., Ackerly, D.D. (1994). Interpreting phenotypic variation in plants. *Trends in Ecology and Evolution* 9: 187–191.
- Connorton, J.M., Balk, J., and Rodríguez-Celma, J. (2017). Iron homeostasis in plants – a brief overview. *Metallomics* 9, 813–823.
- Csardi, G., Nepusz, T. (2006). The igraph software package for complex network research. *Inter Journal Complex Systems* 1695. <http://igraph.org>
- Curie, C., Briat, J.-F. (2003). Iron Transport and Signaling in Plants. *Annual Review of Plant Biology*, 54:183-206
- Curie, C., Mari, S. (2017) New routes for plant iron mining. *The New Phytologist* 214: 521–525.

de Vries, F. T., Williams, A., Stringer, F., Willcocks, R., McEwing, R., Langridge, H., & Straathof, A. L. (2019). Changes in root-exudate-induced respiration reveal a novel mechanism through which drought affects ecosystem carbon cycling. *The New Phytologist*, 224(1), 132–145. <https://doi.org/10.1111/nph.16001>

Dell'Orto, M., De Nisi, P., Pontiggia, A., Zocchi, G. (2003). Fe Deficiency Responses in *Parietaria diffusa*: A calcicole plant. *Journal of Plant Nutrition*, 26 (10-11): 2057–2068.

Díaz, I., Delgado, A., de Santiago, A., del Campillo, M.C., Torrent, J. (2012). Iron deficiency chlorosis in plants as related to Fe sources in soil, EGU General Assembly 2012, 22-27 April, Vienna-Austria, p.4454

Ding, W., Clode, P.L. & Lambers, H. (2019) Is pH the key reason why some *Lupinus* species are sensitive to calcareous soil? *Plant Soil* 434, 185–201. <https://doi.org/10.1007/s11104-018-3763-x>

Donnini S, Dell'Orto M, Zocchi G (2011). Oxidative stress responses and root lignification induced by Fe deficiency conditions in pear and quince genotype. *Tree Physiol*, 31: 102-113.

Donnini S., De Nisi P., Gabotti D., Tato L., Zocchi G. (2012). Adaptive strategies of *Parietaria diffusa* (M.&K.) to calcareous habitat with limited iron availability. *Plant Cell and Environment* 35(6):1171–1184

Donnini S., Castagna A., Ranieri A. & Zocchi G. (2009) Differential responses in pear and quince genotypes induced by Fe deficiency and bicarbonate. *Journal of Plant Physiology* 166, 1181–1193.

Dubrovsky, J.G. and Forde, B.G. (2012). Quantitative analysis of lateral root development: pitfalls and how to avoid them. *Plant Cell* 24(1): 4-14. doi:10.1105/tpc.111.089698.

Edgar, R.C., Haas, B.J., Clemente, Quince, C., Knight, R. (2011). UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics* 27(16): 2194–2200.

Edwards J, Johnson C, Santos-Medellín C, Lurie E, Podishetty NK, Bhatnagar S, Eisen JA, Sundaresan V. (2015). Structure, variation, and assembly of the root-associated microbiomes of rice. *Proc Natl Acad Sci* Feb 24;112(8):E911-20. doi: 10.1073/pnas.1414592112.

Faria JC, Jelihovschi EG, Allaman Bezerra I. (2016). conventional Tukey test. UESC, Ilheus, Brasil.

Fourcroy, P., Sisó-Terraza, P., Sudre, D., Savirón, M., Reyt, G., Gaymard, F., Abadía, A., Abadía, J., Alvarez-Fernández, A., Briat, J.F. (2014). Involvement of the ABCG37 transporter in secretion of scopoletin and derivatives by *Arabidopsis* roots in response to iron deficiency. *New Phytologist* 201: 155–167.

Guerinot M.L., and Yi Y. (1994). Iron: Nutritious, Noxious, and Not Readily Available. *Plant Physiology*, 104: 815–820

Harbort, C.J., Hashimoto, M., Inoue, H., Niu, Y., Guan, R., Rombolà, AD., Kopriva S, Voges M.J.E.E.E., Sattely, E.S., Garrido-Oter, R., Schulze-Lefert, P. (2020). Root-secreted coumarins and the microbiota interact to improve iron nutrition in *Arabidopsis*. *Cell Host Microbe*. Dec 9;28(6):825-837.e6. doi: 10.1016/j.chom.2020.09.006.

- Hartman, K., van der Heijden, M.G.A., Wittwer, R.A. *et al.* (2018). Cropping practices manipulate abundance patterns of root and soil microbiome members paving the way to smart farming. *Microbiome* 6, 14 doi:10.1186/s40168-017-0389-9
- Hell, R., Stephan, U.W. (2003) Iron uptake, trafficking and homeostasis in plants. *Planta* 216: 541–551
- Henkhaus, N., Barlett, M, Gang, D., Grumet, R, Jordan-Thaden, I., Lerence, A., Lyons, E., Miller, S., et al., (2020) Plant Science decadal vision 2020-2030: reimagining the potential of plants for a healthy and sustainable future. *Plant Direct* 00:1-24 doi:10.1002/pld3.252
- Henry, A., Cal, A.J., Batoto, T.C., Torres, R.O., and Serraj, R. (2012). Root attributes affecting water uptake of rice (*Oryza sativa*) under drought. *Journal of Experimental Botany* 63: 4751–4763
- Ho, M.D.; Rosas, J.C.; Brown, K.M., Lynch, J.P. (2005). Root architectural tradeoffs for water and phosphorus acquisition. *Functional Plant Biology* 32: 737–748
- Hu, H., Tang, C, Rengel, Z. (2005). Influence of phenolic acids on phosphorus mobilization in acidic and calcareous soils. *Plant and Soil* 268:173-180.
- Hummel, I., Vile, D., Violle, C., Devaux, J., Ricci, B., Blanchard, A., Garnier, E., Roumet, C. (2007). Relating root structure and anatomy to whole-plant functioning in 14 herbaceous Mediterranean species. *New Phytologist*. 173 (2): 313–321.
- Ihrmark, K., Bödeker, I.T.M., Cruz-Martinez, K., Friberg, H., Kubartova, A., Schenck, J., Strid, Y., Stenlid, J., Brandström-Durling, M., Clemmensen, K.E., Lindahl, B.D. (2012). New primers to amplify the fungal ITS2 region – evaluation by 454-sequencing of artificial and natural communities, *FEMS Microbiology Ecology*, 82 (3): 666-677.
- Isah, T. (2019). Stress and defense responses in plant secondary metabolites production. *Biological Research*. 52: 1–25.
- Ishimaru, Y., Kakei, Y., Shimo, H., Bashir, K., Sato, Y., Sato, Y., Uozumi, N., Nakanishi, H., Nishizawa, N.K. (2011). A rice phenolic efflux transporter is essential for solubilizing precipitated apoplasmic iron in the plant stele. *Journal of Biological Chemistry* 286 (28): 24649–24655.
- Izaguirre, M.M., Mazza, C.A., Svatos, A., Baldwin, I.T., Ballare, C.L., (2007). Solar ultraviolet-B radiation and insect herbivory trigger partially overlapping phenolic responses in *Nicotiana attenuata* and *Nicotiana longiflora*. *Annals of Botany*, 99:103–109
- Jin, C.W., You G.Y., Zheng, S.J. (2008).The iron deficiency-induced phenolics secretion plays multiple important roles in plant iron acquisition underground. *Plant Signaling & Behavior*, 3(1): 60-61
- Jin, CW, You, GY, He, YF, Tang, CX, Wu, P, Zheng , SJ (2007) Iron deficiency-induced secretion of phenolics facilitates the reutilization of root apo-plastic iron in red clover. *Plant Physiology* 144:278–285.
- Jansenn, P.H. (2006)Identifying the Dominant Soil Bacterial Taxa in Libraries of 16S rRNA and 16S rRNA Genes Applied and Environmental Microbiology Vol. 72, No. 3 <https://doi.org/10.1128/AEM.72.3.1719-1728.2006>

Kalayu, G. (2019). Phosphorus solubilising microorganisms: promising approach as biofertilizers. *International Journal of Agronomy* ID 4917256, <https://doi.org/10.1155/2019/4917256>

Kim S.A., Guerinot M.L. (2007). Mining iron: Iron uptake and transport in plants. *FEBS letters*, 581(12): 2273–2280.

Kobayashi, T, Nishizawa, NK. (2012). Iron Uptake, Translocation, and Regulation in Higher Plants. *Annual Review of Plant Biology* 63: 131-152.

Kono, Y, Kashine, S, Yoneyama, T, Sakamoto, Y, Matsui, Y, Shibata, H. (1998). Iron chelation by chlorogenic acid as a natural antioxidant. *Bioscience Biotechnology Biochemistry*.62(1): 22-7. doi: 10.1271/bbb.62.22. PMID: 9501514.

Ladygina, N, Hedlund, K. (2010). Plant species influence microbial diversity and carbon allocation in the rhizosphere. *Soil Biology & Biochemistry* 42: 162–168.

Lattanzio, V. (2019) relationship of phenolic metabolism to growth in plant and cell cultures under stress. In: K.G. Ramawat, H.M. Ekiert, S. Goyal (eds.), plant cell and tissue differentiation and secondary metabolites, reference series in Phytochemistry, pp 1-32. Springer Nature Switzerland AG.

Lattanzio, V., Van Sumere, C.F. (1987). Changes in phenolic compounds during the development and cold storage of artichoke (*Cynara scolymus* L. heads. *Food Chemistry*, 24(1):37

Lattanzio, V., Cardinali, A., Di Venere, D., Linsalata, V., Palmieri, S. (1994). Browning phenomena in stored artichoke (*Cynara scolymus* L.) heads: enzymic or chemical reactions? *Food Chemistry*, 50: 1-7

Lattanzio, V., Di Venere, D., Linsalata, V., Bertolini, P., Ippolito, A., Salerno, M. (2001). Low temperature metabolism of apple phenolics and quiescence of *Phlyctanea vagabunda*. *Journal of Agricultural Food Chemistry*, 49 (12), 5817-5821.

Lindsay, W.L., Schwab, A.P. (1982). The chemistry of iron in soils and its availability to plants. *Journal of Plant Nutrition* 5: 821–840.

Lupini, A, Sorgonà, A, Princi, M.P., Sunseri, F., Abenavoli, M. R. (2016) Morphological and physiological effects of trans-cinnamic acid and its hydroxylated derivatives on maize root types. *Plant Growth Regulation* 78 (2): 263-273. 0.1007/s10725-015-0091-5.

McMurdie, P.J., Holmes, S. (2013). Phyloseq: An R Package for Reproducible Interactive Analysis and Graphics of Microbiome Census Data. *PLoS ONE*. 8(4): e61217.

Mehboob, I., Naveed, M., Zahir, Z.A., Ashraf, M. (2012) Potential of Rhizobia for Sustainable Production of Non-legumes. In: Ashraf M., Öztürk M., Ahmad M., Aksoy A. (eds) *Crop Production for Agricultural Improvement*. Springer, Dordrecht. [https://doi.org/10.1007/978-94-007-4116-4\\_26](https://doi.org/10.1007/978-94-007-4116-4_26)

Miguel, MA, Postma, J.A, Lynch J.P. (2015) phenic synergism between root hair length and basal root growth angle for phosphorus acquisition. *Plant Physiology*, 167 (4) 1430-1439; DOI: 10.1104/pp.15.00145

Mengel, K. (1994). Iron availability in plant tissues—iron chlorosis on calcareous soils. *Plant and Soil*, 165: 275–283

Mimmo, T., Ghizzi, M., Marzadori, C. et al. (2008) Organic acid extraction from rhizosphere soil: effect of field-moist, dried and frozen samples. *Plant Soil* 312, 175–184. <https://doi.org/10.1007/s11104-008-9574-8>

Oksanen, J., Blanchet, F.G., Kindt, R., Legendre, P., Minchin, P.R., O’Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Wagner, H. (2013). Vegan: Community Ecology Package. [WWW document] URL <http://cran.r-project.org/package=vegan>

Osmond, C.B., Austin, M.P., Berry, J.A., Billings, W.D., Boyer, J.S., Dacey, J.W.H., Nobel, P.S., Smith, S.D., Winner, W.E. (1987). Stress physiology and the distributions of plants, *BioScience* 37: 38-48.

Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M., Prettenhofer, P., Weiss, R., Dubourg, V., Vanderplas, J., Passos, A., Cournapeau, D., Brucher, M., Perrot, M., and Duchesnay, E. (2011). Scikit-learn: machine learning in Python. *Journal of machine learning research*, 12:2825–2830

Pérez-Jaramillo, J.E., Mendes, R., Raaijmakers, J.M. (2016). Impact of plant domestication on rhizosphere microbiome assembly and functions. *Plant Molecular Biology* 90(6):635-44. doi: 10.1007/s11103-015-0337-7.

Oburger E. and Jones DL (2018) Sampling root exudates-mission impossible? *Rhizosphere* 6 (2018) 116–133

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., et al. (2012). The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Research* 41: 590-596.

Rajniak, J., Giehl, R.F., Chang, E., Murgiam, I., von Wirén, N., Sattely, E.S. (2018) Biosynthesis of redox-active metabolites as a general strategy for iron acquisition in plants. *Nature Chemical Biology* 442-450. doi:10.1038/s41589-018-0019-2

Rangarajan, H., Postma J.A., Lynch J.P. (2018) Co-optimisation of axial root phenotypes for nitrogen and phosphorus acquisition in common bean. *Annals of Botany* 122:485–499, <https://doi.org/10.1093/aob/mcy092>

Ranjani, A., Dhanasekaran, D., Gopinath, P.M. (2016). An Introduction to Actinobacteria, 11th 2016 DOI: 10.5772/62329

Rodríguez-Celma, J., Vázquez-Reina, S., Orduna, J, Abadía, A., Abadía, J., Álvarez-Fernández, A., López-Millán, A.F. (2011) Characterization of flavins in roots of Fe-deficient strategy I plants, with a focus on *Medicago truncatula*. *Plant Cell Physiology* 52: 2173–2189

Roman-Aviles, B., Snapp, S.S., and Kelly, J.D. (2004) Assessing root traits associated with root rot resistance in common bean. *Field Crops Research*. 86: 147–156.

- Romano, A., Sorgonà, A., Lupini, A., Araniti, F., Stevanato, P., Cacco, G., Abenavoli, M. R. (2013) Morpho-physiological responses of sugar beet (*Beta vulgaris* L.) genotypes to drought stress. *Acta Physiologia Plantarum*. 35: 853-865, DOI: 10.1007/s11738-012-1129-1
- Römheld V. and Marschner, H. (1983) Mechanism of iron uptake by peanut plants. I. Fe<sup>III</sup> reduction, chelate splitting, and release of phenolics. *Plant Physiology* 71: 949-955.
- Römheld, V. (1987). Different strategies for iron acquisition in higher plants. *Physiologia Plantarum*, 70(2): 231–234.
- Ryser, P., Lambers, H. (1995). Root and leaf attributes accounting for the performance of fast- and slow-growing grasses at different nutrient supply. *Plant and Soil* 170: 251–265.
- Sasse, J., Martinoia, E., Northen, T. (2018). Feed Your Friends: Do Plant Exudates Shape the Root Microbiome? *Trends Plant Science*. 23(1):25-41. doi: 10.1016/j.tplants.2017.09.003. Epub 2017 Oct 17. PMID: 29050989.
- Sathya, A., Rajendran Vijayabharathi, R., Gopalakrishnan, S, (2017). Plant growth-promoting actinobacteria: a new strategy for enhancing sustainable production and protection of grain legumes, *Biotechnology* 7(2):102. doi: 10.1007/s13205-017-0736-3.
- Schenkeveld, W.D.C., Kraemer, S.M. (2018). Constraints to synergistic Fe mobilization from calcareous soil by a phytosiderophore and a reductant. *Soil Systems*: 4: 67. <https://doi.org/10.3390/soilsystems2040067>
- Schmid, N.B., Giehl, R.F.H., Döll, S., Mock, H-P, Strehmel, N., Scheel, D., Kong, X., Hider, R.C., von Wirén, N. (2014) Feruloyl-CoA 6'-Hydroxylase1-dependent coumarins mediate iron acquisition from alkaline substrates in *Arabidopsis*. *Plant Physiology* 164: 160-172; DOI: 10.1104/pp.113.228544
- Sorgonà, A., Abenavoli, M.R, Gringeri, P.G. and Cacco, G. (2007). Comparing morphological plasticity of root orders in slow- and fast-growing citrus rootstocks supplied with different nitrate levels. *Annals of Botany*, 100:1287-1296 - doi: 10.1093/aob/mcm207.
- Stevanato, S., Gui, G., Cacco, G., Biancardi, E., Abenavoli, M.R., Romano, A., Sorgonà, A. (2013) Morpho-physiological traits of sugar beet exposed to salt stress. *International Sugar Journal* 115: 800-809.
- Stringlis, I.A., Yu, K., Feussner, K., de Jonge, R., Van Betum, S.V., Van Vek, M.C., Berendsen, R.L., Bakker, P.A.H.M., Feussner, I., Pieterse, C.M.J. (2018) MYB2-dependent coumarin exudation shapes root microbiome assembly to promote plant health. *Proceedings of the National Academy of Sciences* 115: E5213-E5222
- Takahashi, S., Tomita, J., Nishioka, K., Hisada, T., Nishijima, M. (2014). Development of a prokaryotic universal primer for simultaneous analysis of Bacteria and Archaea using next-generation sequencing. *PLoS One* 9 (8): e105592.
- Tato L, De Nisi P, Donnini S, Zocchi G. Low iron availability and phenolic metabolism in a wild plant species (*Parietaria judaica* L.). *Plant Physiology and Biochemistry*. 2013 72:145-53. doi: 10.1016/j.plaphy.2013.05.017

- Tato, L., Islam, M., Mimmo T, Zocchi G and Vigani G (2020) Temporal responses to direct and induced iron deficiency in *Parietaria judaica*. *Agronomy* 10: 1037; doi:10.3390/agronomy10071037
- Tedersoo, L., Bahram, M., Põlme, S., Kõljalg, U., Yorou, N.S., Wijesundera, R., Villarreal-Ruiz, L., ..., Abell S, Abarenkov K. (2014). Global diversity and geography of soil fungi. *Science* 346: 1078.
- Tellah, S., Badiani, M., Trifilò, P., Lo Gullo, M.A., Ounane, G., Ounane, S.M., Sorgonà, A. (2014) Morpho-physiological traits contributing to water stress tolerance in a peanut (*Arachis hypogaea* L.) landraces collection from the Algerian Maghreb. *Agrochimica*. 58: 126-147.
- Terés, J., Busoms, S., Perez Martín, L., et al. (2019) Soil carbonate drives local adaptation in *Arabidopsis thaliana*. *Plant Cell Environment*.;42:2384–2398. <https://doi.org/10.1111/pce.13567>
- Tsai and Schmidt. pH-dependent transcriptional profile changes in iron-deficient *Arabidopsis* roots *BMC Genomics* (2020) 21:694 <https://doi.org/10.1186/s12864-020-07116-6>
- Tyler, G. (2003) Some ecophysiological and historical approaches to species richness and calcicole/calcifuge behaviour—contribution to a debate. *Folia Geobotanica* 38:419–428
- Vigani, G., Murgia, I. (2018). Iron-Requiring enzymes in the spotlight of oxygen. *Trends in Plant Science* 23: 874-882. doi: 10.1016/j.tplants.2018.07.005.
- Voges, M.J.E.E., Bai, Y., Schulze-Lefert, P., Sattely, E.S. (2019). Plant-derived coumarins shape the composition of an *Arabidopsis* synthetic root microbiome. *Proceedings of the National Academy of Sciences* 116: 12558-12565.
- Vives-Peris, V., López-Climent, M.F, Pérez-Clemente, R.M., Gómez-Cadenas, A. (2020) Root involvement in plant responses to adverse environmental conditions. *Agronomy* 10(7):942. <https://doi.org/10.3390/agronomy10070942>
- Wahl, S., Ryser, P. (2000). Root tissue structure is linked to ecological strategies of grasses. *New Phytologist*. 148: 459–471.
- Wang, N., Dong, X., Chen, Y., Ma, B., Yao, C., Ma, F., Liu, Z. 2020. Direct and Bicarbonate-induced iron deficiency differentially affect iron translocation in kiwifruit roots. *Plants* 9, 1578 doi:10.3390/plants9111578
- White, P. J., George, T. S., Dupuy, L. X., Karley, A. J., Valentine, T. A., Wiesel, L., Wishart, J. (2013). Root traits for infertile soils. *Frontiers in Plant Science*, 4: 193. <https://doi.org/10.3389/fpls.2013.00193>
- Yilmaz, P., Parfrey, L. W., Yarza, P., Gerken, J., Pruesse, E., Quast, C., et al. (2013). The SILVA and “all-species living tree project (LTP)” taxonomic frameworks. *Nucleic Acids Research*. 42: 643-648.
- York, L.M., Nord, E.A., Lynch, J.P. (2013) Integration of root phenes for soil resource acquisition. *Frontiers in Plant Science* 4:355. doi: 10.3389/fpls.2013.00355
- Zadworny, M., Eissenstat, D.M. (2011) Contrasting the morphology, anatomy and fungal colonization of new pioneer and fibrous roots. *New Phytologist*, 190: 213-221. <https://doi.org/10.1111/j.1469-8137.2010.03598.x>

Zajc, J., Gunde-Cimerman, N. (2018). The Genus *Walleimia*-From Contamination of Food to Health Threat. *Microorganisms*. May 21;6(2):46. doi: 10.3390/microorganisms6020046. PMID: 29883408; PMCID: PMC6027281.



## Figure legends

**Figure 1.** Venn diagrams of the distribution of prokaryotic (A) and fungal (B) OTUs detected in bulk soil and rhizosphere samples. For each panel the number of identified OTUs in the bulk soil (in orange), and in the rhizosphere (in blue) are reported, as well as the number of overlapping OTUs.

**Figure 2.** Shifts in the microbial composition between the bulk soil and the rhizosphere microbiome at the phylum level. The bars show the relative abundance of each phylum on the overall microbial composition for the fungal (upper panel) and prokaryotic (lower panel) microbiome. Asterisks show significant differences in the bulk soil vs rhizosphere composition for each phylum displayed (p-value < 0.05). Eight independent biological replicates were considered (n=8).

**Figure 3.** Shifts in the microbial composition between the bulk soil and the rhizosphere microbiome at the genus level. The bars show the relative abundance on the overall microbial composition for the fungal and prokaryotic genera considered (n=8). Asterisks show significant differences in the bulk soil vs rhizosphere composition for each genus displayed (p-value < 0.05). Note: the Rhizobium group comprises *Allorhizobium*, *Neorhizobium*, *Pararhizobium* and *Rhizobium*, as they are considered as a single genera according to the Silva taxonomy. Eight independent biological replicates were considered (n=8).

**Figure 4.** Co-occurrence network (meta-network) visualising correlations between prokaryotic and fungal OTUs in the bulk soil and rhizosphere communities. The sensitive OTUs shown in green and blue represent the OTUs identified as indicator species for the rhizosphere and bulk soil condition, respectively (listed in supplemental file 3). Red triangles represent the Keystone OTUs (listed in supplemental file 2, red triangles) are also represented. Modules are defined as areas that show a high density of connections among OTUs. The Gray symbols represent “hot spots” of overlapping OTUs.

**Figure 5.** Phosphorous (P) and iron (Fe) content in root and leaf tissues of *P. judaica* grown in +Fe, -Fe, Bic, and Tric treatments. Different letters correspond to significant differences among means (P<0.05; Tukey test), n=3.

**Figure 6.** Morphological analysis of the whole root system of *P. judaica* grown in +Fe, -Fe, Bic, and Tric media. A) Images captured of root; B) Total root length and total root surface; C) Root length ratio (RLR) and its components, i.e. root mass ratio RMR, fineness F and tissue density ratio TD; D)

Morphological analyses intra-root (LR, lateral roots, AR, adventitious roots); E) Root branching analysis of *P. judaica* (root branching zone's length (BZ), lateral root formation zone (LRFZ)). Different letters correspond to statistically significant differences among mean values ( $P < 0.05$ ; Tukey test),  $n=3$ .

**Figure 7.** A) Score and loading plots of principal component analysis of root traits from *P. judaica* plants exposed to +Fe, -Fe, Bic and Tric treatments. The proportion of variability explained by each PC is given within the bracket. The ellipses denote the grouping of the samples after Hierarchical Cluster Analysis (Ward's method with distance measure by squared Euclidean distance). Correlation between plant dry weight and PC1 (B) and PC2 (C) in *P. judaica* plants exposed to different treatments [+Fe, -Fe, Bic and Tric]. The coefficient of determination and p-values are reported.

**Figure 8.** Profile of caffeoylquinic acid derivatives identified in extracted fraction from tissues (left panel) and exudates fraction (right panel) of root of *P. judaica* grown hydroponically in +Fe, -Fe, Bic, and Tric treatments (A). Profile of caffeoylquinic acid derivatives identified in *P. judaica* harvested from the soil (urban soil) is reported in B. Percentage Pie chart is related to a representative experiment with three independent replicates ( $n=3$ ).

**Figure 9.** A) Fe (III) reduction activity of root exudates, caffeic and citric acids; B) Fe content in the root apoplast fraction and C) lignin visualisation (red color) in root cross sections of *P. judaica* grown in +Fe, -Fe, Bic, and Tric treatments (20x and 40x magnification). Different letters correspond to significant differences among means ( $P < 0.05$ ; Tukey test),  $n=3$ .

## Supplementary figures

**Figure S1.** Rarefaction curves showing the reaching of a satisfactory sequencing depth for each of the sequenced samples. Upper panel: prokaryotic libraries; lower panel: fungal libraries.

**Figure S2.** Alpha indexes showing the OTUs diversities between rhizosphere and bulk soil.

**Figure S3.** Weighted Generalized Unifrac analysis showing the intra-groups diversity of the Prokaryotic (upper diagram) and Fungal (lower diagram) microbial communities between the two conditions considered (BS= Bulk soil; RH= Rhizosphere samples).

**Figure S4.** Co-occurrence networks visualising the positive, significant correlations ( $\rho > 0.7$  and  $p < 0.001$ ) among prokaryotic (upper diagram) and fungal OTUs (lower diagram) from the Rhizosphere and the Bulk soil microbial communities. The sensitive OTUs shown in green and blue represent the OTUs identified as indicator species for the rhizosphere and bulk soil condition, respectively (listed in supplemental file 3). Red squares represent the keystone OTUs (listed in supplemental file 2). Modules are defined as areas that show a high density of connections among OTUs. The Gray symbols represent “hot spots” of overlapping OTUs.

**Figure S5.** Dendrogram of Hierarchical Cluster Analysis of the scores of the PCA using the Ward's method with distance measure by squared Euclidean distance.

**Figure S6.** Characterization of carboxylic acids concentrations in root tissues (upper panel) and root exudates (lower panel) of *P. judaica* grown in +Fe, -Fe, Bic, and Tric treatments. Different letters correspond to significant differences among mean values ( $P < 0.05$ ; Tukey test),  $n=3$ .

**Figure S7.** Quantification of the phenolics compounds by Arnowns' reagent (A) and by HLPC-DAD (B) approaches. Different letters correspond to significant differences among means ( $P < 0.05$ ; Tukey test),  $n=3$

**Figure S8.** Content of phosphate anions in soluble fraction of urban soils after incubation with caffeic and citric acids. Different letters correspond to significant differences among mean values ( $P < 0.05$ ; Tukey test),  $n=3$ .

**Table 1** – Principal components of root traits of rooted cuttings of *Parietaria judaica* exposed to different treatments [+Fe, -Fe, Bic and Tric].

	<b>Attribute loadings</b>	
	<i>PC1</i>	<i>PC2</i>
<i>Statistics</i>		
<i>Eigenvalue and variability</i>		
<b>Eigenvalue</b>	2.474	1.919
<b>Proportion of variability (%)</b>	49	38
<i>Variable</i>		
<i>Eigenvectors</i>		
<b>Tissue density (TD)</b>	<b>-.867</b>	.071
<b>Total length lateral roots (LI)</b>	.192	<b>.959</b>
<b>Average length adventitious roots (aLA)</b>	<b>.926</b>	.073
<b>Average length lateral roots (aLI)</b>	-.074	<b>.981</b>
<b>Root branching zone (BZ)</b>	<b>.907</b>	.165