

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Malignant peritoneal mesothelioma in a boar who lived in Calabria (Italy): wild animal as sentinel system of human health

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1703868> since 2019-06-03T15:45:37Z

Published version:

DOI:10.1016/j.scitotenv.2019.05.254

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 **Malignant peritoneal mesothelioma in a boar who lived in Calabria (Italy): wild animal**
2 **as sentinel system of human health**

3 Elena Colombino^{a*}, Silvana Capella^{b*}, Francesco Casalinuovo^c, Rocco Racco^d, Flavia
4 Pruiti^e, Marco Volante^f, Vincenzo Di Marco Lo Presti^e, Elena Belluso^b, Maria Teresa
5 Capucchio^a

6 *equally contribution.

7 ^a Department of Veterinary Science, University of Torino, Italy

8 ^b Department of Earth Sciences and Interdepartmental Centre for Studies on Asbestos and
9 Other Toxic Particulates G. Scansetti, University of Torino, Italy.

10 ^c Istituto Zooprofilattico Sperimentale del Mezzogiorno, Catanzaro, Italy

11 ^d ASP - Reggio Calabria, Italy

12 ^e Istituto Zooprofilattico Sperimentale della Sicilia, Barcellona P.G. (Messina), Italy

13 ^f Department of Oncology, San Luigi Hospital, University of Torino, Italy

14

15

16 Corresponding author:

17 Elena Colombino, Department of Veterinary Sciences, University of Turin, Largo Paolo

18 Braccini 2, Grugliasco 10095, Turin, Italy. Tel: +39-0116709035. Fax: +390116709031. E-

19 mail: elena.colombino@edu.unito.it

20

21

22

23

24 **Abstract**

25 Mesothelioma is a tumor of the serosal membranes described both in human and veterinary
26 medicine. While in humans the relationship between mesothelioma and exposure to
27 asbestos and some other asbestiform minerals is well known, in animals it is still difficult to
28 establish. In this paper a case of malignant peritoneal mesothelioma probably related to
29 asbestos exposure in a wild boar is described. At post-mortem evaluation the peritoneum,
30 diaphragm and serosal surface of liver and kidneys showed isolated to coalescent multiple
31 nodular lesions. Samples from diaphragm, liver and lung were collected to perform
32 microbiological and histological investigations. To assess the presence of asbestos and/or
33 other asbestiform minerals, SEM-EDS investigations were performed on organs and soil
34 samples collected from the area where the wild boar lived.

35 Microbiological investigations were negative for *Mycobacterium* species. Gross and
36 histological examination were compatible with a biphasic mesothelioma, with nodules
37 composed of epithelioid and sarcomatoid elements with high pleomorphism.
38 Immunohistochemistry revealed only multifocal scattered positivity for WT-1 and D2-40.
39 Asbestos fibres were detected in all samples (organs and soil) by SEM-EDS, demonstrating
40 a potential relationship between the neoplasia and the exposure to naturally occurring
41 asbestos (NOA).

42 In conclusion, the results of the present study are further confirmation that wild animals,
43 such as the boar, are suitable sentinels to indicate the risk of environmental exposure to
44 asbestos for human populations.

45

46

47 **Key words:** mesothelioma, asbestos environmental exposure, boar, sentinel animals

48

49 **Conflict of interest: none**

50

51 **Funding**

52 This research did not receive any specific grant from funding agencies in the public,
53 commercial, or not-for-profit sectors.

54

55 **1. Introduction**

56 Malignant mesothelioma (MM) is a tumor of serosal membranes including pleura,
57 pericardium, peritoneum and tunica vaginalis testis. In humans, MM and its strong
58 correlation with occupational or anthropogenic environmental exposure (at a high dose) to
59 asbestos and other asbestiform minerals has been well documented and confirmed (IARC,
60 2012) and makes it an epidemiological marker (Pavlisko and Sporn, 2014). It is more difficult
61 to establish effects on human health resulting from chronic low-dose exposures as in the
62 case of a natural environmental background, a very common situation.

63 The use of asbestos began in ancient times and reached the height during the second half
64 of the 20th century. Actually mining and use of asbestos minerals remain high in a limited
65 number of states in the world (USGS, 2018). Asbestos minerals are present in many
66 environments, i.e. rocks, soils, air and water. Rocks are their primary source and rocks
67 containing asbestos are widespread worldwide. Owing to natural and/or anthropogenic
68 action asbestos is dispersed from these sources and spread in the environment.

69 The term “asbestos” covers a group of six natural minerals (hydrated silicates), one
70 belonging to the serpentine group and the remaining five to fibrous amphiboles (IARC,
71 2012).

72 Owing to the lack of a recognized definition, according to the dimensional definition by World
73 Health Organization (WHO, 1997) and the literature (Belluso et al., 2017), the meaning of

74 the words used in the paper is shown below. The terms "fibre" and "fibrous" are used to
75 address inorganic particles with a length $\geq 5 \mu\text{m}$, a width $\leq 3 \mu\text{m}$, and a length/width ratio \geq
76 3:1 which present parallel sides when seen in two dimensions, perpendicularly to fibre axis.
77 Asbestiform is an adjective used for non-asbestos-classified fibres with asbestos-fibre-like
78 dimensions and at least one of the asbestos properties such as flexibility or splitting. Fibre
79 bundle is used for a parallel aggregate of mineral fibre.

80 The most widely known fibrous variety of serpentine is chrysotile which is the most
81 commonly used asbestos in industrial production as component of asbestos containing
82 materials (ACM). Chrysotile is abundant as naturally occurring asbestos (NOA), i. e. in rocks
83 . Other asbestiform serpentine varieties are asbestiform antigorite and asbestiform
84 polygonal serpentine, which are difficult to distinguish from chrysotile and often confused
85 with it, depending on the kind of investigation (Belluso *et al.*, 2017). Other commercially
86 common asbestos minerals include the amphiboles crocidolite and amosite (also named
87 cummingtonite-grunerite asbestos), tremolite asbestos, actinolite asbestos and
88 anthophyllite asbestos (NIOSH, 2011).

89 Apart from these five asbestos listed above, there are also certain asbestiform amphibole
90 not asbestos classified as, for example, asbestiform fluoredenite (detected in Italy) and
91 asbestiform winchite (detected in USA). Correlation between a high dose exposure to these
92 minerals and MM was recognized through epidemiological investigations in humans affected
93 by MM (McDonald *et al.*, 2004; Paoletti *et al.*, 2000).

94 The presence of these asbestiform minerals was identified only recently, either because
95 they had not been detected in raw materials or because they had been confused with
96 other minerals, such as tremolite asbestos (Meeker *et al.*, 2003).

97

98 Owing to natural and/or anthropogenic action, asbestos are dispersed from these sources
99 and spread in various types of environments. The same origin and the same fate concern

100 other asbestiform minerals not currently classified as “asbestos” that, at the current state of
101 knowledge, are present only in some areas of the world (Cannata et al., 2018).

102 Diffusion of fibrous minerals (whether classified as asbestos or not) in non-occupational
103 environments has been highlighted in the last twenty years by their detection in lungs of
104 animals. In non-experimental animals, MM is generally rare. It has been more frequently
105 described in dogs (Forbes and Matthews, 1991; Glickman et al., 1983;) and cattle (Girard
106 and Cécire, 1995; Klopfer et al., 1983; Zurwieden et al., 1990) but occasionally also in cats
107 (Umphlet and Bertoy, 1988), lambs (Brown and Weaver, 1981), horses (Mair et al., 1992;
108 Ricketts and Peace, 1976), goats (Campopiano et al., 2017) and pigs (Beytut, 2002; Uzal et
109 al., 2016).

110 Animals have been proposed as sentinel system (SSA) for different aerial pollutants
111 including asbestos to complement human epidemiological studies and to demonstrate the
112 interactive effects of these mineral fibres and their role in biological response (Dumortier et
113 al., 2002; Rey et al., 1994). In fact, animals are not affected by occupational exposure, their
114 lungs are easier to obtain for post-mortem examination than human ones and, finally, they
115 can provide an assessment of the kind and amount of the respirable minerals present in the
116 environment where they lived. Therefore, SSA could be indicators of environmental
117 background exposure to inorganic fibres (asbestos and non-asbestos) (Ardizzone et al.,
118 2014; Capella et al., 2017a; Capella et al., 2017b; De Nardo et al., 2004; Dumortier et al.,
119 2002; Fornero et al., 2009; Glickman et al., 1983).

120 In particular, MM in wild animals can be suitable to provide useful indications for
121 understanding the possible correlation between low-dose exposure and carcinogenesis.
122 Regarding the investigated Italian area (Calabria region, south of Italy), in 2017
123 Campopiano et al. identified tremolite asbestos fibres in the pulmonary tissue of sheep,
124 goats, and two boars that lived near disused quarries. In the Calabria region (Monte

125 Reventino, CZ) several deposits of ophiolite containing mainly tremolite both asbestiform
126 (i.e. asbestos classified) and non-asbestiform (i.e. non asbestos classified) were reported
127 (Zakrzewska, 2008). On the basis of the National Register of mesotheliomas in the same
128 region 6 cases of human tumors related to asbestos exposure were detected in 2012 (INAIL,
129 2015) confirming the presence of asbestos in the environment and the need to activate a
130 suitable method to evaluate the potential human risk. On the basis of these considerations,
131 the aim of this report is to describe the pathological findings of a case of peritoneal
132 mesothelioma detected in a wild boar (from Calabria) and relate them to the environment
133 where the boar lived.

134

135 **2. Material and methods**

136

137 **2.1 The boar**

138 A 3-year-old female wild boar of 70 Kg of weight had been killed during the hunting season
139 in Caulonia (Reggio Calabria, Calabria, Italy) and was submitted to post-mortem inspection.
140 The huntsman referred that the boar did not show any sign of sickness before death and it
141 seemed in good nutritional status.

142

143 **2.2 Microbiological, histological and immunohistochemical investigations**

144 Samples of peritoneal nodules, diaphragm, liver, kidney and lung were collected and frozen
145 to perform microbiological, histopathological and immunohistochemical investigations.

146

147 For microbiological investigations tissue samples were frozen and sent to the Istituto
148 Zooprofilattico Sperimentale of Barcellona P.G. (Messina, Italy). Samples were
149 homogenized, decontaminated with 1 volume of 4% NaOH for 30 min at 37°C, neutralized

150 with 0.067 M of phosphate buffered saline (PBS) at pH 7.2 and centrifuged for 15 min at
151 3,000 *g*. Pellets were suspended in PBS, inoculated into Lowenstein-Jensen medium (LJ)
152 (Biolife®, Italy) and LJ medium without glycerol (Biolife®, Italy), then incubated in CO₂ for
153 8 weeks at 37 °C, according to the OIE *Manual of Diagnostic Tests and Vaccines for*
154 *Terrestrial Animals* (World Organization for Animal Health, 2009).

155

156 After negative response for tuberculosis, frozen tissues were fixed in 10% buffered formalin,
157 and sent to the Department of Veterinary Sciences, University of Torino for histological
158 evaluation. Tissue samples were embedded in paraffin wax blocks, sectioned at 5 µm
159 thickness, mounted on glass slides and stained with Haematoxylin & Eosin (HE) and
160 Periodic Acid Schiff after diastase treatment.

161

162 Immunohistochemistry was performed on selected sections to characterize the neoplasia in
163 an automated system (Omnis Instrument, Dako, Agilent technologies). The characteristics
164 of the antibodies employed and the corresponding working conditions are detailed in Table
165 1. Reactions were visualized using a polymer-conjugated secondary antibody (Envision
166 Flex, Dako). Sections were counterstained with Mayer's hematoxylin. Positive control was
167 represented by human tissues known to express the different markers and negative controls
168 were carried out by omitting the primary antibodies.

169

170 **2.3 SEM-EDS investigation**

171 To assess the presence of asbestos and asbestiform minerals, and to evaluate a
172 relationship between the suspected mesothelioma and the possible exposure to asbestos,
173 the inorganic residue of lung, liver and diaphragmatic nodules samples were investigated
174 by SEM-EDS at the Department of Earth Sciences, University of Torino following the
175 procedure detailed in Belluso *et al.* (2006). In order to remove the organic fraction, 0.25 mg

176 of tissue from each sample have been chemically digested in 30 cc of sodium hypochlorite
177 (NaClO). Subsequently, the solution containing the inorganic material has been filtered on
178 a mixed cellulose ester filter with a diameter of 25 mm and porosity of 0.45 µm. The
179 dissolution of NaClO granules has been completed washing the filter with distilled water
180 preheated at 60° C. The filter has been put on a SEM stub and coated with a thin carbon
181 film.

182 Each membrane has been investigated by SEM (JEOL-IT300LV). Only inorganic particles
183 having aspect ratio >3:1 have been considered and their dimensions have been registered.
184 Their elemental chemical composition has been determined by EDS (Oxford Instrument
185 INCA Energy 2000). The inorganic fibres detected were identified by comparing their EDS
186 spectrum with those collected in the laboratory database. The distinction between chrysotile
187 and asbestiform antigorite, that have the same chemical composition, is possible only by
188 structural determination, but not by SEM-EDS technique (Capella et al., 2017a). As it
189 concerns tremolite and actinolite asbestos, they only differ for a little chemical variation
190 (Hawthorne and Oberti, 2007) that it is not possible to detect on fibres from digested tissue
191 (Goldstein et al., 2012). Therefore, the authors grouped together chrysotile with asbestiform
192 antigorite (i.e. chrysotile/asbestiform antigorite) and tremolite asbestos with actinolite
193 asbestos (i.e. tremolite/actinolite asbestos).

194 The number of detected inorganic fibres has been normalized to 1 g of dry tissue (ff/gdw)
195 as indicated by international guidelines (De Vuyst *et al.*, 1998).

196

197 Moreover, SEM-EDS investigations were performed on 5 samples of soil to assess the
198 presence of asbestos in the environment where the wild boar lived. This area is into the
199 woods, far away from footpaths (latitude: 38.371389; longitude: 16.390556). In the same
200 area there are active/abandoned quarries (Italian Institute for Environmental Protection and
201 Research -ISPRA) with some tunnels which can be used as dens by wild animals. There

202 are also some human settlement (4-5 farms) in the surrounding zone. The collection points
203 were chosen within a circular area of about 3 km² around the site where the animal was
204 felled, considered it as its home range (Figure 1). On the basis of these considerations, more
205 or less equidistant points of about 500 meters were identified from the killing point of the
206 animal. For each site, approximately 1 kg of soil to a depth of 30 cm was collected after
207 elimination of the layer of foliage and shrubs typical of the subsoil.

208 A representative portion of each soil sample was processed following the guidelines of the
209 Italian legislation (Italian Ministerial Decree of September 6, 1994). From the dried fraction
210 with diameter < 2 mm, a portion of soil was ground by using an agate pestle and mortar and
211 5 mg of obtained powder was suspended into a dispersive solution. A share of this solution
212 was filtered on a polycarbonate filter with a diameter of 25 mm and a porosity of 0.8 µm. The
213 filter was put on a SEM stub and coated with a thin carbon film. The presence of inorganic
214 fibres was evaluated on 0.1 mg of soil by SEM-EDS. Their amount was calculated to obtain
215 the value of mg/kg (ppm) as indicated by the Italian legislation (Italian Ministerial Decree of
216 September 6, 1994; the asbestos pollution being with value equal to or more than 100
217 mg/kg).

218

219 **3. Results**

220

221 **3.1 Post-mortem evaluation**

222 At post-mortem evaluation the peritoneum, surface of diaphragm and serosal surface of liver
223 and kidneys showed isolated to coalescent multiple nodular lesions varying from 2 mm to 3
224 cm in diameter from grey-white to red-yellow color depending on the amount of
225 hemorrhages. The nodules had a smooth and translucent surface (Figure 2). On the serosal
226 surface of the liver some nodules were pedunculated with a cauliflower like surface due to

227 many complex fissures. No lesions were found in the thoracic cavity. On the basis of the
228 observed lesions tuberculosis/mesothelioma was suspected.

229

230 **3.2 Microbiological, histological and immunohistochemical investigations**

231 The microbiological exam resulted negative for *Mycobacterium* spp., excluding the
232 hypothesis of Tuberculosis.

233

234 The histological features of the nodules were the same in all the samples (diaphragm, kidney
235 and liver serosal surface) and hepatic and renal parenchyma were not affected. Moreover,
236 in liver hemosiderin accumulations were observed probably suggesting previous
237 hemorrhages. Lungs showed no histological alterations.

238 Nodules were composed by a mixed cell population of epithelioid and sarcomatoid elements
239 with an high pleomorphism. The epithelioid cell population showed oval cells with abundant
240 and light-colored/eosinophilic cytoplasm, round-to-oval nuclei and well-defined cell borders
241 (Figure 3). These cells did not show reactivity with PAS stain indicating the absence of mucin
242 droplets within tumor cells or tubular lamina which were characteristic of adenocarcinoma
243 (Head, 1990). Less commonly binucleate or multinucleate cells were observed. The
244 sarcomatoid pattern was composed by pleomorphic spindled fibroblast-like cells. Areas of
245 necrosis and lymphoplasmacytic infiltrates with rare neutrophils were frequently observed.
246 Both cell populations were mixed in the nodules and tended to dispose in papillary structure,
247 nests or cords.

248

249 Most of the immunohistochemical investigations failed to help in the classification of the
250 tumor masses. In fact, only multifocal scattered WT-1 and D2-40 positivity were detected
251 (Figure 4). All the samples showed negativity for Calretinin, cytokeratin 5, HBME-1 and CEA,

252 but at least for the former three no positivity in internal control cells was observed, thus the
253 reactions were considered inconclusive.

254 Localization and morphological pattern are compatible with the diagnosis of a biphasic
255 mesothelioma containing both epithelioid and sarcomatoid components at histological
256 examination. Malignancy was supported by the presence of necrosis, increased cellularity,
257 pleomorphism and a diffuse proliferation in the connective stromal tissue supporting
258 mesothelial cells.

259

260 **3.3 SEM-EDS investigation**

261 The inorganic fibres detected in different organs at SEM-EDS investigation are reported in
262 Table 2. In lung, mineral fibres attributable to phyllosilicates and amphiboles families, have
263 been detected, both asbestos and asbestiform minerals. Within the asbestos minerals,
264 tremolite/actinolite asbestos (Figure 5–a,b), and chrysotile/asbestiform antigorite (Figure 5-
265 c,d) were detected. Regarding asbestiform minerals, hornblende, Na-Ca amphiboles and
266 illite-smectite have been detected.

267 In the diaphragmatic nodules and in the liver only chrysotile/antigorite asbestiform fibres
268 have been detected.

269 Inorganic fibres detected in soils samples are reported in Table 3. Chrysotile/asbestiform
270 antigorite is present only in 2/5 samples while fibrous amphibole species (both asbestos and
271 non-asbestos classified) have not been detected.

272 Two kinds of fibrous species, chrysotile/asbestiform antigorite and fibrous illite/smectite have
273 been detected both in biological and soil samples. In both types of samples, the amount of
274 chrysotile/asbestiform antigorite is predominant compared to the other fibrous minerals.

275

276 **4. Discussion**

277 Primary peritoneal tumors in animals are rare. One of this is represented by peritoneal
278 mesothelioma even if its diagnosis must be made with caution because of the similarity of
279 mesothelial proliferative lesions with chronic granulomatous peritonitis and metastatic
280 tumors in serous membranes (such as carcinoma) (Head, 1990).

281 In the present case Tuberculosis was suspected during post-mortem inspection as the
282 peritoneum was covered by diffuse small nodules. However, histological examination
283 revealed no granulomatous inflammatory process and microbiological investigations
284 excluded the suspect of mycobacterial infection.

285 Gross and histological examination are compatible with a biphasic mesothelioma as the
286 nodules were composed by both epithelioid and sarcomatoid mixed components (Head,
287 1990). Moreover, immunohistochemistry, although focally, was positive for WT1 e D2-40
288 which are specific markers for mesothelioma (Ordoñez, 2013) supporting the hypothesis of
289 a MM in the boar. Indeed, other tested mesothelioma-specific markers, such as calretinin,
290 cytokeratin 5 and HBME-1, were negative both in tumor cells and in internal control cells
291 (such as normal peritumoral mesothelial cells). This finding appears difficult to explain but
292 the authors assume that it can be due to bad antigen preservation in tissue samples. All the
293 samples have been frozen before chemical fixation and the type of freezing-defrosting
294 process (temperature, delay in tissue allocation) can influence the antigen preservation
295 (Pelstring, 1991).

296 In human medicine mesothelioma usually develops in the pleural cavity (Bianchi and
297 Bianchi, 2014). On the contrary, in veterinary medicine a higher number of cases of
298 peritoneal mesothelioma have been reported in literature (Bacci *et al.*; 2006; Beytut, 2002;
299 Brown and Weaver, 1981; Forbes and Matthews, 1991; Girard and Cécire, 1995; Umphlet
300 and Bertoy, 1988). This different localization could be explained by the capability of
301 pulmonary macrophages with phagocytosed dust particles to move through alveolar walls
302 towards the pleura. In doing so they may penetrate bronchioles and be transported by the

303 mucociliary staircase or they may penetrate blood vessels and be carried to extrapulmonary
304 sites where they excite a reaction for example in the liver, spleen, kidney and abdominal
305 wall (Holt, 1981). In the boar of the present case an ingestion of asbestos fibres cannot be
306 excluded.

307 The evidence of abundant asbestos fibres in lung, liver and diaphragmatic nodules with
308 SEM-EDS methods seems to support this pathological mechanism and the diagnosis of
309 mesothelioma relating to asbestos exposure.

310 The mineral fibres detected in the biological samples are compatible with the geological
311 characteristics of the area where the wild boar lived. In particular chrysotile/asbestiform
312 antigorite was detected both in biological and in soil samples. The authors cannot exclude
313 that the lack of detection of fibrous amphiboles in soil samples is due to a detection limit of
314 SEM-EDS method.

315 To the best of author knowledge, in humans tremolite asbestos accumulates in lung, while
316 chrysotile is quickly cleared. This difference in pulmonary clearance can explain why
317 tremolite asbestos and amphiboles asbestos in general are considered "2-3 orders of
318 magnitude more carcinogenic than chrysotile" (WHO, 2015). Assuming that pulmonary
319 clearance rate in wild boars is similar to human one, the authors can hypothesize that
320 probably this animal respired more chrysotile/asbestiform antigorite than tremolite/actinolite
321 asbestos, but the most persistent fibres are left the most abundant.

322 Moreover, the amount of fibres detected in the boar lungs is lower than that reported in
323 previous studies.

324 As it concerns the amount of fibres in the lung sample (Table 2), the comparison with
325 literature data shows that amphibole asbestos is much lower than that found in lungs of
326 sheep, goats and two boars from more than 100 km far from the provenance site of the
327 examined boar (Campopiano *et al.*, 2017). Similarly, goats from Corsica (Dumortier *et al.*,
328 2002), cows from north-western Italy (Belluso *et al.*, 2017), cats and dogs from California

329 (Abraham *et al.*, 2005) showed lower amount of amphiboles fibres in lung compared to the
330 boar of the present study. Also for chrysotile/asbestos antigorite, the burden in cow lungs
331 from north-western Italy is lower (Belluso *et al.*, 2006) than that found in the investigated
332 boar. These animals did not develop MM even if they had a higher fibre burden compared
333 to the boar.

334 In humans a burden of amphiboles asbestos (fibre with length of $> 5 \mu\text{m}$) higher than $1 \times$
335 $10^5/\text{gdw}$ is considered an indicator of significant asbestos exposure (De Vuyst *et al.*, 1998),
336 able to trigger cancerous pathologies. It is interesting to note that the amphiboles asbestos
337 burden in the boar is even lower than the threshold for humans.

338 This finding, correlated with the mesothelioma diagnosis, seems to support the hypothesis
339 of a cancer risk related to natural environmental exposure (low doses) to asbestos, and the
340 presence of a genetic component of susceptibility to MM (Crovella *et al.*, 2016) also in
341 animals.

342 For these reasons the role of wild boars as SSA has to be considered central in: i)
343 determining the environmental diffusion of asbestos and non-asbestos classified
344 asbestiform minerals; ii) establishing if the natural exposure to asbestos could play a role in
345 the development of human mesotheliomas.

346

347 **5. Conclusion**

348 In conclusion, the results of this study suggest that an active surveillance on regularly
349 slaughtered domestic/wild animals seems suitable to quantify the risk of exposure to
350 asbestos for human population.

351 It would be interesting to carry out diagnostic investigations also on MM cases in other
352 wild/domestic animals to confirm the hypothesis that, at least in animals, the low-dose
353 asbestos respiration, presumably continuous, and for a limited period of time (3 years) may
354 induce MM.

355

356 **References**

357

358 Abraham, J.L., Case, B.W., Burnett, B.R., Trent T., May 2005. Lung-retained fibre in animal
359 lungs confirms environmental exposure to naturally-occurring amphibole asbestos in El
360 Dorado Country, California Am Thorac Soc meeting, San Diego.

361

362 Ardizzone, M., Vizio, C., Bozzetta, E., Pezzolato, M., Meistro, S., Dondo, A., Giorgi, I.,
363 Seghesio, A., Mirabli, D., Capella, S., Vigliaturo, R., Belluso, E., 2014. The wild rat as
364 sentinel animal in the environmental risk assessment of asbestos pollution: a pilot study. *Sci*
365 *Total Environ* 479-480:31-8.

366

367 Bacci, B., Morandi, F., De Meo, M., Marcato, P.S., 2006. Ten cases of feline mesothelioma:
368 an immunohistochemical and ultrastructure study. *J Comp Pathol* 134:347-354.

369

370 Belluso, E., Bellis, D., Fornero, E., Capella, S., Ferraris, G., Coverlizza, S., 2006.
371 Assessment of inorganic fibre burden in biological samples by scanning electron
372 microscopy-energy dispersive spectroscopy. *Mikrochim Acta* 155:95–100.

373

374 Belluso, E., Cavallo, A., Halterman, D., 2017. Crystal habit of mineral fibres. *European*
375 *Mineralogical Union Notes in Mineralogy*. Gualtieri A. Eds. In: *EMU Notes in Mineralogy*,
376 *Mineral fibres: crystal, chemistry, chemical-physical properties, biological interaction and*
377 *toxicity*. 18 (3), 65-109.

378

379 Beytut, E., 2002. Metastatic sclerosing mesothelioma in a cow, *Aust Vet J* 80 (7):409-411.
380 <https://doi.org/10.1111/j.1751-0813.2002.tb10998.x>

381

382 Bianchi, C., Bianchi, T., 2014. Global mesothelioma epidemic: Trend and features. Indian
383 Int J Occup Environ Med 18:82-88. [https://doi.org/ 10.4103/0019-5278.146897](https://doi.org/10.4103/0019-5278.146897).

384

385 Brown, P.J., Weaver A.D., 1981. Malignant mesothelioma in a lamb. Vet Rec 109:159.

386

387 Campopiano, A., Cannizzaro, A., Olori, A., Angelosanto, F., Iannò, A., Bruno, M.R., Bruni,
388 B.M., Casalnuovo, F., Ciambone, L., Esposito, A., Iavicoli, S., Padova 20-22/09/2017.
389 Valutazione indiretta di fibre di amianto respirabili in aree del territorio calabrese con
390 presenza di pietra verde: primi risultati. 80°Congresso Nazionale SiMLi.

391

392 Cannata C.B., Bloise A., De Rosa R., Outdoor fibres air pollution monitoring in the Crotone
393 area (Southern Italy). J Med Earth Sci 10 (2018), 27-36.

394

395 Capella, S., Belluso, E., Bursi Gandolfi, N., Tibaldi, E., Mandrioli, D., Belpoggi, F., 2017a. In
396 vivo biological activity of mineral fibres. Gualtieri A. Ed. In: EMU Notes in Mineralogy, Mineral
397 fibres: crystal, chemistry, chemical-physical properties, biological interaction and toxicity.
398 Vol.18 (9) 307-345.

399

400 Capella, S., Fornero, E., Bellis, D., Belluso, E., 2017b. Analysis of respired amphibole fibres
401 (asbestos and non-asbestos classified): discrimination between natural and anthropogenic
402 sources using sentinel animals. Air Qual Atmos Health. Volume 10 (5): 533-42.

403

404 Crovella, S., Bianco, A.M., Vuch, J., Zupin, L., Moura, R.R., Trevisan E., Schneider, M.,
405 Brollo, A., Nicastro, E.M., Cosenzi, A., Zabucchi, G., Borelli, V., 2017. Iron signature in

406 asbestos-induced malignant pleural mesothelioma: A population-based autopsy study. J
407 Toxicol Environ Health. 79 (3):129-141.

408

409 De Nardo, P., Bruni, B., 2004. Pulmonary fibre burden in sheep living in the Biancavilla area
410 (Sicily): preliminary results. Sci Total Environ 325:51-58. [https://doi.org/
411 10.1016/j.scitotenv.2003.11.018](https://doi.org/10.1016/j.scitotenv.2003.11.018).

412

413 De Vuyst, P., Karjalainen, A., Dumortier, P., Pairon, JC., Monso, E., Brochard, P., Teschler,
414 H., Tossavainen, A., Gibbs, A., 1998. ERS Task Force Report Guidelines for mineral fibre
415 analyses in biological samples: report of the ERS working group. Eur Respir J 11:1416–
416 1426.

417

418 Dumortier, P., Rey, F., 2002. Chrysotile and Tremolite asbestos fibres in the lungs and
419 parietal pleura of Corsican goats. Occup Environ Med 59: 643-646.
420 [https://doi.org/ 10.1136/oem.59.9.643](https://doi.org/10.1136/oem.59.9.643).

421

422 Forbes, DC., Matthews BR., 1991. Abdominal mesothelioma in a dog. Can Vet J 32:176-
423 177.

424

425 Fornero, E., Belluso, E., Capella, S.& Bellis, D., 2009. Environmental exposure to asbestos
426 and other inorganic fibres using animal lung model. Sci Total Environ 407, 1010-1018.

427

428 Girard, CA., Cécyre, A., 1995. Diffuse abdominal epithelioid mesothelioma in a cow. Can
429 Vet J 36:440-441.

430

431 Glikman, L.T., Domanski, L.M., Maguire, T.G., Dubielzig, R.R., Churg, A., 1983. Mesothelioma
432 in pet dogs associated with exposure of their owners to asbestos. *Environ Res* 32; 305-313.
433

434 Goldstein, D. J., Newbury, D. E., Joy, D.C, Lyman, C.E., Echlin, P., Lifshin, E., Sawyer, I.,
435 Michael, J.R., 2012. *Scanning Electron Microscopy and X-Ray Microanalysis: Third Edition*.
436

437 Hawthorne, F.C., Oberti, R., 2007. Classification of the Amphiboles. *Reviews in*
438 *mineralogy & Geochemistry*. Hawthorne F.C, Oberti R., Della ventura G., Mottana A. Eds.
439 In: *Amphiboles: Crystal Chemistry, Occurrence, and Health Issues*. 67:55-85.
440

441 Head, K.H., 1990. Tumors of the alimentary tract. Moulton JE, Ed. In: *Tumors in Domestic*
442 *Animals*. 3rd ed. University of California Press, Los Angeles; pp. 347-435.
443

444 Holt, P.F., 1981. Transport of inhaled dust to extrapulmonary sites. *J Pathol* 133:123-129.
445

446 IARC Working Group on the Evaluation of Carcinogenic Risk to Humans, 2012. *IARC*
447 *Monographs on the Evaluation of Carcinogenic Risks to Humans*, No. 100C, Lyon (FR):
448 International Agency for Research on Cancer.
449

450 Istituto Nazionale Assicurazione Infortuni sul Lavoro (INAIL), 2015. V rapporto Registro
451 Nazionale Mesoteliomi (Re.NaM). (Accessed 04/03/19)
452 [https://www.inail.it/cs/internet/comunicazione/pubblicazioni/catalogo-generale/il-registro-](https://www.inail.it/cs/internet/comunicazione/pubblicazioni/catalogo-generale/il-registro-nazionale-dei-mesoteliomi-v-rapporto.html)
453 [nazionale-dei-mesoteliomi-v-rapporto.html](https://www.inail.it/cs/internet/comunicazione/pubblicazioni/catalogo-generale/il-registro-nazionale-dei-mesoteliomi-v-rapporto.html)
454

455 Italian Institute for Environmental Protection and Research (ISPRA). (Accessed 25/02/19)
456 <http://500.isprambiente.gov.it>

457

458 Klopfer, U., Brenner, G., Nobel, T.A., Perl, S., Yakobson, B., 1983. Mesothelioma in cattle -
459 A rare or an unidentified tumour?. *Zentbl Vet Med Reihe B* 30:785-788.

460

461 Mair, TS., Hillyer, MH., Brown, PJ., 1992. Mesothelioma of the pleural cavity in a horse:
462 diagnostic features. *Equine Vet Educ* 4:59-61.

463

464 McDonald, J.C., Harris, J., Armstrong, B., 2004. Mortality in a cohort of vermiculite miners
465 exposed to fibrous amphibole in Libby, Montana. *Occup Environ Med* 61:363–366.

466

467 Meeker, G.P., Bern, A.M., Brownfild, I.K., Lowers, H.A., Sutley, S.J., Hoefen, T.M., Vance,
468 J.S., 2003. The composition and morphology of amphiboles from the Rainy Creek Complex,
469 near Libby, Montana. *Am Mineral* 88, 1955-1969.

470

471 Ministero Italiano della Sanità, Decreto Ministeriale 06 settembre 1994: Normative e
472 metodologie tecniche di applicazione dell'art. 6, comma 3, e dell'art. 12, comma 2, della
473 legge 27 marzo 1992, n.257, relativa alla cessazione dell'impiego dell'amianto.

474

475 The National Institute for Occupational Safety and Health (NIOSH), 2011. Asbestos Fibres
476 and Other Elongate Mineral Particles: State of the Science and Roadmap for Research,
477 Revised edition. Department of Health and Human Services. DHHS (NIOSH) Publication
478 No. 2011_159, Current Intelligence Bulletin 62, pp. 1-159.

479

480 Ordóñez, NG., 2013. Application of immunohistochemistry in the diagnosis of epithelioid
481 mesothelioma: a review and update. *Hum Pathol* 44(1):1-19. doi:
482 10.1016/j.humpath.2012.05.014

483

484 Paoletti, L., Batisti, D., Bruno, C., Di Paola, M., Gianfagna, A., Mastrantonio, M., et al., 2000.
485 Unusually high incidence of malignant pleural mesothelioma in a town of eastern Sicily: an
486 epidemiological and environmental study. *Arch Environ Occup Health* 55(6):392–8.

487

488 Pavlisko, E.N., Sporn, T.A., 2014. Mesothelioma, Oury T.D., Sporn T.A., Roggli V.L., Eds.
489 In: *Pathology of Asbestos-Associated Disease*. Springer-Verlag, Berlin, Heidelberg, pp 81-
490 140.

491

492 Pelstring, R.J., Allred, D.C., Esther, R.J., Lampkin, S.R., Banks, P.M., 1991. Differential
493 antigen preservation during tissue autolysis. *Hum Pathol.* 1991 Mar;22(3):237-41.

494

495

496 Ricketts, S.W., Peace, C.K., 1976. A case of peritoneal mesothelioma in a thoroughbred
497 mare. *Equine Vet J* 8(2):78-80.

498

499 Rey, F., Boutin, C., Viallat, J. R., Steinbauer, J., Alessandroni, P., Jutisz, P., Di Giambattista,
500 D., Billon-Galland, M A, Hereng, P., Dumortier, P., De Vuyst, P., 1994. Environmental
501 Asbestotic Pleural Plaques in Northeast Corsica: Correlations with Airborne and Pleural
502 Mineralogic Analysis. *Environ Health Perspect* 102(5), 251–252.

503

504 Umphlet, RC., Bertoy, RW., 1988. Abdominal mesothelioma in a cat. *Mod Vet Pract* 69:71-
505 73.

506

507 United States Geological Survey (USGS), 2018. (Accessed 22/02/19)
508 <https://minerals.usgs.gov/minerals/pubs/commodity/asbestos/mcs-2018-asbes.pdf>

509

510 Uzal, F.A., Plattner, BL., Hostetter, JM., 2016. Alimentary system. M. Grant Maxie Ed. in:
511 Jubb, Kennedy, Palmer's Pathology of domestic animals Sixth edition, St. Louis, Missouri,
512 pp 256.

513

514 WHO, 1997. Determination of airborne fibre number concentrations; a recommended
515 method, by phase contrast optical microscopy. World Health Organization, Geneva,
516 Switzerland.

517

518 WHO, 2015. Mesothelial tumours - Diffuse Malignant Mesothelioma - Epithelioid
519 mesothelioma Travis WD, Müller-Hermelink H., Harris C. Eds. In: Classification of Tumours
520 of the Lung, Pleura, Thymus and Heart. pp 130-138

521

522 World Organization for Animal Health, Manual of diagnostic tests and vaccines for terrestrial
523 animals, 2009. Chapter 2.4.7. Bovine tuberculosis. (Accesed 23/02/19)
524 http://www.oie.int/fileadmin/Home/eng/Health_standards/tahm/2.04.07_BOVINE_TB.pdf

525

526 Zakrzewska, AM., Capone, PP., Iannò, A., Tarzia, V., Campopiano, A., Villella, E., Giardino
527 R., 2008. Calabrian ophiolites: dispersion of airborne asbestos fibres during mining and
528 milling operations. *Period Mineral* 77(2), 27-34.

529

530 Zurwieden, NS., Stöber, M., Pohlenz, J., 1990. Morphological characterization of bovine
531 mesothelioma. *Schweiz Arch Tierh* 132:473-474.

532

533

534

535

536 **Figures Legends**

537 **Figure 1.** Geographical area where the boar was killed and soil samples were collected.

538

539 **Figure 2.** Boar, macroscopical findings at post-mortem examination. A: peritoneum,
540 diaphragm and serosal surface of abdominal organs covered by multiple nodules. B: detail
541 of the caudal abdomen characterized by diffuse serosal proliferations.

542

543 **Figure 3.** Boar, histological characterization of the nodular lesions. A: proliferations attached
544 to the diaphragm, Haematoxilin and eosin stain (H-E). B: mixed cell population composed
545 by epithelioid and sarcomatoid elements with disseminated areas of necrosis and
546 lymphoplasmacytic infiltrates. H-E, 100x. C: detail of the tumor masses attached to the
547 serosal surface characterized by pleomorphism and nest or cords organization. H-E, 200x.

548

549 **Figure 4.** Boar, immunohistochemical characterization of the nodular lesions. A: multifocal
550 scattered WT1 positivity. Immunohistochemistry for WT1 detection, haematoxylin
551 counterstaining, 400x. B: multifocal D2-40 positivity. Immunohistochemistry for D2-40
552 detection, haematoxylin counterstaining, 400x.

553

554 **Figure 5.** SEM images (2000X) and relative EDS spectra of chrysotile/asbestiform antigorite
555 (a,b) and tremolite/actinolite asbestos (c,d) detected in lung samples of the boar.

556

557

558 **Table legends**

559 **Table 1.** Immunohistochemical protocols.

560 **Table 2.** Inorganic fibres detected in different kinds of tissue (ff/ gdt: number of fibres per
561 gram of dry tissue; * fibre bundle of 2 fibres minimum; ** fibre bundle of 6 fibres minimum)

562 **Table 3.** Inorganic fibres detected in soil samples of 5 different areas (* fibre bundle)

563

564