



AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Biomarkers of folate and vitamin B12 and breast cancer risk: report from the EPIC cohort

This is the author's manuscript
Original Citation:
Availability:
This version is available http://hdl.handle.net/2318/1766555 since 2021-01-13T09:17:42Z
Published version:
DOI:10.1002/ijc.30536
Terms of use:
Open Access
Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Biomarkers of folate and vitamin B12 and breast cancer risk: report from the EPIC cohort

M. Matejcic¹, J. de Batlle^{1†}, C. Ricci¹, C. Biessy¹, F. Perrier¹, I. Huybrechts¹, E. Weiderpass^{2,3,4,6}, M.C. Boutron-Ruault^{5,7}, C. Cadeau^{5,7}, M. His^{5,7}, D.G. Cox⁸, H. Boeing⁹, R.T. Fortner¹⁰, R. Kaaks¹⁰, P. Lagiou^{11,12,13}, A. Trichopoulou^{11,12}, V. Benetou¹², R. Tumino¹⁴, S. Panico¹⁵, S. Sieri¹⁶, D. Palli¹⁷, F. Ricceri^{18,19}, H.B(as) Bueno-de-Mesquita^{20,21,22}, G. Skeie³, P. Amiano^{23,24}, M.J. Sanchez ^{23,25}, M.D. Chirlaque^{23,26,27}, A. Barricarte^{23,28,29}, J.R. Quiros³⁰, G. Buckland³¹, C.H. van Gils³², P.H. Peeters^{32,33}, T.J. Key³⁴, E. Riboli³³, B. Gylling³⁵, A. Zeleniuch-Jacquotte³⁶, M.J. Gunter¹, I. Romieu¹ and V. Chaje`s¹

1 International Agency for Research on Cancer, Lyon, France

2 Genetic Epidemiology Group, Folkh€alsan Research Center, Helsinki, Finland

3 Department of Community Medicine, University of Tromsø – The Arctic University of Norway, Tromsø, Norway

4 Department of Research, Cancer Registry of Norway, Institute of Population-Based Cancer Research, Oslo, Norway

5 Institut Gustave Roussy, Villejuif, France

6 Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden

7 Universit e Paris-Saclay, Universite Paris-Sud, UVSQ, CESP, INSERM, Villejuif, France 8 Centre Leon B erard, INSERM U1052, Cancer Research Center of Lyon, Lyon, France

9 Epidemiology, German Institute of Human Nutrition Potsdam-Rehbruecke (DIfE), Nuthetal, Germany

10 Division of Cancer Epidemiology, German Cancer Research Center (DKFZ), Heidelberg, Germany

11 Hellenic Health Foundation, Athens, Greece

12 WHO Collaborating Center for Nutrition and Health, Unit of Nutritional Epidemiology and Nutrition in Public Health, Department of Hygiene, Epidemiology and Medical Statistics, School of Medicine, National and Kapodistrian University of Athens, Athens, Greece

13 Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, MA

14 Cancer Registry and Histopathology Unit, Civic - M.P. Arezzo Hospital, ASP Ragusa, Ragusa, Italy

15 Dipartimento di Medicina Clinica e Chirurgia, Universita degli Studi di Napoli Federico II, Naples, Italy

16 Epidemiology and Prevention Unit, Fondazione IRCCS Istituto Nazionale dei Tumori, Milano, Italy

17 Molecular and Nutritional Epidemiology Unit, Cancer Research and Prevention Institute – ISPO, Florence, Italy

18 Unit of Cancer Epidemiology, Department of Medical Sciences, University of Turin, Turin, Italy

19 Unit of Epidemiology, Regional Health Service ASL TO3, Grugliasco, Italy

20 Department of Social & Preventive Medicine, Faculty of Medicine, University of Malaya, Kuala Lumpur, Malaysia

21 Department for Determinants of Chronic Diseases (DCD), National Institute for Public Health and the Environment (RIVM), Bilthoven, The Netherlands

22 Department of Epidemiology and Biostatistics, The School of Public Health, Imperial College London, London, United Kingdom

23 CIBER de Epidemiologia y Salud Publica (CIBERESP), Madrid, Spain

24 Public Health Division of Gipuzkoa, BioDonostia Research Institute, San Sebastian, Spain

25 Escuela Andaluza de Salud Publica, Instituto de Investigaci on Biosanitaria ibs, GRANADA, Hospitales Universitarios de Granada/Universidad de Granada, Granada, Spain

26 Department of Epidemiology, Regional Health Council, IMIB-Arrixaca, Murcia, Spain

27 Department of Health and Social Sciences, Universidad de Murcia, Murcia, Spain

28 Navarra Institute for Health Research (IdiSNA), Pamplona, Spain

29 Navarra Public Health Institute, Pamplona, Spain

30 Public Health Directorate, Asturias, Spain

31 Unit of Nutrition and Cancer, Cancer Epidemiology Research Programme, Catalan Institute of Oncology (ICO-IDIBELL), Barcelona, Spain

32 Department of Epidemiology, Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Utrecht, The Netherlands 33 Department of Epidemiology and Biostatistics, School of Public Health, Imperial College, London, United Kingdom

34 Cancer Epidemiology Unit, Nuffield Department of Population Health, University of Oxford, Oxford, United Kingdom

35 Department of Medical Biosciences, Pathology, Umea° University, Umea°, Sweden

36 Department of Population Health, NYU School of Medicine, New York, NY

Correspondence to:

Marco Matejcic, International Agency for Research on Cancer, 150, Cours Albert-Thomas, 69372 Lyon CEDEX 08, France, Tel.: 133-0-4-72-73-8029, E-mail: <u>matejcicm@fellows.iarc.fr</u>

Abstract

Epidemiological studies have reported inconsistent findings for the association between B vitamins and breast cancer (BC) risk. We investigated the relationship between biomarkers of folate and vitamin B12 and the risk of BC in the European Prospective Investigation into Cancer and Nutrition (EPIC) cohort. Plasma concentrations of folate and vitamin B12 were determined in 2,491 BC cases individually matched to 2,521 controls among women who provided baseline blood samples. Multivariable logistic regression models were used to estimate odds ratios by quartiles of either plasma B vitamin. Subgroup analyses by menopausal status, hormone receptor status of breast tumors (estrogen receptor [ER], progesterone receptor [PR] and human epidermal growth factor receptor 2 [HER2]), alcohol intake and MTHFR polymorphisms (677C > T and 1298A > C) were also performed. Plasma levels of folate and vitamin B12 were not significantly associated with the overall risk of BC or by hormone receptor status. A marginally positive association was found between vitamin B12 status and BC risk in women consuming above the median level of alcohol (OR Q4-Q1 = 1.26; 95% CI 1.00–1.58; Ptrend 5 0.05). Vitamin B12 status was also positively associated with BC risk in women with plasma folate levels below the median value (OR Q4-Q1 = 1.29; 95% CI 1.02 1.62; Ptrend 5 0.03). Overall, folate and vitamin B12 status was not clearly associated with BC risk in this prospective cohort study. However, potential interactions between vitamin B12 and alcohol or folate on the risk of BC deserve further investigation.

What's new?

Does B-vitamin intake play a role in breast cancer (BC) risk? Results have been inconsistent. In this analysis of data from a large, prospective European study, the authors found that, overall, folate and vitamin B12 status were not clearly associated with BC risk. However, the risk did seem to increase somewhat for women who had higher vitamin B12 levels and either low plasma folate or increased alcohol consumption. The authors suggest that this may involve nutrient-nutrient or gene-nutrient interactions, such as changes in DNA methylation, which require further investigation.

Key words:

plasma biomarkers, folate, vitamin B12, alcohol, hormone receptor status, MTHFR polymorphism, breast cancer

Introduction

The etiology of breast cancer (BC) is complex and results from the combination of lifetime reproductive events, genetics, dietary and lifestyle factors. [1] According to the latest breast cancer report from the World Cancer Research Fund (WCRF), there is novel evidence that alcohol intake and factors that lead to a greater adult attained height are positively associated with postmenopausal and probably also premenopausal BC. [2] Among dietary factors, deficiencies of B vitamins related to Western dietary patterns have been suggested to play a role in breast carcinogenesis. [3,4] Vitamin B9 (folate) and vitamin B12 (cobalamin) are two water soluble B vitamins involved in one-carbon metabolism, [5] which generates substrates for DNA methylation and DNA synthesis. [6] Thus, deficiencies of these micronutrients may trigger both genetic and epigenetic procarcinogenic processes. [7] Prospective studies that investigated the association between biomarkers of folate and BC risk have reported either an inverse association [8,9] or no association [10,11] overall. The prospective investigation of the relationship between biomarkers of vitamin B12 and BC risk has also produced mixed results. [8-12] While studies found no evidence for an association between blood levels of vitamin B12 and BC risk in the overall population, [9–11] an inverse association was independently reported among either postmenopausal women [10] or premenopausal women. [9] However, a recent meta-analysis of prospective studies revealed no significant association between biomarkers of vitamin B12 and BC risk in the subgroup analysis by menopausal status. [13] A number of factors have been suggested to influence the association between B vitamins and the risk of BC, including menopausal status, [11,14–16] alcohol consumption, [15,17] nutrient interactions [18] and methylenetetrahydrofolate reductase (MTHFR) gene polymorphisms. [12] MTHFR is a key enzyme in one-carbon metabolism where it balances the folate pool between synthesis and methylation of DNA. [19] Although a number of MTHFR single nucleotide polymorphisms (SNPs) have been reported in the literature, [20-22] only the C677T and A1298C SNPs have been consistently associated with decreased enzyme activity and reduced plasma folate levels compared with the wildtype genotypes. [23,24] A prospective study found a positive association between plasma folate concentration and postmenopausal BC among carriers of the MTHFR 677T allele. [12] Breast tumors are also subdivided into subgroups according to the expression of sex hormone receptors (estrogen receptor [ER], progesterone receptor [PR] and human epidermal growth factor receptor 2 [HER2]), which have been differentially associated with both folate intake [17] and folate status [11] among premenopausal women only.

We conducted a large nested case–control study within European Prospective Investigation into Cancer and Nutrition (EPIC) to evaluate the association between plasma concentrations of folate and vitamin B12 and BC risk overall and stratified by hormone receptor status and potential risk factors.

In addition, we examined the interaction between the MTHFR 677C > T (rs1801133) and 1298A > C (rs1801131) polymorphisms and the two plasma B vitamins on the risk of BC using data from a subsample of this nested case–control population.

Material and Methods

Study design

The EPIC study is an ongoing multicenter European cohort study designed to investigate the role of dietary habits and lifestyle factors on the incidence of cancer of various sites, including BC. [25] The cohort includes over 521,000 participants recruited between 1992 and 2000 from 23 centers in 10 European countries (Denmark, France, Germany, Greece, Italy, the Netherlands, Norway, Spain, Sweden and UK). Of 367,903 women (age 35–70 years) recruited into the EPIC study, the present analysis excluded women with prevalent cancers at recruitment (n = 19,853) and missing diagnosis or censoring date (n = 2,892). A total of 10,713 women with malignant primary BC were identified after a median followup of 11.5 years. The follow-up rate was very high (98.5%: 91.4% alive and 7.1% dead) and only 1.5% of women were lost to follow-up.

Details of the recruitment procedures and data collection in the EPIC study have been previously described in details. [26] Briefly, sociodemographic, lifestyle and dietary data were collected at baseline from all the cohort members by administration of country-specific questionnaires. Anthropometric measurements and peripheral blood samples of the participants were also collected. Methods of blood collection, processing and storage are described in details elsewhere. [27] All participants signed an informed consent for the use of their blood samples and data. The study was approved by the Ethical Review Board of the IARC and those of all national recruiting centers.

Selection of study subjects

A nested case–control study was designed among women who provided a blood sample and completed the lifestyle and dietary questionnaires at recruitment. A total of 2,491 BC cases with a confirmed first diagnosis of invasive BC were identified between 1992 and 2010. Each case was individually matched to at least one control subject chosen randomly among cohort women with available blood samples and free of cancer (except nonmelanoma skin cancer) at the time of diagnosis of the corresponding case. Control subjects were matched to cases for study center, age at blood donation (63 months), exogenous hormone use at blood collection (yes; no; unknown), menopausal status (pre; surgical post; natural post), fasting status (<3, 3-6, >6 hr) and phase of the menstrual cycle (early follicular, late follicular, periovulatory, mid luteal, other luteal) at recruitment.

Dietary and lifestyle data collection

Dietary data were obtained at enrollment using validated country-specific dietary history and foodfrequency questionnaires (FFQs), designed to collect local dietary habits of the participants over the preceding year. [26] Dietary intakes of folate and vitamin B12 were estimated using the updated EPIC Nutrient Database (ENDB), [28] following standardization from country-specific food composition tables (FCT) according to Bouckaert's recommendations. [29] Details on dietary assessment have been discussed previously. [17] Participants also completed a baseline lifestyle questionnaire providing information on anthropometric and sociodemographic characteristics, reproductive history, family history of cancer, physical activity, alcohol use, smoking habits, use of oral contraceptives, hormone replacement therapy and vitamin supplements in the year prior to enrollment date.

Outcome assessment

Participants were followed from the date of enrollment until first cancer diagnosis, death, emigration or end of the followup period, whichever occurred first. Incident cancer cases were identified through population cancer registries (Denmark, Italy except Naples, the Netherlands, Norway, Spain, Sweden and UK) or by a combination of methods including health insurance, cancer and pathology registries and active ollow-up through study subjects and their next-of-kin in three countries (France, Germany, Greece and Naples). Data on clinical and tumor characteristics were coded according to the 10th Revision of the International Statistical Classification of Diseases, Injuries and Causes of Death (ICD). In the present study, 91% of BC cases were confirmed by histological or cytological examination, whereas the remaining 9% was diagnosed through clinical observation, ultrasound, autopsy or death certificate. The most frequent subtype of BC was ductal carcinoma (71.5%), followed by lobular carcinoma (14.1%) and tubular carcinoma (2.7%). The remaining BC cases were classified as mixed (5.0%) or other (6.7%) subtypes.

Hormone receptor status determination

Determination of ER, PR and HER2 status of BC cases was performed within each EPIC center. Information on hormone receptor status as well as on the methods for its determination was retrieved from each EPIC center using the same approaches used for collection of incident cases. To standardize the quantification of the receptor status collected across centers, the following criteria were applied for a positive receptor status: 10% cells stained, any "plus-system" description, 20 fmol/mg, an Allred score of 3, an IRS 2 or an H-score 10.30 ER, PR and HER2 status was available for 98, 84 and 44% of cases, respectively. For the remaining cases, hormone receptor status was not determined because of insufficient amount of tumor tissue available for histopathological evaluation. Furthermore, HER2

status could not be ascertained in the majority of cases because of the lack of a specific test in the nineties.

Laboratory measurements

All biochemical analyses were performed at the Bevital AS laboratory in Bergen, Norway (www.bevital.no). Microbiological assays were used to determine plasma concentrations of folate [31] and vitamin B12. [32] The assays were adapted to a microtiter plate format and carried out by a robotic workstation. Throughout all steps of the biochemical analysis, samples from each case– control set were analyzed within the same batch. The laboratory personnel were blinded to case– control status. To assess the measurement precision, each batch contained six quality control (QC) samples with known biomarker concentrations and four samples without biomarker (blanks). The six QC samples were three samples in parallels. The coefficient of variation calculated from the three duplicate sets of identical QC samples was 8.6% for folate and 5.0% for vitamin B12. Plasma concentrations of folate and vitamin B12 were determined for all study participants.

Genotyping analysis

Determination of the genotype status was carried out only in a subsample of 401 cases and 401 matched control individuals from this nested case–control population. DNA extraction from white blood cells was carried out using Autopure LS kit (Gentra Systems, Minneapolis, MN). DNA concentration was quantified with Quant-iT PicoGreen dsDNA reagent (Thermo Fisher Scientific, Waltham, MA). The MTHFR 677C > T (rs1801133) and 1298A > C (rs1801131) single nucleotide polymorphisms (SNPs) were genotyped by Kaspar allelic discrimination assay using allelespecific probes and fluorescent reporters (LGC Group, UK). Each reaction was carried out according to the manufacturer's instructions using supplied kits. Amplifications and end-point allele determination were performed in 96-well plates using a StepOne Plus system (Applied Biosystems). Each plate contained randomly placed case and control samples, while matched sets were analyzed within the same plate. Genotyping success rates were 98.0 and 96.5% for rs1801131 and rs1801133, respectively. Samples not yielding genotypes were removed from further analyses.

Statistical methods

Lifestyle and dietary baseline characteristics of study participants were described using mean 6 standard deviation (SD) for continuous variables and percentages for categorical variables. Plasma concentrations of folate and vitamin B12 were log natural transformed to normalize their distribution. The paired t test and $\chi 2$ test were used to assess differences between cases and control individuals with regard to continuous and categorical variables, respectively. Multivariable conditional logistic

regression models were used to estimate odds ratios (ORs) and 95% confidence intervals (95% CIs) for overall BC and specific subgroups stratified by menopausal status at recruitment (dichotomized as natural/surgical postmenopausal and premenopausal) and by hormone receptor status (ER1/ER2, PR1/PR2, HER21/HER22). Crude ORs were also presented to observe the effect of confounding on the risk estimates. In addition, the association between each plasma biomarker and the risk of BC was examined using four-knot restricted cubic splines with the midpoint of the fifth decile of plasma vitamin B12 as the reference category. [33] Quartiles and tertiles of plasma levels of biomarkers for the overall and hormone receptor-specific analyses, respectively, were determined on the basis of the distribution among control individuals. Tests for linear trends were performed by entering the median value of each category as continuous term in the multivariable models. All multivariate models were adjusted by BMI, height, alcohol intake, total energy intake, educational attainment (primary school, technical/professional school, secondary school, university degree, 4.2% unknown), physical activity (inactive, moderately inactive, moderately active, active, 6.9% unknown), ever use of hormone replacement therapy (never, ever, 4.1% unknown), parity and age at first full-term birth combined (nulliparous, <30 year and 1-2 children, <30 year and 3 children, >21-30 year, 30 year, 3.6% unknown) and family history of BC (yes, no, 53.1% unknown). These confounders were previously related to BC risk or blood measurements and were chosen based on previous studies in the literature. Unknown categories of the above mentioned variables were included in the model using indicator variables. Multivariate unconditional logistic regression models were used to investigate the association between plasma concentrations of folate and vitamin B12 and BC risk by levels of alcohol intake or plasma folate (low and high levels based on median values) and by MTHFR genotypes. The joint effect of plasma folate (in tertiles) and categories of alcohol intake (0-3, 3-12, >12 g/day) on BC risk was evaluated by using the lowest tertile of plasma folate and highest category of alcohol intake as reference category, as previously assessed. [17] Tests for interaction between each plasma biomarker as continuous variable and potential risk factors were computed by likelihood ratio test. Formal tests of heterogeneity between ORs in menopausal and hormone receptor subgroups were based on χ^2 statistics, calculated as the deviations of logistic beta-coefficients observed in each of the subgroups relative to the overall beta-coefficient. The association between the SNPs and overall BC risk was evaluated by conditional logistic regression. Genotypic (codominant) and dominant models were assumed for SNP effects. A trend test was conducted by treating the genotypes as equally spaced integer weights and entering the variable as a continuous term in the model. Specific sensitivity analyses were carried out by excluding women consuming multivitamin supplements and cases diagnosed within the first 2 years of follow-up (to reduce the chance of reverse causality).

Statistical tests were two-sided, and p values below 0.05 were considered statistically significant. All analyses were performed using STATA 12.1 (StataCorp. 2011, Stata Statistical Software: Release 11, College Station, TX).

Results

Table 1 summarizes the sociodemographic, reproductive and lifestyle characteristics of study participants by case–control status. Cases had slightly older age at menopause (p = 0.026) and at first live birth (p = 0.023) than control individuals. A slightly higher BMI in cases compared with the control group was found among postmenopausal women (p < 0.001), but not among premenopausal women. Cases were also more likely to have had a first-degree relative with BC (p = 0.009), and had higher daily alcohol intake (p = 0.002). Both MTHFR SNPs were in Hardy-Weinberg equilibrium (p = 0.298 for C677T; p = 0.823 for A1298C) and the frequency of the minor allele among control individuals was 30.8% at locus C677T and 37.0% at locus A1298C (data not shown).

There was no significant association between plasma levels of folate and vitamin B12 and the overall risk of BC (Table 2) or by ER, PR and HER2 status (Table 3). No further association emerged after adjustment by MTHFR polymorphisms for the available subsample (data not shown).

A nonlinear modeling of the association between plasma concentrations of vitamin B12 and BC risk showed a borderline significant trend (Ptrend = 0.07) in increased risk associated with plasma concentrations of vitamin B12 higher than 360 pmol/l, while the odds ratio plateaued at levels 500 pmol/l. No dose-dependent effect of plasma folate on the risk of BC was observed (data not shown). Because of the impaired folate absorption and altered onecarbon metabolism due to chronic alcohol consumption, [34] we reported risk estimates by tertiles of plasma folate and categories of alcohol consumption (Fig. 1). The association between plasma folate concentration and BC risk was not significantly modified by levels of alcohol intake (Pinteraction = 0.69). Similarly, no significant association between plasma folate and BC risk was observed by median level of alcohol consumption (data not shown).

The association between plasma levels of vitamin B12 and BC risk stratified by the median intake of alcohol is summarized in Table 4. There was a borderline significant increase in risk associated with the highest quartile of plasma vitamin B12 in women consuming at least 3.36 g/day of alcohol (OR Q4-Q1 = 1.26; 95% CI 1.00–1.58; Ptrend = 0.05), while no significant association emerged in women drinking lower amounts of alcohol (OR Q4-Q1 = 1.08; 95% CI 0.86–1.35; Ptrend = 0.56). However, no significant heterogeneity by alcohol intake was found (Pheterogeneity = 0.14). The multivariable

risk estimates did not change appreciably after further adjustment by plasma folate concentration (data not shown). A statistically significant interaction between plasma concentrations of folate and vitamin B12 on the risk of BC was observed (Pinteraction = 0.04; data not shown). To further explore this interaction, a stratification analysis by the median level of plasma folate was carried out (Table 4). A marginally increased risk of BC associated with increasing concentrations of plasma vitamin B12 was found in women with plasma levels of folate below 13.56 nmol/l (OR Q4-Q1 = 1.29; 95% CI 1.02–1.62; Ptrend = 0.03), while no significant association occurred in women with higher levels of plasma folate (Ptrend = 0.68). A borderline significant heterogeneity by plasma folate levels was also found (Pheterogeneity = 0.05). Exclusion from analyses of women who consumed multivitamin supplements or cases diagnosed within the first two years of follow-up did not change the risk estimates in our study population (data not shown).

The MTHFR 677C > T and 1298A > C SNPs were in low linkage disequilibrium among both the cases (r2 = 0.24) and control individuals (r2 = 0.25). There was no significant association between either C677T (OR TTvs.CC = 0.71; 95% CI 0.42–1.19; Ptrend = 0.38) or A1298C (OR CCvs.AA = 0.97; 95% CI 0.62–1.53; Ptrend = 0.91) and the overall risk of BC. The interaction between plasma folate or vitamin B12 and MTHFR SNPs was not statistically significant (Pinteraction > 0.05). Plasma concentrations of the two B vitamins were not significantly associated with BC risk in any of the genotypic classes (homozygous wild-type, heterozygous, homozygous variant) of each SNP (data not shown). No further association emerged in the dominant models, and adjustment for the alternative SNP did not change the risk estimates.

Discussion

In this large prospective European study, circulating levels of folate and vitamin B12 were not significantly associated with the overall risk of BC. However, we found borderline positive associations between plasma concentrations of vitamin B12 and BC risk restricted to women with either high alcohol intake or low folate status. The MTHFR C677T and A1298C polymorphisms had no effect modification on the association between either plasma B vitamin and BC risk in a subsample of this nested case–control study.

A study was recently conducted to assess the reliability of plasma biomarkers involved in one-carbon metabolism in a subsample from the EPIC study (38 men and 35 women), which was estimated over a period of 2–5 years using an intraclass correlation coefficient (ICC). [35] The study showed that plasma vitamin B12 was a highly reliable biomarker (ICC = 0.75), while a modest reliability was

observed for plasma folate (ICC = 0.45). Because our study was performed on a larger number of subjects and extended over 18 years of follow-up, it is difficult to predict whether and to what extent the single biomarker measurements may have led to attenuation of the risk estimates in our study. However, when the models were adjusted for the regression dilution using the ICCs as adjustment coefficients, no significant change in risk estimates was observed.

Consistent with our findings, a prospective study within EPIC reported a lack of significant association between dietary folate intake and the overall risk of BC. [17] Prospective investigations based on biomarkers of nutrient status reported inconsistent findings between folate and BC risk. [8–12] The mean plasma folate concentration in our study population (14.2 nmol/l) was comparable to that reported in the Malmo o Diet and Cancer cohort (12.8 nmol/l), which also reported a null association between plasma levels of folate and overall BC risk. [12] However, our highest category of plasma folate (>19.8 nmol/l; 609 cases) was substantially lower than that reported in the US population-based cohort from the Nurses' Health Study (>14.0 ng/ml = 31.7 nmol/l; 120 cases) in which a higher consumption of folic-acid containing foods and an inverse association between plasma folate levels and BC risk were observed. [9] Thus, a minimal level of blood folate might be required for observing a beneficial effect of this nutrient on the risk of BC.

The high plasma folate concentrations reported in US based population studies is likely due to folic acid fortification of flour and cereal-grain products, which became mandatory in the United States since 1997 to prevent neural tube defects. [36,37] On the other hand, no policy of folic acid fortification of foods has been implemented in European countries. Folic acid from fortified foods or supplementation is estimated to be approximately 1.7 times more bioavailable than natural folates. [38] Because most of the enzymes that use folate as cofactor cannot use the synthetic form, there might be important perturbations in one-carbon metabolism and cellular processes that rely on this pathway. A recent dose–response meta-analysis of 16 prospective studies including a total of 26,205 BC patients identified a U-shaped relationship between energy-adjusted dietary folate intake and BC risk, [39] supporting prior evidence of an increased risk of BC associated with folic acid fortification. [40] The lack of data on consumption of folic acid intake might have been associated with high levels of plasma folate and an increased BC risk. However, the proportion of vitamin supplement users in our study population was only 23% among cases, suggesting that plasma levels of folate and other B vitamins were primarily attributable to natural food sources.

The lack of a significant interaction between plasma folate levels and alcohol intake on BC risk in our analysis is consistent with results from previous prospective studies that used biomarkers of folate

status. [8–12] However, a recent prospective investigation within the EPIC study reported an inverse association between dietary folate intake and the risk of BC among heavy alcohol drinkers.17 Since alcohol may impair folate absorption, [34,39] alcohol consumption behaviors are more likely to modify the risk of BC associated with dietary folate intake rather than plasma folate levels, which can be affected by a variety of other factors including genetic polymorphisms. [41] Thus, women with high intake of both folate and alcohol may not necessarily have a high folate status and consequently a reduced risk of BC.

The main sources of vitamin B12 are animal products, including meat, fish, dairy products, eggs and liver. Our finding of a positive association between plasma levels of vitamin B12 and BC risk in subgroup analyses is in accordance with two previous prospective studies that measured either dietary intake [42] or plasma levels11 of this nutrient. However, an inverse association between biomarkers of vitamin B12 and the risk of BC has also been reported. [9,10] The median value of plasma vitamin B12 in our study population (377 = 511 pg/ml in cases) was not substantially different from those reported in other population-based prospective studies, ranging between 421 and 467 pg/ml.9–11 Thus, several other factors might have contributed to the inconsistent findings, including differences in alcohol consumption, genetic polymorphisms, and nutrient interactions in one-carbon metabolism. [34,43]

As a cofactor required for the generation of methylgroups, a high vitamin B12 status could result in hypermethylation of CpG island promoters for tumor suppressor genes, [44] which may lead to reduced expression of these cancer-related genes and ultimately promote breast carcinogenesis. [45] These DNA methylation changes may also impair the proper expression and/or function of cell-cycle regulatory genes and thus confer a selective growth advantage to neoplastic cells. [46] A randomized crossover trial suggested that moderate alcohol intake may diminish plasma vitamin B12 concentrations. [47] In contrast, a case–control study found that plasma levels of vitamin B12 in heavy alcohol drinkers were significantly higher than those in light alcohol drinkers. [48] Further studies are needed to clarify the modifying effect of alcohol on the association between vitamin B12 and BC risk.

The positive association between plasma levels of vitamin B12 and BC risk among women with low folate status is unexpected. Previous prospective studies found no evidence of an interaction between these two nutrients on the risk of BC. [16,42,49,50] On the other hand, a prospective analysis within the French E3N cohort reported a strong joint protective effect of high intake of folate and vitamin B12 on BC risk. [18] The almost exclusive form of folate in plasma is 5-methyl THF, which reflects the amount of folate available for DNA methylation.51 5-methyl THF is converted to tetrahydrofolate

(THF) via the vitamin B12-dependent enzyme methionine synthase. A high vitamin B12 status indicates that methionine synthase activity is increased, leading to depletion of 5-methyl THF and thus plasma folate concentration if not replaced by new 5-methyl THF from diet. In this situation, cells lack the substrate needed for methionine synthesis and DNA methylation is impaired. There is evidence that a low folate status may induce carcinogenesis through alteration of DNA methylation pathways.52 Thus, the possibility that low plasma folate concentrations (mainly 5-methyl THF) as a consequence of high vitamin B12 status would impair DNA methylation might be suggested.

Epidemiological studies provide support that the association between the MTHFR C677T polymorphism and BC risk is modified by intakes of some B vitamins, including folate and vitamin B12. [53–55] We observed no significant effect modification of MTHFR SNPs on the association between plasma folate or vitamin B12 and BC risk. The low power of these subgroup analyses prevented us from finding a potential interaction between MTHFR genotypes and B vitamin status on the risk of BC. Furthermore, the effect of MTHFR polymorphisms on plasma levels of B vitamins is highly complex and may depend on the interaction with other dietary and genetic factors. [56]

The present study is the largest prospective investigation to date to have examined the association between biomarkers of folate and vitamin B12 and the risk of BC. The high follow-up rates and large number of cases provided sufficient statistical power for most subgroup analyses. The major strength of our study is, however, the collection of blood samples prior to diagnosis and the use as biomarkers of exposure as reflection of true vitamin status.

Major limitations include the single collection of blood samples at baseline and the measurement of a single biomarker of folate or vitamin B12 status. Folate concentration measured in plasma is considered to be a sensitive biomarker of recent dietary intake, and thus is not very informative for the assessment of long-term folate status. [57] Plasma vitamin B12 is the most widely used biomarker of total cobalamin status, but not the most specific biomarker to characterize adequate vitamin concentrations.58 In order to obtain more reliable information on vitamin status, multiple measurements of plasma biomarkers should be taken over a period of time or a combination of different biomarkers should be used. Additional limitations include (i) the large percentage of missing data for family history of BC (53.1%) and supplement use (54.5%), (ii) the determination of menopausal status at recruitment and not at diagnosis, (iii) the lack of complete hormone receptor status data and (iv) the insufficient statistical power for gene-nutrient interaction analyses. Because controlling for family history of BC and supplement use had minimal effect on the risk estimates, our results are unlikely to be explained by residual confounding by those factors.

In conclusion, no clear support for an association between plasma levels of folate and BC risk was found in this large prospective study. However, potential interactions between vitamin B12 and alcohol or folate on the risk of BC were observed. Our findings suggest a potential role of vitamin B12 in breast carcinogenesis and raise the possibility of important nutrient–nutrient and gene–nutrient interactions in the etiology of BC. The potential deleterious effect of high vitamin B12 status in combination with other risk factors for BC deserves further investigation. Given the inconsistent findings to date and the possibility that associations between folate and BC could be influenced by some factors yet to be identified, further studies based on novel biomarkers that take into account the effect of potential risk factors and genetic polymorphisms are warranted.

Acknowledgements

The authors gratefully acknowledge Deborah Postoly, laboratory technician, for sample processing and genotyping of study samples, as well as chief engineer Gry Kvalheim within the Bevital laboratory, Bergen, Norway, for her outstanding assistance with laboratory measurements of blood B vitamins. We confirm that all authors listed have contributed to the planning, execution and analysis of the submitted manuscript and that there are no conflicts of interest whatsoever.

Funding

International Agency for Research on Cancer; Grant sponsor: European Commission FP7 Marie Curie Actions-PeopleCofunding of regional, national, and international programs (COFUND); Grant Sponsor: Ecumenical Project for International Cooperation; Grant sponsor: Danish Cancer Society (Denmark); Grant sponsor: Ligue Contre le Cancer, Institut Gustave Roussy, Mutuelle Generale de l'Education Nationale, Institut National de la Sante et de la Recherche Medicale (INSERM) (France); Grant sponsor: Deutsche Krebshilfe, Deutsches Krebsforschungszentrum and Federal Ministry of Education and Research (Germany); Grant sponsor: Hellenic Health Foundation (Greece); Grant sponsor: Italian Association for Research on Cancer (AIRC) and National Research Council (Italy); Grant sponsor: Dutch Ministry of Public Health, Welfare and Sports (VWS); Grant sponsor: Netherlands Cancer Registry (NKR); Grant sponsor: LK Research Funds; Grant sponsor: Dutch Prevention Funds; Grant sponsor: Dutch ZON (Zorg Onderzoek Nederland); Grant sponsor: World Cancer Research Fund (WCRF); Grant sponsor: Statistics Netherlands (The Netherlands); Grant sponsor: AGAUR, Generalitat de Catalunya; Grant number: Exp. 2014 SGR 726; Grant sponsor: Health Research Funds; Grant number: RD12/0036/0018; Grant sponsor: Swedish Cancer Society, Swedish Scientific Council and Regional Government of Skåne and Voasterbotten (Sweden); Grant sponsor: Cancer Research UK; Grant sponsor: Medical Research Council; Grant sponsor: Stroke Association; Grant sponsor: British Heart Foundation; Grant sponsor: Department of Health, Food Standards Agency; Grant sponsor: Welcome Trust (United Kingdom); Grant sponsor: World Cancer Research Funds; Grant sponsor: Institut National du Cancer (INCA); Grant sponsor: la Fondation de France (FDF); Grant sponsor: La Ligue Nationale contre le Cancer (LNCC)

References

1. Hankinson SE, Colditz GA, Willett WC. Towards an integrated model for breast cancer etiology: the lifelong interplay of genes, lifestyle, and hormones. Breast Cancer Res 2004;6: 213–8.

Breast Cancer 2010 Report. Food, nutrition, physical activity, and the prevention of breast cancer.
2016.

3. Ames BN, Wakimoto P. Are vitamin and mineral deficiencies a major cancer risk?. Nat Rev Cancer 2002;2:694–704.

4. Vera-Ramirez L, Ramirez-Tortosa MC, SanchezRovira P, et al. Impact of diet on breast cancer risk: a review of experimental and observational studies. Crit Rev Food Sci Nutr 2013;53:49–75.

5. Mason JB. Biomarkers of nutrient exposure and status in one-carbon (methyl) metabolism. J Nutr 2003;133(Suppl.3):9418–7S.

6. Davis CD, Uthus EO. DNA methylation, cancer susceptibility, and nutrient interactions. Exp Biol Med (Maywood) 2004;229:988–95.

7. Szyf M, Pakneshan P, Rabbani SA. DNA methylation and breast cancer. Biochem Pharmacol 2004;68:1187–97.

8. Rossi E, Hung J, Beilby JP, et al. Folate levels and cancer morbidity and mortality: prospective cohort study from Busselton, Western Australia. Ann Epidemiol 2006;16:206–12.

9. Zhang SM, Willett WC, Selhub J, et al. Plasma folate, vitamin B6, vitamin B12, homocysteine, and risk of breast cancer. J Natl Cancer Inst 2003;95:373–80.

10. Wu K, Helzlsouer KJ, Comstock GW, et al. A prospective study on folate, B12, and pyridoxal 50-phosphate (B6) and breast cancer. Cancer Epidemiol Biomarkers Prev 1999;8:209–17.

11. Lin J, Lee IM, Cook NR, et al. Plasma folate, vitamin B-6, vitamin B-12, and risk of breast cancer in women. Am J Clin Nutr 2008;87:734–43.

12. Ericson UC, Ivarsson MI, Sonestedt E, et al. Increased breast cancer risk at high plasma folate concentrations among women with the MTHFR 677T allele. Am J Clin Nutr 2009;90:1380–9.

13. Wu W, Kang S, Zhang D. Association of vitaminB6, vitamin B12 and methionine with risk of breast cancer: a dose-response meta-analysis. Br J Cancer 2013;109:1926–44.

14. Ericson U, Sonestedt E, Gullberg B, et al. High folate intake is associated with lower breast cancer incidence in postmenopausal women in the Malmo Diet and Cancer cohort. Am J Clin Nutr 2007;86:434–43.

15. Stolzenberg-Solomon RZ, Chang SC, Leitzmann MF, et al. Folate intake, alcohol use, and postmenopausal breast cancer risk in the prostate, lung, colorectal, and ovarian cancer screening trial. Am J Clin Nutr 2006;83:895–904.

16. Stevens VL, McCullough ML, Sun J, et al. Folate and other one-carbon metabolism-related nutrients and risk of postmenopausal breast cancer in the Cancer Prevention Study II Nutrition Cohort. Am J Clin Nutr 2010;91:1708–15.

17. de BJ, Ferrari P, Chajes V, et al. Dietary folate intake and breast cancer risk: European prospective investigation into cancer and nutrition. J Natl Cancer Inst 2015;107:dju367.

18. Lajous M, Romieu I, Sabia S, et al. Folate, vitamin B12 and postmenopausal breast cancer in a prospective study of French women. Cancer Causes Control 2006;17:1209–13.

19. Choi SW, Mason JB. Folate and carcinogenesis: an integrated scheme. J Nutr 2000;130:129-32.

20. Koushik A, Kraft P, Fuchs CS, et al. Nonsynonymous polymorphisms in genes in the one-carbon metabolism pathway and associations with colorectal cancer. Cancer Epidemiol Biomarkers Prev 2006;15:2408–17.

21. Sharp L, Little J. Polymorphisms in genes involved in folate metabolism and colorectal neoplasia: a HuGE review. Am J Epidemiol 2004;159: 423–43.

22. Hazra A, Wu K, Kraft P, et al. Twenty-four nonsynonymous polymorphisms in the one-carbon metabolic pathway and risk of colorectal adenoma in the Nurses' health study. Carcinogenesis 2007;28:1510–9.

23. Frosst P, Blom HJ, Milos R, et al. A candidate genetic risk factor for vascular disease: a common mutation in methylenetetrahydrofolate reductase. Nat Genet 1995;10:111–3.

24. van der Put NM, Gabreels F, Stevens EM, et al. A second common mutation in the methylenetetrahydrofolate reductase gene: an additional risk factor for neural-tube defects?. Am J Hum Genet 1998;62:1044–51.

25. Bingham S, Riboli E. Diet and cancer – the European Prospective Investigation into Cancer and Nutrition. Nat Rev Cancer 2004;4:206–15.

26. Riboli E, Hunt KJ, Slimani N, et al. European Prospective Investigation into Cancer and Nutrition (EPIC): study populations and data collection. Public Health Nutr 2002;5:1113–24.

27. Riboli E, Kaaks R. The EPIC project: rationale and study design. European Prospective Investigation into Cancer and Nutrition. Int J Epidemiol 1997;26(Suppl.1):S6–S14.

28. Slimani N, Deharveng G, Unwin I, et al. The EPIC nutrient database project (ENDB): a first attempt to standardize nutrient databases across the 10 European countries participating in the EPIC study. Eur J Clin Nutr 2007;61:1037–56.

29. Bouckaert KP, Slimani N, Nicolas G, et al. Critical evaluation of folate data in European and international databases: recommendations for standardization in international nutritional studies. Mol Nutr Food Res 2011;55:166–80.

30. Ritte R, Tikk K, Lukanova A, et al. Reproductive factors and risk of hormone receptor positive and negative breast cancer: a cohort study. BMC Cancer 2013;13:584.

31. Molloy AM, Scott JM. Microbiological assay for serum, plasma, and red cell folate using cryopreserved, microtiter plate method. Methods Enzymol 1997;281:43–53.

32. Kelleher BP, Broin SD. Microbiological assay for vitamin B12 performed in 96-well microtitre

plates. J Clin Pathol 1991;44:592–5.

33. Durrleman S, Simon R. Flexible regression models with cubic splines. Stat Med 1989;8:551-61.

34. Halsted CH, Villanueva JA, Devlin AM, et al. Metabolic interactions of alcohol and folate. J Nutr 2002;132(Suppl. 8):2367S–72S.

35. Leenders M, Ros MM, Sluijs I, et al. Reliability of selected antioxidants and compounds involved in one-carbon metabolism in two Dutch cohorts. Nutr Cancer 2013;65:17–24.

36. Pfeiffer CM, Johnson CL, Jain RB, et al. Trends in blood folate and vitamin B-12 concentrations

in the United States, 1988 2004. Am J Clin Nutr 2007;86:718-27.

37. MMWR. Trends in wheat-flour fortification with folic acid and iron–worldwide, 2004 and 2007.MMWR Morb Mortal Wkly Rep 2008;57:8–10.

38. Caudill MA. Folate bioavailability: implications for establishing dietary recommendations and optimizing status. Am J Clin Nutr 2010;91:1455S–60S.

39. Chen P, Li C, Li X, et al. Higher dietary folate intake reduces the breast cancer risk: a systematic review and meta-analysis. Br J Cancer 2014;110: 2327–38.

40. Kim YI. Does a high folate intake increase the risk of breast cancer?. Nutr Rev 2006;64(10 Part 1):468–75.

41. Lalouschek W, Aull S, Serles W, et al. Genetic and nongenetic factors influencing plasma homocysteine levels in patients with ischemic cerebrovascular disease and in healthy control subjects. J Lab Clin Med 1999;133:575–82.

42. Bassett JK, Baglietto L, Hodge AM, et al. Dietary intake of B vitamins and methionine and breast cancer risk. Cancer Causes Control 2013;24:1555–63.

43. Powers HJ. Interaction among folate, riboflavin, genotype, and cancer, with reference to colorectal and cervical cancer. J Nutr 2005;135(Suppl. 12): 2960S–6S.

44. Baylin SB, Herman JG, Graff JR, et al. Alterations in DNA methylation: a fundamental aspect of neoplasia. Adv Cancer Res 1998;72:141–96.

45. Widschwendter M, Jones PA. DNA methylation and breast carcinogenesis. Oncogene 2002;21: 5462–82.

46. Gonzalgo ML, Jones PA. Mutagenic and epigenetic effects of DNA methylation. Mutat Res 1997; 386:107–18.

47. Laufer EM, Hartman TJ, Baer DJ, et al. Effects of moderate alcohol consumption on folate and vitamin B(12) status in postmenopausal women. Eur J Clin Nutr 2004;58:1518–24.

48. Cylwik B, Czygier M, Daniluk M, et al. Vitamin B12 concentration in the blood of alcoholics. Pol Merkur Lekarski 2010;28:122–5.

49. Kabat GC, Miller AB, Jain M, et al. Dietary intake of selected B vitamins in relation to risk of major cancers in women. Br J Cancer 2008;99:816–21.

50. Cho E, Holmes M, Hankinson SE, et al. Nutrients involved in one-carbon metabolism and risk of breast cancer among premenopausal women. Cancer Epidemiol Biomarkers Prev 2007;16:2787–90.

51. Fazili Z, Pfeiffer CM. Measurement of folates in serum and conventionally prepared whole blood lysates: application of an automated 96-well plate isotope-dilution tandem mass spectrometry method. Clin Chem 2004;50:2378–81.

52. Crider KS, Yang TP, Berry RJ, et al. Folate and DNA methylation: a review of molecular mechanisms and the evidence for folate's role. Adv Nutr 2012;3:21–38.

53. Chen J, Gammon MD, Chan W, et al. One-carbon metabolism, MTHFR polymorphisms, and risk of breast cancer. Cancer Res 2005;65:1606–14.

54. Shrubsole MJ, Gao YT, Cai Q, et al. MTHFR polymorphisms, dietary folate intake, and breast cancer risk: results from the Shanghai Breast Cancer Study. Cancer Epidemiol Biomarkers Prev 2004;13:190–6.

55. Maruti SS, Ulrich CM, Jupe ER, et al. MTHFR C677T and postmenopausal breast cancer risk by intakes of one-carbon metabolism nutrients: a nested case-control study. Breast Cancer Res 2009;11:R91.

56. Coughlin SS, Piper M. Genetic polymorphisms and risk of breast cancer. Cancer Epidemiol Biomarkers Prev 1999;8:1023–32.

57. Green R. Indicators for assessing folate and vitamin B-12 status and for monitoring the efficacy of intervention strategies. Am J Clin Nutr 2011; 94:6668–728.

58. EFSA Panel on Dietetic Products NaAN. Scientific opinion on dietary reference values for cobalamin (vitamin B12). EFSA J 2015;13:4150.