



# AperTO - Archivio Istituzionale Open Access dell'Università di Torino

## Perfectionism and cognitive rigidity in anorexia nervosa: Is there an association?

This is a pre print version of the following article:				
Original Citation:				
Availability:				
This version is available http://hdl.handle.net/2318/1695466 since 2019-03-26T10:14:04Z				
Published version:				
DOI:10.1002/erv.2591				
Terms of use:				
Open Access				
Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.				

(Article begins on next page)

# Perfectionism and Cognitive Rigidity in Anorexia Nervosa: is there an association?

Buzzichelli Sara, Marzola Enrica, Amianto Federico, Fassino Secondo, Abbate-Daga Giovanni.

Eating Disorder Center, Department of Neuroscience, University of Turin, Via Cherasco 11, 10126 Torino.

Correspondence Prof. Giovanni Abbate-Daga Eating Disorder Center, Department of Neuroscience, University of Turin Via Cherasco 11, 10126, Turin, Italy. Tel: +39 011 5707701 Fax: +39 011 6335749 Email: giovanni.abbatedaga@unito.it

#### Abstract

Little is known about the relationship between neuropsychology, personality and eating psychopathology in anorexia nervosa (AN). We aimed to investigate the interaction between set-shifting and perfectionism in AN and to ascertain the role of perfectionism as a mediator between set-shifting and eating psychopathology. **85** patients with AN and 71 healthy controls completed: Eating Disorder Inventory-2, Beck Depression Inventory, Wisconsin Card Sorting Test, Trail Making Task and Hayling Sentence Completion Task. Our findings support heightened cognitive inflexibility in individuals with AN, particularly in those with high perfectionism traits. Considering the sample as a whole, perfectionism resulted to be a mediator of the relationship between a measure of set-shifting (i.e., Trail Making Task) and drive for thinness but this finding **did not** remain significant when including in the model only those with AN. Taken together, these data suggest a complex and non-exclusive association between set-shifting, eating psychopathology and perfectionism.

#### Keywords

Eating Disorders Anorexia Nervosa Cognitive Rigidity Set Shifting Perfectionism

# Introduction

Anorexia nervosa (AN) is a complex mental illness whose pathogenesis is largely unknown (Brockmeyer, Friederich, & Schmidt, 2017; Bulik, Berkman, Brownley, Sedway & Lohr, 2007).

Several studies explored the possibility that cognitive rigidity could be an endophenotype of AN (Talbot, Hay, Buckett & Touyz, 2015). Set-shifting has been investigated as an endophenotype also because AN patients consistently showed inefficient set-shifting abilities which have been consistently found to be stable in AN (Abbate-Daga et al., 2011; Abbate-Daga, Buzzichelli, Marzola, Amianto & Fassino, 2014; Tchanturia et al., 2012; Aloi et al., 2015; Westwood, Stahl, Mandy & Tchanturia, 2016). Also personality has been investigated to better understand AN pathogenesis; among personality traits, perfectionism is strongly associated with eating disorders (EDs) as both risk and maintaining factor (Egan, Wade & Shafran, 2011). Interestlyngly, research showed perfectionism not only as correlated with drive for thinness (DT) in patients with AN, but also as preceding eating psychopathology in AN (Halmi et al., 2012).

Little is known about the relationship between set-shifting and personality. Theoretically, rigidity, inhibition, and perfectionism could be linked to difficulties in executive functions. Parallels between perfectionism and set-shifting are common in every-day clinical practice and have been already hypothesized in literature (Schmidt & Treasure, 2006). Furthermore, retrospective research showed set-shifting as associated with childhood perfectionism (Tchanturia et al., 2004); therefore, perfectionism could influence AN patients' thinking style and problem solving abilities thus partially influencing their neuropsychological performances.

The interaction between neuropsychological performance and perfectionism in AN received scant attention and only contradictory findings are available. In a recent study, higher personal standards (Frost, Marten, Lahart & Rosenblate, 1990), were associated with better Trial Making Test scores in both affected and non-affected individuals (Vall & Wade, 2015). In another study higher personal standards were found to be correlated with good set-shifting in recovered AN patients and with poor set-shifting in healthy controls (Lindner, Fichter & Quadflieg, 2014). Buhren and collaborators (2012) found a correlation between greater accuracy on neuropsychological test and perfectionism in young AN patients (Shott et al., 2012). In contrast, other studies highlighted the need for clarifying the interactions between perfectionism and set-shifting (Pignatti & Bernasconi, 2013) and no significant correlations between these two constructs have been reported as well (Friederich et al., 2012). The available evidence is overall unsatisfactory also because previous studies used different measures of set-shifting and cognitive flexibility in verbal domains has not been investigated so far in this regard.

In order to bridge the aforementioned gaps in literature, this study has the following goals: a) to analyze the set-shifting performance concerning both verbal and non-verbal domains across subsamples of AN patients and healthy controls (HCs) grouped according to their degrees of perfectionism; b) to ascertain the role of perfectionism as a mediator between cognitive rigidity and eating psychopathology, in particular drive for thinness. When designing this study we hypothesized to find: a) a relationship between suboptimal neuropsychological performance and perfectionism; b) perfectionism as mediator of the link between cognitive rigidity and eating psychopathology.

#### **Materials and Methods**

#### **Participants**

The sample consisted of 156 adult female participants, 85 individuals diagnosed with AN restricting subtype (RAN) and 71 healthy controls (HCs). AN patients were consecutively recruited at the Eating Disorders Center of University of Turin, Italy. All participants provided written informed consent according to the ethical committee of University of Turin. Patients were included in this study who met criteria for AN according to DSM-5 (American Psychiatric Association, 2013), HCs were recruited at the University of Turin through flyers. Exclusion criteria for the entire sample are the following: a) male gender; b) Wechsler Adult Intelligence Scale-Revised Intellectual

Quotient < 85 (Wechsler, 1997); c) history of cranial trauma with loss of consciousness; d) alcohol or substance dependenceor abuse ; e) medical problems; f) suicidal ideation. Moreover, none of the HCs was on psychotropic medications or met criteria for a current or lifetime diagnosis of EDs or other mental disorders, as clinically assessed by an experienced psychiatrist. Participants were all Caucasian, all affected individuals completed the assessment in the first week of treatment to minimize confounders due to the treatment.

#### Measures and procedure

#### Psychiatric assessment

All participants were assessed using the following self-reported measures: Eating Disorders Inventory-2 (EDI-2; Garner, 1991) to evaluate eating psychopatology and Beck Depression Inventory (BDI; Beck et al., 1961) to assess depression. Participants' weight and high were measured by a trained nurse to calculate body mass index (BMI). Perfectionism was measured with the perfectionism sub-scale of the EDI-2, and we set the cut off at 8 according to the EDI-2 manual instructions (Garner, 1991). Three groups emerged: RAN High Perfectionism (RAN-HP; n=21), RAN Low Perfectionism (RAN-LP; n=64) and HCs with low perfectionism scores (n=64). Seven individuals in the HCs group were outliers and thus excluded.

To evaluate cognitive flexibility we used the following pen and paper neuropsychological tests: Wisconsin Card Sorting Test (WCST; Berg, 1948; Heaton, 1981), Trail Making Task (TMTA-B; Reitan, 1958) and the Hayling Sentence Completion task (HSCT; Burgess & Shallice, 1997).

# Statistical Analysis

The SPSS 24.0 statistical software package (IBM SPSS Statistics for Windows, Version 24.0. Armonk, NY: IBM Corp) has been used for data analysis. Sample t-test has been run to evaluate significant differences between RAN versus HCs. Cohen's d effect size has been calculated for neuropsychological tests. A Chi Square analysis has been used to evaluate the distribution of perfectionism score between RAN and HCs groups; seven HCs were considered as

outliers and thus excluded from all subsequent analyses. To check clinical differences between RAN-HP and RAN-LP a t-test has been used. The one way ANOVA with Bonferroni post-hoc analysis evaluated differences between RAN-HP, RAN-LP, and HCs.

Linear regression analyses were used to: a) explore the eventual independent associations between cognitive rigidity and both perfectionism and Drive for Thinness (DT); b) verify the possible association between perfectionism and DT. Multivariate regression analysis and the Sobel test (Baron & Kenny, 1986; Preacher & Hayes, 2004) were then used in order to ascertain the mediation role of perfectionism. Perfect mediation holds if the independent variable has no effect when the mediator is controlled for (Baron & Kenny, 1986). The amount of mediation is usually called the "indirect effect" (Figure 1).

## Results

# Demographics and clinical features of the sample

The sample was composed of 156 individuals, 85 were RAN and 71 HCs. Those with RAN were inpatients (see table 1 for sample charactheristics)

## Neuropsychological performances

As shown in Table 1, the RAN group performed significantly worse than HCs on all WCST subscales.

Regarding the TMT, both time scores, part A and B were significantly higher in RAN patients when compared to HCs; in contrast, no significant differences were found in the number of errors in part B (Table 1).

RAN individuals performed significantly worse on the HSCT with both net score and reaction time significantly higher than those of HCs (p<0.001 and p<0.05, respectively; Table 1).

# Neuropsychological performances across perfectionism groups

Within the RAN group, 21 (24.7%) individuals had high perfectionism scores (RAN-HP) and 64 (75.3%) had low perfectionism scores (RAN-LP); within the HCs group 7 individuals

(9.9%) reported high perfectionism while the majority of this group (n=64, 90.1%) had low perfectionism scores. The distribution of high perfectionism scores significantly differed between RAN and HCs (p=0.013). Also in this light, HCs with high perfectionism have been considered as outliers and excluded from subsequent analysis.

Between RAN-HP and RAN-LP no significant differences were found in BMI,, and years of education but RAN-HP showed a significantly longer duration of illness, more severe eating and depressive symptoms (Table 1).

With respect to the differences in neuropsychological performances across perfectionism groups (Table 2), RAN-LP and RAN-HP performed significantly worse than HCs on the WCST global score although the differences between RAN groups did not reach statistical significance. Perseverative errors did not significantly differ across groups; in contrast, RAN-HP differed from HCs in non-perseverative errors with RAN-HP making considerably more errors than HCs. Although not statistically significant, RAN-LP were found to have intermediate scores between RAN-HP and HCs with respect to non-perseverative errors. Finally, HCs reported a significantly lower number of failures than both AN groups which did not differ from each other.

Time A on the TMT resulted significantly lower for HCs when compared to RAN-LP. Time B resulted significantly lower in HCs when compared to both RAN groups also, RAN-HP showed greater scores than RAN-LP but the differences between RAN groups showed only a trend toward statistical significance. Similarly, the score time B-A was significantly higher in RAN-HP than HCs , with RAN-LP scoring an intermediate value; however, only the difference between RAN-HP and RAN-LP reached significance.

Concerning the HSCT, we found that the net score resulted significantly higher in RAN-HP when compared to HCs. The net score of RAN-LP was intermediate between those of RAN-HP and HCs although not reaching statistically significance.

#### **Mediation model**

We used the whole sample to run the mediation model. As a proxy for cognitive rigidity we chose to carry on in the mediation model those indices of cognitive rigidity that significantly differed across groups of perfectionism (Table 2), namely TMT A, TMT B, TMT B-A, WCST global score, WCST non perseverative errors, WCST failures, and HSCT net score. As a first step, as shown in Table 3, we investigated the association between cognitive rigidity and DT finding that TMT B, TMT B-A, and WCST non perseverative errors resulted as significantly associated with DT (p= 0.024; p=0.027; p=0.015, respectively). Secondly, we tested the effect of cognitive rigidity on perfectionism reporting that TMT A, TMT B, and HSCT net score were significantly associated with perfectionism (p=0.016; p=0.017; p=0.005, respectively). Lastly, we found a significant association between perfectionism and DT (p=0.001). Among cognitive rigidity indices, only TMT B was significantly associated with both DT and perfectionism (p= 0.024 and p=0.017, respectively), so we used TMT B to run the mediation model. TMT B was significantly associated to DT but, when controlling for perfectionism, such an association was no longer significant (Table 3) thus supporting perfectionism as a mediator of the relationship between cognitive rigidity and DT. The Sobel test confirmed this finding (Sobel test p= 0.031; Figure 1).

When including only the AN sample in the mediation model the association between cognitive rigidity and both DT and perfectionism was no longer significant (data not shown) while the association between perfectionism and DT remained significant (p=0.001).

#### Discussion

Findings from this study confirmed heightened cognitive rigidity in AN and showed perfectionism to be a significant mediator of the relationship between set-shifting and eating psychopathology; nevertheless, this finding did not remain significant when only patients with AN were included in the same mediation model.

In line with the a priori hypothesis we found a relationship between neuropsychological performance and perfectionism. Moreover, those with high perfectionism reported the most marked

inefficiencies in cognitive rigidity when compared to patients with low perfectionism and HCs; however, the differences between patients with high versus low perfectionism did not reach statistical significance in all comparisons.

Patients with different levels of perfectionism differed on the majority of the neuropsychological tests performed but not all, providing support to a multifaceted interaction between cognitive rigidity and perfectionism. With more detail, all AN patients scored poorly on perseverative errors of the WCST, while non-perseverative errors resulted more linked to perfectionism. This finding is in line with literature showing perfectionism as influencing negatively recovered individuals' set-shifting (Lindner et al., 2014). Moreover, RAN-HP resulted significantly slower than RAN-LP on the TMT Time B-A; this finding is interesting as Time B-A represents the time required to perform cognitive set-shifting without considering the time required to perform the control task (Time A). RAN-HP showed significantly longer shifting time than RAN-LP, thus indicating greater difficulty in applying an effective and flexible strategy in cognitive shifting with perfectionism potentially complicating and slowing down RAN patients' set-shifting ability.

Our findings are not in line with previous research (Val & Wade, 2015) reporting an association between TMT scores and perfectionism in an opposite direction; however, different measures have been used to assess perfectionismin fact, clinical perfectionism (as measured by the EDI-2) could have a pejorative role on cognitive flexibility while high personal standards might be interpreted as enhancing cognitive performance.

Similarly, also verbal domains of cognitive rigidity seemed to be partially linked with perfectionism even though statistical significance was not reached between RAN-HP and RAN-LP. This is a novel finding because very few studies investigated the verbal domains of cognitive rigidity (Abbate-Daga et al., 2011; Pignatti & Bernasconi, 2013).

From a clinical point of view, RAN-HP and RAN-LP differed in duration of illness and eating psychopathology but not in BMI. According to these findings, perfectionism resulted indeed unrelated to malnutrition but rather to AN patients' history and severity. This is consistent with data proposing perfectionism as maintaining factor in AN (Fairburn, Shafran & Cooper, 1999; Schmidt & Treasure, 2006; Lilenfeld, 2011); relatedly, it could potentially increase AN severity and duration (Pignatti & Bernasconi, 2013) In future l'associazione tra perfezionismo, basse performances cognitive, longere duration of illness e maggir sintomatologia psichiatrica potrebbe essere studiata anche in relazione ai tratti autisitic evidenziati in AN (ref,...).

We tested the role of perfectionism as a possible mediator of the association between cognitive rigidity and drive for thinness This hypothesis was then confirmed when the relationship between cognitive rigidity and DT was controlled for perfectionism and such an association did not remain significant. However, caution is required when analyzing this mediation model because of two main reasons: a) only TMT B resulted to mediate this association; b) the mediation effect resulted significant when considering the whole sample (Sobel test p=0.031) but not if only individuals with AN were carried on into the analysis. Considering only patients with AN, perfectionism was significantly associated with DT but set-shifting was unrelated to both DT and perfectionism. On one hand, the sample size could partially explain this discrepancy so studies with more power are needed; notwithstanding, perfectionism would seem to play a partially independent role in the acute stage of AN.

According to our findings, set-shifting and perfectionism resulted to be related factors and this is in line with every-day clinical practice. However, the mediation role of perfectionism should be targeted by future studies. In fact, earlier research (Tchanturia et al., 2004) found that cognitive rigidity in AN was associated with childhood perfectionism but not with perfectionism in adulthood. Similarly to our findings on the AN sample, when eating psychopathology is taken into account, perfectionism and cognitive rigidity seem to have an independent role.

When analyzing these data, some limitations should be borne in mind: first and foremost, perfectionism has been assessed with a unidimensional scale while other instruments could be more detailed in describing perfectionism in a multidimensional way. also, other factors (e.g., anxiety

**т** 0

symptoms) could be taken into account when investigating perfectionism and eating psychopathology. Finally a larger sample would be preferable and AN patients and HCs are not matched with respect to years of education.

A complex and non-exclusive association between set-shifting, eating psychopathology, and perfectionism emerged from this study with perfectionism having a potential mediation role. Treatment approaches have been developed to target such aspects [e.g., cognitive remediation therapy (ref) and perfectionism-focused treatments (Lloyd, Schmidt, Khondoker & Tchanturia, 2015; Tchanturia, Larsson & Adamson, 2016; )] but further research would be recommended to improve the therapeutic armamentarium in this regard.

#### References

- Abbate-Daga, G., Buzzichelli, S., Amianto, F., Rocca, G., Marzola, E., McClintock, S. M., & Fassino, S. (2011). Cognitive flexibility in verbal and nonverbal domains and decision making in anorexia nervosa patients: a pilot study. *BMC Psychiatry*, 11, 162. DOI:10.1186/1471-244X-11-162.
- Abbate-Daga, G., Buzzichelli, S., Marzola, E., Amianto, F., & Fassino, S. (2014). Clinical investigation of set-shifting subtypes in anorexia nervosa. *Psychiatry Research*, 219(3), 592-597. DOI:10.1016/j.psychres.2014.06.024.
- Aloi, M., Rania, M., Caroleo, M., Bruni, A., Palmieri, A., Cauteruccio, M. A., . . . Segura-García, C. (2015). Decision making, central coherence and set-shifting: a comparison between Binge Eating Disorder, Anorexia Nervosa and Healthy Controls. *BMC Psychiatry*, 15, 6. DOI:10.1186/s12888-015-0395-z.

American Psychiatric Association (APA). (2013) Diagnostic and Statistic Manual of Mental

Disorders (5<sup>th</sup> ed, DSM-5). Washinton DC.

- Bardone-Cone, A. M., Wonderlich, S. A., Frost, R. O., Bulik, C. M., Mitchell, J. E., Uppala, S., & Simonich, H. (2007). Perfectionism and eating disorders: current status and future directions. *Clinical Psychology Review*, 27(3), 384-405. DOI:10.1016/j.cpr.2006.12.005.
- Baron, R. M., & Kenny, D. A. (1986). The moderator-mediator variable distinction in social psychological research: conceptual, strategic, and statistical considerations. *Journal of Personality and Social Psychology*, 51(6), 1173-1182.
- Beck, A.T., Ward, C.H., Mendelson, M., Mock, J., Erbaugh, J. (1961). An inventory for measuring depression. *Archives of General Psychiatry* 4, 561–571.
- Berg, E.A., (1948). A simple objective technique for measuring flexibility in thinking. *The Journal* of General Psychology, 39,15–22.
- Boone, L., Soenens, B., & Luyten, P. (2014). When or why does perfectionism translate into eating disorder pathology? A longitudinal examination of the moderating and mediating role of body dissatisfaction. *Journal of Abnormal Psychoogyl, 123*(2), 412-418. DOI:10.1037/a0036254.
- Brockmeyer, T., Friederich, H. C., & Schmidt, U. (2017). Advances in the treatment of anorexia nervosa: a review of established and emerging interventions. *Psychological Medicine*, 1-37. DOI:10.1017/S0033291717002604.
- Bulik, C. M., Berkman, N. D., Brownley, K. A., Sedway, J. A., & Lohr, K. N. (2007). Anorexia nervosa treatment: a systematic review of randomized controlled trials. *International Journal of Eating Disorders*, 40(4), 310-320. DOI:10.1002/eat.20367.
- Burgess, P., & Shallice, T. (1997). *The Hayling and Brixton Test*.: Test manual Bury St Edmunds: Thames Valley Test Company.
- Egan, S. J., Wade, T. D., & Shafran, R. (2011). Perfectionism as a transdiagnostic process: a clinical review. *Clinical Psychology Review*, 31(2), 203-212. DOI:10.1016/j.cpr.2010.04.009.

- Fairburn, C. G., Shafran, R., & Cooper, Z. (1999). A cognitive behavioural theory of anorexia nervosa. *Behavioural Research Therapy*, 37(1), 1-13. DOI: 10.1016/S0005-7967(98)00102-8.
- Friederich, H. C., Walther, S., Bendszus, M., Biller, A., Thomann, P., Zeigermann, S., . . . Herzog,
  W. (2012). Grey matter abnormalities within cortico-limbic-striatal circuits in acute and weight-restored anorexia nervosa patients. *Neuroimage*, 59(2), 1106-1113. DOI:10.1016/j.neuroimage.2011.09.042.
- Frost, R. O., Marten, P., Lahart, C., & Rosenblate, R. (1990). The dimensions of perfectionism. *Cognitive Therapy and Research*, *14*(5), 449–468. DOI:10.1007/BF01172967.
- Garner, D. (1991). Eating Disorder Inventory 2: Professional Manual. Psychological Assessment Resources, Odessa. Giel.
- Halmi, K. A., Bellace, D., Berthod, S., Ghosh, S., Berrettini, W., Brandt, H. A., . . . Strober, M. (2012). An examination of early childhood perfectionism across anorexia nervosa subtypes. *Internationa Journal of Eating Disorders*, 45(6), 800-807. DOI:10.1002/eat.22019.

Heaton, K.R. (1981). A Manual for Wisconsin Card Sorting Test. FI: P.E.A, Odessa

- Lilenfeld, L. R. (2011). Personality and temperament. *Current Topics in Behavioral Neurosciences*, 6, 3-16. DOI:10.1007/7854\_2010\_86.
- Lindner, S. E., Fichter, M. M., & Quadflieg, N. (2014). Set-shifting and its relation to clinical and personality variables in full recovery of anorexia nervosa. *European Eating Disorders Review*, 22(4), 252-259. DOI:10.1002/erv.2293.
- Lloyd, S., Yiend, J., Schmidt, U., Tchanturia, K. (2014). Perfectionism in Anorexia Nervosa: Novel Performance Based Evidence. *PLoS ONE 9*(10), e111697. DOI:10.1371/journal.pone.0111697.
- Lloyd, S., Schmidt, U., Khondoker, M., Tchanturia, K. (2015). Can Psychological Interventions Reduce Perfectionism? A Systematic Review and Meta-analysis. *Behavioural and Cognitive Psychoterapy*, 43, 705-731. DOI:10.1017/S1352465814000162.

- Pignatti, R., & Bernasconi, V. (2013). Personality, clinical features, and test instructions can affect executive functions in Eating Disorders. *Eating Behaviors*, 14(2), 233-236. DOI:10.1016/j.eatbeh.2012.12.003.
- Preacher, K. J., & Hayes, A. F. (2004). SPSS and SAS procedures for estimating indirect effects in simple mediation models. *Behavior Research Methods, Instrments, & Computers, 36*(4), 717-731. DOI:10.3758/BF03206553.
- Reitan, R.M. (1958). Validity of the Trail Making Test as indicator of organic brain damage. *Perception and Motor Skills*, 8:271-276.
- Rigaud, D., Pennacchio, H., Bizeul, C., Reveillard, V., & Vergès, B. (2011). Outcome in AN adult patients: a 13-year follow-up in 484 patients. *Diabetes & Metabolism, 37*(4), 305-311. DOI:10.1016/j.diabet.2010.11.020.
- Schmidt, U., & Treasure, J. (2006). Anorexia nervosa: valued and visible. A cognitive-interpersonal maintenance model and its implications for research and practice. *British Journal of Clinical Psychology*, 45(Pt 3), 343-366. DOI:10.1348/014466505X53902.
- Shott, M. E., Filoteo, J. V., Bhatnagar, K. A., Peak, N. J., Hagman, J. O., Rockwell, R., . . . Frank,
  G. K. (2012). Cognitive set-shifting in anorexia nervosa. *European Eating Disorders Review*, 20(5), 343-349. DOI:10.1002/erv.2172.
- Sutandar-Pinnock, K., Blake Woodside, D., Carter, J. C., Olmsted, M. P., & Kaplan, A. S. (2003). Perfectionism in anorexia nervosa: a 6-24-month follow-up study. *International Journal of Eating Disorders*, 33(2), 225-229. DOI:10.1002/eat.10127.
- Talbot, A., Hay, P., Buckett, G., & Touyz, S. (2015). Cognitive deficits as an endophenotype for anorexia nervosa: an accepted fact or a need for re-examination? *International Journal of Eating Disorders, 48*(1), 15-25. DOI:10.1002/eat.22332.
- Tchanturia, K., Davies, H., Roberts, M., Harrison, A., Nakazato, M., Schmidt, U., . . . Morris, R. (2012). Poor cognitive flexibility in eating disorders: examining the evidence using the Wisconsin Card Sorting Task. *PLoS One*, 7(1), e28331. DOI:10.1371/journal.pone.0028331.

- Tchanturia, K., Lounes, N., & Holttum, S. (2014). Cognitive remediation in anorexia nervosa and related conditions: a systematic review. *European Eating Disorders Review*, 22(6), 454-462. DOI:10.1002/erv.2326.
- Tchanturia, K., Larsson, E., & Adamson, J. (2016). Brief Group Intervention Targeting Perfectionism in Adults with Anorexia Nervosa: Empirically Informed Protocol. *European Eating Disorders Review*, 24(6), 489-493. DOI:10.1002/erv.2467.
- Tchanturia, K., Morris, R. G., Anderluh, M. B., Collier, D. A., Nikolaou, V., & Treasure, J. (2004). Set shifting in anorexia nervosa: an examination before and after weight gain, in full recovery and relationship to childhood and adult OCPD traits. *Journal of Psychiatry Research*, 38(5), 545-552. DOI:10.1016/j.jpsychires.2004.03.001.
- Vall, E., & Wade, T. D. (2015). Predictors of treatment outcome in individuals with eating disorders: A systematic review and meta-analysis. *International Journal of Eating Disorders*, 48(7), 946-971. DOI:10.1002/eat.22411.

Wechsler, D. (1997). WechslerAdultIntelligenceScale - Revised. GiuntiO.S, Firenze, Italy.

Westwood, H., Stahl, D., Mandy, W., & Tchanturia, K. (2016). The set-shifting profiles of anorexia nervosa and autism spectrum disorder using the Wisconsin Card Sorting Test: a systematic review and meta-analysis. *Psychological Medicine*, 46, 1809-1827. DOI: 10.1017/S0033291716000581.

		RAN	НС	Test st	atistics	
		(n=85)	(n=71)			
		Mean(SD)	Mean(SD)	t	р	
Age, years		23.73 (6.68)	24.82 (3.09)	-1.263	0.209	
Years of education		11.99 (2.97)	15.52 (2.67)	-7.735	0.001**	
Body Mass Index		15.06 (1.89)	20.82 (2.19)	-17.618	0.001**	
Duration of illness		6.24 (6.37)	-	-	-	
EDI-2	DT	12.54 (7.24)	2.07 (3.97)	10.886	0.001**	
	В	2.07 (3.46)	1.01 (2.38)	2.176	0.031*	
	BD	13.34 (6.76)	5.48 (5.41)	7.912	0.001**	
BDI		13.77 (7.85)	3.12 (3.77)	10.446	0.001**	
HSCT	Net Score	3.58 (2.60)	2.32 (1.97)	3.331	0.001**	
	Reaction time A	0.502 (0.238)	0.389 (0.233)	2.967	0.003*	
	Reaction time B	2.110 (1.632)	1.518 (1.130)	2.570	0.011*	
ТМТ	Time A	40.69 (15.53)	34.05 (9.27)	3.162	0.002*	
	Time B	79.12 (31.64)	64.42 (17.24)	3.504	0.001**	
	Time B-A	39.28 (26.23)	30.52 (15.10)	2.489	0.014*	
	Errors part B	0.28 (0.64)	0.17 (0.41)	1.247	0.214	
WCST	Global Score	26.40 (21.14)	16.77 (11.49)	3.434	0.001**	
	Perseverations	6.42 (5.01)	4.62 (2.89)	2.686	0.008*	
	No perseverative errors	10.07 (10.09)	6.61 (6.32)	2.510	0.013*	
	Failure	0.62 (1.32)	0.17 (0.61)	2.676	0.008*	

Table 1. Demographic, clinical and neuropsychological assessment

Female AN-R patients and healthy controls; mean and (standard deviation) are provided; statistically significant differences are shown with \* if p<0.05, and \*\* if p<0.001; BMI: Body Mass Index; BDI: Beck Depression Inventory; EDI-2: Eating Disorders inventory-2; DT: Drive for

Thinnes; B: Bulimia; BD: Body Dissatisfaction; WCST: Wisconsin Card Sorting Test; HSCT: Hayling Sentence Completion Task; TMT: Trail Making Test.

Table 2. Neuropsychological performance across groups comparing AN patients with high versus low perfectionism and healthy controls.

		RAN-HP (n=21)	RAN-LP (n=64)	HC (n=64)	Test statistics		atistics
		Mean(SD)	Mean(SD)	Mean(SD)	F	р	Bonferroni post-hoc
HSCT	Net Score	4.05 (2.99)	3.42 (2.46)	2.45 (2.02)	4.662	0.011*	RAN-HP>HC
	Reaction time A	0.508 (0.141)	0.501 (0.263)	0.399 (0.235)	3.354	0.055	-
	Reaction time B	1.850 (1.358)	2.193 (1.711)	1.563 (1.152)	3.017	0.052	-
TMT	Time A	41.90 (13.61)	40.29 (16.19)	34.05 (9.38)	4.685	0.011*	RAN-LP>HC
	Time B	88 95 (39 56)	75 90 (28 20)	64 40 (17 73)	7 650	0.001**	RAN-HP>HC
	Time D	88.95 (39.50)	75.90 (28.20)	04.40 (17.73)	7.030	0.001	RAN-LP>HC
	Time B-A	50.47 (31.65)	35.61 (23.31)	30.53 (15.45)	6.617	0.002*	RAN-HP>AN-LP RAN-HP>HC
	Errors part B	0.33 (0.79)	0.27 (0.59)	0.17 (0.45)	0.784	0.459	-
WCST	Global Score	27.10(21.52)	26.14(21.19)	17.00(12.01)	4 078	0.008*	RAN-HP>HC
west		27.19 (21.55)	20.14 (21.18)	17.09 (12.01)	4.970		RAN-LP>HC
	Perseverations	6.00 (4.75)	6.56 (5.11)	4.77 (2.93)	2.916	0.057	-
	No perseverative errors	12.67 (16.35)	9.22 (6.92)	6.69 (6.63)	3.974	0.021*	RAN-HP>HC
	Failure	0.52 (0.98)	0.66 (1.42)	0.17 (0.63)	3.302	0.040*	RAN-LP>HC

Female AN-HP (high perfectionism), AN-LP (low perfectionism) patients and healthy controls; mean and (standard deviation) are provided; statistically significant differences are shown with \* if p<0.05, and \*\* if p<0.001; WCST: Wisconsin Card Sorting Test; HSCT: Hayling Sentence Completion Task; TMT: Trail Making Test.

-		DT		
		D1		
		F (1.148)	р	
HSCT	Net Score	3.781	0.054	
TMT	Time A	2.660	0.105	
	Time B	5.166	0.024*	
	Time B-A	4.966	0.027*	
WCST	Global Score	3.732	0.055	
	No perseverative errors	6.061	0.015*	
	Failure	1.690	0.196	
TMT	Time B <sup>§</sup>	27.384	0.230	

 Table 3. Linear regressions of the unmediated and mediated model investigating the relationship between set-shifting and drive for thinness

WCST: Wisconsin Card Sorting Test; HSCT: Hayling Sentence Complemion Task; TMT: Trail Making Test. F and p value of the association between Cognitive Rigidity and Drive for Thinness are given. Statistically significant differences are shown in bold under p values. <sup>§</sup>After correcting the model for perfectionism.

# Figure 1. Mediation model

A)Unmediated model



Representation of the mediation model. A) The direct effect of cognitive rigidity on drive for Thinness is represented, B) The mediator role of perfectionism is shown and Sobel Test is given. <sup>§</sup>After correcting the model for perfectionism.