

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Anti-inflammatory and anti-diabetic effects of anthocyanins from colored cereals.**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1830649> since 2022-01-06T18:44:12Z

*Publisher:*

BOKU University

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

# ANTI-INFLAMMATORY AND ANTI-DIABETIC EFFECTS OF ANTHOCYANINS FROM COLORED CEREALS

Abbasi Parizad, Parisa<sup>1</sup>; Capraro, Jessica<sup>1</sup>; Scarafoni, Alessio<sup>1</sup>; Bonomi, Francesco<sup>1</sup>; Blandino, Massimo<sup>2</sup>; Giordano, Debora<sup>2</sup>; Carpen, Aristodemo<sup>1</sup>; Iametti, Stefania<sup>1</sup>

<sup>1</sup>Department of Food, Environmental and Nutritional Sciences (*DeFENS*), University of Milan, Italy; <sup>2</sup>Department of Agricultural, Forest and Food Sciences (*DISAFA*), University of Turin, Italy

[parisa.abbasi@unimi.it](mailto:parisa.abbasi@unimi.it)

**Introduction:** Anthocyanins are located in aleuronic or pericarp layers of various types of pigmented cereal grains. Many studies have been reported on the health-promoting effects of anthocyanins and on their anti-inflammatory and antidiabetic activities. This study was aimed at investigating the anthocyanin profile of pigmented grains, and at scrutinizing bioactive molecules by assessing their effects on the expression of inflammatory markers and on the activity of pancreatic and brush border enzymes.

**Material & Methods:** The present study compares five different colored grains: three corn samples and two wheat samples. A pH-differential method was used to measure total anthocyanin content, while anthocyanin profiling was carried out by means of RP-HPLC/DAD. Antioxidant capacity was determined by means of the FRAP assay. Immunomodulating effects were evaluated by using cytokine-stimulated biosynthesis of inflammation factors in Caco-2 cells transfected with a pNiFty2-Luc vector. In these studies, cyanidin-3-glucoside was used as positive control. Inhibitory activity against intestinal  $\alpha$ -glucosidase and pancreatic  $\alpha$ -amylase was also assessed.

**Results:** Four major anthocyanins were identified by HPLC profiling: cyanidin-3-glucoside was predominant in purple wheat, and delphinidin in blue corn. The highest total anthocyanin content was detected in purple wheat (0.74 mg/g), followed by some of the blue corn varieties (0.66 mg/g and 0.53 mg/g in cultivars T and MF, respectively), whereas non-blue corn had very low levels (0.18mg/g). Conversely, the antioxidant capacity was the highest in the non-blue Rostrato corn variety (47.2  $\mu$ molFe(II)/g), comparable to that of the purple wheat (47.5  $\mu$ molFe(II)/g). The highest ferulic acid concentration was detected in blue corn (cultivar T) and in the red-colored Rostrato corn, whereas specific fractions derived from de-branning of blue wheat (cultivar Skorpion) had the highest rutin content.

The immune response of Caco-2 cells decreased significantly in a dose-responsive manner for all grains extracts. Extracts from blue corn (cultivar SK) and purple wheat were effective in decreasing inflammatory response at anthocyanin concentrations comparable to those of pure cyanidin-3-glucoside. Purple wheat and blue corn (cultivar T) extracts also displayed significant inhibition of  $\alpha$ -amylase activity. Values ranged between 78% (wheat)

and 95.4% (corn) at final concentrations of extract solids in the 100-130 µg/ml range. Under comparable conditions, brush border α-glucosidase inhibition from the same extracts was 57.8 % (wheat) and 65.4% (corn).

**Conclusion:** This study highlights the promising antioxidant activity, anti-inflammatory, and antidiabetic effects of anthocyanins-rich extracts from corn and wheat. These cereals can be incorporated into staple grain-based foods such as pasta, bread, noodles, or snacks, but also in other types of foods, such as dairy products or fiber-rich drinks. In this frame, they could be regarded as both a natural food colorant and as a functional food ingredient. Their use as natural dietary supplements deserves further attention, given their ability to control the carbohydrate metabolism and to modulate inflammation. Also worth of further studies are their reported effects on cardiovascular risk factors, as well as their effects on the intestinal microbiota.

*References:*

*Nankar, Amol N., et al. "Quantitative and qualitative evaluation of kernel anthocyanins from southwestern United States blue corn." Journal of the Science of Food and Agriculture 96.13 (2016): 4542-4552.*

*Huang, Bo, et al. "Anti-diabetic effect of purple corn extract on C57BL/KsJ db/db mice." Nutrition research and practice 9.1 (2015): 22-29.*

*Tsuda, Takanori, et al. "Dietary cyanidin 3-O-β-D-glucoside-rich purple corn color prevents obesity and ameliorates hyperglycemia in mice." The Journal of nutrition 133.7 (2003): 2125-2130.*

*Kim, Eun Ok, et al. "Anti-inflammatory activity of hydroxycinnamic acid derivatives isolated from corn bran in lipopolysaccharide-stimulated Raw 264.7 macrophages." Food and Chemical Toxicology 50.5 (2012): 1309-1316.*

*Miguel, Maria Graça. "Anthocyanins: Antioxidant and/or anti-inflammatory activities." (2011).*