

27 showed a high frequency of *Aeromonas spp* contamination in *Sparus aurata* reared on the Italian
28 coast and an elevated biodiversity in isolated bacterial strains. *Aeromonas* isolates comprise
29 potentially pathogenic species for humans, often resistant to several antibiotics and able to transfer
30 the genes responsible for antibiotic resistance to microorganisms pathogenic for humans throughout
31 the food chain. The few ECV studies available on many antibiotics against *Aeromonas spp* strains
32 isolated from the aquaculture environment highlight the need for further research in this area, while
33 regular monitoring programmes should be stepped up to check for antibiotic resistance.

34

35 **Keywords:** gilthead seabream; *Aeromonas spp* bacteria; minimum inhibitory concentration;
36 antibiotics; epidemiological cut-off values.

37

38

39 1. Introduction

40 World fish consumption has been growing in the last thirty years reaching 20 kg per capita in 2014.

41 For the first time, the aquaculture production of fish for human consumption has overtaken the
42 supply of wild-caught fish and is expected to rise to 62% by 2030 (FAO, 2016). Gilthead sea bream
43 (*Sparus aurata*) is a very suitable species for mariculture in the Mediterranean basin and has
44 become one of Europe's main fish species in aquaculture. Greece, Turkey and Spain are the main
45 producers worldwide while Italy is the third main producer in the EU (EC, 2017). Large-scale
46 aquaculture is characterized by the intensive and semi-intensive production systems with high
47 stocking density, which leads to poor hygiene conditions and the emergence of infectious diseases
48 (Diana et al., 2013).

49 The genus *Aeromonas* comprises a group of bacteria with a ubiquitous distribution in natural
50 habitats, including the aquatic environment (Janda & Abbott, 2010) where species such as *A.*
51 *hydrophila*, *A. caviae*, *A. salmonicida* and *A. veronii* biovar *sobria* cause disease in marine fish
52 (Radu et al., 2003). *Aeromonas spp* are also important human opportunistic pathogens able to cause

53 intestinal, blood, skin and soft tissue and trauma-related infections, particularly in young children
54 and **the** elderly (Janda and Abbott, 2010; Real et al., 1994). *Aeromonas* species have been
55 frequently isolated from fish and other foods (Callister and Agger, 1987; Gobat and Jemmi, 1993).
56 **These bacteria are responsible for food spoilage and may serve as vectors for disease transmission**
57 **to humans** (Tsai and Chen, 1996). Infection can **also occur** after contact with contaminated **water** or
58 fish (Janda and Abbott, 2010). ***Aeromonas* pathogenicity is linked** to the production of a number of
59 extracellular hydrolytic enzymes such as lipases and proteases, which **aid in bacterial invasion and**
60 **the establishment of infection** (Galindo et al., 2006). Among an array of other virulence factors, the
61 **biological activities of** cytotoxic enterotoxin (Act) **include haemolysis**, cytotoxicity, enterotoxicity
62 and lethality (Chopra et al., 1991).

63 The **worldwide** expansion of intensive fish farming has increased the use of antibiotics to treat
64 bacterial **infections** (Díaz-Cruz et al., 2003). In aquaculture, antimicrobials are generally added to
65 the feed or directly to the water to prevent the spread of infectious fish disease (Defoirdt et al.,
66 2011) and in some circumstances **to promote fish growth illegally** (Serrano, 2005). **Regulations**
67 **governing the use of antibiotics in aquaculture differ widely** with little to no enforcement in many
68 of the world's major **aquaculture-producing** countries (Pruden et al., 2013). The extensive use of
69 antibiotics in aquaculture has in turn resulted in the emergence of antibiotic resistance in both
70 foodborne and opportunistic human **pathogens** (Marshall and Levy, 2011). The resistance of
71 *Aeromonas* species to diverse groups of antibiotics **is** a major concern for human health (Figueira et
72 al., 2011) **as resistant bacteria can spread** from the aquatic environment **to humans via the** food
73 chain or direct contact (Taylor et al., 2011). **In addition, resistance genes can be transferred** by
74 mobile genetic elements such as plasmids, phages and transposons (Levy and Marshall, 2004).

75 Janda and Abbott (2010) reviewed the general susceptibility profiles of Aeromonads to various
76 antimicrobial classes, showing resistance to **sulfamethoxazole**, **cephalosporins**, **penicillins**
77 (**amoxicillin**, **ampicillin**, **ampicillin-sulbactam**, **ticarcillin**, **oxacillin** and **penicillin**) and **macrolides**
78 (**clarithromycin**). ***Aeromonas* species resistant to penicillins** and first generation **cephalosporins** are

79 associated with the production of chromosomally encoded beta-lactamases (Janda and Abbott,
80 2010). Other important resistance determinants to beta-lactam antimicrobials and tetracyclines are
81 *bla* genes and *tet* genes respectively encoded in mobile genetic elements (Agersø et al., 2007; Wu et
82 al., 2011) or integrons, responsible for resistance to tetracyclines, aminoglycosides,
83 chloramphenicol and trimethoprim (Chang et al., 2007; Kadlec et al., 2011). Indeed, acquired
84 antibiotic resistance among fish pathogens could determine serious therapeutic problems in humans
85 following the use of molecules whose class and structure are similar or, in some cases, identical to
86 those used in mariculture (Cabello, 2006). Despite recent efforts by international agencies such as
87 the European Centre for Disease Prevention and Control and the National Antimicrobial Resistance
88 Monitoring System (EFSA, 2014), the role of antibiotic usage in aquaculture in the development
89 and dissemination of antibiotic resistance genes is still poorly understood. The potential risk of
90 transferring such resistance from the aquaculture environment to humans is underestimated
91 (Cabello et al., 2013) so the effectiveness of antibiotics used in fish farming should be carefully
92 monitored.

93 Little information is available on the susceptibility of *Aeromonas* spp isolated from mariculture to
94 antibiotics used in both fish farming and human therapy. Antimicrobial susceptibility is generally
95 tested by measuring the drug's minimum inhibitory concentration (MIC). MIC breakpoints are the
96 MICs at which an organism should be considered susceptible, intermediate or resistant. Breakpoint
97 values are published by organizations such as the European Committee on Antimicrobial
98 Susceptibility Testing (EUCAST) and the American Clinical Laboratory Standard Institute (CLSI),
99 based on pharmacokinetic/pharmacodynamic data and clinical studies. MIC₅₀ and MIC₉₀ indicate
100 the lowest concentrations of the antimicrobial agent inhibiting visible growth of 50% and 90% of
101 the bacterial population respectively. However, few interpretation criteria for *Aeromonas* spp have
102 been published to date. The only available breakpoints proposed by the CLSI are from clinical
103 isolates adapted from *Enterobacteriaceae*, while no criteria have been established by EUCAST.
104 Epidemiological cut-off values (ECVs) must be set to discriminate wild-type strains (with no

105 acquired resistance mechanism to the tested antibiotic) from non-wild-type strains (with one or
106 more acquired resistance mechanisms) (Kahlmeter et al., 2003). These cut-off values are the upper
107 limit of the MIC distribution of fully susceptible strains.

108 The purpose of the present study was to estimate the MICs of *Aeromonas* spp strains isolated from
109 *Sparus aurata* against 15 antimicrobial agents, and to determine the ECVs for *Aeromonas* spp.

110

111 2. Materials and methods

112 2.1 Fish sampling

113 The study was conducted on gilthead sea bream (*Sparus aurata*) collected from six offshore
114 mariculture farms in three Italian regions (Sardinia, Sicily and Tuscany). All fish farms were
115 characterized by intensive rearing systems in sea cages. Water salinity was ca. 33‰ and the
116 temperature ranged between 16°C and 22°C. Twenty commercial size (~250 g) *Sparus aurata*
117 specimens were randomly collected at each farm during two different visits conducted four months
118 apart. After collection, fish were slaughtered by immersion in fusing ice, placed in expanded
119 polystyrene boxes and covered with a plastic film then transported to the laboratory under
120 refrigeration and processed within three hours after collection.

121 2.2 Microbiological analysis

122 Samples of skin, gills, muscle and intestinal content were aseptically collected from each specimen
123 for microbiological analysis. The initial suspension and decimal dilution for microbiological
124 examination were prepared according to ISO 6887–1:1999. Each matrix was tested for *Aeromonas*
125 species (presence/absence) inoculating 0.1 mL of homogenized PBS (pH 7.4) on plates of
126 *Aeromonas* Medium Base (Ryan's medium) (Oxoid, Basingstoke, UK) supplemented with
127 ampicillin selective supplement at 5 mg/L. The agar plates were incubated at +30°C for 48 hours.
128 Colonies with typical growth characteristics, opaque dark green with darker centres, were picked
129 and subcultured on brain heart infusion (BHI) agar plates (BHI, Oxoid, Basingstoke, UK). After
130 incubation, isolates were tested as follows: morphology in Gram staining, cytochrome oxidase,

131 amylase and trealose fermentation. After presumptive genus identification, strains were stored at -
132 80°C for subsequent genetic confirmation and species identification.

133 2.3 Bacterial identification

134 Genus identification of isolates was confirmed by PCR (Khan et al., 2009). To avoid over-
135 representation of clones, 16S ribosomal DNA sequencing was conducted on a selection of strains to
136 identify bacterial species. A hierarchical method was used to select up to three strains from each of
137 the following nested criteria: region of collection (three levels), fish farm (two levels), fish
138 specimens (40 levels) and fish matrix (4 levels). For species identification, colonies with
139 morphological and biochemical features of *Aeromonas* spp were grown overnight at 37°C in
140 tryptone soya broth (Oxoid). DNA was extracted using the following protocol: 1mL of broth culture
141 (10^8 CFU/mL) was centrifuged at 12,000 g for five minutes, then the pellet was resuspended in 1mL
142 of phosphate-buffered saline, boiled for five minutes, and centrifuged again (Bottero et al., 2004).
143 The supernatant was stored at -20°C until use. The DNA was quantified using a spectrophotometer
144 (Nanodrop 2000, Thermo Fisher Scientific). All extracted DNA were subjected to sequencing
145 analysis with the MicroSeq 500 16S rDNA bacterial sequencing kit (Thermo Fisher Scientific). 16S
146 rDNA amplicons were purified by Exo-Sap treatment according to the manufacturer's
147 recommendations (USB Europe, Staufeu, Germany). Forward and reverse sequencing reactions
148 were performed using the ABI Prism BigDye Terminator Cycle Sequencing Ready Reaction Kit,
149 version 1.1 (Thermo Fisher Scientific). The extended products were purified with DyeEx 2.0 Spin
150 kit (Qiagen, Valencia, CA, USA) and resolved by capillary electrophoresis using an ABI 310
151 Genetic Analyzer (Thermo Fisher Scientific). The electropherograms were analyzed using Chromas
152 2.22 software (Technelysium, Epoch Life Science Inc.) and the sequences were submitted to the
153 BLAST similarity search software on the National Center for Biotechnology Information (NCBI)
154 website.

155 2.4 Antibiotic susceptibility

156 Antibiotic susceptibility was determined for *Aeromonas* strains at the genus level. MICs of 15
157 antibiotics were measured by the broth microdilution method (CLSI, 2011). The antimicrobial
158 agents chosen among those mainly used in aquaculture and human therapy were: oxolinic acid
159 (OXA), ampicillin (AM), amoxicillin (AMX), cephalothin (CF), cloramphenicol (CL),
160 erythromycin (E), florfenicol (FF), flumequine (FM), gentamicin (GM), kanamycin (K),
161 oxytetracycline (OT), streptomycin (S), sulfadiazine (SZ), tetracycline (TE) and trimethoprim
162 (TMP). To obtain stock solutions, the antibiotic powders (Sigma Aldrich, MI, Italy) were weighed
163 and dissolved in the following solvents (Sigma Aldrich): phosphate buffer, pH 8.0, 0.1 mol/L (AM),
164 phosphate buffer, pH 6.0, 0.1 mol/L (AMX and CF), ethanol 95% (CL and E), methanol 96% (FF),
165 aqueous alkaline solution NaOH 0.1M + ethanol 2:1 (FM), aqueous alkaline solution NaOH 0.1M,
166 pH 10 (OXA), methanol-water 2:1 (OT and TE), aqueous acidic solution HCl 10% (TMP), and
167 deionized water (GM, K, S and SZ). Once dissolved, each stock solution (2,560 µg/mL) was
168 dispensed in 1.5 mL aliquots into polypropylene vials and frozen at - 80 °C until use. Each
169 microtitre plate was prepared with 12 serial twofold dilutions of each antibiotic stock solution
170 (Work Station - Micro Star, Hamilton, Bonaduz GR, Switzerland) with deionized water (phosphate
171 buffer, pH 6.0, 0.1 mol/L, only for AMP and AMX antibiotics). The antibiotic concentrations
172 obtained ranged between 0.06 µg/mL and 128 µg/mL (0.12-256 µg/mL for SZ antibiotic). Strains
173 were subcultured twice in BHI plates before preparation of the inoculum. After overnight
174 incubation at 37 °C, two or more colonies were picked from BHI plates and dissolved in salt
175 solution (0.85% w/v) to obtain 0.5 McFarland turbidity, measured using a portable photometric
176 reader (Densimat, bioMérieux, Lyon, France). Each bacterial suspension was further diluted (1:100)
177 in cation-adjusted Mueller Hinton broth (CAMHB, Oxoid, Basingstoke, UK) supplemented with
178 NaCl (1%) to obtain an inoculum concentration of ca. 10^6 cfu/mL. Fifty µL of the final suspension
179 were transferred into microtitre wells (one strain for each row of the microplate) containing 50 µL
180 of each antimicrobial agent. The density of the final inoculum in each well was ca. 5×10^5 cfu/mL.
181 The reference strain *E. coli* ATCC 25922 was used as quality control. Each microplate was

182 subsequently incubated under aerobic conditions for 20 hours at 35 °C. The MIC of each antibiotic
183 was compared with breakpoint values to determine resistance (CLSI, 2005, 2007, 2011, 2016;
184 NCCLS, 1998, 1999, 2002). The MIC range and mode, MIC₅₀ and MIC₉₀ of each antimicrobial
185 agent were also determined. Multiple antibiotic resistance (MAR) among *Aeromonas* spp strains
186 was evaluated applying the MAR index defined as a/b , where “a” was the number of antimicrobials
187 the isolate was resistant to and “b” was the number of antibiotics against which the isolate was
188 tested. According to Krumperman (1985), a MAR index below 0.2 is interpreted as strains
189 originating from animals in which antibiotics are seldom or never used, while a MAR index above
190 0.2 is interpreted as strains originating from an elevated selective pressure environment where
191 antibiotics are frequently used.

192 2.5 Epidemiological cut-off values

193 The distribution of MIC values served to determine the ECVs. The statistical determination of ECV
194 values for each antimicrobial agent at genus level was conducted according to Turnidge et al.
195 (2006) using the freely available ECOFFinder Microsoft Excel spreadsheet calculator
196 (<https://clsi.org/education/microbiology/ecoffinder/>). The spreadsheet is designed to estimate the
197 ECVs based on the observed MIC of the tested bacterial population, i.e. it estimates the MIC value
198 best describing where wild type distribution ends.

199

200 3. Results

201 3.1 Isolation and identification of *Aeromonas* spp

202 *Aeromonas* spp were observed in 98 skin samples (30.6%), 154 gills (48.2%) and 40 gut contents
203 (12.5%) whereas the bacteria were never detected in muscle. One hundred and four *Aeromonas* spp
204 strains isolated from *Sparus aurata* were speciated by 16S ribosomal DNA sequencing. Of the total
205 strains, 59 originated from Tuscany, 40 from Sicily and five from Sardinia. The *Aeromonas* strains
206 were isolated from skin (n. 48), gut content (n. 20) and gills (n. 36). Sequencing identified 23
207 different *Aeromonas* species or species-complex. The most frequently recovered species were

208 *Aeromonas media* (15 strains, 14.4%), *Aeromonas salmonicida/bestiarium/hydrophila/caviae*
209 species-complex (12 strains, 11.5%), *A. molluscorum* (11 strains, 10.6%) and *A. bivalvum* (10
210 strains, 9.6%). Table 1 reports a complete list of *Aeromonas* species and species-complex identified
211 and the relative number of strains. Fig. 1 shows their distribution by region of origin.

212 3.2 Antimicrobial susceptibility

213 For some of the selected antimicrobial agents, the reference strain used as quality control for the
214 MIC determination assay indicated the *Aeromonas* strains in compliance with CLSI
215 recommendations (CLSI, 2005). Over 90% of the speciated *Aeromonas* strains showed
216 susceptibility to CL, FF and GM antibiotics. Table 1 lists the resistance profile for each *Aeromonas*
217 species. All tested *Aeromonas* strains showed resistance to two or more antibiotics. One strain of *A.*
218 *bivalvium* and one strain of the *A. punctata/hydrophila/enteropelogenes* species-complex were
219 resistant to 11 antibiotics while one *A. molluscorum* strain was resistant to 12 antibiotics. Table 2
220 reports the MAR index indicating the multiple antibiotic resistance of *Aeromonas* spp strains by
221 region of origin. Table 3 shows the MIC₅₀, MIC₉₀, mode and range and cut-off of MICs for each
222 tested antibiotic, and the number of sensitive, intermediate and resistant strains with reference to the
223 CLSI breakpoints. The most frequent combination of antibiotic resistance profiles was AM, AMX,
224 CF, E, S, SZ and TMP. For CF and SZ, the MIC₅₀ and MIC₉₀ values were above the tested range
225 (128 and 256 µg/mL respectively).

226 3.3 Determination of wild-type strains

227 The ECVs were computed for 12 out of 15 antimicrobial agents. In addition to CF and SZ, no
228 values were computed for TMP due to a high number of isolates with MIC values greater than the
229 upper limit of the tested dilutions (128 µg/mL).

230 The wild-type strains ranged between 59.6% and 96.2% of the tested strains. Thirty-one strains
231 (29.8%) were wild-type for all antibiotics with a computable ECV. One strain (*A. bivalvium*)
232 resulted wild-type exclusively for E, one strain for GM and K (*A.*
233 *punctata/hydrophila/enteropelogenes* species-complex) and one strain (*A. molluscorum*) for GM, K

234 and S. The remaining 70 strains were wild-type for five up to 11 different antibiotics, yielding 31
235 different combinations of antibiotic wild-type profiles. More than 80% of wild-type strains were
236 resistant to eight antibiotics (CL, E, FF, GM, K, OT, S, TE). Table 3 reports the complete results on
237 the ECVs and percentage of wild-type strains.

238

239 4. Discussion

240 The worldwide growth of aquaculture has seen the development of intensive fish farming. This in
241 turn has been associated with an extensive use of antibiotics to treat or prevent bacterial infections.
242 Regulations governing the antimicrobial agents authorized in fish farming differ from country to
243 country. The selective pressure exerted by intensive fish farming has resulted in the emergence of
244 antibiotic-resistant food-borne pathogens, opportunistic pathogens and human commensal flora of
245 food animals (Sorum, 2006; Teuber, 2001; Witte, 2000). The potential transfer of antibiotic
246 resistance from the aquatic environment to humans through direct contact or via the food chain is a
247 serious concern for human health (Marshall and Levy, 2011). Antibiotic resistance monitoring fails
248 to collect extensive information of the classes of antimicrobials used in aquaculture and the efficacy
249 of antibiotics (Cabello et al., 2013).

250 The present study provided useful information on the resistance of *Aeromonas* spp isolated from
251 gilthead sea bream (*Sparus aurata*) reared in Italian fish farms. *Aeromonas* spp were widely
252 distributed in skin, gills and intestinal content of *Sparus aurata* whereas they were never detected in
253 muscle. *Aeromonas* spp can potentially cause human illness by direct contact or through the
254 ingestion of contaminated fish (Janda & Abbott, 2010).

255 Clinical breakpoints are useful to assess the efficacy of antibiotics during treatments, while the
256 determination of ECVs will establish the emergence of antibiotic resistance mechanisms within a
257 bacterial population. Based on these values, the present study documented high resistance rates for
258 β -lactams, erythromycin, sulfadiazine and trimethoprim. The MIC₉₀ of ampicillin, amoxicillin and
259 cephalothin (>128 μ g/mL) were higher than the reference breakpoints for resistance and the number

260 of resistant strains ranged between 40.4% and 86.5%. The literature reports resistance rates for
261 these antibiotics as high as 100% (Hatha et al., 2005; Snoussi et al., 2011). The MIC₅₀ for ampicillin
262 and amoxicillin was 16 µg/mL, an intermediate value between the reference breakpoints for
263 susceptibility and resistance, while the MIC₅₀ for cephalothin was greater than the reference value
264 for resistance (>128 µg/mL). Amoxicillin and ampicillin are susceptible to β-lactamase and to rapid
265 onset antibiotic resistance especially when the antibiotic is repeatedly used in a short time period,
266 typical of intensive fish farming systems. Three different types of β-lactamase have been observed
267 in *Aeromonas* spp (Walsh et al., 1997), but little information is available on the ECV_{ST} for
268 *Aeromonas* spp and limited to few antibiotics, hampering a comparison with the MIC distribution
269 observed in our microbial population. Despite the high resistance rates observed for β-lactam
270 antibiotics, based on the ECVs computed for amoxicillin and ampicillin, an elevated percentage of
271 strains could be considered wild-type.

272 Among the quinolone antibiotics, various countries have authorised oxolinic acid (a first generation
273 quinolone) for therapeutic use in aquaculture, while flumequine is the only one of the five
274 fluoroquinolone antibiotics listed in Reg. EC 37/2010 authorised for fish farming. These antibiotics
275 are used in mariculture for the treatment of furunculosis caused by *Aeromonas salmonicida* (Giraud
276 et al., 2004). Oxolinic acid and flumequine showed resistance in 32.7% and 22.1% of the tested
277 *Aeromonas* spp strains respectively. The antibiotic resistance of *Aeromonas* spp in the present study
278 is in agreement with previous investigations conducted in mariculture farms where resistance was
279 between 25% and 50% (Inglis et al., 1991; Snoussi et al., 2011; Cattoir et al., 2008).

280 The ECV computed in the present study was 0.25 µg/mL for both oxolinic acid and flumequine
281 while the literature reported values of 0.031 µg/mL and 0.06 µg/mL respectively (Baron et al.,
282 2017; Smith and Kronvall, 2015). However, these results are not comparable as the values obtained
283 in our study coincided with the lowest dilution tested.

284 Due to their broad-spectrum activity, low toxicity and cost, tetracyclines are the most commonly
285 used antibiotics in both human and veterinary medicine. In mariculture, oxytetracycline is

286 authorized for therapeutic immersion in Europe, while elsewhere (USA and Asian countries) it is
287 also administered with medicated foods. The widespread use of tetracycline has resulted in the
288 dissemination of resistance to many marine bacteria (Furushita et al., 2003) with the number of
289 resistant strains ranging from 7.7% (TE) to 11.5% (OT). These results are comparable with previous
290 studies where *Aeromonas* spp strains showed sensitivity to tetracycline and oxytetracycline (Awan
291 et al., 2009). *Aeromonas* spp strains showed high in vitro sensitivity against both oxytetracycline
292 (80.8%) and tetracycline (85.6%) in the tetracycline class with MIC₅₀ values below the reference
293 breakpoint of susceptibility ($\leq 1\mu\text{g/mL}$) and MIC₉₀ of 4 $\mu\text{g/mL}$ and 16 $\mu\text{g/mL}$ for tetracycline and
294 oxytetracycline, respectively. The MIC₅₀ and MIC₉₀ observed in our study were within the range
295 reported in previous investigations conducted on Aeromonads isolated from freshwater fish (Baron
296 et al., 2017; Čížek et al., 2010). The ECV of *Aeromonas* spp was 2 $\mu\text{g/mL}$ for both tetracycline and
297 oxytetracycline, values greater than those reported by Barone et al. (2017).

298 Among the Macrolides, erythromycin is the bacteriostatic drug of choice against Gram-positive
299 bacteria. Although it is not approved for aquaculture use in most European countries, the EU has
300 established maximum residue limits (MRLs) (Reg. EC 37/2011). In the present study, erythromycin
301 showed little effectiveness against *Aeromonas* spp. The MIC₉₀ of erythromycin was higher than the
302 reference breakpoints (8 $\mu\text{g/mL}$) with 84.6% of the tested *Aeromonas* strains showing resistance.
303 This high resistance rate and the MIC₅₀, MIC₉₀ and ECV for erythromycin are in agreement with
304 other studies (Mejdi et al., 2010; Baron et al., 2017).

305 Trimethoprim is mainly used in fish culture and often combined with sulfadiazine in commercial
306 preparations. Because of the potential carcinogenic effect of both antibacterial agents, the EU set
307 MRLs in fish muscle. The present study tested the two antimicrobials independently. Low efficacy
308 was obtained for sulfonamides with resistance rates of 92.3% and 69.2% of strains for sulfadiazine
309 and trimethoprim, respectively. For both sulfadiazine and trimethoprim the MIC₅₀ and MIC₉₀ were
310 above the reference breakpoints, so the ECVs could not be estimated. These values could not be

311 compared with other studies on *Aeromonas* spp as these antibacterials are generally used in
312 combination.

313 Aminoglycoside antibiotics showed intermediate MIC₉₀ values for gentamicin (8 µg/mL) and
314 kanamycin (32 µg/mL), whereas they were above the breakpoint for streptomycin (64 µg/mL) to
315 which 39.4% of strains were resistant. The ECVs for gentamicin and streptomycin were greater than
316 those observed by Baron et al. (2017), while the MIC₅₀ for streptomycin was comparable with data
317 obtained by Goñi-Urriza et al. (2000).

318 In the present study, chloramphenicol and florfenicol MIC₉₀ were lower than the reference
319 breakpoints. These results were expected for chloramphenicol as it has been banned from use in
320 animal food production since 1994 (EC 1430/94) due to its serious side effects on human health
321 (irreversible aplastic anaemia). While florfenicol is registered for use in aquaculture only in some
322 European countries, resistance to the fenicol category ranged between 2.9 % and 3.8% of the tested
323 strains. The MIC₅₀ and ECVs for these two antimicrobials were in agreement with values reported
324 by Baron et al. (2017).

325 Overall, *Aeromonas* spp showed elevated multiple resistance to the antibiotics tested. Most of the
326 strains (82.7%) showed a MAR index between 0.3 and 0.5 (corresponding to resistance to four to
327 eight different antibiotics) while six strains were resistant to nine to 11 antibiotics. One strain, *A.*
328 *punctata/hydrophila/enteropelogenes*, was resistant to 11 out of 15 antibiotics tested, indicating that
329 the isolates were exposed to high-risk sources of contamination with broad use of antibiotics, as in
330 intensive fish farming. This result is in agreement with previous studies indicating the high
331 antibiotic resistance of *Aeromonas* spp (Dumontet et al., 2000; Nguyen et al., 2014). The antibiotics
332 most frequently associated with multiple resistance were amoxicillin, ampicillin, cephalothin,
333 erythromycin, streptomycin, sulfadiazine and trimethoprim.

334

335 5. Conclusions

336 The present study confirms that selective pressure in the aquatic environment of intensive fish farms
337 leads to acquired antibiotic resistance by *Aeromonas* spp in gilthead sea bream reared in Italy.
338 Compared to clinical breakpoints, measuring epidemiological cut-off values allows a better
339 distinction between wild-type strains and strains which have acquired drug resistance due to
340 selective pressure. The multiple antibiotic resistance of almost all strains raises serious concerns due
341 to the possible transfer via food of antibiotic-resistant bacteria to humans or the acquisition of
342 antibiotic resistance by human pathogens. In the light of these findings, regular monitoring
343 programmes should be stepped up to check for antibiotic resistance in the aquaculture production of
344 fish for human consumption.

345 **Acknowledgements**

346 Anne Collins edited the English text.

347 **References**

- 348 1. Agersø, Y., Bruun, M. S., Dalsgaard, I., & Larsen, J. L. (2007). The tetracycline resistance
349 gene tet (E) is frequently occurring and present on large horizontally transferable plasmids
350 in *Aeromonas* spp. from fish farms. *Aquaculture*. 266(1), 47-52.
- 351 2. Awan, M. B., Maqbool, A., Bari, A., & Krovacek, K. (2009). Antibiotic susceptibility
352 profile of *Aeromonas* spp. isolates from food in Abu Dhabi, United Arab Emirates. *New*
353 *Microbiol.* 32(1), 17.
- 354 3. Baron, S., Granier, S. A., Larvor, E., Jouy, E., Cineux, M., Wilhelm, A., Gassilloud, B., Le
355 Bouquin, S., Kempf, I., & Chauvin, C. (2017). *Aeromonas* Diversity and Antimicrobial
356 Susceptibility in Freshwater-An Attempt to Set Generic Epidemiological Cut-Off Values.
357 *Front. Microbiol.* 8.
- 358 4. Bottero, M. T., Dalmaso, A., Soglia, D., Rosati, S., Decastelli, L., & Civera, T. (2004).
359 Development of a multiplex PCR assay for the identification of pathogenic genes of
360 *Escherichia coli* in milk and milk products. *Mol. Cell Probes*. 18(4), 283-288.

- 361 5. Cabello, F. C. (2006). Heavy use of prophylactic antibiotics in aquaculture: a growing
362 problem for human and animal health and for the environment. *Environ Microbiol.* 8(7),
363 1137-1144.
- 364 6. Cabello, F. C., Godfrey, H. P., Tomova, A., Ivanova, L., Dölz, H., Millanao, A., &
365 Buschmann, A. H. (2013). Antimicrobial use in aquaculture re-examined: its relevance to
366 antimicrobial resistance and to animal and human health. *Environ. Microbiol.* 15(7), 1917-
367 1942.
- 368 7. Callister, S. M., & Agger, W. A. (1987). Enumeration and characterization of *Aeromonas*
369 *hydrophila* and *Aeromonas caviae* isolated from grocery store produce. *Appl. Environ.*
370 *Microbiol.* 53(2), 249-253.
- 371 8. Cattoir, V., Poirel, L., Aubert, C., Soussy, C. J., & Nordmann, P. (2008). Unexpected
372 occurrence of plasmid-mediated quinolone resistance determinants in environmental
373 *Aeromonas* spp. *Emerg. Infect. Dis.* 14(2), 231.
- 374 9. Chang, Y. C., Shih, D. Y. C., Wang, J. Y., & Yang, S. S. (2007). Molecular characterization
375 of class 1 integrons and antimicrobial resistance in *Aeromonas* strains from foodborne
376 outbreak-suspect samples and environmental sources in Taiwan. *Diagn. Micr. Infec. Dis.*
377 59(2), 191-197.
- 378 10. Chopra, A. K., Houston, C. W., & Kurosky, A. (1991). Genetic variation in related cytolytic
379 toxins produced by different species of *Aeromonas*. *FEMS Microbiol. Lett.* 78(2-3), 231-
380 238.
- 381 11. Čížek, A., Dolejská, M., Sochorová, R., Strachotová, K., Piačková, V., & Veselý, T. (2010).
382 Antimicrobial resistance and its genetic determinants in *Aeromonads* isolated in ornamental
383 (koi) carp (*Cyprinus carpio koi*) and common carp (*Cyprinus carpio*). *Vet. Microbiol.*
384 142(3), 435-439.

- 385 12. CLSI (2005). Methods for Antimicrobial Dilution and Disk Susceptibility Testing of
386 Infrequently Isolated or Fastidious bacteria. 3rd ed. CLSI guideline M45. Clinical and
387 Laboratory Standards Institute, Wayne, PA.
- 388 13. CLSI (2007). Performance Standards of Antimicrobial Susceptibility Testing; Seventeenth
389 Informational Supplement. CLSI document M100-S17. Clinical and Laboratory Standards
390 Institute, Wayne, PA.
- 391 14. CLSI (2011). Performance Standards of Antimicrobial Susceptibility Testing of Bacteria
392 Isolated From Aquatic Animals: First Informational Supplement. CLSI document
393 M42/M49-S1. Clinical and Laboratory Standards Institute, Wayne, PA.
- 394 15. CLSI (2016). Performance Standards of Antimicrobial Susceptibility Testing; Seventeenth
395 Informational Supplement. CLSI document M100-S26. Clinical and Laboratory Standards
396 Institute, Wayne, PA.
- 397 16. Council Regulation (EEC) No. 2377/90 laying down a community procedure for the
398 establishment of maximum residue limits of veterinary medicinal products in foodstuffs of
399 animal origin, amending regulation no. 1430/94 of 22 June 1994, Off. J. Eur. Commun.,
400 L156, 1994, 23.
- 401 17. CVM. (2014). National Antimicrobial Resistance Monitoring System Integrated Report.
402 Center for Veterinary Medicine, Food and Drug Administration.
403 <http://www.fda.gov/AnimalVeterinary/SafetyHealth/AntimicrobialResistance/NationalAntimicrobialResistanceMonitoringSystem/ucm059103.htm>. Accessed 024 July 2017.
- 404
- 405 18. Defoirdt, T., Sorgeloos, P., & Bossier, P. (2011). Alternatives to antibiotics for the control
406 of bacterial disease in aquaculture. *Curr. Opin. Microbiol.*, 14(3), 251-258.
- 407 19. Diana, J. S., Egna, H. S., Chopin, T., Peterson, M. S., Cao, L., Pomeroy, R., Verdegem, M.,
408 Slack, W.T., Bondad-Reantaso, M.G., & Cabello, F. (2013). Responsible aquaculture in
409 2050: valuing local conditions and human innovations will be key to success. *BioScience*,
410 63(4), 255-262.

- 411 20. Díaz-Cruz M.S., Lopez De Alda M.J., Barcelò D. (2003). Environmental behaviour and
412 analysis of veterinary and human drugs in soils, sediments and sludge. *Trends Anal. Chem.*
413 22, 340-351.
- 414 21. Dumontet, S., Krovacek, K., Suenson, S.S., Pasquale, V., Baloda, S.B., Figliuolo, G. (2000).
415 Prevalence and diversity of *Aeromonas* and *Vibrio* spp. in coastal water of Southern Italy.
416 *Comp. Immunol. Microb.* 23, 53–72.
- 417 22. EC. (2017).
418 https://ec.europa.eu/fisheries/marine_species/farmed_fish_and_shellfish/seabream_en.
419 Accessed 04 July 2017.
- 420 23. EFSA (European Food Safety Authority), ECDC (European Centre for Disease Prevention
421 and Control) (2014) The European Union Summary Report on antimicrobial resistance in
422 zoonotic and indicator bacteria from humans, animals and food in 2012. *EFSA J*
423 12(3):3590–336. doi:10.2903/j.efsa.2014.3590.
- 424 24. FAO. (2016). *The State of World Fisheries and Aquaculture 2016. Contributing to food*
425 *security and nutrition for all.* Rome. 200 pp. Available at: [http://www.fao.org/3/a-](http://www.fao.org/3/a-i5555e.pdf)
426 [i5555e.pdf](http://www.fao.org/3/a-i5555e.pdf). Accessed 15 July 2017.
- 427 25. Figueira, V., Vaz-Moreira, I., Silva, M., & Manaia, C. M. (2011). Diversity and antibiotic
428 resistance of *Aeromonas* spp. in drinking and waste water treatment plants. *Water Res.*
429 45(17), 5599-5611.
- 430 26. Furushita, M., Shiba, T., Maeda, T., Yahata, M., Kaneoka, A., Takahashi, Y., Torii, K.,
431 Hasegawa, T., Ohta, M. (2003). Similarity of tetracycline resistance genes isolated from fish
432 farm bacteria to those from clinical isolates. *Appl. Environ. Microbiol.* 69, 5336–5342.
- 433 27. Galindo, C. L., Sha, J., Fadl, A. A., Pillai, L. L., & Chopra, A. K. (2006). Host immune
434 responses to *Aeromonas* virulence factors. *Current Immunology Reviews* 2(1), 13-26.

- 435 28. Giraud, E., Blanc, G., Bouju-Albert, A., Weill, F. X., & Donnay-Moreno, C. (2004).
436 Mechanisms of quinolone resistance and clonal relationship among *Aeromonas salmonicida*
437 strains isolated from reared fish with furunculosis. *J. Med. Microbiol.* 53(9), 895-901.
- 438 29. Gobat, P. F., & Jemmi, T. (1993). Distribution of mesophilic *Aeromonas* species in raw and
439 ready-to-eat fish and meat products in Switzerland. *Int. J. Food Microbiol.* 20(2), 117-120.
- 440 30. Goñi-Urriza, M., Pineau, L., Capdepuy, M., Roques, C., Caumette, P., & Quentin, C.
441 (2000). Antimicrobial resistance of mesophilic *Aeromonas* spp. isolated from two European
442 rivers. *J. Antimicrob. Chemoth.* 46(2), 297-301.
- 443 31. Hatha, M., Vivekanandhan, A. A., & Joice, G. J. (2005). Antibiotic resistance pattern of
444 motile *Aeromonads* from farm raised fresh water fish. *Int. J. Food Microbiol.* 98(2), 131-
445 134.
- 446 32. Inglis, V., Frerichs, G. N., Millar, S. D., & Richards, R. H. (1991). Antibiotic resistance of
447 *Aeromonas salmonicida* isolated from Atlantic salmon, *Salmo salar* L., in Scotland. *J. Fish*
448 *Dis.*, 14(3), 353-358.
- 449 33. Janda, J. M., & Abbott, S. L. (2010). The genus *Aeromonas*: taxonomy, pathogenicity, and
450 infection. *Clin. Microbiol. Rev.* 23(1), 35-73.
- 451 34. Kadlec, K., von Czapiewski, E., Kaspar, H., Wallmann, J., Michael, G. B., Steinacker, U., &
452 Schwarz, S. (2011). Molecular basis of sulfonamide and trimethoprim resistance in fish-
453 pathogenic *Aeromonas* isolates. *Appl. Environ. Microbiol.* 77(20), 7147-7150.
- 454 35. Kahlmeter, G., Brown, D. F., Goldstein, F. W., MacGowan, A. P., Mouton, J. W.,
455 Österlund, A., ... & Vatopoulos, A. (2003). European harmonization of MIC breakpoints for
456 antimicrobial susceptibility testing of bacteria. *J. Antimicrob. Chemoth.* 52(2), 145-148.
- 457 36. Khan, I. U. H., Loughborough, A., and Edge, T. A. (2009). DNA-based real-time detection
458 and quantification of *Aeromonads* from fresh water beaches on Lake Ontario. *J. Water*
459 *Health* 7, 312–323. doi: 10.2166/wh.2009.041

- 460 37. Krumperman, P. H. (1983). Multiple antibiotic resistance indexing of *Escherichia coli* to
461 identify high-risk sources of fecal contamination of foods. *Appl. Environ. Microbiol.* 46(1),
462 165-170.
- 463 38. Levy, S. B., & Marshall, B. (2004). Antibacterial resistance worldwide: causes, challenges
464 and responses. *Nat. Med.* 10, S122-S129.
- 465 39. Marshall, B. M., & Levy, S. B. (2011). Food animals and antimicrobials: impacts on human
466 health. *Clin. Microbiol. Rev.* 24(4), 718-733.
- 467 40. Mejdi, S., Emira, N., Ali, M., Hafedh, H., & Amina, B. (2010). Biochemical characteristics
468 and genetic diversity of *Vibrio* spp. and *Aeromonas hydrophila* strains isolated from the Lac
469 of Bizerte (Tunisia). *World J. Microbiol. Biotechnol.* 26(11), 2037-2046.
- 470 41. Nguyen, H. N. K., Van, T. T. H., Nguyen, H. T., Smooker, P. M., Shimeta, J., & Coloe, P. J.
471 (2014). Molecular characterization of antibiotic resistance in *Pseudomonas* and *Aeromonas*
472 isolates from catfish of the Mekong Delta, Vietnam. *Vet. Microbiol.* 171(3), 397-405.
- 473 42. Pruden, A., Larsson, D. J., Amézquita, A., Collignon, P., Brandt, K. K., Graham, Lazorchak,
474 J.M., Suzuki, S., Silley, P., Snape, J. R., Topp, E., Zhang, T., Zhu, Y.G. (2013).
475 Management options for reducing the release of antibiotics and antibiotic resistance genes to
476 the environment. *Environ. Health Persp.* 121(8), 878.
- 477 43. Radu, S., Ahmad, N., Ling, F. H., & Reezal, A. (2003). Prevalence and resistance to
478 antibiotics for *Aeromonas* species from retail fish in Malaysia. *Int. J. Food Microbiol.* 81(3),
479 261-266.
- 480 44. Real, F., Acosta, B., Deniz, S., Oros, J., Rodriguez, E. (1994). *Aeromonas salmonicida*
481 infection in *Sparus aurata* in the Canaries. *Bull. Eur. Assoc. Fish Pathol.* 14, 148-153.
- 482 45. Serrano, P.H. (2005). Responsible use of antibiotics in aquaculture, FAO Fisheries
483 Technical Paper no. 469. FAO, Rome.

- 484 46. Smith, P., & Kronvall, G. (2015). How many strains are required to set an epidemiological
485 cut-off value for MIC values determined for bacteria isolated from aquatic animals?.
486 *Aquacult. Int.* 23(2), 465-470.
- 487 47. Snoussi, M., Noumi, E., Lajnef, R., Belila, A., Yazidi, N., Bakhrouf, A. (2011). Phenotypic
488 characterization and enterobacterial repetitive intergenic consensus polymerase chain
489 reaction (eric-pcr) of *Aeromonas* spp. and *Vibrio* spp. strains isolated from *Sparus aurata*
490 fish farm (Khenis, Tunisia). *Afr. J. Microbiol. Res.* 5, 2920-2928.
- 491 48. Sorum, H. (2006). Antimicrobial drug resistance in fish pathogens. In *Antimicrobial*
492 *Resistance in Bacteria of Animal Origin*. Aarestrup, F.M., ed. Washington, DC: ASM Press.
493 pp. 213-238.
- 494 49. Teuber, M. (2001). Veterinary use and antibiotic resistance. *Curr. Opin. Microbiol.* 4, 493-
495 499.
- 496 50. Tsai, G. J., & Chen, T. H. (1996). Incidence and toxigenicity of *Aeromonas hydrophila* in
497 seafood. *Int. J. Food Microbiol.* 31(1-3), 121-131.
- 498 51. Turnidge J., Kahlmeter G., Kronvall G. (2006). Statistical characterisation of bacterial wild-
499 type MIC value distributions and the determination of epidemiological cut-off values. *Clin.*
500 *Microbiol Infect.* 12(5):418-425.
- 501 52. Walsh, T.R., Stunt, R.A., Nabi, J.A., MacGowan, A.P., Bennett, P.M. (1997). Distribution
502 and expression of β -lactamase genes among *Aeromonas* spp. *J. Antimicrob. Chemother.*, 40,
503 171-178.
- 504 53. Witte, W. (2000). Selective pressure by antibiotic use in livestock. *Int. J. Antimicrob.*
505 *Agents. Suppl* 1, S19-24.
- 506 54. Wu, C. J., Chuang, Y. C., Lee, M. F., Lee, C. C., Lee, H. C., Lee, N. Y. Chang, C.M., Chen,
507 P.L., Lin, Y.T., Yan, J.J., & Ko, W. C. (2011). Bacteremia due to extended-spectrum- β -
508 lactamase-producing *Aeromonas* spp. at a medical center in Southern Taiwan. *Antimicrob.*
509 *Agents Ch.* 55(12), 5813-5818.

1 Table 1 – Resistance of *Aeromonas* species and species-complex to 15 antimicrobial agents

Species	N (%)	OXA	AMX	AM	CF	CL	E	FM	FF	GM	K	OT	S	SZ	TE	TMP
<i>A. media</i>	15 (14.4%)	8	8	6	13	1	13	5	-	1	3	1	6	13	1	12
<i>A. enteropelogenes</i>	1 (0.9%)	-	1	1	1	-	1	-	-	-	-	-	-	1	-	1
<i>A. bivalvium</i>	10 (9.6%)	2	6	6	10	-	10	1	-	1	1	2	6	10	1	8
<i>A. media/veronii</i>	4 (3.8%)	2	4	4	4	-	4	2	1	-	-	-	-	4	-	4
<i>A. salmonicida/bestiarium/hydrophila/caviae</i>	12 (11.5%)	1	7	8	12	1	11	-	1	-	-	1	7	12	1	9
<i>A. punctata/hydrophila/enteropelogenes</i>	1 (0.9%)	1	1	1	1	1	1	1	-	-	-	1	1	1	-	1
<i>A. salmonicida/bestiarium</i>	8 (7.7%)	-	4	5	8	-	8	-	-	-	-	1	6	8	1	3
<i>A. popoffii</i>	3 (2.9%)	2	1	2	3	-	3	-	-	-	-	-	1	2	-	2
<i>A. molluscorum</i>	11 (10.6%)	7	1	1	4	1	3	6	1	1	2	5	4	11	3	6
<i>A. encheleia</i>	2 (1.9%)	1	1	1	2	-	2	-	-	-	-	-	2	2	-	2
<i>A. hydrophila/salmonicida/bestiarum</i>	4 (3.8%)	1	-	-	4	-	3	-	-	-	-	-	-	4	-	4
<i>A. punctata</i>	6 (5.8%)	1	3	1	6	-	6	1	-	-	-	-	1	4	-	6
<i>A. bivalvium/ popoffii</i>	3 (2.9%)	-	-	-	2	-	3	-	-	-	-	-	-	3	-	1
<i>A. salmonicida/bestiarum/popoffii</i>	2 (1.9%)	1	-	-	1	-	2	1	-	-	-	-	-	2	-	-
<i>A. media/hydro</i>	2 (1.9%)	1	2	2	2	-	2	1	-	-	-	-	-	2	-	2
<i>A. allosacrophila</i>	1(0.9%)	-	-	-	1	-	1	-	-	-	-	-	-	1	-	1

<i>A. tasmaniensis/ hydro/ punctata</i>	2(1.9%)	1	2	-	2	-	2	1	-	-	-	1	1	2	1	2
<i>A. media/ punctata</i>	5 (4.8%)	4	3	2	3	-	5	3	-	1	2	-	2	3	-	4
<i>A. encheleia/ molluscorum</i>	3 (2.9%)	-	2	2	3	-	3	-	-	-	-	-	1	2	-	1
<i>A. salmonicida/ sobria/popoffii</i>	4 (3.8%)	-	-	-	4	-	3	-	-	-	-	-	1	4	-	2
<i>A. salmonicida</i>	3 (2.9%)	-	-	-	3	-	1	-	-	-	-	-	-	3	-	1
<i>A. salmonicida/sobria</i>	1 (0.9%)	-	-	-	1	-	1	-	-	-	-	-	1	1	-	-
<i>A. molluscorum /eucrenophila</i>	1 (0.9%)	1	-	-	-	-	-	1	-	-	-	-	1	1	-	-
Total	104 (100%)	34	46	42	90	4	88	23	3	4	8	12	41	96	8	72

- 2 oxolinic acid (OXA), amoxicillin (AMX), ampicillin (AM), cephalothin (CF), cloramphenicol (CL), erythromycin (E), flumequine (FM), florfenicol
- 3 (FF), gentamicin (GM), kanamycin (K), oxytetracycline (OT), streptomycin (S), sulfadiazine (SZ), tetracycline (TE) and trimethoprim (TMP).

4 Table 2. Multiple antibiotic resistance (MAR) index of *Aeromonas* spp strains isolated from gilthead sea bream reared in 3 Italian regions.

MAR index	Region			Total
	Sardinia	Sicily	Tuscany	
0.1	-	<i>n</i> = 1	<i>n</i> = 2	<i>n</i> = 3
0.2	-	-	<i>n</i> = 13	<i>n</i> = 13
0.3	-	<i>n</i> = 14	<i>n</i> = 31	<i>n</i> = 45
0.4	<i>n</i> = 3	<i>n</i> = 7	<i>n</i> = 4	<i>n</i> = 14
0.5	<i>n</i> = 1	<i>n</i> = 14	<i>n</i> = 7	<i>n</i> = 22
0.6	-	<i>n</i> = 1	<i>n</i> = 1	<i>n</i> = 2
0.7	<i>n</i> = 1	<i>n</i> = 2	<i>n</i> = 1	<i>n</i> = 4
0.8	-	<i>n</i> = 1	-	<i>n</i> = 1
0.9	-	-	-	-

5

6

7

8

9

10 **Table 3.** MIC ($\mu\text{g/mL}$) and antimicrobial susceptibility of *Aeromonas* spp strains isolated from *Sparus aurata*. *=M45-P (CLSI, 2005); a=M100-
 11 S26 (CLSI, 2016); b= M42/49 (CLSI, 2011); c=M31A2 (NCCLS, 2002); d=M31A (NCCLS, 1998).


Antibiotic	Breakpoints	MIC₅₀	MIC₉₀	Moda	Range	S (%)	I (%)	R (%)	ECV	WT strains (%)
OXA	≤ 0.12 - $\geq 1^b$	0.06	16	0.06	0.06- ≥ 128	65 (62.5)	5 (4.8)	34 (32.7)	0.25	69 (66.3)
AMX	≤ 8 - $\geq 32^*$	16	≥ 128	8	0.06- ≥ 128	44 (42.3)	14 (13.5)	46 (44.2)	32	66 (63.5)
AM	≤ 8 - $\geq 32^*$	16	≥ 128	16- ≥ 128	0.06- ≥ 128	42 (40.4)	20 (19.2)	42 (40.4)	64	77 (74.0)
CF	≤ 8 - $\geq 32^*$	≥ 128	≥ 128	≥ 128	0.06- ≥ 128	14 (13.5)	-	90 (86.5)	ND	ND
CL	≤ 8 - $\geq 32^*$	0.5	4	0.5	0.25-64	98 (94.8)	2 (1.9)	4 (3.8)	2	91 (87.5)
E	≤ 0.5 - $\geq 8^c$	16	64	8	0.06- ≥ 128	9 (8.7)	7 (6.7)	88 (84.6)	32	93 (89.4)
FM	≤ 2 - $\geq 4^c$	0.12	16	0.06	0.06- ≥ 128	81 (77.9)	-	23 (22.1)	0.25	62 (59.6)
FF	≤ 4 - $\geq 8^b$	1	4	0.5	0.06-64	101 (97.1)	-	3 (2.9)	2	93 (89.4)
GM	≤ 4 - $\geq 16^*$	2	8	2	0.12-32	92 (88.5)	8 (7.7)	4 (3.8)	8	100 (96.2)
K	≤ 16 - $\geq 64^a$	8	32	16-32	0.5- ≥ 128	89 (85.6)	7 (6.7)	8 (7.7)	32	96 (92.3)
OT	≤ 1 - $\geq 8^b$	0.5	16	0.5	0.12- ≥ 128	84 (80.8)	8 (7.7)	12 (11.5)	2	84 (80.8)
S	≤ 6 - $\geq 25^d$	16	64	16	1- ≥ 128	9 (8.7)	54 (51.9)	41 (39.4)	64	90 (86.5)
SZ	≤ 38 - $\geq 76^a$	≥ 256	≥ 256	≥ 256	0.12- ≥ 256	8 (7.7)	-	96 (92.3)	ND	ND
TE	≤ 1 - $\geq 8^b$	0.5	4	0.5	0.12- ≥ 128	89 (85.6)	7 (6.7)	8 (7.7)	2	85 (81.8)
TMP	≤ 8 - $\geq 16^a$	32	64	64	0.12- ≥ 128	32 (30.8)	-	72 (69.2)	ND	ND

12 MIC (Minimum Inhibitory Concentrations), S (susceptible strains), I (intermediate strains), R (resistant strains), **ECV** (epidemiological cut-off
 13 value) ($\mu\text{g/mL}$), WT strains (wild-type strains), oxolinic acid (OXA), amoxicillin (AMX), ampicillin (AM), cephalothin (CF), cloramphenicol (CL),

- 14 erythromycin (E), flumequine (FM), florfenicol (FF), gentamicin (GM), kanamycin (K), oxytetracycline (OT), streptomycin (S), sulfadiazine (SZ),
15 tetracycline (TE) and trimethoprim (TMP). N.D.= not determined.

1 Figure 1. Percentages of *Aeromonas* species - species complex isolated from 6 mariculture farms in 3 Italian regions.

2



3

4 1=*A. media*; 2=*A. enteropelogenes*; 3=*A. bivalvium*; 4=*A. media/veronii*; 5=*A. salmonicida/bestiarium/hidrophila/caviae*; 6=*A. punctata/hydrophila/enteropelogenes*; 7=*A.*
 5 *salmonicida/bestiarium*; 8=*A. popoffii*; 9=*A. molluscorum*; 10=*A. encheleia*; 11=*A. hydrophila/salmonicida/bestiarum*; 12=*A. punctata*; 13=*A. bivalvium/popoffii*; 14=*A.*
 6 *salmonicida/bestiarium/popoffii*; 15=*A. media/hydro*; 16=*A. allosacarophila*; 17=*A. tasmaniensis/hydro/punctata*; 18=*A. media/punctata*; 19=*A. encheleia/molluscorum*; 20=*A.*
 7 *salmonicida/sobria/popoffii*; 21=*A. salmonicida*; 22=*A. salmonicida/sobria*; 23=*A. molluscorum /eucrenophila*.

8