

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Detection of rat hepatitis E virus in wild Norway rats (*Rattus norvegicus*) and Black rats (*Rattus rattus*) from 11 European countries

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1726177> since 2022-02-01T16:19:09Z

Published version:

DOI:10.1016/j.vetmic.2017.07.001

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

**Detection of rat hepatitis E virus in wild Norway rats (*Rattus norvegicus*) and
Black rats (*R. rattus*) from 11 European countries**

René Ryll^a, Samuel Bernstein^a, Elisa Heuser^a, Mathias Schlegel^{a,1}, Paul Dremsek^{a,2},
Maxi Zumpe^{a,3}, Sandro Wolf^{b,4}, Michel Pépin^c, Daniel Bajomi^d, Gabi Müller^e, Ann-
Charlotte Heiberg^f, Carina Spahr^g, Johannes Lang^h, Martin H. Groschup^a, Hermann
Ansorgeⁱ, Jona Freise^j, Sebastian Guenther^k, Kristof Baert^l, Francisco Ruiz-
Fonscisco^m, Jiri Pikula^{n,o}, Nataša Knap^p, Ioannis Tsakmakidis^q, Chrysostomos
Dovas^r, Stefania Zanet^s, Christian Imholt^t, Gerald Heckel^{u,v}, Reimar Johnen^g, Rainer
G. Ulrich^{a,w,*}

^a Friedrich-Loeffler-Institut, Institute of Novel and Emerging Infectious Diseases,
Südufer 10, 17493 Greifswald-Insel Riems, Germany

^b Institute for Microbiology, Technische Universität Dresden, Zellescher Weg 20b,
01217 Dresden, Germany

^c VetagroSup (Microbiologie/Immunologie/Pathologie infectieuse), Coordinateur de
l'Unité d'Enseignement de MEDECINE PREVENTIVE, USC INRA-VAS 1233 /
Equipe PERS ("Pathogènes Emergents et Rongeurs Sauvages"), 1 avenue
Bourgelat, 69280 Marcy l'étoile, France

^d Bábolna Bio Ltd., Szállás u. 6, H-1107 Budapest, Hungary

^e Stadt Zürich, Umwelt- und Gesundheitsschutz Zürich, Walchestrasse 31, 8021
Zürich, Switzerland

^f AC Heiberg Rådgivning, Åmosevej 69, Skellingsted, 4440 Mørkøv, Denmark

24 ^g German Federal Institute for Risk Assessment, Max-Dohrn-Str. 8-10, 10589 Berlin,
 25 Germany

26 ^h Working Group for Wildlife Biology, Justus Liebig University Giessen, Leihgesterner
 27 Weg 217, 35392 Giessen, Germany

28 ⁱ Senckenberg Museum of Natural History, Am Museum 1, 02826 Görlitz, Germany

29 ^j Niedersächsisches Landesamt für Verbraucherschutz und Lebensmittelsicherheit,
 30 Fachbereich Schädlingsbekämpfung, Task-Force Veterinärwesen, Röverskamp 5,
 31 26203 Wardenburg, Germany

32 ^k Institute of Microbiology and Epizootics, Veterinary Faculty, Freie Universität Berlin,
 33 Robert-von-Ostertag-Str. 7-13, 14163 Berlin, Germany

34 ^l Research Institute for Nature and Forest (INBO), Research Group Wildlife
 35 Management, Brussels, Belgium

36 ^m Spanish Wildlife Research Institute IREC (CSIC-UCLM-JCCCM), Ronda de Toledo
 37 12, 13071 Ciudad Real, Spain

38 ⁿ Department of Ecology and Diseases of Game, Fish and Bees, University of
 39 Veterinary and Pharmaceutical Sciences Brno, Brno, Czech Republic

40 ^o CEITEC - Central European Institute of Technology, University of Veterinary and
 41 Pharmaceutical Sciences Brno, Brno, Czech Republic

42 ^p Institute of Microbiology and Immunology, Faculty of Medicine, Zaloška 4, 1000
 43 Ljubljana, Slovenia

44 ^q Laboratory of Parasitology and Parasitic Diseases, School of Veterinary Medicine,
 45 Faculty of Health Sciences, Aristotle University of Thessaloniki, Greece

46 ^r Diagnostic Laboratory, School of Veterinary Medicine, Faculty of Health Sciences,
 47 Aristotle University of Thessaloniki, Greece

48 ^s Università degli Studi di Torino Dipartimento di Scienze Veterinarie Largo Paolo
49 Braccini, 2 (già Via L. DaVinci, 44)- 10095 Grugliasco (TO), Italy

50 ^t Julius Kühn-Institute, Federal Research Centre for Cultivated Plants, Institute for
51 Plant Protection in Horticulture and Forestry, Vertebrate Research, 48161 Münster,
52 Germany

53 ^u University of Bern, Institute of Ecology and Evolution, Baltzerstrasse 6, 3012 Bern,
54 Switzerland

55 ^v Swiss Institute of Bioinformatics, Quartier Sorge - Batiment Genopode, 1015
56 Lausanne, Switzerland

57 ^w German Center for Infection Research (DZIF), partner site Hamburg-Luebeck-
58 Borstel-Insel Riems, Germany

59

60 Current addresses:

¹ Seramun Diagnostica GmbH, 15754 Heidesee, Germany

² Labor Doz. DDr. Stefan Mustafa, 1030 Wien, Austria

³ University Medicine Greifswald, Pediatric Hematology and Oncology, Ferdinand-
Sauerbruch-Straße, 17475 Greifswald, Germany

⁴ Labor Ostsachsen, Bremer Str. 57, 01067 Dresden, Germany

61 *Corresponding author at: Institute of Novel and Emerging Infectious Diseases,
62 Friedrich-Loeffler-Institut, Federal Research Institute for Animal Health, Südufer 10,
63 D-17493 Greifswald-Insel Riems, Germany. Tel. +49 38351 7 1159; fax +49 38351 7
64 1192.

65 *E-mail address:* rainer.ulrich@fli.de (R.G. Ulrich)

66 **Keywords:** *Hepeviridae*, hepatitis E virus, Genotype 3, Norway rat, Black rat, Europe

67

ABSTRACT

Rat hepatitis E virus (ratHEV) is genetically only distantly related to hepeviruses found in other mammalian reservoirs and in humans. It was initially detected in Norway rats (*Rattus norvegicus*) from Germany, and subsequently in rats from Vietnam, the USA, Indonesia, China, Denmark and France.

Here, we report on a molecular survey of Norway and Black rats from 12 European countries for ratHEV and human pathogenic hepeviruses. RatHEV-specific real-time and conventional RT-PCR investigations revealed the presence of ratHEV in 63 of 508 (12.4%) rats at the majority of sites in 11 of 12 countries. In contrast, a real-time RT-PCR specific for human pathogenic HEV genotypes 1-4 and a nested broad-spectrum (NBS) RT-PCR with subsequent sequence determination did not detect any infections with these genotypes. Only in a single Norway rat from Belgium a rabbit HEV-like genotype 3 sequence was detected. Phylogenetic analysis indicated a clustering of all other novel Norway and Black rat-derived sequences with ratHEV sequences from Europe, the USA and a Black rat-derived sequence from Indonesia within the proposed ratHEV genotype 1. No difference in infection status was detected related to age, sex, rat species or density of human settlements and zoological gardens.

In conclusion, our investigation shows a broad geographical distribution of ratHEV in Norway and Black rats from Europe and its presence in all settlement types investigated.

1. Introduction

The family *Hepeviridae* comprises an increasing number of viruses in mammals, birds and fish (Johne et al., 2014, Pérez-Gracia et al., 2015). Initially, hepatitis E virus (HEV) was the only member of this virus family, which was divided into four genotypes. The genotypes 1 and 2 are supposed to exclusively infect humans, whereas genotypes 3 and 4 are zoonotic with wild boar, domestic pig and deer representing animal reservoirs (Meng et al., 2013). In chicken, additional divergent genotypes were discovered and designated as Avian HEV, which can be associated with the diseases Big Liver and Spleen Disease and Hepatitis-Splenomegaly Syndrome (Handler and Williams 1988, Ritchie and Riddell 1991; Gerber et al., 2015). The International Committee on Taxonomy of Viruses (ICTV) currently classifies the human pathogenic HEV genotypes 1-4 into species *Orthohepevirus A*, avian HEV into *Orthohepevirus B*, batHEV into *Orthohepevirus D* and the carnivore and ratHEV into *Orthohepevirus C* (<http://ictvonline.org/virusTaxonomy.asp>, accessed 07.04. 2017).

The hepevirus genome is a positive stranded RNA of approximately 6.7 to 7.3 kilobases (Meng et al., 2012). The genome contains the typical sequence elements of an eukaryotic mRNA with a cap structure at its 5'-end and a polyadenylation at its 3'-end (Tam et al., 1991). For all hepeviruses, three major open reading frames (ORF) were identified with almost the same organization, but differences in the junction or overlapping region of ORF1 and ORF2/ORF3 (Johne et al., 2014). The ORF1 of 4.6 to 5.2 kb is located at the 5'-end of the genome and codes for a polyprotein comprising several nonstructural proteins including regions with similarity to methyltransferases, papain-like proteases, helicases and RNA-dependent RNA

115 polymerases (Koonin et al., 1992). The capsid protein of 600-675 amino acid
116 residues is encoded by ORF2 and contains three domains with the carboxyterminal
117 domain being exposed on the surface of the virion (Yamashita et al., 2009). The
118 overlapping ORF3 codes for a small phosphoprotein of strongly varying length in
119 avian, mammalian and fish hepeviruses (Zafrullah et al., 1997; Holla et al., 2013;
120 Johne et al., 2014). This protein is essential for virus egress and found to be
121 associated with lipid membranes (Okamoto, 2013). Interestingly, ratHEV as well as
122 ferretHEV contains an additional putative open reading frame (ORF4), overlapping
123 ORF1 at its 5'-end, of still unknown function (Johne et al., 2010a, Raj et al., 2012).

124 Using a broad-spectrum RT-PCR assay, a novel, only distantly-related hepevirus was
125 identified in 2010 in Norway rats (*Rattus norvegicus*) from Hamburg, Germany
126 (Johne et al., 2010a; Johne et al., 2010b). This initial finding was confirmed by
127 detection of closely related sequences in Norway rats from other cities in Germany
128 (Johne et al., 2012). Detection of related sequences in rats from the USA, Vietnam,
129 Denmark, France, China and Indonesia suggests a host specificity of ratHEV for rats
130 of the genus *Rattus* and indicated its broad geographical distribution (Li et al., 2013b;
131 Li et al., 2013d; Mulyanto et al., 2013; Mulyanto et al., 2014; Purcell et al., 2011;
132 Widen et al., 2014; Wolf et al., 2013). The host specificity of this virus was also
133 demonstrated by infection experiments using laboratory rats and other mammals
134 (Cossaboom et al., 2012; Li et al., 2013c). However, recent studies in China
135 suggested a broader host range of the virus or frequent spillover infections of
136 bandicoot rats and even shrews (Guan et al., 2013; Li et al., 2013d). The genotypes
137 G1, G2 and G3 of ratHEV were previously defined on the basis of a complete
138 genome sequence comparison; a further comparison of 31 ORF 2-derived

sequences of 281-bp length revealed two additional sequences of a non-designated clade (ND), which clustered with G1 (Mulyanto et al., 2014). All G1 ratHEV sequences in previous studies originated from *R. norvegicus* or *R. rattus*, whereas ratHEV sequences of G3 originated exclusively from *R. rattus*. In contrast, genotype G2 was detected in *R. rattus*, *R. tanezumi*, *R. rattoides losea* and the shrew *Suncus murinus* (Li et al., 2013b; Li et al., 2013d; Mulyanto et al., 2013).

The zoonotic potential of ratHEV is currently controversially discussed. Serological studies in forestry workers showed a few seropositive individuals (Dremsek et al., 2012). In addition, febrile patients from China showed a stronger reactivity with ratHEV antigen than with genotype 1 and 3 antigens (Shimizu et al., 2016). Furthermore, ratHEV was shown to replicate in a human-derived cell line (Jirintai et al., 2014; Li et al., 2015). In contrast, experimental infection of monkeys and domestic pigs with ratHEV failed (Cossaboom et al., 2012; Purcell et al., 2011). Reproducible experimental infections of nude rats and Wistar rats with ratHEV (Li et al., 2013c; Purcell et al., 2011) and the availability of a recently developed reverse genetics system for ratHEV (Li et al., 2015) led to the suggestion to use ratHEV-infected laboratory rats as an infection model for hepeviruses. On the other hand, Norway rats were found to be infected with human pathogenic genotype 3 associated strains, suggesting a potential role for zoonotic transmission (Lack et al., 2012; Kanai et al., 2012).

Here, we describe a molecular survey of Norway and Black rats from 12 European countries for ratHEV and human pathogenic HEV genotypes, and evaluated influences of sex, age, rat species and human settlement type on ratHEV prevalence.

2. Material and methods

2.1. Rat collection, dissection and sample collection

The collection of Norway rats in Copenhagen and Berlin has been already described previously (Sachsenroder et al., 2014; Wolf et al., 2013). Additional Norway rats were collected in Germany, Denmark, Austria, Switzerland, Czech Republic, Belgium, France, Slovenia and Greece; Black rats (*R. rattus*) were collected in Italy, Slovenia, Greece and Spain (Fig. 1).

The dissection and collection of tissue and chest cavity fluid samples followed standard protocols. For the evaluation of the influence of sex, age, reservoir species and human settlement type on ratHEV prevalence, previously published results for rats from Hamburg, Berlin, Stuttgart, Esslingen and Copenhagen (Johne et al., 2012; Johne et al., 2010a; Johne et al., 2010b; Wolf et al., 2013) were also included.

2.2. RNA isolation, real-time and conventional RT-PCR and sequencing

After homogenizing rat liver tissue using a TissueLyser (Qiagen, Hilden, Germany), RNA was extracted with the RNeasy Mini Kit (Qiagen). A ratHEV-specific real-time RT-PCR (Johne et al., 2012, RTD, see Fig. 2) and a real-time RT-PCR specific for HEV genotypes 1-4 (Jothikumar et al., 2006) were performed as previously published. The QuantiTect Probe RT-PCR Kit (Qiagen) was used in a 7500 Real Time PCR System (Applied Biosystems Life Technologies, Darmstadt, Germany) and

the data were evaluated using 7500 Software v2.0.1 (Applied Biosystems Life Technologies, Darmstadt, Germany).

A one-step RT-PCR (designated SW-RT-PCR; see Fig. 2) was then performed using a SuperScriptIII One-Step RT-PCR with PlatinumTaq Kit (Invitrogen Life Technologies, Carlsbad, CA, USA) in a C1000 Thermal Cycler (Bio-Rad Laboratories, Munich, Germany). Reverse transcription was conducted at 42 °C for 50 min, followed by a denaturation step at 94 °C for 2 min. A total of 45 PCR cycles each consisting of 30 s at 94 °C, 30 s at the primer-specific annealing temperature (Table 1), 1 min at 68 °C and a final incubation at 68 °C for 10 min were performed.

Additionally, a slightly modified nested broad-spectrum (NBS) RT-PCR was performed to test the samples for all possible HEV strains, including ratHEV and human pathogenic genotypes as described (Johne et al., 2010b; see Fig. 2). A first RT-PCR was performed using a One-Step RT-PCR kit (Qiagen) with primers HEV-cs and HEV-cas in a 2720 thermal cycler (Applied Biosystems). The thermal profile comprised 42 °C for 60 min and 95 °C for 15 min, followed by 40 cycles of 94 °C for 30 s, 50 °C for 30 s and 74 °C for 45 s, with a final incubation at 74 °C for 5 min. An aliquot of the RT-PCR product (5 µl) was used in a nested PCR with a GoTaq kit (Promega) and the primers HEV-csn and HEV-casn. The thermal profile consisted of 95 °C for 5 min and 35 cycles of 94 °C for 30 s, 50 °C for 30 s and 72 °C for 45 s, with a final incubation at 72 °C for 5 min.

To generate a longer sequence stretch, overlapping the SW-/NBS-RT-PCR products and including the 3'-end of ORF 1, the 5'-region of ORF 2 and a partial or complete ORF 3, selected samples were analyzed by a primer walking-based attempt using

two different primer pairs and following the protocols of the SW-RT-PCR (see Fig. 2 and Table 1; Primer-walking RT-PCR-I/II).

RT-PCR products were purified using a MiniElute PCR Purification Kit (Qiagen) or a NucleoSpin Gel and PCR Clean-up Kit (Machery-Nagel, Düren, Germany), separated by agarose gel electrophoresis and visualized by ethidium bromide staining.

For sequencing, the purified RT-PCR product was amplified by PCR using the same primers and the following temperature profile: 96 °C for 1 min, followed by 30 cycles of 96 °C for 15 s, 50 °C for 15 s and 60 °C for 90 s. Amplicons were purified using a Sigma Spin Post-Reaction Clean-up Column Kit (Sigma-Aldrich, Hamburg, Germany) and sequenced on an ABI 3100 Avant DNA-Sequencer (Applied-Biosystems, Darmstadt, Germany). Sequences were assembled and aligned using BioEdit 7.2.0 (Hall, 1999) and MEGA 7 (Kumar et al., 2016), respectively. The novel HEV sequences were deposited at GenBank (for accession numbers see Fig. 3A).

2.3. Phylogenetic analysis

The General Time Reversible + discrete Gamma distribution (GTR+G) model was the best suited substitution model determined by MEGA 7 for both regions spanning nucleotides (nt) 4,105-4,387 (numbering based on strain R63, acc. no. GU345042) and nucleotides 4,105-5,226. The phylogenetic analyses were performed by Bayesian algorithms via MrBayes v.3.2.2 and CIPRES online portal (Ronquist et al., 2012) and by Maximum likelihood algorithm performed via MEGA7 (Kumar et al., 2016).

2.4. Evaluation of demographic, rat species and human settlement type influence

The statistical evaluation of demographic, rat species and human settlement type influences on individual ratHEV infection status was performed similarly to the previously described methodology for other infectious agents on a sub-sample (Heuser et al., 2016). Briefly, generalized linear modelling (GLM) with a binomial error distribution was applied using individual infection status as the response variable, with sex and age classes (≤ 200 g (juvenile) and > 200 g (adult) (Webster et al., 1995)) as demographic predictors as well as the association of ratHEV with a particular *Rattus* species (*R. norvegicus* vs. *R. rattus*) and human settlement type, based on human population density (urban ($> 1,500$ inhabitants/km²), small town (300-1,500 inhabitants/km²), rural (< 300 inhabitants/km²)) (database: Geostat, 2012). Rats collected in zoological gardens were put in a separate category. Model selection was performed using the *drop1* function. Goodness of fit of all performed regression models was assessed using the Le Cessie-van Houwelingen test statistic implemented in the *rms*-package. All analyses were performed in R (R Core Team, 2015).

3. Results

3.1. Collection of rats and initial real-time RT-PCR screening of rats

From 2005 to 2016 a total of 508 rats were collected in 12 European countries (Fig. 1). This sample contained 420 Norway rats from trapping sites in Germany (23 sites,

156 rats), Denmark (1 site, 11 rats), Austria (1 site, 43 rats), Switzerland (3 sites, 29 rats), Czech Republic (3 sites, 58 rats), Belgium (2 sites, 60 rats), France (1 site, 28 rats), Slovenia (1 site, 1 animal) and Greece (3 sites, 16 rats) and 88 Black rats from trapping sites in Italy (1 site, 17 rats), Slovenia (1 site, 17 rats), Greece (2 sites, 4 rats) and Spain (1 site, 50 rats). Initially, liver-derived RNA preparations of a Norway rat sample subset were tested in parallel by real-time RT-PCR assays either targeting ratHEV or HEV genotypes 1 to 4. The ratHEV-specific real-time RT-PCR (RTD) resulted in the detection of 5 out of 145 (3.4%) samples from Germany (Table 2). Norway rat samples from Hungary, Denmark, Switzerland and France were also positive for ratHEV-RNA by ratHEV-specific real-time RT-PCR with a detection range of 5.5% (1/18) to 18.1% (2/11; see Table 2). The Ct values of positive samples ranged between 20 and 34. In the real-time RT-PCR targeting the human pathogenic genotypes 1-4 none of the Norway rat samples showed a Ct value <35, used as cut-off (Table 2).

3.2. Conventional SW-RT-PCR and NBS-RT-PCR analysis

A conventional RT-PCR approach using ORF1-specific SW-RT-PCR (nt positions 4,105-4,387, prototype strain R63, accession number GU345042, see Fig. 2) and NBS RT-PCR (nt positions 4,000-4,423, see Fig. 2) resulted in the detection of HEV-specific RNA in 17 of 156 (10.8%) samples from Germany (Table 2). The prevalence for samples from the sites in the other ten countries reached from 4% (2/50) to 27.2% (3/11; Table 2). The prevalences in Norway rats and Black rats were 10%-27.2% (2/20 and 3/11) and 4%-5.8% (2/50 and 1/17), respectively. None of the single Norway rat and 17 Black rats from Slovenia was HEV-RNA positive (Table 2).

Using a primer-walking based approach for thirteen samples from nine sites in Germany, France, Spain, Belgium, Austria and Denmark, a 1,122/1,125-base pair (bp) long region including parts of ORF1, ORF2 and partial or entire ORF3 (see Fig. 2) was RT-PCR amplified and sequenced (see Table 4). The different lengths of the sequences B1 and B4 from France were caused by a triplet indel, i.e., insertion/deletion of three nucleotides (data not shown).

3.3. Sequence comparison and phylogenetic analysis

Phylogenetic analysis of the 280 bp fusion-product of the SW-/NBS-RT-PCR assays showed that almost all novel sequences, independently whether from Norway or Black rats, clustered together with ratHEV sequences, species *Orthohepevirus C1*, well separated from sequences of species *Orthohepevirus C2* (Figs. 3A and 3B). In one Norway rat sample from Belgium (KS/16/825) a sequence with 88.8% sequence similarity to genotype 3 HEV sequences was found (see below); in no other sample human pathogenic genotype-related sequences were found. This HEV genotype 3 - like sequence from the single Norway rat sample from Belgium clustered in the phylogenetic tree with three rabbit HEV strains from China and a human rabbit HEV sequence from France within species *Orthohepevirus A* (Fig. 3C); attempts to generate a longer sequence failed. The phylogenetic analysis of the concatenated 1,122/1,125 bp product of the coding sequences revealed clustering of all novel sequences within the ratHEV genotype G1 defined by Mulyanto et al. (2014), in sister clade relationship with ratHEV genotypes G2 and G3 (Fig. 3D). Genotype G1 contains the prototype sequence R63 from a Norway rat from Hamburg, Norway rat-derived sequences from different European countries and the USA, Black rat-derived

301 sequences from Spain and Italy and one sequence originating from a Black rat
 302 collected in Solo, Indonesia (Figs. 3B and D).

303 A novel sequence from rats in Berlin, detected in five animals, clustered with a
 304 previously determined sequence from Berlin and two novel sequences from rats in
 305 Esslingen, with one found in four animals, clustered with a sequence detected
 306 previously in Stuttgart, a site close to Esslingen (Johne et al., 2012; see Fig. 3B, and
 307 legend to Fig. 3). Similarly, two sequences from Warburg formed a well-separated
 308 subclade and all sequences from Czech Republic were highly related (Fig. 3B). Most
 309 novel ratHEV sequences from Vienna formed a well-supported cluster but one
 310 sequence (KS12-1338) was highly divergent. Both sequences from Spain are closely
 311 related, independently if the 280 bp or 1,222 bp products were analyzed (Figs. 3B
 312 and D). Interestingly, ratHEV sequences from three trapping sites close to Lyon (B
 313 and E/A) formed two well-separated subclades and sequences from Zurich belonged
 314 also to two subclades (Fig. 3B). Sequences from Norway rats from Belgium were
 315 found at highly divergent positions within the tree (Fig. 3B).

316 Comparison of ORF1-derived sequences from the fusion product of SW-/NBS-RT-
 317 PCR from the same site resulted in an intra-cluster sequence similarity of 79.6% to
 318 100% for the nucleotide and 86.8% to 100% for the corresponding amino acid
 319 sequences (Table 3). When analyzing the nucleotide sequence similarity within
 320 partial ORF1 or the overlapping ORF1/ORF2/ORF3 regions between different sites,
 321 the values reached similar levels of 81.0% to 96.1% and 87.2% to 91.5%,
 322 respectively (Supplementary Table and Table 4). The corresponding aa sequence
 323 similarities of ORF1-encoded protein and concatenated ORF1- and ORF2-encoded

proteins ranged between 93.4% and 100% and 95.9% and 98.6%, respectively
(Supplementary Table and Table 4).

3.4. Association of ratHEV infections with age, sex, rat species and human settlement density

For a total of 668 rats, including those of this study (n=508) and those investigated previously (n= 160; Johne et al., 2012; Johne et al., 2010a; Wolf et al., 2013), no association with age, sex or the *Rattus* species and the individual ratHEV infection status could be detected (Table 5). In addition, ratHEV was detected in Norway rats from all four settlement types investigated. Human population density did not seem to have an effect on ratHEV occurrence, as prevalences in small towns and rural sites did not differ significantly from high density urban areas. The prevalence in zoological gardens was lower compared to urban areas, though not formally significant (Table 5). For all models goodness of fit analysis did not provide any evidence of a lack of fit.

4. Discussion

In this study, we investigated Norway and Black rat samples from 12 European countries for the presence of ratHEV and other hepeviruses using ratHEV-specific real-time RT-PCR (Johne et al., 2010a) and human HEV genotype 1-4-specific real-time RT-PCR (Jothikumar et al., 2006) as well as conventional RT-PCR assays (SW-RT-PCR (Wolf et al., 2013) and NBS RT-PCR (Johne et al., 2010b)). Using these four methods, almost exclusively ratHEV was detected in Norway and Black rats from

11 of 12 countries. This finding is in line with the previously demonstrated inability in experimentally infecting Norway rats with human pathogenic genotypes ((Li et al., 2013a; Li et al., 2013c; Purcell et al., 2011) and results from earlier field studies in Norway rats (Johne et al., 2012; Johnne et al., 2010a). Similar to previous studies reporting the human pathogenic HEV genotype 3 in Norway rats (Lack et al., 2012; Kanai et al., 2012), in one Norway rat from Belgium a short rabbit HEV-like genotype 3 sequence was detected. This might be explained by a spillover infection of this strain from a rabbit reservoir. [Rabbits and rats may share their habitats in this region of Belgium, either in wildlife habitats or when wild \(pest\) rats search for food close to private rabbit husbandry.](#)

This study demonstrates the occurrence of ratHEV not only in Norway rats, as previously reported for Germany, France and Denmark, but for the first time in Europe also in Black rats, namely from Italy and Spain. This finding is in line with studies in Asia, where ratHEV has been demonstrated in different *Rattus* species and in Bandicoot rats (Guan et al., 2013, Li et al., 2013d).

In addition, in our study ratHEV was not only detected in rats from urban areas, but also in rats from small towns and rural areas. The detection of ratHEV in rural areas complements our previous finding of a local absence of ratHEV in a rural area close to Ahlen (Johne et al., 2012), which may suggest site-specific differences and a heterogeneous distribution of ratHEV not primarily driven by human settlement. In addition, ratHEV was identified in pest rats from zoological gardens raising questions on the potential transmission of this virus to zoo animals. In fact, serological investigations have detected HEV-specific antibodies in captive macaques and HEV-RNA in different mammalian and avian species in a wildlife rescue center in China

(Korzaia et al., 2007; Zhang et al., 2008). The recently developed in-house ELISA technology based on ratHEV- and HEV genotype 3-derived recombinant capsid protein derivatives (Dremsek et al., 2012; Johne et al., 2012) may be used in the future for differentiation of antibodies raised against these viruses in zoo animals.

The phylogenetic analysis of the novel ratHEV sequences showed for almost all a high similarity to ratHEV genotype 1 defined recently (Mulyanto et al., 2014), independently whether the sequences originated from Norway or Black rats. In line with a previous investigation (Purdy and Sue, 2017), the resolution of the phylogenetic analysis using the short-sized ORF1 region was lower than the resolution for the larger segment of ORF1/ORF2/ORF3. The observed phylogenetic clustering of many sequences from the same or neighbouring sites may indicate the persistence of ratHEV strains within the local populations. The separate clustering of sequences from the same geographical origin might be caused by an incursion (and perhaps establishment) of additional, highly divergent ratHEV strains by invading rats. In line with this assumption, sequences from the USA (strain LA-8350) and Indonesia (strain SOLO-006SF) cluster also within genotype 1 of ratHEV (Figs. 3B and D).

The previous finding of the majority of rats being only HEV RNA or anti-ratHEV antibody positive suggested non-persistent infections in individual rats (Johne et al., 2012). In line with this assumption, we did not find here a significantly higher RNA prevalence in adult rats compared to juvenile animals. These findings of non-persistent infections of rats are also in line with results of experimental infection studies in Norway rats (Purcell et al., 2011). At this time we cannot exclude age-dependent differences in susceptibility and mortality of rats for ratHEV infection,

possibly associated with co-infections with other pathogens or genetic or environmental factors.

5. Conclusion

The detection of ratHEV in Norway and Black rats from 11 European countries indicates a broad geographical distribution of ratHEV suggesting an (almost) continent-wide occurrence and no specific association with human population density. Phylogenetic investigations indicated clustering of all European ratHEV sequences within ratHEV genotype 1. Well-separated subclades of sequences from the same or neighbouring sites might indicate the incursion of novel ratHEV strains into local Norway rat populations with a parallel persistence of a local ratHEV strain. This necessitates future studies on the population structure and potential invasion of individuals into existing rat populations and their association with ratHEV incursion. In addition, the finding of ratHEV infections in zoological gardens may allow future studies on the zoonotic potential of ratHEV based on the investigation of putative natural ratHEV transmission to non-human primates. Finally, the finding of a rabbit HEV-like sequence in a single Norway rat necessitates further studies, especially in habitats with sympatric occurrence of rabbits or pigs and rats, to evaluate potential spillover infections of human pathogenic genotype(s) and their potential public health impact.

Conflict of Interest

The authors declare that they have no competing interests.

418

419 **Authors' contributions**

420 Designed the study: RGU, RJ, GH. Performed the experiments: SB, RR, EH, MS,
421 PD, MZ, SW. Analyzed the data: SB, RR, CI, RJ, GH, RGU. Contributed materials:
422 MP, DB, GM, ACH, JL, HA, JF, SG, KB, FRF, JP, NK, JT, CD, SZ. Wrote the
423 manuscript: RR, SB, CI, GH, RJ, RGU. All authors read and approved the
424 manuscript.

425

426 **Acknowledgments**

427 The authors are very grateful to Dörte Kaufmann and Günter Strebelow for excellent
428 technical assistance, Patrick Wysocki for generation of Figure 1 and Hanan Sheikh
429 Ali, Kathrin Baumann, Angele Breithaupt, Thomas Büchner, Stephan Drewes, Ulrike
430 Duve, Julie Elkins, Bärbel Hammerschmidt, Dörte Kaufmann, Robert Klopffleisch,
431 Nastasja Kratzmann, Christian Kretzschmar, Denny Maaz, Grit Möwert, Josephin
432 Schröder, Ulrike M. Rosenfeld, Josephine Schlosser, Sabrina Schmidt, Ramona
433 Spließ, Annemarie Steiner, André Schütte, Kerstin Tauscher, Franziska Thomas,
434 Konrad Wanka and Ute Wessels for support of the dissections. The authors kindly
435 acknowledge the collection of rats by Marianne Jäger (Vienna) and support of the rat
436 trapping by Zoltán Kiss (Báblona Bio Ltd. Budapest, Hungary), Ezio Ferroglio
437 (Università degli Studi di Torino Dipartimento di Scienze Veterinarie Largo Paolo
438 Braccini, Grugliasco, Italy) and Diego Bünter (Anticimex, Zurich, Switzerland).
439 Additional rats were kindly provided by Anne Balkema-Buschmann, Dan Balkema,
440 Lothar Stitz, Lieselotte Minke (Friedrich-Loeffler-Institut, Greifswald-Insel Riems and

441 Wusterhausen, Germany), Marion Saathoff (LAVES, Wardenburg, Germany),
442 Richard Kruczewski (Dortmund, Germany), Tomi Trilar (Slovenian Museum of
443 Natural History, Slovenia), Diana Žele (Veterinary Faculty, University of Ljubljana,
444 Slovenia) and the Flanders Environment Agency (VMM, Belgium). The investigations
445 in the laboratory of Rainer G. Ulrich were supported by the German Center for
446 Infection Research (DZIF).

447

Figure legends

Fig. 1. Geographical map representing the rat collection sites in Denmark (1, Copenhagen), Germany (2, Hamburg; 3, Elmenhorst; 4, Stahlbrode; 5, Osnabrück; 6, Wolbrechtshausen; 7, Magdeburg; 8, Kampehl; 9, Berlin; 10, Neschwitz; 11, Königshain; 12, Görlitz; 13, Niederoderwitz; 14, Zittau; 15, Aachen; 16, Köln; 17, Oer-Erkenschwick; 18, Münster; 19, Ahlen; 20, Warburg; 21, Heidelberg; 22, Stuttgart; 23, Esslingen; 24, Möggingen), Switzerland (25, Gränichen, 26, Dübendorf, 27, Zurich), Austria (28, Vienna), Hungary (29, Budapest), France (30, five sites close to Lyon), Belgium (31 Dender, 32 Ijzer), Italy (33 Pianosa Island), Slovenia (34 close to Ljubljana), Spain (35 Cadiz), Czech Republic (36 Prague, 37 Brno, 38 Northern Moravia), Greece (39 Thessaloniki, 40 Kilkis, 41 Chalkidiki). All or some of the rats from sites 1, 2, 9, 19, 22 and 23 were investigated for ratHEV previously (Johne et al., 2010, 2012; Wolf et al., 2013; indicated by empty or half-filled circles, respectively) and were included here for analysis of demographic, reservoir and human settlement type association of ratHEV infections (see Heuser et al., 2016).

Fig. 2. Genome organization of rat HEV, prototype strain R63 (accession number GU345042), and location of primer binding sites for real-time (RTD) and conventional screening SW-/NBS-RT-PCR and primer-walking RT-PCRs as well as the corresponding amplification products.

Fig. 3. Consensus phylogenetic trees based on Bayesian and Maximum-Likelihood analyses of a part of ORF 1 with all species within genus *Orthohepevirus* (A) and a

471 zoom-in for species *Orthohepevirus C1* and *Orthohepevirus C2* (B) and
472 *Orthohepevirus A* (C), and the concatenated region of ORF1 and partial ORF2/ORF3
473 overlapping region (D).

474 Consensus phylogenetic trees based on Bayesian analyses were done with
475 8,000,000 or 6,000,000 generations and a burn-in of 25%, and Maximum-Likelihood
476 analysis with 1,000 bootstraps and 50% cut-off, of a part of ORF 1 (nt positions
477 4,105-4,387, counting according prototype strain R63, accession number GU345042)
478 (A-C) and the concatenated region of ORF1 (4,105-4,921) and partial ORF2/ORF3
479 overlapping region (nt positions 4,949-5,226) (D) of ratHEV. Posterior
480 probability/bootstrap values of >50 are given at the supported nodes.

481 The ratHEV genotypes G1, G2 and G3 were defined previously (Mulyanto et al.,
482 2014); the two sequences of clade ND (not designated) were found in this previous
483 study, based on a partial region of ORF2, to be clustering with G1.

484 Novel sequences are given in bold and labeled by a star. Identical sequences were
485 omitted from the analysis and only different sequence types are presented (Berlin
486 KS11/573 = KS11/576, /578, /580, /587, Esslingen Mu10/1564= Mu10/1567, /1568,
487 /1571, Warburg Mu10/697= Mu10/698, Zurich KS12/1361=KS12/1363, Czech
488 Republic KS14/73 = KS14/75, /76, /99, and KS14/70 = KS14/80, /98).

489

References

- Batts, W., Yun, S., Hedrick, R., Winton, J., 2011. A novel member of the family *Hepeviridae* from cutthroat trout (*Oncorhynchus clarkii*). *Virus Res* 158, 116-123.
- Bodewes, R., van der Giessen, J., Haagmans, B.L., Osterhaus, A.D., Smits, S.L., 2013. Identification of multiple novel viruses, including a parvovirus and a hepevirus, in feces of red foxes. *J Virol* 87, 7758-7764.
- Cossaboom, C.M., Cordoba, L., Sanford, B.J., Pineyro, P., Kenney, S.P., Dryman, B.A., Wang, Y., Meng, X.J., 2012. Cross-species infection of pigs with a novel rabbit, but not rat, strain of hepatitis E virus isolated in the United States. *J Gen Virol* 93, 1687-1695.
- Dremsek, P., Wenzel, J.J., Johne, R., Ziller, M., Hofmann, J., Groschup, M.H., Werdermann, S., Mohn, U., Dorn, S., Motz, M., Mertens, M., Jilg, W., Ulrich, R.G., 2012. Seroprevalence study in forestry workers from eastern Germany using novel genotype 3- and rat hepatitis E virus-specific immunoglobulin G ELISAs. *Med Microbiol Immunol* 201, 189-200.
- Drexler, J.F., Seelen, A., Corman, V.M., Fumie Tateno, A., Cottontail, V., Melim Zerbinati, R., Gloza-Rausch, F., Klose, S.M., Adu-Sarkodie, Y., Oppong, S.K., Kalko, E.K., Osterman, A., Rasche, A., Adam, A., Muller, M.A., Ulrich, R.G., Leroy, E.M., Lukashev, A.N., Drosten, C., 2012. Bats worldwide carry hepatitis E virus-related viruses that form a putative novel genus within the family *Hepeviridae*. *J Virol* 86, 9134-9147.

512 Gerber, P. F., Trampel, D. W., Willingham, E. M., Billam, P., Meng, X. J., &
 513 Opriessnig, T. (2015). Subclinical avian hepatitis E virus infection in layer
 514 flocks in the United States. *Vet. J.*, 206(3), 304-311.

515 Geostat initiative, Eurostat. [http://ec.europa.eu/eurostat/web/gisco/gisco-](http://ec.europa.eu/eurostat/web/gisco/gisco-activities/integrating-statistics-geospatial-information/geostat-initiative)
 516 [activities/integrating-statistics-geospatial-information/geostat-initiative](http://ec.europa.eu/eurostat/web/gisco/gisco-activities/integrating-statistics-geospatial-information/geostat-initiative) 2012).

517 Guan, D., Li, W., Su, J., Fang, L., Takeda, N., Wakita, T., Li, T.C., Ke, C., 2013.
 518 Asian musk shrew as a reservoir of rat hepatitis E virus, China. *Emerg Infect*
 519 *Dis* 19, 1341-1343.

520 Hall, T.A., 1999. BioEdit: a user-friendly biological sequence alignment editor and
 521 analysis program for Windows 95/98/NT. *Nucleic acids symposium series* Vol.
 522 41.

523 Handler, J. H., & Williams, W. (1988). An egg drop associated with splenomegaly
 524 in broiler breeders. *Avian Dis*, 32(4): 773-778.

525 Heuser, E., Fischer, S., Ryll, R., Mayer-Scholl, A., Hoffmann, D., Spahr, C., Imholt,
 526 C., Alfa, D.M., Frohlich, A., Luschow, D., Johne, R., Ehlers, B., Essbauer, S.,
 527 Nockler, K., Ulrich, R.G., 2016. Survey for zoonotic pathogens in Norway rat
 528 populations from Europe. *Pest Manag Sci*, 73(2), 341-348.

529 Holla RP, Ahmad I, Ahmad Z, Jameel S. Molecular virology of hepatitis E virus.
 530 *Semin Liver Dis*. 2013 Feb;33(1):3-14.

531 Jirintai, S., Tanggis, Mulyanto, Suparyatmo, J.B., Takahashi, M., Kobayashi, T.,
 532 Nagashima, S., Nishizawa, T., Okamoto, H., 2014. Rat hepatitis E virus
 533 derived from wild rats (*Rattus rattus*) propagates efficiently in human
 534 hepatoma cell lines. *Virus Res* 185, 92-102.

535 Johne, R., Dremsek, P., Kindler, E., Schielke, A., Plenge-Bonig, A., Gregersen, H.,
 536 Wessels, U., Schmidt, K., Rietschel, W., Groschup, M.H., Guenther, S.,

537 Heckel, G., Ulrich, R.G., 2012. Rat hepatitis E virus: geographical clustering
 538 within Germany and serological detection in wild Norway rats (*Rattus*
 539 *norvegicus*). Infect Genet Evol 12, 947-956.

540 Johne, R., Dremsek, P., Reetz, J., Heckel, G., Hess, M., Ulrich, R.G., 2014.
 541 *Hepeviridae*: an expanding family of vertebrate viruses. Infect Genet Evol 27,
 542 212-229.

543 Johne, R., Heckel, G., Plenge-Bonig, A., Kindler, E., Maresch, C., Reetz, J., Schielke,
 544 A., Ulrich, R.G., 2010a. Novel hepatitis E virus genotype in Norway rats,
 545 Germany. Emerg Infect Dis 16, 1452-1455.

546 Johne, R., Plenge-Bonig, A., Hess, M., Ulrich, R.G., Reetz, J., Schielke, A., 2010b.
 547 Detection of a novel hepatitis E-like virus in faeces of wild rats using a nested
 548 broad-spectrum RT-PCR. J Gen Virol 91, 750-758.

549 Jothikumar, N., Cromeans, T.L., Robertson, B.H., Meng, X.J., Hill, V.R., 2006. A
 550 broadly reactive one-step real-time RT-PCR assay for rapid and sensitive
 551 detection of hepatitis E virus. J Virol Methods 131, 65-71.

552 Kanai, Y., Miyasaka, S., Uyama, S., Kawami, S., Kato-Mori, Y., Tsujikawa, M.,
 553 Yunoki, M., Nishiyama, S., Ikuta, K., Hagiwara, K. 2012. Hepatitis E virus in
 554 Norway rats (*Rattus norvegicus*) captured around a pig farm. BMC research
 555 notes, 5(1), 4.

556 Koonin EV, Gorbalenya AE, Purdy MA, Rozanov MN, Reyes GR, Bradley DW.
 557 Computer-assisted assignment of functional domains in the nonstructural
 558 polyprotein of hepatitis E virus: delineation of an additional group of positive-
 559 strand RNA plant and animal viruses. Proc Natl Acad Sci U S A. 1992 Sep
 560 1;89(17):8259-63.

561 Korzaia, L.I., Lapin, B.A., Keburia, V.V., Lazareva, I., 2007. [Hepatitis E virus
 562 antibodies in the macaques and in the personnel serving the macaques of the
 563 Adler apery]. *Vopr Virusol* 52, 36-40.

564 Krog, J.S., Breum, S.O., Jensen, T.H., Larsen, L.E., 2013. Hepatitis E virus variant in
 565 farmed mink, Denmark. *Emerg Infect Dis* 19, 2028-2030.

566 Kumar, S., Stecher, G., Tamura, K., 2016. MEGA7: Molecular Evolutionary Genetics
 567 Analysis Version 7.0 for Bigger Datasets. *Mol Biol Evol* 33, 1870-1874.

568 Lack, J.B., Volk, K., Van Den Bussche, R.A., 2012. Hepatitis E virus genotype 3 in
 569 wild rats, United States. *Emerg Infect Dis* 18, 1268-1273.

570 Li, T.C., Ami, Y., Suzaki, Y., Takeda, N., Takaji, W., 2013a. No evidence for hepatitis
 571 E virus genotype 3 susceptibility in rats. *Emerg Infect Dis* 19, 1343-1345.

572 Li, T.C., Ami, Y., Suzaki, Y., Yasuda, S.P., Yoshimatsu, K., Arikawa, J., Takeda, N.,
 573 Takaji, W., 2013b. Characterization of full genome of rat hepatitis E virus strain
 574 from Vietnam. *Emerg Infect Dis* 19, 115-118.

575 Li, T.C., Yang, T., Yoshizaki, S., Ami, Y., Suzaki, Y., Ishii, K., Haga, K., Nakamura,
 576 T., Ochiai, S., Takaji, W., Johne, R., 2015. Construction and characterization
 577 of an infectious cDNA clone of rat hepatitis E virus. *J Gen Virol* 96, 1320-1327.

578 Li, T.C., Yoshizaki, S., Ami, Y., Suzaki, Y., Yasuda, S.P., Yoshimatsu, K., Arikawa, J.,
 579 Takeda, N., Wakita, T., 2013c. Susceptibility of laboratory rats against
 580 genotypes 1, 3, 4, and rat hepatitis E viruses. *Vet Microbiol* 163, 54-61.

581 Li, W., Guan, D., Su, J., Takeda, N., Wakita, T., Li, T.C., Ke, C.W., 2013d. High
 582 prevalence of rat hepatitis E virus in wild rats in China. *Vet Microbiol* 165, 275-
 583 280.

584 Lin, J., Norder, H., Uhlhorn, H., Belak, S., Widen, F., 2014. Novel hepatitis E like
 585 virus found in Swedish moose. *J Gen Virol* 95, 557-570.

586 Meng, X. J., Anderson, D. A., Arankalle, V. A., Emerson, S. U., Harrison, T. J.,
 587 Jameel, S., Okamoto, H. (2012). Hepeviridae. Virus Taxonomy 9th Report of
 588 the ICTV, Elsevier Academic Press, London (2012), pp. 1021–1028.
 589 Meng, X. J. (2013). Zoonotic and foodborne transmission of hepatitis E virus. In
 590 Seminars in liver disease (Vol. 33, No. 01, pp. 041-049). Thieme Medical
 591 Publishers.
 592 Mulyanto, Depamede, S.N., Sriasih, M., Takahashi, M., Nagashima, S., Jirintai, S.,
 593 Nishizawa, T., Okamoto, H., 2013. Frequent detection and characterization of
 594 hepatitis E virus variants in wild rats (*Rattus rattus*) in Indonesia. Arch Virol
 595 158, 87-96.
 596 Mulyanto, Suparyatmo, J.B., Andayani, I.G., Khalid, Takahashi, M., Ohnishi, H.,
 597 Jirintai, S., Nagashima, S., Nishizawa, T., Okamoto, H., 2014. Marked
 598 genomic heterogeneity of rat hepatitis E virus strains in Indonesia
 599 demonstrated on a full-length genome analysis. Virus Res 179, 102-112.
 600 Okamoto, H., 2013. Culture systems for hepatitis E virus. J Gastroenterol 48, 147-
 601 158.
 602 Purcell, R.H., Engle, R.E., Rood, M.P., Kabrane-Lazizi, Y., Nguyen, H.T.,
 603 Govindarajan, S., St Claire, M., Emerson, S.U., 2011. Hepatitis E virus in rats,
 604 Los Angeles, California, USA. Emerg Infect Dis 17, 2216-2222.
 605 Pérez-Gracia, M. T., García, M., Suay, B., & Mateos-Lindemann, M. L. (2015).
 606 Current knowledge on hepatitis E. J Clin Transl Hepatol 3(2), 117.
 607 R Foundation for Statistical Computing, R: A language and environment for statistical
 608 computing. <https://www.R-project.org/> (2015).

609 Raj, V.S., Smits, S.L., Pas, S.D., Provacia, L.B., Moorman-Roest, H., Osterhaus,
 610 A.D., Haagmans, B.L., 2012. Novel hepatitis E virus in ferrets, the
 611 Netherlands. *Emerg Infect Dis* 18, 1369-1370.

612 Ritchie, S. J., & Riddell, C. (1991). British Columbia. "Hepatitis-splenomegaly"
 613 syndrome in commercial egg laying hens. *Can. Vet. J.*, 32(8), 500.

614 Ronquist, F., Teslenko, M., van der Mark, P., Ayres, D.L., Darling, A., Hohna, S.,
 615 Larget, B., Liu, L., Suchard, M.A., Huelsenbeck, J.P., 2012. MrBayes 3.2:
 616 efficient Bayesian phylogenetic inference and model choice across a large
 617 model space. *Syst Biol* 61, 539-542.

618 Sachsenroder, J., Braun, A., Machnowska, P., Ng, T.F., Deng, X., Guenther, S.,
 619 Bernstein, S., Ulrich, R.G., Delwart, E., Johne, R., 2014. Metagenomic
 620 identification of novel enteric viruses in urban wild rats and genome
 621 characterization of a group A rotavirus. *J Gen Virol* 95, 2734-2747.

622 Shimizu, K., Hamaguchi, S., Ngo, C. C., Tian-Cheng, L. I., Yoshimatsu, K., Yasuda,
 623 S. P., Fujita, H., Pham, T.T., Le, M.Q., Dang, A.D., Nguyen, T.Q., Yoshida,
 624 L.M., Ariyoshi, K., Arikawa, J.. (2016). Serological evidence of infection with
 625 rodent-borne hepatitis E virus HEV-C1 or antigenically related virus in
 626 humans. *J. Vet. Med. Sci.*, 78(11), 1677-1681.

627 Smith, D.B., Simmonds, P., International Committee on Taxonomy of Viruses
 628 *Hepeviridae* Study, G., Jameel, S., Emerson, S.U., Harrison, T.J., Meng, X.J.,
 629 Okamoto, H., Van der Poel, W.H., Purdy, M.A., 2014. Consensus proposals
 630 for classification of the family *Hepeviridae*. *J Gen Virol* 95, 2223-2232.

631 Tam, A. W., Smith, M. M., Guerra, M. E., Huang, C. C., Bradley, D. W., Fry, K. E., &
 632 Reyes, G. R. (1991). Hepatitis E virus (HEV): molecular cloning and
 633 sequencing of the full-length viral genome. *Virology*, 185(1), 120-131.

634 Webster, J.P., Ellis, W.A., Macdonald, D.W., 1995. Prevalence of *Leptospira* spp. in
 635 wild brown rats (*Rattus norvegicus*) on UK farms. Epidemiol Infect 114, 195-
 636 201.

637 Widen, F., Ayrat, F., Artois, M., Olofson, A.S., Lin, J., 2014. PCR detection and
 638 analysis of potentially zoonotic Hepatitis E virus in French rats. Virol J 11, 90.

639 Wolf, S., Reetz, J., Johne, R., Heiberg, A.C., Petri, S., Kanig, H., Ulrich, R.G., 2013.
 640 The simultaneous occurrence of human norovirus and hepatitis E virus in a
 641 Norway rat (*Rattus norvegicus*). Arch Virol 158, 1575-1578.

642 Yamashita, T., Mori, Y., Miyazaki, N., Cheng, R.H., Yoshimura, M., Unno, H., Shima,
 643 R., Moriishi, K., Tsukihara, T., Li, T.C., Takeda, N., Miyamura, T., Matsuura,
 644 Y., 2009. Biological and immunological characteristics of hepatitis E virus-like
 645 particles based on the crystal structure. Proc Natl Acad Sci U S A 106, 12986-
 646 12991.

647 Zafrullah M, Ozdener MH, Panda SK, Jameel S. The ORF3 protein of hepatitis E
 648 virus is a phosphoprotein that associates with the cytoskeleton. J Virol. 1997
 649 Dec;71(12):9045-53.

650 Zhang, W., Shen, Q., Mou, J., Yang, Z.B., Yuan, C.L., Cui, L., Zhu, J.G., Hua, X.G.,
 651 Xu, C.M., Hu, J., 2008. Cross-species infection of hepatitis E virus in a zoo-like
 652 location, including birds. Epidemiol Infect 136, 1020-1026.