

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Effects of dietary *Tenebrio molitor* meal inclusion in free-range chickens

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1560985> since 2022-04-01T11:21:33Z

Published version:

DOI:10.1111/jpn.12487

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 **Effects of dietary *Tenebrio molitor* meal inclusion in free-range chickens**

2

3 I. Biasato¹, M. De Marco¹, L. Rotolo², M. Renna², C. Lussiana², S. Dabbou², M.T.
4 Capucchio¹, E. Biasibetti¹, P. Costa¹, F. Gai³, L. Pozzo³, D. Dezzutto⁴, S. Bergagna⁴, S.
5 Martínez⁵, M. Tarantola^{1,6}, L. Gasco^{2,3}, and A. Schiavone^{1,6}

6

7 ¹Dipartimento di Scienze Veterinarie, Università degli Studi di Torino, Italy

8 ²Dipartimento di Scienze Agrarie, Forestali e Alimentari, Università degli Studi di Torino,
9 Italy

10 ³Istituto di Scienze delle Produzioni Alimentari (ISPA), CNR, Italy

11 ⁴Istituto Zooprofilattico Sperimentale del Piemonte, Liguria e Valle d'Aosta, Torino, Italy.

12 ⁵Departamento de Producción Animal, Universidad de Murcia, Spain

13 ⁶Istituto di Ricerche Interdisciplinari sulla Sostenibilità, Università degli Studi di Torino, Italy

14

15 Correspondence: Prof. Achille Schiavone, Dipartimento di Scienze Veterinarie, Università
16 degli Studi di Torino, largo Paolo Braccini 2, 10095 Grugliasco (TO), Italy. Tel: +39-011-
17 6709208; Fax: +39-011-6709240; E-mail: achille.schiavone@unito.it

18

19

20

21 **Running head:** Dietary *Tenebrio molitor* meal in free-range chicken diets

22

23

24

25 **Summary**

26 Insects are currently being considered as a novel protein source for animal feeds, because they
27 contain a large amount of protein. The larvae of *Tenebrio molitor* (TM) have been shown to
28 be an acceptable protein source for broiler chickens in terms of growth performance, but till
29 now no data on histological or **intestinal** morphometric features have been reported. The
30 present study has had the aim of evaluating the effects of dietary TM inclusion on the
31 performance, welfare, intestinal morphology and histological features of free-range chickens.
32 A total of 140 medium-growing hybrid female chickens were free-range reared and randomly
33 allotted to two dietary treatments: i) a control group and ii) a TM group, in which TM meal
34 was included at 75 g/kg. Each group consisted of five pens as replicates, with 14 chicks per
35 pen. Growth performance, hematological and serum parameters and welfare indicators were
36 evaluated and the animals were slaughtered at the age of 97 days. Two birds per pen (10
37 birds/treatment) were submitted to histological (liver, spleen, thymus, bursa of Fabricius,
38 kidney, heart, glandular stomach and gut) and morphometric (duodenum, jejunum and ileum)
39 investigations. The inclusion of TM did not affect the growth performance, hematological or
40 serum parameters. The morphometric and histological features were not significantly affected
41 either, thus suggesting no influence on nutrient metabolization, performance or animal health.
42 Glandular stomach alterations (chronic flogosis with epithelial squamous metaplasia) were
43 considered paraphysiological in relation to free-range farming. The observed chronic
44 intestinal flogosis, with concomitant activation of the lymphoid tissue, was probably due to
45 previous parasitic infections, which are very frequently detected in free-range chickens. In
46 conclusion, the findings of this study show that yellow mealworm inclusion does not affect
47 the welfare, productive performances or morphological features of free-range chickens, thus
48 confirming that TM can be used safely in poultry diets.

49 **Keywords**

50 Poultry, insect meal, yellow mealworm, histology, morphometry.

51 **Introduction**

52 Insects are currently considered a novel protein source for animal feeds (Sánchez-Muros et
53 al., 2014). Insect meals exhibit a great potential for becoming a standard ingredient in animal
54 feeding, because of the high quality and quantity of protein (Ramos-Elorduy, 1997).
55 **Furthermore**, the low competitiveness with human food (Ballitoc and Sun, 2013) and the
56 reduction in environmental impact in terms of energy cost, land area utilization and footprints
57 (Oonincx and de Boer, 2012; Makkar et al., 2014; Sánchez-Muros et al., 2014), **make insects**
58 **promising in an ecological perspective**. Invertebrates are included in the European Union
59 Feed Material Register as a feed material, even though they are currently only authorized for
60 pets. However, insect-derived feeds could also represent a possible ingredient for livestock
61 animals, such as poultry, pigs and fish (Veldkamp et al., 2012; Van Huis, 2013; Makkar et al.,
62 2014; Henry et al., 2015). In particular, the most promising insect species for industrial
63 production are *Hermetia illucens* (black soldier fly), *Musca domestica* (common house fly),
64 *Tenebrio molitor* (yellow mealworm), *Bombyx mori* (silkworm) and several grasshoppers
65 (Van Huis, 2013).

66 Considering that insects are consumed naturally by wild birds and free-range poultry (Zuidhof
67 et al., 2003), some studies have evaluated the feasibility of using insects as an alternative feed
68 source for poultry (Khatun et al., 2003; Wang et al., 2005; Oyegoke et al., 2006; Adeniji,
69 2007; Hwangbo et al., 2009; Ijaiya and Eko, 2009; Ballitoc and Sun, 2013). Some authors
70 have observed no differences in growth performance (in terms of feed intake, body weight
71 gain and feed conversion efficiency) in broilers fed a control diet and an insect-based diet
72 (Wang et al., 2005; Oyegoke et al., 2006; Adeniji, 2007; Ijaiya and Eko, 2009). Other studies
73 have reported that insect meal inclusion in chicken diets improved animal growth indexes
74 (Khatun et al., 2003; Hwangbo et al., 2009; Ballitoc and Sun, 2013). The same studies also
75 observed an improvement in carcass yield characteristics, such as dressing percentage, breast

76 muscle, thigh muscle, slaughter, dressed carcass and eviscerated weights (Khatun et al., 2003;
77 Hwangbo et al., 2009; Ballitoc and Sun, 2103).

78 The larvae of *Tenebrio molitor* (TM) are easily bred, because of their efficient growth on
79 dried and cooked waste materials from fruit, vegetables and cereals in various combinations.
80 For this reason, they are already industrially produced as feeds for pets and zoo animals,
81 including birds, reptiles, small mammals, amphibians and fish (Makkar et al., 2014). On a dry
82 matter basis, the meal derived from TM larvae contains a large amount of crude protein (440
83 to 690 g/kg) and fat (230 to 470 g/kg) (Veldkamp et al., 2012). In livestock, TM has been
84 shown to be an acceptable protein source for broiler chickens (Ramos-Elorduy et al., 2006;
85 Ballitoc and Sun, 2013) and fish (Roncarati et al., 2015; Belforti et al., in press).

86 Intestinal morphology is the main indicator of gut health and functioning (Kristy et al., 2005).
87 Dietary protein level and digestibility have been reported to significantly affect the intestinal
88 development and the mucosal architecture of the gastrointestinal tract of broilers (Laudadio et
89 al., 2012; Qaisrani et al., 2014). Intestinal development can be assessed through morphometric
90 measurements of the villus height (to determine the area available for digestion and
91 absorption) and crypt depth (the region in which new intestinal cells are formed) (Franco et
92 al., 2006). The villus height/crypt depth ratio can also be evaluated, because it generally gives
93 an indication of the likely maturity and functional capacity of the enterocytes (Hampson,
94 1986).

95 The nutrient profile and structure of protein sources may also vary significantly, and thus have
96 different effects on nutrient utilization and metabolism, and consequently on some blood
97 constituents, such as serum cholesterol, triglycerides and uric acid (Wang et al., 2015). Some
98 blood parameters have been used as physiological indicators of the stress response of
99 chickens. The heterophils-to-lymphocytes (H/L) ratio is affected by stress factors and could
100 be used as an indicator of stress in poultry (De Marco et al., 2013; Salamano et al., 2010). The

101 H/L ratio is correlated to a bird's health status and responds to stimuli associated with diet,
102 chronic bacterial infections, stress, light and trauma, and it varies according to the change in
103 the percentage of heterophils and lymphocytes in the blood (Gross and Siegel, 1983;
104 Maxwell, 1993; Maxwell and Robertson, 1998). Finally, diet composition can also affect the
105 litter quality (Lynn and Elson, 1990; Jones et al., 2005), which is directly related to the
106 development of footpad dermatitis (FPD) (Bessei, 2006). FPD is a condition that involves
107 inflammation and necrotic lesions on the plantar surface of the bird's feet and is considered to
108 be an important welfare indicator in broilers (Ekstrand et al., 1997; Meluzzi et al., 2008;
109 Welfare Quality[®], 2009).

110 Although insect meals are considered a suitable ingredient for poultry feeding (Veldkamp et
111 al., 2012; Van Huis, 2013; Makkar et al., 2014), there is currently a lack of data on their
112 utilization. Apart from the evaluation of growth performance and carcass yield characteristics,
113 no anatomopathological or morphometric investigations, blood traits analysis or welfare
114 assessments have been carried out on chickens fed diets with insect meal inclusion. Therefore,
115 the aim of the present study was to evaluate the effects of TM dietary inclusion on the
116 productive performance, intestinal morphology, histological features, hematochemical
117 parameters and welfare of free-range chickens.

118

119

120 **Materials and Methods**

121 The study was performed by the Department of Veterinary Sciences and the Department of
122 Agricultural, Forest and Food Sciences of the University of Torino (Italy) in collaboration
123 with an external farm called "Fattoria La Fornace", located in Montechiaro d'Asti (At - Italy).
124 The experimental protocol was designed according to the guidelines of the current European
125 and Italian laws on the care and use of experimental animals (Directive 2010/63/EU, put into

126 force in Italy with D.L. 2014/26). The experiment was carried out between November-
127 December 2014, when the photoperiod was 9-10L:14-15D.

128

129 *Birds and diets*

130 In this experiment, a total of 140 female Label Hubbard hybrid chickens (female: JA 57 ×
131 male: S77CN), a medium-growing genotype, were used. All the birds were free-range reared,
132 in identical environmental conditions, throughout the experimental trial. At the age of 43 days
133 (average weight 715 g), the birds were randomly allotted to two dietary treatments, each
134 consisting of five pens as replicates, with 14 chicks per pen. Each pen had an indoor area
135 (2.5×3.5 m) and an outdoor paddock of the same dimension (2.5×3.5 m). The floor was
136 covered, to a height of 10 cm, with wood shaving litter.. The birds were only exposed to
137 natural light. Two diets were formulated: a control diet (C), normally used by the breeder, and
138 an experimental diet (TM), in which TM meal (Gaobeidian Shannong Biology Co. Ltd.,
139 Gaobeidian, Hebei province - China) was included at 75 g/kg in substitution of corn gluten
140 meal (Table 1). The diets were designed to meet or exceed the current poultry requirements
141 (NRC, 1994). Both diets were isonitrogenous and isoenergetic and were formulated using the
142 apparent metabolizable energy (AMEn) values for TM calculated for broiler chickens (De
143 Marco et al., 2015). Feed and water were provided *ad libitum*.

144 The experiment lasted 54 days. The average chicken weight and feed intake were recorded at
145 the beginning and at the end of the experiment on a pen basis. The final body weight was
146 recorded on day 97, and the feed conversion ratio was calculated for the 43-97 day period.

147 The chicks were vaccinated at hatching against coccidiosis, Newcastle disease and infectious
148 bronchitis. All the birds were individually identified with a shank ring.

149

150 *Chemical analyses of the diets*

151 The diets were subsequently ground to pass through a 0.5-mm sieve and stored in airtight
152 plastic containers for DM, ash, CP, crude fibre (AOAC, 2005) and EE (Folch et al., 1957)
153 determination. The fatty acid composition of the control and TM diets was assessed using the
154 method described by Alves *et al.* (2008). Fatty acid methyl esters were separated, identified
155 and quantified on the basis of the chromatographic conditions reported by Renna *et al.* (2014).
156 Heptadecanoic acid (C17:0) was used as the internal standard. The results were expressed in
157 absolute values as g/kg DM.

158 In order to perform the AA determination, samples of the diets were prepared using a 22 h
159 hydrolysis step in 6 HCl at 112°C under a nitrogen atmosphere. Performic acid oxidation
160 occurred prior to acid hydrolysis for methionine and cystine. The AA in the hydrolysate was
161 determined by means of HPLC after postcolumn derivatization, according to the procedure
162 described by Madrid et al. (2013). Tryptophan was not determined.

163 All the analyses were performed in duplicate.

164

165 *Hematological and serum parameters*

166 At the end of the experiment (day 97), blood samples were collected at slaughtering from 4
167 birds per pen: 2.5 mL was placed in an EDTA tube and 2.5 mL in a serum-separating tube. A
168 blood smear was prepared, using one glass slide for each bird, from a drop of blood without
169 anticoagulant. The smears were stained using May-Grünwald and Giemsa stains (Campbell,
170 1995). The total red and white blood cell counts were determined in an improved Neubauer
171 haemocytometer on blood samples previously treated with a 1:200 Natt-Herrick solution. One
172 hundred leukocytes, including granular (heterophils, eosinophils and basophils) and non-
173 granular (lymphocytes and monocytes) leukocytes, were counted on the slide and the H/L
174 ratio was calculated. The tubes without anticoagulant were left to clot in a standing position at
175 room temperature for approximately two hours to obtain serum. The serum was separated by

176 means of centrifugation at $700 \times g$ for 15 minutes and frozen at -80°C until analysis. The total
177 proteins were quantified by means of the “biuret method” (Bio Group Medical System kit;
178 Bio Group Medical System, Talamello, Italy); the electrophoretic pattern of the serum was
179 obtained using a semi-automated agarose gel electrophoresis system (Sebia Hydrasys®,
180 Norcross, GA, USA). The alanino-aminotransferase (ALT), aspartate-aminotransferase
181 (AST), triglycerides, cholesterol, glucose, phosphorus, magnesium, iron, uric acid and
182 creatinine serum concentrations were measured by means of enzymatic methods in a clinical
183 chemistry analyzer (Screen Master Touch, Hospitex diagnostics Srl., Florence, Italy).

184

185 *Slaughtering procedures, footpad dermatitis lesion assessment, histological investigations*
186 *and morphometric analysis*

187 On day 97, all the chickens were individually marked, weighed and slaughtered in a
188 commercial abattoir. The plucked and eviscerated carcasses were obtained and the head, neck,
189 feet and abdominal fat were removed to obtain carcass-for-grilling. The weight of the breasts,
190 thighs, deboned thighs, spleen, bursa of Fabricius, liver, gizzard and abdominal fat were
191 recorded.

192 In order to evaluate the FPD lesions, the feet collected at the slaughterhouse were assessed
193 macroscopically using the so-called Swedish footpad scoring system (Ekstrand et al., 1997).
194 According to this system, 0 = no lesion, slight discoloration of the skin or healed lesion; 1 =
195 mild lesion, superficial discoloration of the skin and hyperkeratosis; 2 = severe lesion,
196 affected epidermis, blood scabs, hemorrhage and severe swelling of the skin.

197 After slaughtering, 2 birds per pen (10 birds/treatment) were sampled for
198 anatomopathological investigations. Intestinal segment samples (approximately 5 cm in
199 length) of the duodenum, jejunum, ileum and caecum were excised and flushed with 0.9%
200 saline to remove all the contents. The collected intestine segments were the loop of the

201 duodenum, the tract before Meckel's diverticulum (jejunum), the tract before the ileocolic
202 junction (ileum) and the apex of the caeca (caecum). Samples of the liver, spleen, thymus,
203 bursa of Fabricius, kidneys, heart and glandular stomach were also collected. The gut
204 segments were fixed in both 10% buffered formalin (for the histological examination) and
205 Carnoy's solution (for the morphometric analysis), while the other organ samples were only
206 fixed in 10% buffered formalin solution. The tissues were routinely embedded in paraffin wax
207 blocks, sectioned at a thickness of 5 μm , mounted onto glass slides and stained with
208 Haematoxylin & Eosin (H&E). Morphometric analyses (Image Pro-Plus software) (Fig. 1A)
209 were performed on 10 well-oriented and intact villi and 10 crypts chosen from the duodenum,
210 jejunum and ileum (Qaisrani et al., 2014). The evaluated morphometric indices were villus
211 height (Vh; from the tip of the villus to the crypt), crypt depth (Cd; from the base of the villi
212 to the submucosa) and their ratio (Vh/Cd) (Laudadio et al., 2012). Histological changes were
213 scored using a semi-quantitative scoring system as follows: absent or minimal (score 0), mild
214 (score 1) and severe (score 2). The semi-quantitative scoring system was assessed evaluating
215 the defined parameters of each organ (Table 2).

216

217 *Statistical analysis*

218 The statistical analysis was performed using SPSS 17 for Windows (SPSS, Inc., Chicago, IL,
219 USA; SPSS, 2008). The experimental unit was the pen. The influence of diet on the
220 performance parameters, hematological and serum biochemical traits and intestinal
221 morphometric measurements were analyzed using Student's *t*-tests for independent samples.
222 Morphometric data were also analyzed by means of one-way ANOVA (post hoc test:
223 Duncan's multiple range test) to evaluate the influence of the intestinal segment within each
224 dietary treatment. Mann-Whitney *U* tests were used to compare the histological scores
225 between treatments for each considered organ (spleen, thymus, bursa of Fabricius, liver,

226 glandular stomach, intestine, heart and kidney). Kruskal-Wallis tests were used to compare
227 intestine alterations among the considered segments (duodenum, jejunum, ileum and caecum),
228 within each dietary treatment. The results were considered statistically significant when
229 associated with a lower probability than 5%, and highly significant if the probability was
230 lower than 1%. The results were expressed as mean and pooled standard error of the mean
231 (SEM).

232

233 **Results**

234 The inclusion of TM in the medium-growing diet in the 43-97d period did not affect the
235 growth and slaughtering performances (Table 3), the blood and serum traits (Table 4) or the
236 FPD lesion incidence. The blood parameters fell within the physiological range.

237 The morphometric data are summarized in Table 5. There was no significant difference in the
238 morphology of the small intestine between the two dietary treatments ($P > 0.05$). In the C
239 group, the Vh was greater ($P < 0.01$) in the duodenum than in the other gut segments, while
240 the Vh/Cd ratio was only significantly different ($P = 0.01$) between the duodenum and the
241 ileum. In the TM group, Vh and Cd were greater ($P < 0.01$ and $P = 0.03$) in the duodenum
242 than in the other gut segments.

243 The glandular stomach, intestine, spleen, thymus, bursa of Fabricius and liver were the most
244 frequently affected organs, while the heart and kidneys showed no significant alterations.

245 Chronic inflammation, with lymphoid tissue activation, was observed in the intestinal
246 segments (Fig. 1B). The glandular stomach showed lymphoplasmacytic flogosis, with focal to
247 multifocal lymphoid tissue activation and different degrees of severity of epithelial squamous
248 metaplasia (Fig. 1C-D). The jejunum and the caecum showed the most severe alterations in
249 both dietary treatments ($P < 0.01$). White pulp hyperplasia or depletion was identified in the
250 spleen, while cortical depletion was detected in the thymus. Follicular depletion, with or

251 without intrafollicular cysts, was observed in the bursa of Fabricius (Fig. 2A-B). Finally, the
252 liver showed different degrees of lymphoid tissue activation (Fig. 2C-D). The histological
253 changes were not significantly different between the dietary treatments ($P > 0.05$).

254

255 **Discussion**

256 The trial was set up to study the effect of TM inclusion in the diet of medium-growing
257 chickens reared in free-range conditions. The inclusion of TM did not affect the performance,
258 the blood and serum traits or the welfare parameters of the birds. These results confirm that
259 TM can be used safely in poultry diets. Ramos-Elorduy et al. (2006) showed that up to 100
260 g/kg of dried yellow mealworms can be included in a broiler starter diet based on sorghum
261 and soybean meal, without negative effects on either performance or palatability. Ballitoc and
262 Sun (2013), including different levels of TM meal (5, 10, 20 and 100 g/kg, respectively) in a
263 standard commercial broiler diet, pointed out that the inclusion level of 10 g/kg TM had a
264 great impact on the growth performance of the broilers and improved the final body weight,
265 feed intake and feed efficiency, as well as the slaughter yield. Bovera et al. (2015) showed
266 that 30% inclusion of TM meal in the diet of Shaver brown broilers did not affect the final
267 body weight or blood traits, except for uric acid, albumin/globulin ratio, AST and ALT. In
268 another study, it was noted that TM could be successfully used to replace 4% soyabean meal
269 in laying hens diets (Wang et al., 2015).

270 FPD is a condition that involves inflammation and necrotic lesions on the plantar surface of
271 the bird's feet. It is considered to be an important welfare indicator in broilers (Ekstrand et al.,
272 1997; Meluzzi et al., 2008). It is histologically validated and is easy to use for the routine
273 assessment of broiler welfare in processing plants (Michela et al., 2012). A low prevalence
274 and severity of FPD is highly desirable as far as the health of birds and product quality are
275 concerned (Abd El-Wahab et al., 2012). Furthermore, the lesions can be a gateway for

276 bacteria that may spread hematogenously and impair product quality. Birds with severe
277 lesions may also show reduced weight gains, due to pain-induced decreases in feed intake
278 (Martland, 1985). FDP can be caused by several factors, the most important being the
279 condition of the litter in the broiler house (Bessei, 2006). Litter quality can be affected by diet
280 composition, as well as by the number and design of the drinkers in the pen (Lynn and Elson,
281 1990; Jones et al., 2005). The score obtained in the present study was zero for all of the birds,
282 thus showing that they were in a good welfare condition.

283 Intestinal morphology is the main indicator of gut health and functioning (Kristy et al., 2005).
284 In the present study, mealworm inclusion did not affect the morphology of the small intestine,
285 thus suggesting no influence on nutrient metabolization or performance. The greater
286 development of the duodenum, in relation to the other intestinal segments, is in agreement
287 with the results of Uni et al. (1999), Kondo (2003) and Murakami et al. (2007). In fact, the
288 duodenum is the intestinal tract with the fastest cell renewal, and is also the first segment of
289 the small intestine to receive physical, chemical and hormonal stimuli provoked by the
290 presence of the diet in the lumen (Macari, 1998).

291 In the present study, mealworm inclusion did not induce histological changes, thus suggesting
292 no negative influence on animal health. Glandular stomach alterations were considered
293 paraphysiological because, in the authors' opinion, they were probably related to the free-
294 range farming system. Outdoor access in fact entails less control of animal feeding, with the
295 possible ingestion of foreign bodies. However, the activation of the lymphoid tissue, in both
296 dietary treatments, represents the most interesting result. It has been reported that keeping
297 birds in free-range systems may favour the occurrence of gastrointestinal parasitic diseases,
298 with a high prevalence of coccidiosis and helminthiasis (Magwisha et al., 2002; Tomza-
299 Marciniak et al., 2014). Although all the birds were vaccinated against coccidiosis, the
300 immunological variation between strains of the same species might not have conferred total

301 protection (Chapman, 2014). Furthermore, the chronic flogosis observed in the intestinal
302 segments could suggest previous parasitic infections, with subsequent activation of the
303 lymphoid tissue. Tong et al. (2015) also evaluated the effects of outdoor access on the
304 lymphoid organ index in a local chicken breed and found that the spleen was the only
305 lymphoid organ that responded to outdoor access. The thymus, bursa of Fabricius and liver
306 were not altered, but the possibility of an adaption of the birds to the environment, in terms of
307 immune system modification, cannot be excluded.

308 In conclusion, the findings of this study suggest that the inclusion of yellow mealworm in
309 free-range chicken diets does not affect the welfare, productive performance or morphological
310 features of the birds. These results confirm previous data concerning the safety of TM meal
311 use in poultry diets. A key point in future studies would be to establish the point of view of
312 European consumers regarding the use of insects as feed for livestock. Legislative issues will
313 also have to be addressed at a European level, in order to allow insect meal to be used as
314 transformed animal protein to feed monogastric farm animals.

315

316 **Acknowledgments**

317 The authors are grateful to Mr. Alessandro Varesio and Mrs. Roberta Lacopo - “Fattoria La
318 Fornace”, Montechiaro d'Asti (At - Italy) - for their technical support. The research was
319 supported by the University of Torino (2014) and by the Regione Piemonte, Italy (PSR-PIAS
320 no. 08000558869).

321

322 **References**

- 323 1. Abd El-Wahab, A.; Visscher, C.F.; Wolken, S.; Reperant, J.-M.; Beineke, A.;
324 Beyerbach, M.; Kamphues, J., 2012: Foot-pad dermatitis and experimentally
325 induced coccidiosis in young turkeys fed a diet without anticoccidia. *Poultry Science*
326 **91**, 627–635.
- 327 2. Adeniji, A.A., 2007: Effect of replacing groundnut cake with maggot meal in the
328 diet of broilers. *International Journal of Poultry Science* **6**, 822–825.
- 329 3. Alves, S.P.; Cabrita, A.R.J.; Fonseca, A.J.M.; Bessa, R.J.B., 2008: Improved
330 method for fatty acid analysis in herbage based on direct transesterification followed
331 by solid-phase extraction. *Journal of Chromatography A* **1209**, 212–219.
- 332 4. AOAC, 2005. Official methods of analysis, 18th ed. Association of Official
333 Analytical Chemists, Washington, DC, USA.
- 334 5. Ballitoc, D.A.; Sun, S., 2013: Ground yellow mealworms (*Tenebrio molitor* L.) feed
335 supplementation improves growth performance and carcass yield characteristics in
336 broilers. *Open Science Repository Agriculture* (open-access), e23050425.
- 337 6. Belforti, M.; Gai, F.; Lussiana, C.; Renna, M.; Malfatto, V.; Rotolo, L.; De Marco,
338 M.; Dabbou, S.; Schiavone, A.; Zoccarato, I.; Gasco, L., in press: *Tenebrio molitor*
339 meal in rainbow trout (*Oncorhynchus mykiss*) diets: effects on animal performance,
340 nutrient digestibility and chemical composition of fillets. *Italian Journal of Animal*
341 *Science*.
- 342 7. Bessei, W., 2006: Welfare of broilers: a review. *World's Poultry Science Journal*
343 **62**, 455–466.
- 344 8. Bovera, F.; Piccolo, G.; Gasco, L.; Marono, S.; Loponte, R.; Vassalotti, G.;
345 Mastellone, V.; Lombardi, P.; Attia, Y.A.; Nizza, A., in press: Yellow mealworms

- 346 larvae (*Tenebrio molitor*, L.) as protein source for broilers: effects on growth
347 performance and blood profiles. *British Poultry Science*.
- 348 9. Campbell, T.W., 1995: *Avian Hematology and Cytology*, 2nd edn. Iowa State
349 University Press, Ames, Iowa, pp. 104.
- 350 10. Chapman, H.D., 2014: Milestones in avian coccidiosis research: A review. *Poultry*
351 *Science* **93**, 501–511.
- 352 11. De Marco, M.; Martínez, S.; Hernandez, F.; Madrid, J.; Gai, F.; Rotolo, L.; Belforti,
353 M.; Bergero, D.; Katz, H.; Dabbou, S.; Kovitvadhi, A.; Zoccarato, I.; Gasco, L.;
354 Schiavone, A., 2015: Nutritional value of two insect meals (*Tenebrio molitor* and
355 *Hermetia illucens*) for broiler chickens: apparent nutrient digestibility, apparent ileal
356 amino acid digestibility and apparent metabolizable energy. *Animal Feed Science*
357 *and Technology* **209**, 211–218.
- 358 12. De Marco, M.; Martinez, S.; Tarantola, M.; Bergagna, S.; Mellia, E.; Gennero, M.
359 S.; Schiavone, A., 2013: Effect of genotype and transport on tonic immobility and
360 heterophil/lymphocyte ratio in two local Italian breeds and Isa Brown hens kept
361 under free-range conditions. *Italian Journal of Animal Science* **12**, 481–485.
- 362 13. Ekstrand, C.; Algers, B.; Svedberg, J., 1997: Rearing conditions and foot-pad
363 dermatitis in Swedish broiler chickens. *Preventive Veterinary Medicine* **31**, 167–
364 174.
- 365 14. Folch, J.; Lees, M.; Sloane-Stanley, H., 1957: A simple method for the isolation and
366 purification of total lipids from animal tissue. *The Journal of Biological Chemistry*
367 **226**, 479–509.
- 368 15. Franco, J.R.G.; Murakami, A.E.; Natali, M.R.M.; Garcia E.R.M.; Furlan, A.C.,
369 2006: Influence of delayed placement and dietary lysine levels on small intestine

- 370 morphometrics and performance of broilers. *Brazilian Journal of Poultry Science* **8**,
371 233–241.
- 372 16. Gross, W.B.; Siegel, H.S., 1983: Evaluation of the heterophil/lymphocyte ratio as a
373 measure of stress in chickens. *Avian Disease* **27**, 972–979.
- 374 17. Hampson, D.J., 1986: Alterations in piglet small intestinal structure at weaning.
375 *Research in Veterinary Science* **40**, 32–40.
- 376 18. Henry, M.; Gasco, L.; Piccolo, G.; Fountoulaki, E., 2015: Review on the use of
377 insects in the diet of farmed fish: Past and future. *Animal Feed Science and*
378 *Technology* **203**, 1–22.
- 379 19. van Huis, A., 2013: Potential of insects as food and feed in assuring food security.
380 *Annual Review of Entomology* **58**, 563–583.
- 381 20. Hwangbo, J.; Hong, E.C.; Jang, A.; Kang, H.K.; Oh, J.S.; Kim, B.W.; Park, B.S.,
382 2009: Utilization of house fly-maggots, a feed supplement in the production of
383 broiler chickens. *Journal of Environmental Biology* **30**, 609–614.
- 384 21. Ijaiya, A.T.; Eko, E.O., 2009: Effect of replacing dietary fish meal with silkworm
385 (*Anaphe infracta*) caterpillar meal on growth, digestibility and economics of
386 production of starter broiler chickens. *Pakistan Journal of Nutrition* **8**, 845–849.
- 387 22. Jones, T.A.; Donnelly, C.A.; Dawkins, M.S., 2005: Environmental and management
388 factors affecting the welfare of chickens on commercial farms in the United
389 Kingdom and Denmark stocked at five densities. *Poultry Science* **84**, 1155–1165.
- 390 23. Khatun, R.; Howlider, M.A.R.; Rahman, M.M.; Hasanuzzaman, M., 2003:
391 Replacement of fish meal by silkworm pupae in broiler diets. *Pakistan Journal of*
392 *Biological Sciences* **6**, 955–958.
- 393 24. Kondo, N., 2003: Estudo das características morfométricas de diferentes regiões do
394 intestino delgado e índices zootécnicos em quarto linhagens de frangos de corte.

- 395 2003. Tese (Doutorado) Fac. Med. Vet. Zoot. Univ. Estadual Paulista, Botucatu,
396 Brazil.
- 397 25. Kristy, N.K.; Kelly, S.; Swanson, K.A.; Tappenden, L.B.S.; George, C.F.J., 2005:
398 Diet and age affect intestinal morphology and large bowel fermentative end-product
399 concentrations in senior and young adult dogs. *The Journal of Nutrition* **135**, 1940–
400 1945.
- 401 26. Laudadio, V.; Passantino, L.; Perillo, A.; Lopresti, G.; Passantino, A.; Khan, R.U.;
402 Tufarelli, V., 2012: Productive performance and histological features of intestinal
403 mucosa of broiler chickens fed different dietary protein levels. *Poultry Science* **91**,
404 265–270.
- 405 27. Lynn, N.; Elson, A., 1990: Which drinkers reduce possible downgrades. *Misset*
406 *World Poultry* **1**, 11–12.
- 407 28. Macari, M., 1998: Aspectos fisiológicos do sistema digestivo das aves. In 8^a.
408 Semana Acadêmica de Medicina Veterinária. São Paulo, Brazil. Pp. 4–18.
- 409 29. Madrid, J.; Martínez, S.; López, C.; Orengo, J.; López, M.J.; Hernández, F., 2013:
410 Effects of low protein diets on growth performance, carcass traits and ammonia
411 emission of barrows and gilts. *Animal Production Science* **53**, 146-153.
- 412 30. Magwisha, H.; Kassuku, A.; Nyvsgaard, N.; Permin, A., 2002: A comparison of the
413 prevalence and burdens of helminth infections in growers and adult free-range
414 chickens. *Tropical Animal Health Production* **34**, 205–214.
- 415 31. Makkar, H.P.; Tran, G.; Heuzé, V.; Ankers, P., 2014: State-of-the-art on use of
416 insects as animal feed. *Animal Feed Science and Technology* **197**, 1–33.
- 417 32. Maxwell, M.H., 1993: Avian blood leucocyte responses to stress. *World's Poultry*
418 *Science Journal* **49**, 34–43.

- 419 33. Maxwell, M.H.; Robertson, G.W., 1998: The avian heterophil leucocyte: a review.
420 *World's Poultry Science Journal* **54**, 155–178.
- 421 34. Martland, M.F., 1985: Ulcerative dermatitis in broiler chickens: The effects of wet
422 litter. *Avian Pathology* **14**, 353–364.
- 423 35. Meluzzi, A.; Fabbri, C.; Folegatti, E.; Sirri, F., 2008: Effect of less intensive rearing
424 conditions on litter characteristics, growth performance, carcass injuries and meat
425 quality of broilers. *British Poultry Science* **49**, 509–515.
- 426 36. Michela, V.; Pramparta, E.; Mirabitob, L.; Allaina, V.; Arnouldc, C., Huonnica,
427 D.; Le Bouquina, S.; Albaricd, O., 2012: Histologically-validated footpad dermatitis
428 scoring system for use in chicken processing plants. *British Poultry Science* **93**,
429 275–281.
- 430 37. Murakami, A.E.; Sakamoto, M.I.; Natali, M.R.M.; Souza, L.M.G.; Franco, J.R.G.,
431 2007: Supplementation of glutamine and vitamin E on the morphometry of the
432 intestinal mucosa in broiler chickens. *Poultry Science* **86**, 488–495.
- 433 38. National Research Council (NRC), 1994: *Nutrient Requirements of Poultry*, 9th
434 revised edn. National Academy Press, Washington, DC.
- 435 39. Oonincx, D.G.A.B.; de Boer, I.J.M., 2012: Environmental impact of the production
436 of mealworms as a protein source for humans - A Life Cycle Assessment. *PLoS*
437 *ONE* **7**:e51145.
- 438 40. Oyegoke, O.O.; Akintola, A.J.; Fasoranti, J.O., 2006: Dietary potentials of the
439 edible larvae of *Cirina forda* (westwood) as a poultry feed. *African Journal of*
440 *Biotechnology* **5**, 1799–1802.
- 441 41. Qaisrani, S.N.; Moquet, P.C.; van Krimpen, M.M.; Kwakkel, R.P.; Verstegen,
442 M.W.; Hendriks, W.H., 2014: Protein source and dietary structure influence growth

- 443 performance, gut morphology, and hindgut fermentation characteristics in broilers.
444 *Poultry Science* **93**, 3053–3064.
- 445 42. Ramos-Elorduy, J., 1997: Insects: a sustainable source of food? *Ecology of Food*
446 *and Nutrition* **36**, 247–276.
- 447 43. Ramos-Elorduy, J.; Medeiros-Costa, E.; Ferreira-Santos, J.; Pino-Moreno, J.M.;
448 Landero-Torres, I.; Ángeles-Campos, S.C.; García-Pérez, A., 2006: Estudio
449 comparativo del valor nutritivo de varios coleoptera comestibles de México y
450 *Pachymerus nucleorum* (Fabricius, 1792) (Bruchidae) de Brasil. *Interciencia* **31**,
451 512–516.
- 452 44. Renna, M.; Gasmi-Boubaker, A.; Lussiana, C.; Battaglini, L.M.; Belfayez, K.;
453 Fortina, R., 2014: Fatty acid composition of the seed oils of selected *Vicia* L. taxa
454 from Tunisia. *Italian Journal of Animal Science* **13**, 308–316.
- 455 45. Roncarati, A.; Gasco, L.; Parisi, G.; Terova, G., 2015. Growth performance of
456 common catfish (*Ameiurus melas* Raf.) fingerlings fed *Tenebrio molitor* meal diet.
457 *Journal of Insects as Food and Feed* **1**, 233–240.
- 458 46. Salamano, G.; Mellia, E.; Tarantola, M.; Gennero, M.S.; Doglione, L.; Schiavone,
459 A., 2010: Acute phase proteins and heterophil:lymphocyte ratio in laying hens in
460 different housing systems. *Veterinary Record* **167**, 749–751.
- 461 47. Sánchez-Muros, M.J.; Barroso, F.G.; Manzano-Agugliaro, F., 2014 : Insect meal as
462 renewable source of food for animal feeding: a review. *Journal of Cleaner*
463 *Production* **65**, 16–27.
- 464 48. SPSS, 2008. Statistical Package for the Social Sciences. Mc Graw-Hill, version
465 17.0, New York, USA.
- 466 49. Tomza-Marciniak, A.; Pilarczyk, B.; Tobiańska, B.; Tarasewicz, N., 2014:
467 Gastrointestinal parasites of free-range chickens. *Annals of Parasitology* **60**, 305-08.

- 468 50. Tong, H.B.; Cai, J.; Lu, J.; Wang, Q.; Shao, D.; Zou, J.M., 2015: Effects of outdoor
469 access days on growth performance, carcass yield, meat quality, and lymphoid organ
470 index of a local chicken breed. *Poultry Science* **94**, 1115–121.
- 471 51. Uni, Z.; Platin, R.; Sklan, D., 1999: Functional development of the small intestine in
472 domestic birds: Cellular and molecular aspects. *Poultry and Avian Biology Reviews*
473 **10**, 167–79.
- 474 52. Veldkamp, T.; Van Duinkerken, G.; Van Huis, A.; Iakemond, C.M.M.; Ottevanger,
475 E.; Bosch, G.; Van Boekel, M.A.J.S., 2012. Insects as a sustainable feed ingredient
476 in pig and poultry diets - a feasibility study. Wageningen UR Livest. Res., Report
477 638.
- 478 53. Wang, D.; Zhai, S.W.; Zhang, C.X.; Bai, Y.Y., An, S.H.; Xu, Y.N., 2005:
479 Evaluation of nutritional value of field crickets as a poultry feedstuff. *Asian -*
480 *Australasian Journal of Animal Sciences* **18**, 667–670.
- 481 54. Wang, X.C.; Zhang, H.J.; Wu, S.G.; Yue, H.Y.; Li, J.; Qi, G.H., 2015: Effect of
482 dietary protein sources on egg performance and egg quality of Jinghong laying hens
483 at peak production. *Scientia Agricultura Sinica* **48**, 2049–2057.
- 484 55. Welfare Quality[®] 2009: Welfare Quality[®] Assessment Protocol for Poultry
485 (Broilers, Laying Hens). Welfare Quality[®] Consortium: Lelystad, the Netherlands.
- 486 56. Zuidhof, M.J.; Molnar, C.L.; Morley, F.M.; Wray, T.L.; Robinson, F.E.; Khan,
487 B.A.; Al-Ani, L.; Goonewardene, L.A., 2003: Nutritive value of house fly (*Musca*
488 *domestica*) larvae as a feed supplement for turkey poults. *Animal Feed Science and*
489 *Technology* **105**, 225–30.
- 490
- 491
- 492

493 **Figures captions**

494

495 **Figure 1.** Histological and morphometric evaluation of the gastrointestinal tract of the free-
496 range chickens. A) **TM group.** Morphometric measurements of the villus height (Vh) and the
497 crypt depth (Cd) in the jejunum segment. 2.5× H&E. B) **TM group.** Severe and diffuse
498 mucosal lymphoplasmacytic and macrophagic flogosis with activation of the lymphoid tissue
499 (*) in the caecum. 5× H&E. C) **C group.** A normal glandular stomach. 5× H&E. D) **C group.**
500 Severe and diffuse mucosal lymphoplasmacytic flogosis with epithelial squamous metaplasia
501 (arrow) and activation of the lymphoid tissue (*) in the glandular stomach. 5× H&E.

502

503 **Figure 2.** Histological examination of the organs of the free-range chickens. A) **TM group.** A
504 normal follicle in the bursa of Fabricius. 10× H&E. B) **TM group.** Severe and diffuse
505 follicular depletion (*) in the bursa of Fabricius. 20× H&E. C) **C group.** A normal liver. 10×
506 H&E. D) **C group.** Mild and multifocal perivascular activation of the lymphoid tissue (arrow)
507 in the liver. 10× H&E.

508

510 **Table 1.** Ingredients and chemical composition of the experimental diets.

Ingredients (g/kg as fed)	Control diet	TM diet
Corn meal	720.0	720.0
Soybean meal	170.0	170.0
<i>Tenebrio molitor</i> meal	-	75.0
Gluten meal	75.0	-
Vitamin-mineral premix	35.0	35.0
Metabolizable energy (MJ/kg)	12.18	12.22
<i>Analized composition</i>		
Chemical composition (g/kg DM)		
Dry Matter	868	867
Crude Protein	169	168
Crude Fat	31	50
Crude Fiber	23	22
Indispensable aminoacids (g/kg DM)		
Arginine	9.3	10.0
Histidine	4.3	4.7
Isoleucine	7.1	6.9
Leucine	19.4	14.6
Lysine	7.0	9.0
Methionine	4.0	3.0
Phenylalanine	9.1	7.8
Threonine	7.0	7.0
Valine	8.1	8.2
Fatty acid composition (g/kg DM)		

511	C16:0	3.13	6.24
512	C18:0	0.47	1.06
513	C18:1 n9	4.00	10.05
514	C18:2 n6	10.42	20.41
515	C18:3 n3	0.52	0.73
516	Other fatty acids	0.35	1.12
517	Total SFA	3.66	7.75
518	Total MUFA	4.29	10.71
519	Total PUFA	10.94	21.15

520 *The mineral-vitamin premix (Trevit Volatili 3.5 - Trei - Rio Saliceto (RE) Italy) given values
521 are supplied per kg: 650.000 IU of vitamin A; 65.000 IU of vitamin D3; 650 IU of vitamin E;
522 80 mg of vitamin K; 80 mg of vitamin B1; 150 mg of vitamin B2; 770 mg of vitamin B3; 80
523 mg of vitamin B6; 0.5 mg of vitamin B12; 240 mg of pantothenic acid; 4700 mg of betaine;
524 1750 mg of Iron (II) carbonate; 1835 mg of Magnesium oxide; 1612 mg of Zinc oxide; 178
525 mg of Copper (II) oxide; 18.3 mg of Potassium iodide; 6.6 mg of Sodium selenite; 4100 mg
526 of DL-methionine; 5500 mg of L-lysine; 120 g Calcium carbonate; 450 g Calcium phosphate;
527 11.5 g of Sodium chloride. †SFA: saturated fatty acids; ‡MUFA: monounsaturated fatty
528 acids; §PUFA: polyunsaturated fatty acids. ¶Other FAs (all less than 0.40 g/kg DM): C12:0,
529 C14:0, C14:1 *cis*9, C16:1 *cis*9, C18:1 *cis*11, C20:0, C18:3 n6, C20:1 *cis*9, C20:1 *cis*11.

530

531

532

533

534

535 **Table 2.** Parameters evaluated for the histological semi-quantitative scoring system.

Organ	Alterations	Parameters evaluated
Spleen	White pulp hyperplasia	Number and dimension of follicles
	White pulp depletion	Number of apoptosis
		Cell types
Thymus	Cortical depletion	Cortico-medullar ratio
Bursa of Fabricius	Follicular depletion	Number of apoptosis
	Intrafollicular cysts	Cell types
Liver	Lymphoid tissue activation	Number of lymphoid aggregates
Glandular stomach	Lymphoplasmacytic flogosis	Distribution of inflammatory infiltrates
	Lymphoid tissue activation	Number and dimension of follicles
	Epithelial squamous metaplasia	
Intestine	Lymphoplasmacytic flogosis	Distribution of inflammatory infiltrates
	Lymphoid tissue activation	Number and dimension of follicles

536

537

538 **Table 3.** Growth and slaughtering performance of the free-range chickens (mean \pm s.d.)

	Control diet	TM diet	<i>P</i>
Initial body weight (d 43) (g)	720.2 \pm 26.81	712.3 \pm 19.62	0.612
Final body weight (d 97) (g)	2131.0 \pm 134.1	2162.5 \pm 306.8	0.845
Average daily intake (g)	112.75 \pm 9.9	111.6 \pm 11.6	0.873
Feed conversion ratio	4.4 \pm 0.7	4.4 \pm 0.6	0.982
Mortality rate (%)	0	0	-
Footpad dermatitis score	0	0	-
Chilled carcass (g)	1459.3 \pm 116.2	1544.8 \pm 106.5	0.104
Breasts (g)	347.1 \pm 41.6	370.8 \pm 51.3	0.275
Thighs (g)	479.2 \pm 45.3	502.9 \pm 45.2	0.273
Thigh muscle (g)	349.3 \pm 41.1	353.7 \pm 27.7	0.793
Thigh bone (g)	83.8 \pm 7.9	89.3 \pm 9.2	0.186
Spleen (g)	4.1 \pm 1.2	3.8 \pm 1.1	0.615
Bursa of Fabricius (g)	4.3 \pm 1.4	4.2 \pm 2.1	0.873
Liver (g)	36.7 \pm 4.0	39.4 \pm 6.5	0.278
Gizzard (g)	66.7 \pm 13.7	69.5 \pm 9.8	0.603
Abdominal fat (g)	40.6 \pm 35.4	45.1 \pm 23.7	0.752

539

540

541 **Table 4.** Hematological and serum biochemical traits of the free-range chickens (mean \pm s.d.).

	Control diet	TM diet	<i>P</i>
Erythrocytes (10^6 cell/ μ l)	2.40 \pm 0.34	2.63 \pm 0.38	0.641
Leukocytes (10^3 cell/ μ l)	9.42 \pm 1.90	9.94 \pm 1.74	0.659
H/L* ratio	0.55 \pm 0.21	0.51 \pm 0.22	0.798
Albumin (g/dl)	1.04 \pm 0.24	1.31 \pm 0.43	0.174
Total protein (g/dl)	3.98 \pm 0.58	4.05 \pm 0.79	0.293
AST † (UI/l)	190.49 \pm 20.29	197.97 \pm 15.16	0.649
ALT ‡ (UI/l)	14.49 \pm 8.01	13.99 \pm 5.64	0.411
Uric Acid (mg/dl)	4.99 \pm 1.43	3.93 \pm 1.34	0.894
Creatinine (mg/dl)	0.44 \pm 0.05	0.46 \pm 0.02	0.384
Triglycerides (mg/dl)	43.45 \pm 17.98	47.81 \pm 33.82	0.296
Cholesterol (mg/dl)	78.61 \pm 17.09	78.90 \pm 20.37	0.664
Glucose (mg/dl)	262.30 \pm 39.45	243.60 \pm 19.40	0.085
Phosphorus (mg/dl)	7.31 \pm 4.99	6.45 \pm 1.16	0.148
Magnesium (mEq/l)	0.90 \pm 0.53	1.02 \pm 0.53	0.979
Iron (μ g/dl)	54.83 \pm 37.03	61.54 \pm 46.84	0.361

542 *H/L: heterophils/lymphocytes; †AST: aspartate-aminotransferase; ‡ALT: alanino-
 543 aminotransferase.

544

545 **Table 5.** Intestinal morphometric measurements of the free-range chickens.

	Control diet	TM diet	SEM	<i>P</i>
Duodenum				
1. Villus height (µm)	2.29 ^a	2.49 ^a	0.06	0.126
2. Crypt depth (µm)	0.18	0.21 ^d	0.01	0.119
3. Villus height / crypt depth ratio	13.09 ^f	12.17	0.64	0.491
Jejunum				
1. Villus height (µm)	1.94 ^b	1.93 ^b	0.08	0.945
2. Crypt depth (µm)	0.17	0.17 ^e	0.01	0.813
3. Villus height / crypt depth ratio	11.29 ^{fg}	11.14	0.44	0.875
Ileum				
1. Villus height (µm)	1.60 ^c	1.72 ^b	0.07	0.405
2. Crypt depth (µm)	0.17	0.17 ^e	0.01	0.848
3. Villus height / crypt depth ratio	9.44 ^g	10.27	0.31	0.180

546 **Within each dietary treatment**, different letters (a, b, c) in the same column mean significant
 547 differences ($P < 0.05$) among the intestinal segments (duodenum, jejunum and ileum) for
 548 villus height.

549 **Within each dietary treatment**, different letters (d, e) in the same column mean significant
 550 differences ($P < 0.05$) among the intestinal segments (duodenum, jejunum and ileum) for
 551 crypt depth.

552 **Within each dietary treatment**, different letters (f, g) in the same column mean significant
 553 differences ($P < 0.05$) among the intestinal segments (duodenum, jejunum and ileum) for
 554 villus height/crypt depth ratio.