



AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Gastric and Large Colon Impactions Combined With Aggressive Enteral Fluid Therapy May Predispose to Large Colon Volvulus: 4 Cases

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1832466> since 2023-04-05T07:35:22Z

Published version:

DOI:10.1016/j.jevs.2021.103617

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 **Gastric and large colon impactions combined with aggressive enteral fluid**
2 **therapy may predispose to large colon volvulus: 4 cases**

3

4

5 Gessica Giusto¹, Anna Cerullo^{1*}, Marco Gandini¹

6 ¹Department of Veterinary Sciences, University of Turin, L.go Paolo Braccini 2-5,
7 Grugliasco (TO), Italy.

8 **Corresponding author email: anna.cerullo@unito.it*

9 **Abstract**

10 **Background:** To resolve large colon impactions, frequent enteral administration of
11 large volumes of water (8-10 liters in a 500 kg horse) has been suggested.
12 Furthermore, in large colon volvulus, the simultaneous presence of gastric impaction
13 has been described as a possible predisposing factor .

14 **Objectives:** To describe the clinical and surgical features of horses with large colon
15 volvulus associated with aggressive enteral fluid therapy performed to resolve an initial
16 large colon impaction.

17 **Methods:** Records of horses that underwent exploratory laparotomy at the Veterinary
18 Teaching Hospital of Turin between 2012-2019 were reviewed. Clinical and surgical
19 features of cases initially diagnosed with large colon impaction that developed a large
20 colon volvulus after enteral administration of large volumes of fluids were retrieved.

21 **Results:** Four horses met the criteria. In all horses an initial diagnosis of large colon
22 impaction was made by the referring veterinarian. In all cases a sudden increase in
23 pain was noticed shortly after enteral administration of large volumes (8-10 liters) of
24 water. Administration of analgesic didn't resolve the pain in any of the horses that were
25 then referred. A large colon volvulus was found during exploratory laparotomy in all
26 cases. A moderate gastric impaction was also found intraoperatively, which was
27 confirmed by gastroscopy postoperatively.

28 **Conclusions:** Nasogastric administration of large quantity of fluids with a pre-existing
29 gastric and colon impaction, may reduce the available space in the abdominal cavity
30 and potentially predispose the colon volvulus. More frequent enteral administration of
31 smaller volumes of fluids might be preferred when treating large colon impactions.

32 **Keywords:** horse; colic; colon impaction; enteral fluid therapy; large colon volvulus
33 Large colon impaction is considered the second most common cause of colic in horses
34 as well as the most common example of simple intestinal obstruction [1,2]. It has a
35 multifactorial aetiology and is often associated with moderate pain; generally, it is
36 treated medically with the administration of enteral and parenteral fluids [3,4]. In
37 several studies, enteral fluid therapy was more effective than intravenous fluid therapy
38 to resolve large colon impaction [5-8]. An experimental study demonstrated that the
39 administration of up to 10 l of water in a 500-kg horse (20 ml/kg) via a nasogastric tube
40 was safe and effective in hydrating faeces in the colon [7]. It is important to consider
41 that the administration of excessive volumes of enteral water through gravity flow may
42 be associated with a worsening of colic symptoms, following excessive distension of
43 the gastric wall. This latter condition can, in fact, represent an important pain stimulus,

44 especially if gastric emptying is hindered and the contents flow slowly [8]. This
45 condition is also associated with an increase in intra-abdominal pressure, that could
46 predispose to large colon dislocation or volvulus (LCV) [9,10] alone or in combination
47 with other predisposing factors. The following report describes the clinical signs and
48 surgical characteristics of horses with large colon volvulus which developed after
49 aggressive enteral fluid therapy to resolve an initial large colon impaction. In all horses,
50 a moderate gastric impaction was also found intraoperatively, which was confirmed by
51 gastroscopy postoperatively.

52

53 **1. Materials and methods**

54 **1.1 Case selection**

55 The study included horses hospitalised for acute abdomen and undergoing
56 exploratory laparotomy at the Veterinary Teaching Hospital of Turin from 2012 to
57 2019. All horses were diagnosed with colon impaction that developed a large colon
58 volvulus after enteral administration of large volumes of fluids.

59 **1.2 Case signalment**

60 Four horses met the criteria for the study; ages ranged from 16 to 19 years, with a
61 mean of 17.5 years, and mean weight was 563 kg (500–675). Two horses were
62 females, one was gelding, and one was entire male.

63 **2. Results**

64 **2.1 Case presentation** We report the history and clinical findings of four horses
65 initially seen by four different referring veterinarians for inappetence and mild
66 colic symptoms. Horses had mild to moderate colic pain on the first visit; the

referring veterinarians carried out the clinical examination and rectal exploration and issued a diagnosis of large colon impaction. Treatment was initiated with spasmolytics and with the administration of intravenous fluid therapy. Approximately 8 to 10 litres of water were administered to each horse via a nasogastric tube, following which the patients began to show a worsening of the clinical signs, with an increase in heart rate and colic pain not respondent to analgesics.

2.1.1 Case 1 A 16-year-old 500-kg Warmblood mare was referred for colic syndrome. The referring veterinarian had diagnosed a large colon impaction and treatment was initiated. Spasmolytics, analgesics and intravenous fluid therapy were administered. Approximately 10 litres of water (20 ml/kg) were also administered via a nasogastric tube, following which the patient began to show a worsening of the clinical signs with an increase in heart rate and severe colic pain not respondent to treatment with flunixin meglumine¹ (1.1 mg/kg IV) and detomidine hydrochloride² (0.2 ml/100 kg IV). The horse was then referred. Upon arrival, the horse was in severe pain and difficult to handle and explore. A dose of xylazine³ (1 mg /kg IV) and butorphanol⁴ (0.01 mg/kg IV) was administered to perform a quick preoperative examination and then used for anaesthesia premedication. At physical examination, increased heart rate (70 beats/min), absent gut sounds, increased values of PCV (45%) and TP (7 g/dl) were found. A severe large colon impaction was palpated on abdominal examination *per rectum*. Passage of a nasogastric tube did not reveal any net reflux. Because of the unrelentless pain, ultrasonography was not performed. Based on clinical findings, surgical

treatment was advised. After induction of general anaesthesia and aseptical preparation of the operating field, a ventral midline laparotomy was performed. This revealed a 360° counterclockwise colon volvulus, which was then corrected by reverse rotation. A pelvic flexure enterotomy was also performed to resolve the initial colon impaction. The large colon presented just mild oedematous intestinal walls, whose colour improved during surgery. Palpation revealed the presence of a distended stomach with solid content. At the end of the surgery, the abdomen was closed routinely with a double-layer closure on the *linea alba* and on the skin. Surgery lasted 125 min, and the horse recovered from anaesthesia uneventfully. In the first postoperative day, a gastroscopy was performed, and gastric impaction confirmed (Fig.1 A, B). The horse was then administered 2 litres of water with 240 mg of dioctyl sodium sulfosuccinate (DSS)⁵ and 36 g of Sorbitol⁵ twice a day for 3 days. Follow-up gastroscopy on the fourth postoperative day confirmed resolution of the impaction. No complications were encountered during hospitalisation, and the patient was discharged after eight days. A telephone follow-up was obtained by owner interview, and the horse was alive after 40 months.

2.1.2 Case 2 A 17-year-old 535-kg Warmblood entire male was referred for severe colic pain. The referring veterinarian carried out the clinical examination and rectal exploration, and a severe large colon impaction was diagnosed. Treatment was initiated with intravenous fluid therapy, and approximately 10 litres of water (18 ml/kg) were administered via a nasogastric tube. Approximately 10 minutes later, the horse began to show

worsening of the clinical signs, with colic pain not respondent to treatment with analgesics. At presentation, the horse had severe abdominal pain, heart rate was increased, and there was decreased intestinal motility. A distended small intestine was palpated on rectal examination. Exploratory laparotomy was recommended due to the severity of abdominal pain. A 270° counterclockwise colon volvulus was found and corrected by reverse rotation; additionally, pelvic flexure enterotomy was performed. The intra-abdominal palpation revealed the concomitant presence of gastric impaction which was not surgically resolved. Postoperative gastroscopy confirmed the presence of the gastric impaction (Fig. 1C) that was treated as per case #1 and was resolved at follow-up gastroscopy 4 days later. The horse was discharged on the 9th day postoperation and was alive at 20 months follow-up.

2.1.3 Case 3 An 18-year-old 675-kg Warmblood broodmare was referred for abdominal pain. The referring veterinarian had diagnosed a large colon impaction which was treated medically with administration of parenteral fluids, analgesics and abundant enteral fluids (approximately 15 litres- “one full bucket”- according to the referring veterinarian) via a nasogastric tube. After gastric intubation, the horse began to show severe colic pain and was referred. Upon arrival, the horse had moderate abdominal pain, a slightly increased heart rate, decreased intestinal motility and mild dehydration. The large colon was distended, and a taenial band, painful on traction, was palpated on rectal examination; exploratory laparotomy was recommended. A 360° counterclockwise colon volvulus was found and resolved with

reverse rotation; the large colon was exteriorised, and a pelvic flexure enterotomy was performed to resolve the initial colon impaction. Intra-abdominal palpation revealed the concomitant presence of a moderate gastric impaction. The abdomen was closed routinely. The surgery lasted approximately 80 minutes, and the horse recovered from anaesthesia uneventfully. No complications were encountered during hospitalisation, the patient was discharged after 8 days and resulted alive after 12 months.

2.1.4 Case 4 A 19-year-old 540-kg Warmblood gelding was referred for acute abdominal pain, which started after the administration of about 10 litres of water via a nasogastric tube to resolve an initial colon impaction. Upon arrival, the horse showed moderate abdominal pain not responding to analgesics and had an increased heart rate and decreased intestinal motility. A large colon displacement was suspected on rectal examination, and an exploratory laparotomy was performed. A 360° counterclockwise colon volvulus, which was resolved by reverse rotation, and a gastric impaction were found. A pelvic flexure enterotomy was performed, and the abdomen was closed routinely. Treatment with DSS and sorbitol was performed as in the previous cases, and the horse was discharged on postoperative day 10. The horse was alive at follow-up at 5 months.

3. Discussion

All horses included in this study had an initial large colon impaction and began to show a worsening of their clinical condition following the enteral administration of large quantities of water (18–22 ml/kg) through gravity flow. Pain demonstrated by the

163 subjects after administration of water was well more than what is normally shown by
164 horses with colon impaction. Administration of analgesic did not resolve the pain in
165 any of the horses, which were then referred because of the unrelentless pain. Based
166 on clinical findings, surgical treatment was advised and colon torsion was diagnosed
167 in all patients during exploratory laparotomy Torsion of the colon is one of the most
168 painful and rapidly fatal causes of colic in horses, with a high mortality rate associated
169 with the onset of postoperative complications. Therefore, recognising the risk factors
170 could be beneficial to prevent the onset of this problem. Receiving medication in the
171 previous 7 days, drastic changes in diet, reduced exercise, increased height,
172 management changes and parturition in mares are the main risk factors [11]. However,
173 the horses included in our study had not been exposed to these risk factors for LCV,
174 if not an anatomical predisposition relating to the height of the animals, as they are all
175 Warm-blood horses. Gastric impaction, which is a rare condition characterised by
176 persistent and progressive accumulation of dehydrated ingesta in the stomach [12], is
177 also hypothesised to be considered among the predisposing factors of large colon
178 volvulus [13].

179 Gastric impaction also leads to a reduction in the passage of ingesta, which causes
180 an alteration of the colon contents and, therefore, a change in the colonic microbiota
181 and pH that can predispose to the onset of dislocation or volvulus [11]. However, the
182 connection between the two problems it not well known, since gastric impaction could
183 cause variations in intra-abdominal pressure that predispose to variations in the
184 position of the large colon. However, it is possible that the displaced colon compresses
185 the pylorus, causing an arrest of the passage of gastric contents into the small
186 intestine; therefore, gastric impaction is not a cause, but a consequence [13]. Also, a

187 reduced gastrointestinal motility could explain the concurrent presence of large colon
188 impactions and gastric impactions. Changes in intra-abdominal pressure have been
189 studied mainly in post-partum mares [14]. Both in horses with gastric impaction and in
190 mares after parturition, pressure changes, albeit opposite, occur in the abdominal
191 cavity, which could make the colon more prone to dislocation or torsion [10]. Also, in
192 our study, we hypothesised that gastric distension could predispose to the
193 development of volvulus of the large colon; however, while McGovern [13] describes
194 cases of severe gastric impaction, in our cases, gastric distension could have been
195 caused by enteral administration of large volumes of fluids in conjunction with the pre-
196 existing condition of mild or moderate gastric impaction, found intraoperatively and
197 confirmed at gastroscopy. In both cases, however, it is not possible to say with
198 certainty that this is a causal association rather than a coincidence.

199 The presence of a gastric impaction must be considered when initiating an enteral fluid
200 therapy to resolve the large colon impaction because the gastric distension could
201 cause a drastic increase in intra-abdominal pressure, with a consequent greater
202 predisposition of the colon to torsion. There are several risk factors, as previously
203 mentioned, for the development of large colon volvulus. Even severe large colon
204 impaction could physically displace the large colon due to the weight of the ingesta,
205 which could propagate into a large colon volvulus, could worsen the clinical conditions
206 of horses with colic syndrome. It is advisable to assess the severity of intestinal
207 impaction by rectal examination, and it is also advisable, whenever possible, to
208 perform a transabdominal ultrasound exam to evaluate gastric distension before
209 starting enteral therapy to avoid worsening a pre-existing gastric distension condition.

210 An experimental study has shown that intra-abdominal pressure increases already
211 after the enteral administration of 5 litres of fluids [9]; the clinical significance of this
212 condition is unknown. It is not possible, however, to establish whether the change in
213 intra-abdominal pressure also occurred in our cases, since intra-abdominal pressure
214 was not measured before and after the administration of fluids. However, the number
215 of cases considered is small, and it is impossible to know if there was some partial
216 rotation of the colon before the torsion and, at the same time, it is not clear whether
217 the gastric impaction, found intraoperatively, was present at the first visit by the
218 referring vet. The effectiveness of enteral fluid therapy can be enhanced, rather than
219 by increasing the amount of water administered, by combining the administration of
220 water with the use of other products such as DSS. It is an anionic surfactant, and
221 several studies have been conducted using extremely high dosages of DSS as
222 laxatives and to define its toxic dose in horses [15-17]. Diethyl sodium sulfosuccinate
223 is also a surfactant and used in humans mainly to promote hydration of faecal content
224 by reducing the surface tension and allowing fluids to penetrate ingesta or faecal mass
225 [18]. We hypothesised that water alone is not able to permeate the gastric ingesta.
226 Administration of a low quantity of water with a low DSS amount twice daily for 3 days
227 resulted in a resolution of the gastric impaction, possibly by increasing water
228 permeability of the ingesta. In the study by McGovern et al., cases of gastric impaction
229 associated with colon dislocation and torsion were subjected to intraoperative or
230 postoperative euthanasia [13]. Although the number of cases is smaller, we report a
231 favourable short- and long-term survival. Further, in our cases, the colon resulted
232 oedematous and hyperhaemic, no resection was considered necessary, and simple
233 reposition and emptying were performed. This is probably due to the short-time

234 intercourse between the onset of symptoms and surgical treatment, with a prompt
235 referral by the attending veterinarians.

236 **4. Conclusions**

237 It is possible that the enteral administration of large quantities of water may lead to a
238 worsening of colic symptoms and predispose to further problems such as colon
239 displacement or torsion. Although enteral fluid therapy is the most effective treatment
240 for resolving episodes of colon impaction, it must be calibrated to hydrate the gastro-
241 enteric content and stimulate motility, but not to cause excessive distension of the
242 stomach that can be a pain stimulus and a predisposing factor to the development of
243 further problems such as large colon volvulus.

244 Based on what has been observed in these cases, we have modified our approach for
245 the therapy of gastric and large colon impaction referred to the hospital in cases like
246 those described in our study. Therapy consists of combining parenteral administration
247 of crystalloids with continuous enteral administration of fluids (through a nasogastric
248 tube connected to fluid bags). This method may reduce the risk of a sudden distension
249 of the gastric wall, as could happen by administration of bolus fluids through gravity
250 flow or via a pump. Another alternative could be the frequent enteral administration of
251 small volumes of water (4–10 ml/kg) through gravity flow with low doses of DSS. We
252 therefore hypothesise that this therapy may resolve impactions whilst reducing the risk
253 of colon volvulus.

254 **Authors' declaration of interest:** no conflict of interest have been declared.

255 **Ethical animal research:** the owners were informed and signed an informed consent
256 form. Animal welfare was respected throughout the hospital stay.

257 **Authorship:** all authors contributed to this case and preparation of the manuscript. All
258 authors gave their final approval.

259 **Source of funding:** None.

260 **Manufacturers' addresses:**

261 ¹IZO s.r.l. a socio unico Via San Zeno 99/A – 25124, Brescia (BS) - Italia

262 ²Ecuphar Italia S.R.L. Viale Francesco Restelli, 3/7, piano 1, 20124 Milano (MI) – Italia

263 ³P.H. Farmaceutici SpA – Via Aguggiari, 4, 20900 Monza (MB) – Italia

264 ⁴ACME s.r.l. – Via Portella della Ginestra, 9, 42025, Cavriago (RE) – Italia

265 ⁵SIT s.r.l. – ^aVia Cavour, 70, 27035 Mede (PV) - Italia

266

267 **References:**

268 [1] Cohen ND, Matejka PL, Honnas CM, et al. Case-control study of the association
269 between various management factors and development of colic in horses. Texas
270 Equine Colic Study Group. J Am Vet Med Assoc. 1995 Mar 1; 206 (5): 667-73.

271 [2] Proudman CJ. A two-year, prospective survey of equine colic in general practice.
272 Equine Vet J. 1992 Mar; 24 (2): 90-3.

273 [3] Hillyer MH, Taylor FG, Proudman CJ, et al. Case control study to identify risk
274 factors for simple colonic obstruction and distension colic in horses. Equine Vet J.
275 2002 Jul; 34 (5): 455-63.

276 [4] Jennings K, Curtis L, Burford J, et al. Prospective survey of veterinary practitioners'
277 primary assessment of equine colic: clinical features, diagnoses, and treatment of
278 120 cases of large colon impaction. BMC Vet Res. 2014; 10 (Suppl 1): S2.

- 279 [5] Hallowell GD. Retrospective study assessing efficacy of treatment of large colonic
280 impactions. Equine Vet J. 2008 Jun; 40 (4): 411-3.
- 281 [6] Lopes MA, Walker BL, White NA 2nd, et al. Treatments to promote colonic
282 hydration: enteral fluid therapy versus intravenous fluid therapy and magnesium
283 sulphate. Equine Vet J. 2002 Jul; 34 (5): 505-9.
- 284 [7] Lopes MA, White NA 2nd, Donaldson L. et al. Effects of enteral and intravenous
285 fluid therapy, magnesium sulfate, and sodium sulfate on colonic contents and feces
286 in horses. Am J Vet Res. 2004 May; 65 (5): 695-704.
- 287 [8] Montreal L, Navarro M, Armengou L, et al. Enteral fluid therapy in 108 horses with
288 large colon impactions and dorsal displacements. Vet Rec. 2010 Feb 27; 166 (9):
289 259-63.
- 290 [9] Barrett EJ, Munsterman AS, Hanson RR. Effects of gastric distension on
291 intraabdominal pressures in horses. J Vet Emerg Crit Care (San Antonio). 2013
292 Jul-Aug; 23 (4): 423-8.
- 293 [10] Blikslager, A.T. Gastric impaction and large colon volvulus: Can one lead to the
294 other? Equine Vet Educ. 2015; 27: 460-461.
- 295 [11] Suthers JM, Pinchbeck GL, Proudman CJ, et al. Risk factors for large colon
296 volvulus in the UK. Equine Vet J. 2013 Sep; 45 (5): 558-63
- 297 [12] Freeman, D. E. Gastric impaction. Equine Vet. Educ. 2011; 23 (4): 174-176
- 298 [13] McGovern, K. F., Suthers, J. M., James, F. M., et al. Gastric impaction
299 associated with displacement and volvulus of the large colon in seven mature
300 horses. Equine Vet. Educ. 2015; 27 (9), 453–459.

- 301 [14] Weese JS, Holcombe SJ, Embertson RM, et al. Changes in the faecal
302 microbiota of mares precede the development of post-partum colic. Equine Vet J.
303 2015 Nov; 47 (6): 641-9
- 304 [15] Freeman DE, Ferrante PL, Palmer JE. Comparison of the effects of intragastric
305 infusions of equal volumes of water, dioctyl sodium sulfosuccinate, and magnesium
306 sulfate on fecal composition and output in clinically normal horses. Am J Vet Res.
307 1992 Aug; 53 (8): 1347-53
- 308 [16] Moffatt RE, Kramer LL, Lerner D, et al. Studies on dioctyl sodium sulfosuccinate
309 toxicity: clinical, gross and microscopic pathology in the horse and guinea pig. Can
310 J Comp Med. 1975 Oct; 39 (4): 434-41
- 311 [17] Southwood L.L., Schott H.C., Wunn D. et al. Effects of dioctyl sodium
312 sulfosuccinate on hematological values in normal horses. J Equine Vet Sci. 1999
313 Apr; 19 (4): 266-270
- 314 [18] David H. Shaw. Drugs Acting on the Gastrointestinal Tract. In: Frank J. Dowd,
315 Barton S. Johnson, Angelo J. Mariotti, editors. Pharmacology and Therapeutics for
316 Dentistry. 7th ed. Mosby, 2017; 404-416.
- 317

318 **Figure legends**

- 319 Figure 1: Gastric impaction of Case 1 at gastroscopy in the first postoperative day(
- 320 A, B). Gastroscopy of Case 2 obtained during the second postoperative day (C).
- 321