

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Nasal and Tracheal Cytological Changes After Total Laryngectomy in Long-Term Survivors

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1617575> since 2022-03-01T15:23:25Z

Published version:

DOI:10.1177/0003489416676500

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This work is licensed under a Creative Commons-AttributionNonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) <https://creativecommons.org/licenses/by-nc-nd/4.0/>



Nasal and tracheal cytological changes after total laryngectomy in long-term survivors

Giuseppe Riva¹, MD, Monica Boita², PhD, Andrea Corvino¹, MD, Matteo Sensini¹, MD,
Daniela Peruzzetto¹, MD, Luigi Chiusa³, MD, Giancarlo Pecorari¹, MD, Massimiliano
Garzaro¹, MD

1 1st ENT Division, Surgical Sciences Department, University of Turin, Via Genova 3, Turin
- Italy

2 Allergology and Clinical Immunology, Medical Science Department, University of Turin,
Via Genova 3, Turin - Italy

3 Department of Biomedical Sciences and Human Oncology, University of Turin, Via
Genova 3, Turin - Italy

The authors declare that they have no Financial Support/Funding.

The authors declare that they have no conflict of interest.

Corresponding Author and Reprints Request:

Giuseppe Riva, MD

1st ENT Division, Surgical Sciences Department, University of Turin

Via Genova, 3 - 10126 Turin, Italy. E-mail: giuseppe.riva84@gmail.com

Tel: +39.011.633.66.88 - Fax +39.011.633.66.50

Abstract

Objective. Complete separation of upper and lower respiratory tract after total laryngectomy results in permanent effects on nasal cavities and tracheo-bronchial airways. Aim of this study is evaluating nasal and tracheal cytological alterations of mucosa in laryngectomy long-term survivors, analyzing the feasibility of scraping for cytological examination of tracheal mucosa.

Methods. Twenty-five laryngectomy patients underwent symptoms' evaluation, endoscopic fiber optic examination, prick tests and nasal and tracheal scraping for cytological exam. Twenty-five healthy subjects underwent the same assessment, except for tracheal scraping. Eleven laryngectomy patients accepted inferior turbinate biopsy for histological examination.

Results. Nasal cytological analysis demonstrated mucous cell metaplasia in 20% of laryngectomized patients, but it was absent in all healthy subjects; no squamous cell metaplasia was found in both groups. In 15 patients (60%) bacteria were present, without inflammatory infiltrate. Tracheal cytological analysis demonstrated a quite high rate of squamous cell metaplasia (24%), neutrophilic infiltrate (32%) and presence of bacteria (40%). Histological examination of inferior turbinate showed submucosal stromal fibrosis in all patients and submucosal inflammatory infiltrate in one case (9%).

Conclusion. Nasal cavities and trachea of laryngectomy patients undergo long-term cytological and histological changes of mucosa and submucosa, probably due to airflow modifications.

Keywords: Nasal cytology; Tracheal cytology; Laryngectomy; Histological changes; Cytological changes; Total laryngectomy

Introduction

Complete separation of upper and lower respiratory tract after total laryngectomy results in permanent effects on nasal cavities and tracheo-bronchial airways. In laryngectomy patients the air comes directly into trachea. Therefore, nasal cavities are excluded from respiration and cannot carry out their physiological functions, such as filtering, heating and moistening of the inspired air and smell. Hyposmia, impaired mucociliary function, cytological and histological alterations of nasal mucosa arise.¹⁻⁴ Moreover, the "unconditioned" inspired air may induce chronic inflammation of trachea and bronchi, with cough, excessive secretions and sometimes crusting, especially in the first three months after total laryngectomy.

Very few studies evaluated cytological and histological changes of nasal mucosa in laryngectomy individuals¹⁻² and none investigated such alterations in tracheal mucosa. Previous studies focused on smell disorders of these patients.⁴

Nasal airflow has been proposed to be necessary for normal functioning of nasal epithelium. However, there is no convincing evidence that airflow changes are a causative factor in nasal pathology.⁵ Epidemiological studies have showed an increased incidence of lower respiratory tract infections in patients with a tracheostomy.^{6,7} Conversely, rhinosinusitis has not been generally mentioned as associated with laryngectomy.⁵ Previous studies have showed a microbial colonization of nasal cavities and trachea after total laryngectomy, without clinical signs of infection.^{1,8,9,10}

Aim of this study was to evaluate nasal and tracheal cytological alterations of mucosa in laryngectomy long-term survivors, due to airflow changes. Furthermore, we analyzed the feasibility of scraping for cytological examination of tracheal mucosa.

Materials and methods

Twenty-five healthy subjects and twenty-five consecutive long-term survivor patients who underwent total laryngectomy for advanced laryngeal cancer between 2005 and 2011 at our Department were evaluated in this retrospective cross-sectional comparative study. All laryngectomy patients included in the study were disease free at the time of engagement. Moreover, they had no clinically evident viral, bacterial and/or mycotic diseases involving nasal cavities or lower respiratory tract. No antimicrobial therapy was administered to patients for at least one week before the evaluation. Enrolled patients had at least a 2-year follow-up period (mean 52.97 ± 28.96 months, range 26-97 months), in order to evaluate long-term mucosal alterations. No patient had tracheal cannula and/or tracheo-esophageal voice prostheses at enrolment. They were previously evaluated for nasal and tracheal microbial colonization at our Department.¹⁰ Healthy subjects were patients with vocal fold nodules or polyps and without any clinical and endoscopic sign of sinonasal disease. Written informed consent was obtained in every case. Institutional Review Board approval was obtained.

At enrolment, laryngectomy patients underwent symptoms evaluation (rhinorrhea, hyposmia, cough), endoscopic fiber optic nasal and tracheal examination, skin prick tests and nasal and tracheal scraping for cytological exam. Healthy subjects underwent the same assessment, except for tracheal cytological, because they had no tracheostomy. Eleven laryngectomy patients accepted inferior turbinate biopsy for histological examination. **No control subject underwent nasal biopsy.**

Cytological samples were obtained by scraping with a Rhino-pro® Curette (ASI Arlington Scientific Inc, Springville, UT, USA). Nasal samples were collected from the medial portion of the inferior turbinate, while tracheal samples from posterior wall of the

1
2
3
4 trachea 2 cm below the tracheostomy, preventing contamination from the tracheostomy
5
6 border. After sampling, the material is laid on a microscope slide, fixed in 95° alcohol for 4 s
7
8 and stained by the May-Grünwald-Giemsa method. Observation is performed by a common
9
10 optical microscope, provided it is able of a 1000x magnification, analyzing at least 50
11
12 microscopic fields. Scraping and cytological analysis were performed by an otolaryngologist
13
14 expert in these procedures and not knowing whether the subjects underwent laryngectomy or
15
16 not. A pathologist re-analyzed and interpreted the tissue samples. All subjects underwent skin
17
18 prick tests to correctly evaluate cytological data, in particular inflammatory infiltrate, when an
19
20 allergic aetiology is suspected. All cytological samples were collected in November, when the
21
22 majority of pollens were absent. Specimen evaluation was performed using a checklist for
23
24 cytological findings.¹¹ Respiratory tracheal mucosa is similar to nasal mucosa, therefore we
25
26 used the same staining method. Mucous cell metaplasia was defined as the presence of
27
28 mucous cells > 25% of epithelial cells, while squamous cell metaplasia as the presence of
29
30 squamous cells. Cytological atypias were defined as nuclear enlargement associated with
31
32 increased nuclear-cytoplasmic ratio, hyperchromatism, chromatin clumping with moderately
33
34 prominent nucleoli, irregular nuclear membranes and multinucleation, and variation in size
35
36 and/or shape of the cells and nuclei.¹¹

37
38
39
40
41
42 Statistical Package for Social Sciences (SPSS), version 17.0 was employed for data
43
44 analysis. A descriptive analysis of all data was performed and they were reported as means or
45
46 percentages and standard deviations. The Kolmogorov-Smirnov test demonstrated a non
47
48 Gaussian distribution of variables, so non parametric tests were used. The chi-square test was
49
50 used to assess differences between groups in the mean of categorical variables. A p value less
51
52 than 0.05 was considered statistically significant.
53
54
55
56
57
58
59
60

Results

Patients

Mean age was 62.64 ± 9.53 years (range 48-76 years) for healthy subjects and 68.76 ± 8.10 years (range 50-83 years) for laryngectomy patients. Table 1 reports socio-demographic and clinical characteristics, such as age and sex, tobacco, alcohol consumption, previous exposure to irritant substances (i.e. asbestos dust) and allergies, and tumor related factors, such as histological type, TNM classification, grade and stage. There were no clinical or demographic differences (sex, tobacco, alcohol consumption, previous exposure to irritant substances and allergies) between control group and treatment group ($p > 0.05$).

All patients underwent bilateral selective neck dissection, associated to total laryngectomy. Total laryngectomy was performed as salvage surgery for local recurrence in 20% of cases: in 3 patients after previous laser cordectomy and in 2 patients after partial supracricoid laryngectomy. Adjuvant radiation therapy or chemo-radiotherapy was administrated in 8 and 2 patients respectively, because of histopathological adverse features (extracapsular nodal spread, positive margins, pT4 primary, N2 or N3 nodal disease, perineural invasion, vascular embolism).

Concerning symptoms evaluation, anterior rhinorrhea was reported by 56% of patients, dry or productive cough by 48% and hyposmia by all patients.

At nasal fiber optic endoscopic evaluation, no statistically significant difference was observed between laryngectomy patients and healthy subjects ($p > 0.05$). The main findings were turbinate hypertrophy (40%), pale nasal mucosa (40%) and serous nasal secretions (68%). Polypoid degeneration of nasal mucosa was seen in one case (4%). Regarding tracheal endoscopic exam, 64% of patients had dry mucosa, while one case (4%) had hyperaemia. Table 2 shows symptoms and endoscopic findings of laryngectomy patients and healthy

1
2
3
4 subjects. No patient reported specific treatments for nasal and/or tracheal complaints. We did
5
6 not observe endoscopic findings of acute or chronic rhinosinusitis. All patients wore a
7
8 protective bib anteriorly to tracheostomy.
9

10 11 12 *Nasal cytology*

13
14
15 Nasal cytological analysis demonstrated a mucous cell metaplasia (Figure 1A) in 20%
16
17 of laryngectomy patients, but it was absent in all healthy subjects ($p < 0.05$); no squamous cell
18
19 metaplasia was found in both groups (Table 3). No statistically significant difference between
20
21 healthy subjects and laryngectomy patients was observed for neutrophilic and eosinophilic
22
23 infiltration ($p > 0.05$). One healthy subject had rare eosinophils, due to allergic perennial
24
25 rhinitis. No mast cell and lymphocyte were found in both groups. In 15 patients (60%)
26
27 bacteria were present, without inflammatory infiltrate (Figure 1B). Figure 1C shows normal
28
29 nasal cytology from a control subject.
30
31

32
33 Nasal cytology did not show atypia. Concerning laryngectomy patients,
34
35 hyperchromatic supranuclear stria was absent in ciliated cells.¹²
36

37
38 Analyzing nasal cytological changes of treated patients, statistical analysis did not
39
40 show significant correlation with symptoms, endoscopic findings (turbinate hypertrophy,
41
42 mucosal hyperemia, nasal secretions), patients' age, tobacco smoking, tumor stage and
43
44 adjuvant radiotherapy ($p > 0.05$).
45
46
47

48 49 *Tracheal cytology*

50
51 Tracheal cytological analysis demonstrated a quite high rate of squamous cell
52
53 metaplasia (24% of patients), neutrophilic infiltrate (32%) and presence of bacteria (40%)
54
55 (Figure 2A-B). Four out of ten patients with bacteria had no inflammatory infiltrate. Mucous
56
57
58
59
60

1
2
3
4 cell metaplasia was observed in 12% of cases. We did not observe tracheal cytological atypia.
5
6 Sample material was insufficient for examination in 20% of cases, due to dry tracheal mucosa
7
8 (Table 4). Statistical analysis did not show significant correlation with symptoms, endoscopic
9
10 findings (mucosal hyperemia, tracheal secretions), patients' age, tobacco smoking, tumor
11
12 stage and adjuvant radiotherapy ($p>0.05$). Insufficient sampling was associated with dry
13
14 tracheal mucosa ($p<0.05$). A normal tracheal cytology in a patient without laryngectomy
15
16 showing ciliated cells was added from an independent bronchoscopy patient whose tracheal
17
18 brushing was banked in the pathology department (Figure 2C).
19

20
21
22 Normal nasal and tracheal cytology was observed in only 4 (16%) and 2 (8%)
23
24 laryngectomy subjects (8%), respectively.
25

26 27 28 **Histology**

29
30
31 Histological examination of inferior turbinate showed submucosal stromal fibrosis in
32
33 all laryngectomy patients and submucosal neutrophilic and eosinophilic infiltrate in one case
34
35 (9%) (Figure 3).
36

37 38 39 **Discussion**

40
41
42 Permanent changes of airflow in nasal cavities and trachea, due to complete separation
43
44 of upper and lower airways in laryngectomy patients, result in loss of physiological nasal
45
46 functions and presence of "unconditioned" inspired air in lower airways. Therefore,
47
48 laryngectomy patients represent a model to study nasal mucosal reaction to inactivity and
49
50 tracheal reaction to unconditioned inspired air. In literature, there is little evidence that
51
52 cessation of nasal airflow after laryngectomy has any harmful effect on nasal cavities. Only
53
54 two studies have evaluated cytological and histological changes on nasal mucosa.^{1,2} In this
55
56
57
58
59
60

1
2
3
4 study we reported cytological alterations of nasal and tracheal mucosa, likely due to airflow
5
6 changes after total laryngectomy.
7

8
9 Experiments on unilateral naris closure in the rabbit demonstrated an increase of
10 goblet cells number and changes of transitional epithelium into ciliated epithelium.^{13,14}
11
12 Human respiratory nasal epithelium after laryngectomy appears to undergo similar changes to
13 those observed in rabbits with unilateral naris closure. An early hypersecretory phase with an
14 increased speed of mucociliary clearance occurs after total laryngectomy, while on the
15 contrary a long term decrease of mucociliary clearance appears.^{3,13,15,16} Early findings could
16
17 be explained in some patients abolishing the harmful effects of cigarette smoke on the nasal
18 epithelium, as many laryngectomy subjects are heavy smokers before surgical treatment. Late
19 alterations may be due to atrophy of nasal mucosa, chronic inflammatory changes and
20 secondary infections by saprophytic bacteria.³ Similar studies concerning lower airways in
21 laryngectomy patients are not present in literature. Moreover, the presence of tracheal mucosa
22 alterations in these subjects has never been studied.
23
24
25
26
27
28
29
30
31
32
33
34

35 Nasal cytology is a safe, not invasive, low-cost method for evaluating mucosal
36 changes. It is already a currently used tool for the diagnosis of allergic and non-allergic
37 rhinitis and other pathological alterations of nasal mucosa. The nasal mucosa of healthy
38 subjects is constituted by four cytotypes (ciliated, goblet, striated and basal), which are
39 arranged in a pseudostratified pattern on an underlying basement membrane. It does not show
40 other cells except, rarely, neutrophils; therefore, the detection of a cell type different from
41 these is a sign of possible pathology.^{11,17}
42
43
44
45
46
47
48
49

50
51 Skoloudik et al. showed that main cytological alterations of nasal mucosa after total
52 laryngectomy were hyperplasia of the basal zone cells and presence of bacteria without any
53 inflammatory changes.¹ Karaca et al. showed a high percentage of histopathologic alterations
54
55
56
57
58
59
60

1
2
3
4 in laryngectomy subjects: destruction of goblet cells, destruction of cilia, destruction of
5
6 submucosal glands and fibrosis in stroma, focal or total atrophy of the mucosa, myxoid
7
8 degeneration, neovascularization and congestion.² Signs of damage, consisting mainly of
9
10 various degree of epithelial degeneration, were also observed for olfactory neuroepithelium,
11
12 associated to structural disorders of Bowman's glands.¹⁸
13
14

15 According to literature, our findings confirmed a high rate of stromal fibrosis in nasal
16
17 submucosa (all laryngectomy patients in our series). On the contrary, we did not see any
18
19 hyperplasia of the basal zone cells at nasal cytological examination and we found an higher
20
21 percentage of laryngectomy patients with mucous cell metaplasia, compared to healthy
22
23 subjects. Moreover, there was not a statistically significant difference for inflammatory
24
25 infiltrate between laryngectomy patients and healthy subjects. No squamous cell metaplasia
26
27 was observed in both groups. Based on these results, we could hypothesise that the absence of
28
29 nasal airflow may induce some cytological and histological long-term changes, not only in the
30
31 mucosa, but also in submucosa, with the tendency to a mucous cell metaplasia in nasal
32
33 mucosa and to a fibrotic status in submucosa. Probably different molecular mechanisms are
34
35 the reason why our findings are divergent from the previously published works on this topic.
36
37
38

39 Different molecular mechanisms could be related to a different individual genetic
40
41 predisposition to fibrosis and mucous cell proliferation, e.g. production of Fibroblast Growth
42
43 Factor (FGF) and/or mucins. Then, further studies on larger samples are necessary to
44
45 overcome bias due to different genetic predisposition and/or to better understand the
46
47 molecular reasons for the findings observed in our and other studies, such as mucous cell
48
49 metaplasia, submucosal fibrosis, and hyperplasia of the basal zone cells.
50
51

52
53 Nasal cytological examination demonstrated the presence of bacteria without
54
55 inflammatory infiltrate, according to previous studies that analyzed the bacterial flora of nasal
56
57
58
59
60

1
2
3
4 cavities in laryngectomy patients. Their nasal cavities and trachea are generally colonized by
5
6 non-pathogenic and/or potentially pathogenic bacteria, without signs and symptoms of
7
8 infection.^{1,8,10} The absence of a statistically significant difference in cytological findings
9
10 between patients who underwent adjuvant radiotherapy and patients without such treatment
11
12 can be explained because the nose was not in the radiation fields.

13
14
15 Bronchoalveolar lavage and bronchial brushing under flexible bronchoscopy are
16
17 generally used for cytological examination of lower respiratory tract mucosa in several
18
19 patients and represent minimally invasive techniques. Since the tracheal mucosa is similar to
20
21 respiratory nasal one, we evaluated the feasibility of scraping with a curette for cytological
22
23 examination of tracheal mucosa in laryngectomy patients, using the same staining method.
24
25 Transient cough during procedure was observed in only five patients (20%) and was the
26
27 unique slight complaint reported by patients. No bleeding due to procedure occurred. May-
28
29 Grünwald-Giemsa staining method allowed a good identification of all cell types, epithelial
30
31 and inflammatory ones. Therefore, tracheal scraping could be considered a safe, not invasive,
32
33 low-cost method to assess alterations of tracheal mucosa in laryngectomy subjects. However,
34
35 it is important to take into account that sampling material could be insufficient in 20% of
36
37 cases, because of dry tracheal mucosa.
38
39
40

41
42 Tracheal cytological examination showed squamous cell metaplasia in 24% of cases,
43
44 and mucous cell metaplasia in 12% of patients. Possible causes of these alterations could be
45
46 the passage of unconditioned air with exposure to irritants and cold temperature, radiotherapy
47
48 effects on tracheal mucosa, and smoke. However, our statistical analysis did not show
49
50 significant correlation with tobacco smoking and adjuvant radiotherapy. Similarly to nasal
51
52 cavities and according to previous studies,^{7,9,10,19} trachea was colonized by bacteria in 40% of
53
54 cases without clinical signs of infection. Only a few cases recognized the presence of bacteria
55
56
57
58
59
60

1
2
3
4 as a cause of neutrophilic inflammation; nevertheless, these patients had not endoscopic
5
6 findings of tracheitis. Further studies on larger samples are necessary to investigate reasons of
7
8 cytological tracheal alterations in these patients, assessing the role of airflow and bacterial
9
10 colonization. Moreover, stability or change of nasal and tracheal cytology over time after total
11
12 laryngectomy should be analyzed in future studies in order to better understand how airflow
13
14 alterations interacts with upper end lower respiratory tract in laryngectomy patients.
15
16

17
18 In conclusion, the present study demonstrated that nasal cavities and trachea of
19
20 laryngectomy patients undergo long-term cytological and histological changes of mucosa and
21
22 submucosa, probably due to modifications of airflow. These changes must be kept in mind
23
24 when a laryngectomy patient is evaluated for nasal or tracheal complaints. Possible roles of
25
26 tracheal cytology in laryngectomy patients in clinical daily practice should be assessed with
27
28 further studies on larger samples. Our data set is small and has current little clinical relevance,
29
30 but it should be a foundation for future works. In future, detailed knowledge of cytological
31
32 changes in patients' tracheal mucosa could represent a key prerequisite for the choice of
33
34 effective interventions for tracheal pathology, such as saline solution aerosol, antibiotics,
35
36 corticosteroids and/or hyaluronic acid. Further studies are necessary in order to investigate
37
38 whether different treatments for tracheal complaints in patients who underwent total
39
40 laryngectomy may be effective for different cytological findings.
41
42
43
44
45

46 **Acknowledgments**

47
48 The authors declare that they have no Financial Support/Funding.
49
50
51
52
53
54
55
56
57
58
59
60

References

1. Skoloudik L, Vokurra J, Zborayova K, Celakovsky P, Kucera M, Ryska A. Cytology of the nasal mucosa after total laryngectomy. *Acta Oto-Laryngologica* 2009;129:1262-1265.
2. Karaca CT, Gültekin E, Yelken MK, Igdem AA, Külekçi M. Long-term histologic changes in nasal mucosa after total laryngectomy. *Int J Otolaryngol* 2010;137128. doi: 10.1155/2010/137128. Epub 2010 Apr 6.
3. Deniz M, Uslu C, Ogredik EA, Akduman D, Gursan SO. Nasal mucociliary clearance in total laryngectomized patients. *Eur Arch Otorhinolaryngol* 2006;263:1099-1104.
4. Caldas AS, Facundes VL, Melo TM, Dourado Filho MG, Pinheiro Júnior PF, Silva HJ. Modifications and evaluation of smell and taste functions in total laryngectomy: systematic review. *J Soc Bras Fonoaudiol* 2011;23(1):82-88.
5. Boyce J, Eccles R. Do chronic changes in nasal airflow have any physiological or pathological effect on the nose and the paranasal sinuses? A systematic review. *Clin Otolaryngol* 2006;31(1):15-19.
6. Kelsey MC, Mitchell CA, Griffin M, Spencer RC, Emmerson AM. Prevalence of lower respiratory tract infections in hospitalized patients in the United Kingdom and Eire. Results from the Second National Prevalence Survey. *J Hosp Infect* 2000;46:12-22.
7. Harlid R, Andersson G, Frostell CG, Jörbeck HJ, Ortqvist AB. Respiratory tract colonization and infection in patients with chronic tracheostomy. *Am J Respir Crit Care Med* 1996;154:124-129.

- 1
- 2
- 3
- 4 8. Cvetnic V, Skrlin J, Rak I. Bacterial flora of the nasal cavity in
- 5 laryngectomized patients. *Infection* 1996;24(1):26-28.
- 6
- 7
- 8 9. Kramp B, Donat M, Dommerich S, Pau HW, Podbielski A. Prospective
- 9 controlled study of microbial colonization of the trachea in tracheotomized and
- 10 laryngectomized patients with HME (heat and moisture exchanger). *Acta Oto-*
- 11 *Laryngologica* 2009;129:1136-1144.
- 12
- 13 10. Riva G, Garzaro M, Zaccaria T, et al. Nasal and tracheal microbial colonization
- 14 in laryngectomized patients. *Ann Otol Rhinol Laryngol*. 2015 Nov 3. pii:
- 15 0003489415613802 [Epub ahead of print] PMID: 26530093
- 16
- 17 11. Gelardi M. Atlas of nasal cytology for the differential diagnosis of nasal
- 18 diseases. 2nd Ed. Milano: Edi-Ermes, 2012; 1-208.
- 19
- 20 12. Gelardi M, Cassano P, Cassano M, Fiorella ML. Nasal cytology: description of
- 21 hyperchromatic supranuclear stria as a possible marker for the anatomical and
- 22 functional integrity of the ciliated cell. *Am J Rhinol* 2003;5:263-268.
- 23
- 24 13. Maurizi M, Paludetti G, Almadori G, et al. Mucociliary clearance and mucosal
- 25 surface characteristics before and after total laryngectomy. *Acta Otolaryngol*
- 26 1986;102:136-145.
- 27
- 28 14. Meseguer HD, Galvez ME, Ortega GF, Arias RM. The mucociliary function
- 29 and the morphology of the nasal mucous membrane in laryngectomees
- 30 compared with normal subjects. *An Otorrinolaringol Ibero Am* 1996;23:81-90.
- 31
- 32 15. Moore-Gillon V. The nose after laryngectomy. *J R Soc Med* 1985;78:435-439.
- 33
- 34 16. Dixon FW, Hoerr NL, McCall JW. The nasal mucosa in laryngectomized
- 35 patients. *Ann Otol Rhinol Laryngol* 1949;58:535-547.
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60

- 1
2
3
4 17. Riva G, Boita M, Ravera M, et al. Nasal cytological changes as late effects of
5
6 radiotherapy for nasopharyngeal cancer. *Am J Rhinol Allergy* 2015;29(2):41-5.
7
8
9 18. Miani C, Ortolani F, Bergamin Bracale AM, Petrelli L, Staffieri A, Marchini
10
11 M. Olfactory mucosa histological findings in laryngectomees. *Eur Arch*
12
13 *Otorhinolaryngol* 2003;260:529-535.
14
15
16 19. Cabello H, Torres A, Celis R, et al. Bacterial colonization of distal airways in
17
18 healthy subjects and chronic lung disease in a bronchoscopic study. *Eur Respir*
19
20 *J* 1997;10:1137-1144
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 **Tables**
5
6
7

8 **Table 1.** Patients and tumor characteristics
9

Characteristic	Number of subjects (%)	
	Laryngectomy group (25 subjects)	Control group (25 subjects)
Sex		
Male	22 (88)	19 (76)
Female	3 (12)	6 (24)
Smoker or former smoker		
Yes	22 (88)	18 (72)
No	3 (12)	7 (28)
Previous alcohol consumption		
Yes	21 (84)	20 (80)
No	4 (16)	5 (20)
Previous exposure to irritant substances		
Yes	1 (4)	0 (0)
No	24 (96)	25 (100)
Allergies		
Yes	4 (16)	6 (24)
No	21 (84)	19 (76)
Tumor site		
Supraglottic larynx		7 (28)
Glottic larynx	17 (68)	-
Subglottic larynx	1 (4)	-
Histological type		
Squamous cell carcinoma	25 (100)	-
Tumor (pTNM VI ed.)		
T1	0 (0)	-
T2	3 (12)	-
T3	16 (64)	-
T4	6 (24)	-
Nodes (pTNM VI ed.)		
N0	16 (64)	-
N1	4 (16)	-
N2	5 (20)	-
N3	0 (0)	-
Distant Metastasis (pTNM VI ed.)		
M0	25 (100)	-
M1	0 (0)	-
Grade		
G1	1 (4)	-
G2	7 (28)	-
G3	17 (68)	-
Stage		
I	0 (0)	-
II	1 (4)	-
III	13 (52)	-
IV	11 (44)	-

55
56
57
58
59
60

Table 2. Symptoms and endoscopic findings

Characteristic	Number of subjects (%)		p value
	Laryngectomy group (25 subjects)	Control group (25 subjects)	
Symptoms			
Rhinorrhea			
Anterior	14 (56)	5 (20)	0.02
Posterior	2 (8)	1 (4)	0.26
Hyposmia			
Yes	25 (100)	2 (8)	0.03
No	0 (0)	23 (92)	
Cough			
Dry	3 (12)	3 (12)	0.03
Productive	9 (36)	1 (4)	
No	13 (52)	21 (84)	
Endoscopic findings - Nasal cavity			
Nasal septal deviation			
Yes	17 (68)	15 (60)	0.82
No	8 (32)	10 (40)	
Turbinates			
Hypertrophic	10 (40)	7 (28)	0.33
Normotrophic	14 (56)	18 (72)	
Atrophic	1 (4)	0 (0)	
Nasal mucosa			
Pale	10 (40)	8 (32)	0.67
Pink	15 (60)	17 (68)	
Hyperemic	0 (0)	0 (0)	
Secretions			
Dry nose	6 (24)	1 (4)	0.24
Serous	17 (68)	22 (88)	
Thick	2 (8)	2 (8)	
Purulent	0 (0)	0 (0)	
Polypoid degeneration of mucosa			
Yes	1 (4)	0 (0)	0.85
No	24 (96)	25 (100)	
Endoscopic findings - Trachea			
Tracheal mucosa			
Pale	2 (8)	-	
Pink	22 (8)	-	
Hyperemic	1 (4)	-	
Secretions			
Dry trachea	16 (64)	-	
Serous	4 (16)	-	
Thick	5 (20)	-	
Purulent	0 (0)	-	

Table 3. Nasal cytology findings.

Characteristics	Number of subjects (%)		p value
	Laryngectomy group (25 subjects)	Control group (25 subjects)	
<i>Normal nasal cytology</i>	4 (16)	21 (84)	0.02
<i>Squamous cell metaplasia</i>	0 (0)	0 (0)	0.99
<i>Mucous cell metaplasia</i>	5 (20)	0 (0)	0.03
<i>Neutrophilic infiltration</i>	2 (8)	3 (8)	0.95
<i>Eosinophilic infiltration</i>	0 (0)	1 (4)	0.64
<i>Presence of bacteria</i>	15 (60)	0 (0)	0.02

Table 4. Tracheal cytology findings.

Characteristics	Number of patients (%) (25 subjects)
<i>Normal tracheal cytology</i>	2 (8)
<i>Squamous cell metaplasia</i>	6 (24)
<i>Mucous cell metaplasia</i>	4 (12)
<i>Neutrophilic infiltration</i>	8 (32)
<i>Eosinophilic infiltration</i>	0 (0)
<i>Presence of bacteria</i>	10 (40)
<i>Insufficient sampling</i>	5 (20)

Figure legends

Figure 1. Nasal cytology (May-Grünwald-Giemsa staining): (A) mucous cell metaplasia (black arrows) in laryngectomy patients (magnification x400); (B) presence of bacteria (white arrows) without inflammatory infiltrate in laryngectomy patients (magnification x1000); (C) normal ciliated cells in control patients (black stars), without mucous cells, inflammatory cells and bacteria (magnification x400).

Figure 2. Tracheal cytology (May-Grünwald-Giemsa staining): (A-B) squamous cell metaplasia (black arrows), neutrophilic infiltrate (white stars) and presence of bacteria (white arrows) in laryngectomy patients (magnification x1000); (C) tracheal cytology in a patient without laryngectomy showing normal ciliated cells (black stars) (magnification x400).

Figure 3. Histological examination of inferior turbinate (Hematoxylin-Eosin staining, magnification x100): submucosal stromal fibrosis (black arrows) in laryngectomy patients.




Figure 1. Nasal cytology (May-Grünwald-Giemsa staining): (A) mucous cell metaplasia (black arrows) in laryngectomy patients (magnification x400); (B) presence of bacteria (white arrows) without inflammatory infiltrate in laryngectomy patients (magnification x1000); (C) normal ciliated cells in control patients (black stars), without mucous cells, inflammatory cells and bacteria (magnification x400).

321x705mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 2. Tracheal cytology (May-Grünwald-Giemsa staining): (A-B) squamous cell metaplasia (black arrows), neutrophilic infiltrate (white stars) and presence of bacteria (white arrows) in laryngectomy patients (magnification x1000); (C) tracheal cytology in a patient without laryngectomy showing normal ciliated cells (black stars) (magnification x400).

329x736mm (300 x 300 DPI)




Figure 3. Histological examination of inferior turbinate (Hematoxylin-Eosin staining, magnification x100): submucosal stromal fibrosis (black arrows) in laryngectomy patients.

485x366mm (72 x 72 DPI)