IRIS AperTO

UNIVERSITÀ DEGLI STUDI DI TORINO

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Development of old-growth characteristics in uneven-aged forests of the Italian Alps

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available http://hdl.handle.net/2318/149666 since 2022-04-04T21:28:48Z

Published version:

DOI:10.1007/s10342-014-0830-6

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

12 October 2024

Renzo Motta[1*], Matteo Garbarino[2], Roberta Berretti[1], Fabio Meloni[1], Antonio Nosenzo[1] and Giorgio Vacchiano[1]

Development of old-growth characteristics in uneven-aged forests of the Italian Alps

[1] *Dipartimento di Scienze Agrarie, forestali e alimentari (DISAFA), Università di Torino, Grugliasco (TO), 10095, Italy*

[2] *Dipartimento D3A, Università Politecnica delle Marche, Ancona (AN), 60131, Italy*

*Corresponding author: Tel: +39 011 670 5538; Fax: +39 011 670 5556; Email: renzo.motta@unito.it

Abstract

During the last millennia, all forests of the Italian Alps have been heavily affected by human land-use. Consequently, forest structures have been modified and there are no old-growth remains. In the last decades, however, many forests have been withdrawn from regular management, because wood production was unprofitable, and left to develop naturally. At the same time, in currently managed forests, silvicultural systems able to develop or maintain old-growth characteristics are being required. The aim of this paper is to assess the status and developmental dynamics of old-growth characteristics in mixed beech, silver fir, and Norway spruce montane forests of the eastern Italian Alps.

We selected along a naturalness gradient (a) three old-growth forests in Bosnia and Montenegro (due to the lack of old-growth forests in the Italian Alps), (b) two forests withdrawn from regular management for at least 50 years, and (c) three currently managed forests. In each forest we analyzed 17 structural attributes, in order to assess their value as indicators of old-growth condition. Old-growth forests were characterized by significantly higher amounts of live and dead biomass, share of beech in the dominant and regeneration layers, and number of large trees. The diameter distribution was best described as a rotated sigmoid, differently from currently and formerly managed forest. We discuss the differences in old-growth characteristics across the management gradient, and use our results to evaluate the effectiveness of retention prescriptions currently applied in the studied regions in maintaining or promoting old-growth structural attributes in managed forests.

Keywords: forest structure, coarse woody debris, selection system, rotated sigmoid, PCA, retention forestry

Introduction

Old-growth forests are "later stages in forest development that are often compositionally and always structurally distinct from earlier successional stages" (Franklin and Spies 1991). Old-growth forests host plant, fungi and animals that are rare or absent in earlier developmental stages, and are important for regional biodiversity (Bauhus et al. 2009; Burrascano et al. 2013; Keeton 2006; Keeton and Franklin 2005). Moreover, they represent an important reference point for evaluating human impacts on forest ecosystems, and for developing silvicultural systems able to emulate natural processes and fulfil socio-economic goals while maintaining a full range of ecosystem services (Wirth et al. 2009). As the definition implies, old-growth stages can occur also in managed forests that have developed for long periods without human influences and without stand-replacing natural disturbances. The time required for the development of old-growth characteristics in previously managed forests may vary from a few decades to a few centuries, depending on site features (e.g., disturbance regime and history, productivity), and tree species functional traits (e.g. growth rate, longevity, wood decay rate, etc.) (Humphrey 2005; Motta et al. 2010).

Starting from the 19th century, silvicultural treatments in central and southern Europe were focused mainly on timber production. "Industrial" production forestry was based mainly on pure, even-aged stands subjected to short rotation periods, i.e., 10-40% of the potential life-span of the main tree species (Bauhus et al. 2009). Only a small part of the forests was subjected to systems based on mixed uneven-aged stands but, also in these systems, wood production was the main purpose. As a consequence, structural attributes typical of later developmental stages are strongly under-represented and, if present, restricted to small and remote forest reserves (Barbati et al. 2012; Peterken 1996).

In the last decades a gradual reversal of this trend has been observed, with a decrease in timber removal, and many forests have been withdrawn from management (Castagneri et al. 2010;

Motta et al. 2010; Vandekerkhove et al. 2009). At the same time, the scientific community has been advocating for forest management strategies that aim at the restoration of old-growth attributes that would benefit a wealth of taxa that are lacking or threatened in managed forests, even where close-to-nature forestry is applied (Lindenmayer and Franklin 2002; Lindenmayer et al. 2006). Policy frameworks have been set up to facilitate the attainment of this goal (e.g., the Helsinki process) and have in some case led to measurable results (Ammer 1991; Kohler 2010; Vandekerkhove et al. 2011).

Despite being an apparent oxymoron, the development of "old-growthness", i.e., the degree to which structural and functional attributes are similar to those associated to old-growth forests (Bauhus et al. 2009) can be facilitated by appropriate forest management measures. These can be described as either ecological restoration practices (Stanturf and Madsen 2002) e.g., extending thinning rotation periods, regulating stem density and regeneration to match wildlife habitat requirements and mimic the effects of natural disturbances (Bergeron et al. 2007; Silver et al. 2013) or retention forestry (Beese et al. 2003; Gustafsson et al. 2010).

Gustafsson et al. (2012) defined retention forestry as "an approach to forest management based on the long-term retention of structures and organisms, such as live and dead trees and small areas of intact forests, at the time of harvest". Retention forestry aims at improving elements of old-growthness pertaining to stand structure, i.e., the quantity and quality of coarse woody debris (CWD), large old trees, and the amount of structural diversity. These elements are not exhaustive of old-growth characters, but provide surrogates for other ecosystem attributes, such as plant or animal diversity (Halpern and Spies 1995; Heilmann-Clausen and Christensen 2003; Kirby and Drake 1993; Lähde et al. 1999; Ódor et al. 2006; Siitonen 2001) with the advantage of being directly manipulated during forest management.

The aim of this paper was to identify the structural attributes that contribute to better differentiate between old-growth forests, actively managed forests, and forests withdrawn from

regular management (FWRM) in mixed beech (*Fagus sylvatica* L.), silver fir (*Abies alba* Mill.) and Norway spruce (*Picea abies* (L.) Karst.) stands of the Alps and Dinaric mountain ranges. In light of our results, we discuss the effectiveness of current retention forestry aimed at promoting old-growthness in managed stands of the Italian Alps.

Material and methods

Study sites

We focused our research on mixed, uneven-aged mountain forests belonging to the association *Piceo-Abieti-Fagetum*. These forests exhibit continuous canopy cover and multilayered vertical structure. In the European temperate forests, old-growth remnants are still present in remote protected areas of Eastern (Parviainen 2005) and South-eastern Europe (Korpel 1995; Nagel et al. 2012). They are highly significant for European silviculture, not only because of their natural value, but also because their structure served as a model to develop the silvicultural selection system (Korpel 1995; Leibundgut 1982; Susmel 1980).

This study was carried out in eight mixed montane forests of the Italian Alps and of the Dinaric mountains, co-dominated by silver fir, beech and Norway spruce (Fig. 1; Tab. 1). We selected three actively managed forests and two FWRM in the eastern Italian Alps. Ordinary silvicultural management is carried out by harvesting single-tree to small-group maintaining a multilayered and unevenaged stand, and has a limited impact on the landscape. Selection cuttings are repeated every 8-10 years and the main goal of the management is the conservation of the desired diameter distribution and species composition. Santo Stefano di Cadore has been managed by single tree selection since the age of the Republic of Venice (Volin and Buongiorno 1996). In Croviana and Amblar, application of single tree selection is more recent (Wolynski 1998). The two FWRM Val Navarza and Ludrin have developed without direct human influence since 1953

and 1962, respectively (Castagneri et al. 2010). Before the abandonment, these forests were managed using a silvicultural selection system but due to their remoteness the cuts were probably less frequent but more intense than in stands closer to a road system. Due to the absence of old-growth forests in the Italian Alps (Motta 2002) we selected three old-growth forests in Bosnia-Herzegovina and Montenegro (Motta et al. In press; Motta et al. 2011; Nagel and Svoboda 2008) about 400 km south-east of the Italian border. Forest cover type, site conditions, and altitudinal range of old-growth forests are similar to the remaining five.

Field sampling

In each forest, located within a narrow elevation range (1080 – 1570 m a.s.l.), we selected a sector of 5 – 40 ha (Tab.1), chosen with the help of local foresters in compliance to the following criteria: (a) mixed species composition, i.e., no species with > 70% contribution to total basal area; (b) for old-growth forests, core area or sector of core area in the optimum stage (Korpel 1995); (c) for managed forests, management unit(s) in the vicinity of the forest road system; and (d) for FWRM, unmanaged for the longest possible period according to available management plans.

In each forest we surveyed the structure by a regular quadrat grid. Grid size varied from 70 to 120 m based on the size of the stand, resulting in one sampling point every 0.4 – 1.2 ha of forest. At each sampling point, three types of measurements were carried out (Motta et al. 2011): (1) species, diameter at breast height (dbh) to the nearest 0.01 m, and height (h) to the nearest 0.5 m of all living trees (dbh ≥ 7.5 cm) within a 615.5 m^2 circular plot (radius = 14 m); (2) species of each regeneration tree (h > 10 cm and dbh < 7.5 cm, i.e., seedlings and saplings) within a concentric, 113.1 m^2 circular plot (radius = 6 m); (3) the diameter of each log crossing a 50 m line (Motta et al. 2006) oriented northward from the centre of each sampling point (orientation bias was assumed to be negligible); and (4) base and top diameter of stumps (height <130 cm),

142 and dbh of snags (height >130 cm) in a 50 x 8 m rectangular plot centred on the previous line.
143 In each site we have measured the height of the overtopping trees in order to estimate the
144 Maximum height (the average of the 5 highest tree ha^{-1}) sensu Susmel (1980).
145
146
147 *Statistical analyses*
148 Volume of living trees, dead standing trees and snags was calculated using local yield tables.
149 Volume of logs was calculated according to line intercept sampling method and the volume of
150 stumps was estimated as a frustum of a cone using diameter at the top and diameter at the ground
151 level (Motta et al. 2006).
152 In each plot we computed 17 structural attributes for living trees and CWD (Tab. 2). We
153 statistically compared structural attributes between management categories (managed, FWRM,
154 old growth) by means of a non-parametric Kruskal-Wallis test, followed by Dunn's post-hoc
155 multiple comparison. In order to identify key gradients of structural variation under different
156 management categories, we carried out a Principal Component Analysis (PCA) on stand
157 structural attributes. Highly correlated attributes (Pearson's R >0.75) were excluded from further
158 analysis (Tab. 2). The statistical significance of PCA axes was tested by random average under
159 permutation, i.e., the average eigenvalue obtained under a randomization of the data matrix,
160 acquired from a randomization test based on eigenvalues. The randomization test was carried out
161 as follows: (1) randomize the values within variables in the data matrix; (2) conduct a PCA on
162 the reshuffled data matrix; and (3) repeat steps 1 and 2 a total of 10000 times. The *P*-value for
163 each axis and each test statistic was then estimated as: (number of random values equal to or
164 larger than the observed +1) / 1000. If the observed exceeds the average random value, that
165 particular axis is considered to be non-trivial (Peres-Neto et al. 2005). PCA was performed using
166 the function PRCOMP of the R statistical package, version 3.0 (R Core Team 2013).

In order to classify and compare dbh distributions, we computed a pooled frequency distribution of tree diameters (midpoints of 5 cm dbh classes) by forest management categories (MAN, FWRM and OG), using data from all living trees and plots (Goodburn and Lorimer 1999). Following $\log_{10}$ transformation, we fitted each frequency distribution by concurrent models based on linear combinations of dbh, dbh^2 and dbh^3, in order to define its shape (Janowiak et al. 2008; Leak 1996). Because the logarithm of zero is undefined, we added 1.0 to the frequency of all dbh classes before log transformation. Among significant models, we selected the model with the highest adjusted R^2 and lowest root mean square error (RMSE) as the best fitting one. The shape of each diameter distribution was determined by examining the significance and sign of model parameters (Alessandrini et al. 2011; Janowiak et al. 2008).

Results

Old-growth exhibited significantly higher quadratic mean diameter (QMD), volume of living trees (VOL) (763 – 1031 m^3 ha^{-1} on average) and volume of coarse woody debris (CWD) (327 – 420 m^3 ha^{-1}), density of trees with dbh>50 and dbh>80 (D50 and D80), volume of snags with dbh >50 cm (VSNAG50) and volume of logs with diameter > 50 cm (VLOG50) (Tab. 2, Fig. 2). FWRM had the highest tree densities (TPH) and lowest standard deviation of tree diameters (SDD), together with managed forests. The latter had the highest frequency of stumps (STUMP), and lowest basal area (BA), CWD, percent basal area by beech (PERCFASY), percent regeneration by beech (REGFASY), and Shannon diversity index for dbh (HD).

PCA of uncorrelated attributes evidenced two significant principal components ($p < 0.001$, Monte Carlo test), that accounted for a cumulative 51,9 % of total variation (Tab. 3; Fig. 3). The type of management gradient was aligned along a direction from managed to old-growth forests of increasing deadwood (VSAG50, VLOG50), increasing incidence of beech (PERCFASY,

REGENFASY), density of large tree (D80), and decreasing anthropogenic disturbance (STUMP, VSTUMP), and was relatively orthogonal to HD, QMD and BA .

The log_{10} diameter cumulative distribution of old-growth forests (Fig. 4) was best described as a rotated sigmoid ($p < 0.05$), while it exhibited a concave and a variable shape in managed forests and FWRM, respectively (Tab. 4).

The maximum height was between 37-45 m for managed forests, 40-41 m for FWRM and between 46 and 55 m in the old-growth forests.

Discussion

Forest structure

Structural attributes were effective in discriminating managed and old-growth mixed mountain forests of central-southern Europe, consistently with previous research (Boncina 2000; Franklin and Van Pelt 2004; Kuuluvainen et al. 1996; Siitonen et al. 2000). The most significant differences arose from the amount and quality of large woody debris (VSNAG50, VLOG50), and the frequency of very large trees (D80), i.e., variables directly influenced by management intensity.

Other indicators associated to the management gradient were maximum height and species composition. The difference in maximum height between managed/FWRM and old-growth forests, i.e., about 5 m, is probably due to the fact that management prevents the trees from a living their complete life-span (Susmel 1980). Species composition, on the other hand, differed between FWRM/old growth and managed forests, especially due to a significantly lower presence of beech in the latter. Even if all stands belonged to the *Abieti-picenion* Br.-Bl. 1939 (group F5: mixed beech forests according to the map of potential natural vegetation in Europe) (Bohn et al. 2003), past management in the Alps resulted in a negative selection on beech, due to a preference for spruce and fir that had a more valuable timber (Poldini and Bressan 2007). This

is in contrast to the observed current trends observed in most old-growth forests of central-eastern Europe where there is long-term quantitative increment of beech and a quantitative and qualitative reduction of the conifers (Diaci et al. 2011; Keren et al. 2014). In forests left unmanaged for some decades, beech has started to increase its share in the lower diameter classes, even if large individuals are still lacking compared to old-growth forests (Fig. 5). A similar trend occurred in the regeneration layer (regeneration density, Tab. 2). Current silvicultural management protocols have abandoned negative selection on beech, but the low number of seed trees, as a legacy of past management, implies that these stands still experience a scarcity of beech in the intermediate and regeneration layers as well. It is possible, however, that the quantity and quality of coarse woody debris would differ between beech-rich (old-growth) and beech-devoid forests (FWRM and managed), due to the highest production of large branches and fastest decay rate of the broadleaf species (Chirici et al. 2014) as opposed to the conifers.

Old-growth forests were represented as a distinct group on the ordination plot, that expressed gradients of large tree density, biomass, specific composition, quantity of deadwood (logs and snags) and incidence of beech. The scattered distribution of old-growth sampling points on the ordination plot revealed a high heterogeneity of forest structure, as opposed to the smaller variability of structural attributes in formerly and currently managed forests.

On the other hand, there was a wide overlap of the structural data coming from managed forests and FWRM. This is mainly due to the relatively short period that FWRM have spent without direct human influence, about 50 and 60 years respectively. Mature managed stands are expected to take about one century to reach amounts of deadwood and large living trees about half of those found in old-growth (Christensen et al. 2005; Nilsson et al. 2002; Vandekerkhove et al. 2009). FWRM differed from managed forests mainly for increased tree density and CWD volume (Sitzia et al. 2012). The amount of CWD in FWRM was in line with what is expected in a forest that was left unmanaged for about 50 years (Meyer and Schmidt 2011; Motta et al. 2006;

241 Vandekerkhove et al. 2009), and twice as much as in managed forests, even though CWD in the
242 studied managed forests was already much higher than the Italian average for similar forests, i.e.,
243 7 m^3 ha^{-1} for beech, 16 m^3 ha^{-1} for spruce, and 21 m^3 ha^{-1} for silver fir-dominated forests (INFC
244 2005). At the current developmental stage, mortality in FRWM was mainly concentrated in
245 small and intermediate diameters, and only sporadically affected the dominant layer.
246 Consequently, the total amount of CWD and the density of large snags and logs were still very
247 low if compared with old-growth forests (for FWRM and old-growth 70 – 85 vs 327 – 420 m^3
248 ha^{-1} of CWD and 1 – 2 vs 19 - 28 snags >50 cm dbh ha^{-1}). However, CWD in both managed and
249 abandoned forests in this study are higher than 20 – 30 m^3 ha^{-1}, i.e. the proposed threshold to
250 safeguard the complete spectrum of species that rely on deadwood (Angelstam et al. 2003;
251 Müller and Bütler 2010). The density of very large trees (dbh>80 cm) was higher in the managed
252 forests than in FWRM (0-10 vs 0-3 ha^{-1}). This is due both to recent retention forestry policies in
253 managed forests and to the fact that before the abandonment, due to their remoteness, the last
254 cuts were probably in between from a selection and a high-grading system. In any case the
255 density of very large trees of both managed and FWRM was very low if compared with old-
256 growth forests (18-37 ha^{-1})
257 Another important difference between the three management categories was the shape of the
258 diameter distributions (Fig. 4; Tab. 4). Traditionally, the diameter distribution of both managed
259 and unmanaged uneven-aged forests has been represented as reverse J-shaped, or negative
260 exponential (de Liocourt 1898; O'Hara and Gersonde 2004). Recent research has revealed that,
261 for primary forests, it can be better modeled by a rotated sigmoid (Alessandrini et al. 2011; Goff
262 and West 1975; Janowiak et al. 2008). Our results confirmed the latter hypothesis, but not the
263 former, as the log-transformed diameter distribution of managed forests was found to fit a
264 concave function. The concave shape is the result of a deficit in intermediate size classes, with
265 the addition of which it would be classify as a rotated sigmoid. Emergence of a concave

distribution in our managed forests may be explained by retention forestry prescriptions, albeit very recent, that aimed at the conservation of cavity-nesting birds by increasing the number of large diameter trees. Similar studies have pointed out that the concave distribution is evident at small spatial scales (<0.8 ha) (Janowiak et al. 2008) as a result of small gap dynamics or, in our case, to single tree selection practices. Old-growth stands are characterized by an ongoing gap-phase dynamics (Bottero et al. 2011; Kucbel et al. 2010) that can explain the coexistence of multiple structural stages in the same forest (hence the large scatter of OG points on Fig. 3). Conversely, in forests currently or formerly managed for timber, natural disturbances have been replaced by anthropogenic intervention, which may have led to a much greater structural homogeneity (Attiwill 1994) and to a different shape of the diameter distribution. Besides the studied structural parameters the tree spatial pattern, analysed in large permanent plots (Lingua et al. 2011) is another important indicator of old-growthness, differentiating more or less random structures resulting from active management, from a gradient of clumping and subsequent mortality-induced regularization from ongoing competition in natural gaps typical of unmanaged and old-growth forests (Moeur 1997).

Finally, the variable shape of FWRM diameter distribution may be a consequence of the structural re-organisation that forests managed for a long period of time experience after the abandonment.

Retention forestry policies

Notwithstanding some concerns about costs, unwanted negative impacts, and social acceptability, foresters and stakeholders have recently increased their awareness about the importance of conserving biodiversity in regular forest management activities and it has been common practice to leave dead and living trees for biodiversity purposes (Gustafsson et al. 2012; Larrieu et al. 2012).

All the managed forests were, from a silvicultural point of view, overstocked. The increment maximization and long-term structural stability in mixed spruce-fir-beech forests should be attained at around 350 – 400 m^3 ha^{-1} (Schütz 1996), while the observed volumes (453 – 622 m^3 ha^{-1}) were much higher. The management regime in selection forests does not depend only on theoretical approaches but it is strongly influenced by social and economical conditions.

Like most aspects of forest legislation in Italy, the strictness of retention rules in the alpine region varies on a regional/autonomous province basis, and whether or not forests are within a Natura 2000 protected area (Tab. 5). The rules are stricter in regions, like Piedmont or Lombardy, where most traditional silvicultural systems are coppices based on a rotation of 20-30 years. The range of required retention is wider, but at the same time the rules are not so strict, in regions rich in high forests, where there is a long tradition of close-to-nature and sustainable forest management.

However the quantity and quality of retention of old-growth elements that would be adequate for conservation purposes is not known. On one hand, current prescriptions require the retention of minimal densities (0 – 5 ha^{-1}) of ageing or standing dead trees (Tab. 5). This figure is comparable to what is being prescribed in actively managed forests of western and central Europe (e.g., Flanders: 4% of total stock; Baden-Wurttenberg: 15 trees every 3 ha) (Spielmann et al. 2013; Vandekerkhove 2013) but far below the levels detected by this study in unmanaged and old-growth forest, and lower than what is recommended for the conservation of specific organisms such as woodpeckers (Bütler 2003).

Moreover, the prescriptions ignore key components of habitat and ecosystem functioning, such as structural diversity or deadwood on the ground and it is not known what quantity and quality of retention of old-growth elements would be adequate for the set conservation purposes, i.e., the habitat requirements of plant and animal species to be protected at each site are often unclear (Paillet et al. 2010)

Finally, retention strategies at the tree or stand scale should be supported by prescriptions on a larger scale, such as restoration of less natural forest ecosystems and establishment of set-aside areas (forest reserves), aiming at restoring landscape connectivity and building an effective biodiversity network (Bengtsson et al. 2000; Fahrig 2003; Lindenmayer and Franklin 2002).

Conclusions

Both managed and FWRM mixed spruce-fir-beech forests of the Italian Alps have currently few old-growth structural elements. An increase of such elements is desirable in order to improve ecosystem functioning and biodiversity conservation.

Live biomass and large-sized living trees, CWD biomass in different decay stages, large snags, and basal area of beech were the most important variables discriminating the three management categories. This result is aligned to the current body of literature, but nonetheless contains valuable information for future studies on indicators of biodiversity.

The establishment of retention forestry measures is relatively recent, but despite their limited scope and strictness, some effects may be achieved relatively quickly, as we observed in forests where management was abandoned in the last decades, i.e., an increment of large trees and CWD, and a potential shift from negative exponential to concave diameter distribution. On the other hand, 50 years of withdrawal from management are obviously not enough to approximate the degree of old-growthness found in pristine reference forests. According to Nilsson et al. (2002), in boreal forests 100 year without management may be enough to hide the traces of past cutting (even if the structure of the forests may be still influenced by the past human disturbances). This is not the case in the studied mixed forests of the Alps, where the development of old-growth structural characteristics is very slow, and where past management has probably introduced a temporal lag in this process. At the same time, our findings highlight additional stand attributes to be targeted by future retention forestry policies. While in the past the traditional selection

management was oriented mainly towards maintaining the targeted structure and composition, our results evidenced that tree and CWD retention and the CWD decay processes are important for the restoration of the biological diversity. Monitoring and research need to be enhanced in order to modify existing silvicultural measures based on this evidence, and assess their effect through time.

Funding

This work was partially supported by the Autonomous Province of Trento and by the Friuli Venezia Giulia Autonomous Region (Research on forests withdrawn from regular managements). Spot Image Inc., through Planet Action Program domain "Forest and deforestation," provided funding for the research in the Bosnia-Herzegovina and Montenegro old-growth forests by donating satellite imagery and image processing systems.

Acknowledgements

The authors wish to thank Alessandro Wolynski (Autonomous Province of Trento), Franca de Ferrari (Piemonte Region), Damiano Penco and Luca Iacopi (Liguria Region), Maurizio Dissegna, Giovanni Carraro and Ivan Davide Orlando (Veneto Region), Enrico Calvo (ERSAF, Lombardy Region) for useful information and comments regarding retention forestry, Giuseppe Menegus (Regione Veneto), Fabio Angeli, Paolo Zorer and Maurizio Mezzanotte (Autonomous Province of Trento) for useful information and for the field support in the three managed forests, Ivan Bjelanovic, Milic Čurović, Srdjan Keren and Vojislav Dukić for the field support in the old-growth forests. We thank the anonymous reviewers for their careful reading of our manuscript and their many insightful comments and suggestions.

Conflict of interest statement

None declared.

References

Alessandrini A, Biondi F, Di Filippo A, Ziaco E, Piovesan G (2011) Tree size distribution at increasing spatial scales converges to the rotated sigmoid curve in two old-growth beech stands of the Italian Apennines. Forest Ecology and Management 262:1950-1962

Ammer U (1991) Konsequenzen aus den Ergebnissen der Totholzforschung für die forstliche Praxis. Forstwissenschaftliches Centralblatt 110:149-157

Angelstam PK, Bütler R, Lazdinis M, Mikusinski G, Roberge JM (2003) Habitat thresholds for focal species at multiple scales and forest biodiversity conservation - dead wood as an example. Annales Zoologici Fennici 40:473-482

Attiwill PM (1994) The disturbance of forest ecosystems: the ecological basis for conservative management. Forest Ecology and Management 63:247-300

Barbati A et al. (2012) Assessing and promoting old-growthness of forest stands: Lessons from research in Italy. Plant Biosystems - An International Journal Dealing with all Aspects of Plant Biology 146:167-174 doi:10.1080/11263504.2011.650730

Bauhus J, Puettmann K, Messier C (2009) Silviculture for old-growth attributes. Forest Ecology and Management 258:525-537

Beese WJ, Dunsworth BG, Zielke K, Bancroft B (2003) Maintaining attributes of old-growth forests in coastal B.C. through variable retention. Forestry Chronicle 79:570-578

Bengtsson J, Nilsson SG, Franc A, Menozzi P (2000) Biodiversity, disturbances, ecosystem function and management of European forests. Forest Ecology and Management 132:39-50

Bergeron Y, Drapeau P, Gauthier S, Lecomte N (2007) Using knowledge of natural disturbances to support sustainable forest management in the northern Clay Belt. The Forestry Chronicle 83:326-337

Boncina A (2000) Comparison of structure and biodiversity in the Rajhenav virgin forest remnant and managed forest in the Dinaric region of Slovenia. Global Ecology and Biodiversity 9:201–211

Bottero A, Garbarino M, Dukić V, Govedar Z, Lingua E, Nagel TA, Motta R (2011) Gap-phase dynamics in the old growth forest of Lom (Bosnia-Herzegovina). Silva Fennica 45:865-873

Burrascano S, Keeton WS, Sabatini FM, Blasi C (2013) Commonality and variability in the structural attributes of moist temperate old-growth forests: A global review. Forest Ecology and Management 291:458-479 doi:http://dx.doi.org/10.1016/j.foreco.2012.11.020

Bütler R (2003) Dead wood in managed forests: how much and how much is enough? Development of a snag quantification method by remote sensing and GIS and snag targets based on the three-toed woodpecker's habitat requirements. Ecole polytechnique fédérale de Lausanne (EPFL)

Castagneri D, Garbarino M, Berretti R, Motta R (2010) Site and stand effects on coarse woody debris in montane mixed forests of Eastern Italian Alps. Forest Ecology and Management 260:1592-1598

Chirici G, Puletti N, Salvati R, Arbi F, Zolli C, Corona P (2014) Is randomized branch sampling suitable to assess wood volume of temperate broadleaved old-growth forests? Forest Ecology and Management 312:225-230 doi:http://dx.doi.org/10.1016/j.foreco.2013.10.001

Christensen M et al. (2005) Dead wood in European beech (Fagus sylvatica) forest reserves. Forest Ecology and Management 210:267-282 doi:10.1016/j.foreco.2005.02.032

de Liocourt F (1898) De l'aménagement des sapinières. Bull Soc For Franche-Comté et Belfort 4:396-409

Diaci J et al. (2011) Structural dynamics and synchronous silver fir decline in mixed old-growth mountain forests in Eastern and Southeastern Europe. Forestry 84:479-491 doi:10.1093/forestry/cpr030

Fahrig L (2003) Effects of habitat fragmenation on biodiversity. Annual Reviews of Ecology, Evolution and Sistematics 34:487-515

Franklin JF, Spies TA (1991) Ecological definitions of old-growth Douglas-fir forests. In: Ruggiero LF, Aubry KB, Carey AB, Huff MH (eds) Wildlife and vegetation of unmanaged Douglas-fir forests. USDA Forest Service, Pacific Northwest Research Station, Portland, pp 61-69

Franklin JF, Van Pelt R (2004) Spatial Aspects of Structural Complexity in Old-Growth Forests. Journal of Forestry 102:22-28

Goff FG, West D (1975) Canopy-understory Interaction Effects on Forest Population Structure. Forest Science 21:98-108

Goodburn JM, Lorimer CG (1999) Population structure in old-growth and managed northern hardwoods: an examination of the balanced diameter distribution concept. Forest Ecology and Management 118:11-29 doi:http://dx.doi.org/10.1016/S0378-1127(98)00478-2

Gustafsson L et al. (2012) Retention Forestry to Maintain Multifunctional Forests: A World Perspective. BioScience 62:633-645 doi:10.1525/bio.2012.62.7.6

Gustafsson L, Kouki J, Sverdrup-Thygeson A (2010) Tree retention as a conservation measure in clear-cut forests of northern Europe: a review of ecological consequences. Scandinavian Journal of Forest Research 25:295-308 doi:10.1080/02827581.2010.497495

Halpern CB, Spies TA (1995) Plant species diversity in natural and managed forests of the Pacific Northwest. Ecological Applications 5:913-934

Heilmann-Clausen J, Christensen M (2003) Fungal diversity on decaying beech logs-implications for sustainable forestry Biodiversity and Conservation 12:953-973

Humphrey JW (2005) Benefits to biodiversity from developing old-growth conditions in British upland spruce plantations: a review and recommendations. Forestry 78:33-53

Inventario Nazionale delle Foreste e dei Serbatoi Forestali di Carbonio (2005) Ministero delle Politiche Agricole Alimentari e Forestali, Ispettorato Generale - Corpo Forestale dello Stato. CRA - Unità di ricerca per il Monitoraggio e la Pianificazione forestale.

Janowiak MK, Nagel LM, Webster CR (2008) Spatial Scale and Stand Structure in Northern Hardwood Forests: Implications for Quantifying Diameter Distributions. Forest Science 54:497-506

Keeton WS (2006) Managing for late-successional/old-growth characteristics in northern hardwood-conifer forests. Forest Ecology and Management 235:129-142

Keeton WS, Franklin JF (2005) Do remnant old-growth trees accelerate rates of succession in mature Douglas-fir forests? Ecological Monographs 75:103-118

Keren S, Motta R, Govedar Z, Lucic R, Medarevic M, Diaci J (2014) Comparative Structural Dynamics of the Janj Mixed Old-Growth Mountain Forest in Bosnia and Herzegovina: Are Conifers in a Long-Term Decline? Forests 5:1243-1266 doi:10.3390/f5061243

Kirby KJ, Drake CM (1993) Dead wood matters: the ecology and conservation of saproxylic invertebrates in Britain. English Nature Science 7:1-105

Kohler F (2010) Die klimabedingte Veränderung der Totholzkäferfauna (Coleoptera) des nördlichen Rheinlandes. Analysen zur Gesamtfauna und am Beispiel von Wiederholungsuntersuchungen in ausgewählten Naturwaldzellen. Report for Ministerium für Umwelt und Naturschutz Landwirtschaft und Verbraucherschutz des Landes Nordrhein-Westfalen, Bornheim

Korpel S (1995) Die Urwälder der Westkarpaten. Fischer, Jena

Kucbel S, Jaloviar P, Saniga M, Vencurik J, Klimaš V (2010) Canopy gaps in an old-growth fir-beech forest remnant of Western Carpathians. European Journal of Forest Research 129:249-259

Kuuluvainen T, Penttinen A, Leinonen L, Nygren M (1996) Statistical opportunities for comparing stand structural heterogeneity in managed and primeval forests: an example from boreal Spruce forest in Southern Finland. Silva Fennica 30:315-328

Lähde E, Laiho O, Norokorpi Y, Saksa T (1999) Stand structure as the basis of diversity index. Forest Ecology and Management 115:213-220 doi:10.1016/s0378-1127(98)00400-9

Larrieu L, Cabanettes A, Delarue A (2012) Impact of silviculture on dead wood and on the distribution and frequency of tree microhabitats in montane beech-fir forests of the Pyrenees. European Journal of Forest Research 131:773-786 doi:10.1007/s10342-011-0551-z

Leak WB (1996) Long-term structural change in uneven-aged northern hardwoods. Forest Science 42:160-165

Leibundgut H (1982) Europäische Urwälder der Bergstufe. Paul Haupt, Bern

Lindenmayer DB, Franklin JF (2002) Conserving Forest Biodiversity: A Comprehensive Multiscaled Approach. Island Press, Washington, DC

Lindenmayer DB, Franklin JF, Fischer J (2006) General management principles and a checklist of strategies to guide forest biodiversity conservation. Biological Conservation 131:433-445 doi:http://dx.doi.org/10.1016/j.biocon.2006.02.019

Lingua E, Garbarino M, Borgogno Mondino E, Motta R (2011) Natural disturbance dynamics in an old-growth forest: from tree to landscape. Procedia Environmental Sciences 7:365-370 doi:10.1016/j.proenv.2011.07.063

Meyer P, Schmidt M (2011) Accumulation of dead wood in abandoned beech (Fagus sylvatica L.) forests in northwestern Germany. Forest Ecology and Management 261:342-352

Motta R (2002) Old-growth forests and silviculture in the Italian Alps: the case-study of the strict reserve of Paneveggio (TN). Plant biosystems 136:223-232

Motta R et al. (In press) Analysis of the spatio-temporal dynamics of mixed beech-silver fir-Norway spruce old-growth forests of Biogradska Gora (Montenegro) and Perućica (Bosnia-Herzegovina). Plant Biosystems

Motta R et al. (2011) Toward a definition of the range of variability of central European mixed Fagus-Abies-Picea forests: the nearly steady-state forest of Lom (Bosnia and Herzegovina). Canadian Journal of Forest Research 41:1871-1884 doi:10.1139/x11-098

Motta R, Berretti R, Castagneri D, Lingua E, Nola P, Vacchiano G (2010) Stand and coarse woody debris dynamics in subalpine Norway spruce forests withdrawn from regular management. Ann For Sci 67:803p801_803p808

Motta R, Berretti R, Lingua E, Piussi P (2006) Coarse woody debris, forest structure and regeneration in the Valbona Forest Reserve, Paneveggio, Italian Alps. Forest Ecology and Management 235:155-163 doi:10.1016/j.foreco.2006.08.007

Müller J, Bütler R (2010) A review of habitat thresholds for dead wood: a baseline for management recommendations in European forests. European Journal of Forest Research 129:981-992-992 doi:10.1007/s10342-010-0400-5

Nagel TA, Diaci J, Rozenbergar D, Rugani T, Firm D (2012) Old-growth forest reserves in Slovenia: the past, present, and future. Schweizerische Zeitschrift für Forstwesen 163:240-246 doi:10.3188/szf.2012.0240

Nagel TA, Svoboda M (2008) Gap disturbance regime in an old-growth *Fagus-Abies* forest in the Dinaric Mountains, Bosnia-Herzegovina. Canadian Journal of Forest Research 38:2728-2737

Nilsson SG et al. (2002) Densities of large living and dead trees in old-growth temperate and boreal forest. Forest Ecology and Management 161:189-204

O'Hara KL, Gersonde RF (2004) Stocking control concepts in uneven-aged silviculture. Forestry 77:131-143

Ódor P et al. (2006) Diversity of dead wood inhabiting fungi and bryophytes in semi-natural beech forests in Europe. Biological Conservation 131:58-71 doi:10.1016/j.biocon.2006.02.004

Paillet Y et al. (2010) Biodiversity Differences between Managed and Unmanaged Forests: Meta-Analysis of Species Richness in Europe. Conservation Biology 24:101-112 doi:10.1111/j.1523-1739.2009.01399.x

Parviainen J (2005) Virgin and natural forests in the temperate zone of Europe. For Snow Landsc Res 79:9-18

Peterken GF (1996) Natural woodland. Ecology and conservation in northern temperate regions. Cambridge University Press, Cambridge

Schütz JP (1996) Sylviculture 2. La gestion des forêts irrégulières et mélangéés. Presses Polytechniques et Universitaires Romandes, Lausanne

Siitonen J (2001) Forest management, coarse woody debris and saproxylic organisms: Fennoscandian boreal forests as an example. Ecological Bulletins 49:11-41

Siitonen J, Martikainen P, Punttila P, Rauh J (2000) Coarse woody debris and stand characteristics in mature managed and old-growth boreal mesic forests in southern Finland. Forest Ecology and Management 128:211-225

Silver EJ, D'Amato AW, Fraver S, Palik BJ, Bradford JB (2013) Structure and development of old-growth, unmanaged second-growth, and extended rotation Pinus resinosa forests in Minnesota, USA. Forest Ecology and Management 291:110-118 doi:http://dx.doi.org/10.1016/j.foreco.2012.11.033

Sitzia T, Trentanovi G, Dainese M, Gobbo G, Lingua E, Sommacal M (2012) Stand structure and plant species diversity in managed and abandoned silver fir mature woodlands. Forest Ecology and Management 270:232-238 doi:10.1016/j.foreco.2012.01.032

Spielmann M, Bücking W, Quadt V, Krumm F (2013) Integration of nature protection in forest policy in Baden-Württenberg (Germany). INTEGRATE country report. EFICENT-OEF. Freiburg

Stanturf JA, Madsen P (2002) Restoration concepts for temperate and boreal forests of North America and Western Europe. Plant Biosystems 136:143-158

Susmel L (1980) Normalizzazione delle foreste alpine. Liviana Editrice, Padova

Vandekerkhove K (2013) Integration of nature protection in Belgian forest policy: The case of Flanders. INTEGRATE country report for Belgium. EFICENT-OEF. Freiburg

Vandekerkhove K, De Keersmaeker L, Menke N, Meyer P, Verschelde P (2009) When nature takes over from man: Dead wood accumulation in previously managed oak and beech woodlands in North-western and Central Europe. Forest Ecology and Management 258:425-435

Vandekerkhove K et al. (2011) Reappearance of old-growth elements in lowland woodlands in northern Belgium: Do the associated species follow? Silva Fennica 45:909-935

Volin VC, Buongiorno J (1996) Effects of alternative management regimes on forest stand structure, species composition, and income: a model for the Italian Dolomites. Forest Ecology and Management 87:107-125 doi:http://dx.doi.org/10.1016/S0378-1127(96)03828-5

Wirth C, Gleixner G, Heimann M (eds) (2009) Old-growth forests. Function, fate and value. vol 207. Ecological Studies. Springer, Berlin

Wolynski A (1998) Evoluzione storica della selvicoltura naturalistica. Sherwood 40:5-11

Table and Figure captions

Table 1 Characteristics of the eight study sites.

Table 2 Forest structural characteristics of the eight study sites.

Table 3. Principal component loadings for the first two principal components for the eight study areas. Codes in Table 2.

Table 4 Signs (+/-) of significant coefficients (dbh, dbh^2 and dbh^3) in polynomial regression models used to determine diameter distribution shape (Janowiak et al. 2008) in uneven-aged forests. All the models grouped in the management categories were significant ($p < 0.001$) and their performance was compared through goodness of fit indices: adjusted R^2 (Adj. R^2), root mean square error (RMSE), Akaike information criterion (AIC) and Sum of squares due to error (SSE).

Table 5 Retention forestry in northern Italian regions and autonomous provinces according to current regional/provincial laws

Figure 1 Map of the study area. Location of sites (AMB - Amblar, CRO - Croviana, CAD - Santo Stefano di Cadore, LUD – Ludrin, NAV - Navarza, LOM - Lom, BIO - Biogradska gora, PER – Perućica) and management category (triangles: old growth; circles: FWRM; squares: managed).

Figure 2 Boxplots of 17 structural attributes in currently managed (MAN) forests, forests withdrawn from regular management (FWRM), and old-growth (OG). Groups with identical letters are not significantly different from each other ($p < 0.05$).

Figure 3 Principal component analysis (PCA of 207 plots) of structure surveyed in the study sites. Arrows represent selected forest structure descriptors (see Tab. 2 for codes). Triangles: old growth (og); circles: withdrawn from regular management (fwrm); squares: managed forests (man).

Figure 4 Diameter distributions in managed forests (MAN), forests withdrawn from regular management (FWRM), and old-growth (OG).

Figure 5 Relative frequency (average number of trees ha^{-1}) for increasing size classes in managed forests (MAN), forests withdrawn from regular management (FWRM), and old-growth (OG).

Tables

Tab. 1

Site	Code	Location	Area (ha)	n. plots	Altitude (m a.s.l.)	Slope (°)	Management category
Amblar	AMB	Italy, TN	5	13	1150-1220	5-25	Managed (single tree selection)
Croviana	CRO	Italy, TN	9	14	1220-1290	0-20	Managed (single tree selection)
Santo Stefano Cadore	CAD	Italy, BL	15	21	1080-1190	0-29	Managed (single tree selection)
Ludrin	LUD	Italy, TN	27	22	1250-1350	0-30	Withdrawn from regular management since 1962
Val Navarza	NAV	Italy, UD	37	33	1300-1570	12-34	Withdrawn from regular management since 1953
Lom	LOM	Bosnia-Herzegovina	40	40	1250-1520	0-40	Old-growth forest
Perućica	PER	Bosnia-Herzegovina	32	32	1340-1510	4-40	Old-growth forest
Biogradska gora	BIO	Montenegro	30	30	1210-1450	0-35	Old-growth forest

Tab. 2

Attached file

Statistical parameters	PC1	PC2
% of variance	*34.32*	*17.57*
cumulative % of variance	*34.32*	*51.89*
p	*0.001*	*0.001*
BA	-0.36	0.41
D80	-0.40	0.20
HD	-0.27	0.32
QMD	-0.39	0.30
REGEN	-0.01	-0.01
STUMP	0.39	0.31
VLOG50	-0.31	-0.25
VSNAG50	-0.27	-0.12
VSTUMP	0.26	0.21
PERCFASY	-0.13	-0.56
REGFASY	-0.28	-0.25

Tab. 3

Tab. 4

Management category	SSE	AIC	RMSE	Adj. R^2	dbh	dbh^2	dbh^3	Distribution shape
AMB	15.901	224.432	0.195	0.933	ns	ns	+	Unclassified
CAD	19.417	226.948	0.193	0.946	-	ns	ns	Negative Exp.
CRO	13.747	222.598	0.204	0.915	ns	+	+	Unclassified
Managed	13.601	222.464	0.121	0.971	-	ns	+	Concave
LUD	19.791	227.188	0.214	0.935	ns	ns	+	Unclassified
NAV	17.300	225.494	0.157	0.960	ns	+	+	Unclassified
FWRM	14.692	223.436	0.132	0.967	ns	+	+	Variable
LOM	11.997	220.884	0.148	0.949	-	ns	ns	Negative Exp.
PER	8.889	217.107	0.127	0.949	-	+	-	Rotated Sig.
BIO	8.054	215.865	0.168	0.902	-	ns	ns	Negative Exp.
Old-growth	8.737	216.890	0.086	0.976	-	+	-	Rotated Sigmoid

Tab. 5

Region/Province	Retention forestry measures All forests	Retention forestry measures Natura 2000 forests	Other	Notes
Piemonte	Some ageing trees (no definite number and no marking) adviced as a “good practice”	Living ageing trees (quantity not defined) and 2 dead trees ha^{-1} (snags or logs) if present. Additional retention measures scheduled locally for Natura 2000 forests		Conservation of dead or decaying habitat trees is advised in Natura 2000 forests
Valle d'Aosta Autonomous Region		Retention measures scheduled locally for Natura 2000 forests		Conservation of dead or decaying habitat trees is advised in Natura 2000 forests
Liguria		5 dead trees ha^{-1}, if present. Other additional measures scheduled locally for Natura 2000 forests		Conservation of "biodiversity islands", old trees, biodiversity trees is advised in Natura 2000 forests
Lombardia	2 living ageing trees ha^{-1} (marked, the trees must be replaced in case of death)	Retention measures scheduled locally for Natura 2000 forests		Conservation of dead or decaying habitat trees is advised in Natura 2000 forests
Bolzano Autonomous Province	Some ageing trees (no definite number and no marking).	Retention measures scheduled locally for Natura 2000 forests		According to the provincial forest rules some ageing trees, should be retained for biodiversity purposes in all the forests. Conservation of dead or decaying habitat trees is advised in Natura 2000 forests
Trento Autonomous Province	Some ageing trees (no definite number and no marking).	Retention measures scheduled locally for Natura 2000 forests		According to the provincial forest rules habitat trees, old or monumental trees, rare species should be retained for biodiversity purposes in all the forests. Conservation of dead or decaying habitat trees is advised in Natura 2000 forests
Veneto		Retention measures scheduled locally for Natura 2000 forests	Public funding (up to 200 € ha^{-1}) for owners that agree to preserve large living trees and dead trees (marked)	Conservation of dead or decaying habitat trees is advised in Natura 2000 forests
Friuli Venezia Giulia Autonomous Region	Some ageing trees (no definite number and no marking).	2 living trees (dbh > 50 cm if present) and 1 dead tree (dbh>40 cm if present) ha^{-1}. Other additional measures scheduled locally for Natura 2000 forests		According to the regional forest rules some ageing trees, should be retained for biodiversity purposes. Conservation of dead or decaying habitat trees is advised in Natura 2000 forests