

# In vitro screening of Latvian isolates of *Bjerkandera adusta* and *Sistotrema brinkmannii* as potential biocontrol agents against *Heterobasidion* root and butt rots

NATĀLIJA BURŅEVIČA<sup>1\*</sup>, DĀRTA KĻAVIŅA<sup>1</sup>, GUGLIELMO LIONE<sup>2</sup>, MARTINA PELLICCIARO<sup>2</sup>, LAUMA SILBAUMA<sup>1</sup>, ASTRA ZAĻUMA<sup>1</sup>, VIZMA NIKOLAJEVA<sup>3</sup> AND PAOLO GONTHIER<sup>2</sup>

<sup>1</sup> Latvian State Forest Research Institute "Silava", Rigas str. 111, Salaspils, LV-2169, Latvia

<sup>2</sup> University of Torino, Department of Agricultural, Forest and Food Sciences (DISAFA), Largo Paolo Braccini str. 2, I-10095 Grugliasco, Italy

<sup>3</sup> University of Latvia, Faculty of Biology, Department of Microbiology and Biotechnology, Jelgavas str. 1, Riga, LV-1004, Latvia

\* Corresponding author: [natalija.burnevica@silava.lv](mailto:natalija.burnevica@silava.lv); phone: +371 25951553

Burņeviča, N., Kļaviņa, D., Leone, G., Pellicciaro, M., Silbauma, L., Zaļuma, A., Nikolajeva, V. and Gonthier, P. 2022. In vitro screening of Latvian isolates of *Bjerkandera adusta* and *Sistotrema brinkmannii* as potential biocontrol agents against *Heterobasidion* root and butt rots. *Baltic Forestry* 28(1): article id 637. <https://doi.org/10.46490/BF637>.

Received 21 February 2022 Revised 28 April 2022 Accepted 29 April 2022

## Abstract

Root and butt rots caused by *Heterobasidion* spp. are destructive diseases of conifer forest stands that may be controlled by spraying freshly cut stumps with chemical or biological treatments, including the effective and widely used biological control agent *Phlebiopsis gigantea*. In this study, we tested *in vitro* different Latvian isolates of the fungal species *Bjerkandera adusta* and *Sistotrema brinkmannii* for their antagonistic activity against both *Heterobasidion* species present in Northern Europe and Baltic countries, namely *H. annosum* and *H. parviporum*, using Latvian isolates of *P. gigantea* and the Rotstop® strain as controls. The growth rate of isolates on agar media and their ability to produce oidia or conidia were also assessed. In general, *B. adusta* performed better than *S. brinkmannii*. Some isolates of *B. adusta* ranked similarly to those of *P. gigantea* in terms of oidia production, growth rate and ability to overgrow the mycelium of both *H. annosum* and *H. parviporum*. These isolates will be further tested in field conditions.

**Keywords:** biological control, antagonistic fungi, fungal isolates

## Introduction

*Heterobasidion* root and butt rots are some of the most significant conifer diseases in the Northern Hemisphere, causing losses of approximately 800 million euros to the European forestry annually (Woodward et al. 1998). These fungal diseases are caused by basidiomycetes included in the *Heterobasidion annosum* species complex, namely *H. abietinum*, *H. annosum sensu stricto*, hereafter referred to as *H. annosum*, and *H. parviporum* in Eurasia, and *H. irregulare* and *H. occidentale* in North America (Garbelotto and Gonthier 2013). In Northern Europe and the Baltic States, particularly in Latvia, both *H. annosum* and *H. parviporum* are present and abundant, the former mainly associated with Scots pine (*Pinus sylvestris*) and the latter with Norway spruce (*Picea abies*) (Korhonen et al. 1992, Korhonen and Piri 1994, Gaitnieks et al. 2021).

*Heterobasidion* species spread in two ways: 1) by airborne basidiospores produced in fruit bodies and infecting freshly cut conifer stumps or tree wounds; 2) by mycelium spreading through root contacts to adjacent trees (Korhonen and Stenlid 1998, Gonthier and Thor 2013). To prevent spore infection of freshly cut stumps, stump treatment using biological or chemical control agents is widely used (Nicolotti and Gonthier 2005, Thor and Stenlid 2005, Gonthier and Thor 2013). If a forest stand is already severely infested, stump removal or planting tree species resistant or less susceptible to the disease could be considered (Vasaitis et al. 2008, Garbelotto and Gonthier 2013).

The most effective biological control agent (BCA) for stump treatments against *Heterobasidion* spp. is the basidiomycete *Phlebiopsis gigantea* (Pratt et al. 2000, Garbelotto and Gonthier 2013), although the bacterium *Pseudomo-*

*nas protegens* has recently been proved as effective both *in vitro* and in forest conditions (Pellicciaro 2021a, b). In Europe, the biological product based on *P. gigantea* most widely employed in practical forestry is Rotstop® (Thor 2003). In the Baltic countries, Rotstop® is registered for use in Estonia and Latvia since 2004 and 2007, respectively (Drenkhan et al. 2008, Kenigvalde et al. 2016). Rotstop® is extensively used by Latvia's State Forests: each year it is applied to an area of more than 5,000 hectares using at least 300 thousand litres of prepared suspension for stump treatment (J. Mālakalns, personal communication). Rotstop® was developed decades ago and it contains a single *P. gigantea* strain native to Finland (Korhonen et al. 1994). In the long term, such a large-scale application of one isolate could negatively affect populations of native *P. gigantea* and of other stump colonizing fungi, as previously suggested (Varese et al. 2001, Vasiliauskas et al. 2004). Although the abundance of the Rotstop® strain of *P. gigantea* in Scots pine and Norway spruce stumps has been reported to decrease within a period of three to six years after treatments (Vainio et al. 2001, Vasiliauskas et al. 2005), from a biological diversity perspective it would be beneficial to use also other BCAs, including native *P. gigantea* strains and other fungi. Moreover, *P. gigantea* is better adapted to Scots pine stumps than to Norway spruce stumps (Korhonen 2003, Webber and Thorpe 2003) making treatments with this BCA more effective when applied on stumps of the former tree species than on stumps of the latter one (Korhonen et al. 1994, Berglund and Rönnerberg 2004, Rönnerberg et al. 2006). For instance, data from Latvia showed that the efficacy of Rotstop® was on average 82% on Scots pine stumps but only 64% on Norway spruce stumps (Kenigvalde et al. 2016). Altogether the above findings point to the opportunity of identifying additional antagonistic fungi effective against *Heterobasidion* spp., especially on Norway spruce stumps. Various saprotrophic fungi have been tested against *Heterobasidion* spp. in laboratory and field conditions (Hyppel 1968, Rishbeth 1970, Aufsess 1976b, Lundborg and Unestam 1980, Holdenrieder 1982, 1984, Negrutskii 1986, Vasiliauskas 1989, Capretti and Mugnai 1989a, b, 1993, Hanso and Hanso 1985, 1992, Holmer and Stenlid 1993, 1996, Negrutskii 1993, Anselmi and Nicolotti 1997, Nicolotti et al. 1999, Roy et al. 2003, Woods et al. 2005, Hamberg et al. 2012). Among them, the basidiomycete *Bjerkandera adusta* was identified as one of the most promising species after *P. gigantea* (Holdenrieder and Greig 1998). *Bjerkandera adusta* is common in wood of deciduous trees (Holdenrieder and Greig 1998), but several studies indicate that it is also associated with Norway spruce stumps (Pechmann and Aufsess 1971, Kallio and Hallaksela 1979, Vasiliauskas et al. 2002). Interestingly, this fungus has been reported to be able to replace other fungi both *in vitro* and in stem sections of *Platanus* sp. (Holdenrieder and Greig 1998 and literature therein). Another fungal species showing the ability to replace other fungi both in laboratory and field experi-

ments is *Sistotrema brinkmannii* (Holdenrieder and Greig 1998 and literature therein), a fungus reported to colonize wood both at the early and at the final stage of decay (Morrison and Johnson 1978, Vasiliauskas et al. 2005, Stokland and Larsson 2011). *Sistotrema brinkmannii* frequently colonizes wood already decayed by *P. gigantea* (Vasiliauskas et al. 2005).

In Latvia, analyses of wood samples collected from 1182 decayed Norway spruce stumps revealed that *H. parviporum* and *Stereum sanguinolentum* are the most common stump-inhabiting basidiomycetes (Arhipova et al. 2011a). The next most abundant basidiomycetes were right *B. adusta* and *S. brinkmannii*, showing approximately the same isolation frequency (Arhipova et al. 2011a).

While the antagonism of *B. adusta* and *S. brinkmannii* towards wood decay fungi, including *Heterobasidion* spp., *Armillaria* spp. and *S. sanguinolentum* has been documented (Rishbeth 1970, Aufsess 1976a, b, Kallio and Hallaksela 1979, Capretti and Mugnai 1989a, b, Nicolotti et al. 1993), as noted by Holdenrieder and Greig (1998) studies on the efficacy of these two fungi against *Heterobasidion* spp. are still scarce to determine their level of antagonistic potential.

From a biotechnological perspective, oidia production by a fungus is an important feature for the production of a biological product (Pratt et al. 2000). Similarly, to *P. gigantea*, *B. adusta* produces oidia, although little is known about intraspecific variation of the oidia production between different *B. adusta* isolates. *Sistotrema brinkmannii*, to our knowledge, can produce both basidiospores and Ptychogaster-like conidia, but information about this production are scarce (Carris and Glawe 1989).

The aim of this study was to screen *B. adusta* and *S. brinkmannii* isolates *in vitro* for their antagonistic potential against both *H. annosum* and *H. parviporum*, using *P. gigantea* as a control. The specific aims were: i) to assess the mycelial growth of *B. adusta* and *S. brinkmannii* isolates on malt agar media; ii) to evaluate their level of antagonism against *H. annosum* and *H. parviporum* using dual culture assays; iii) to appraise the oidia production potential of isolates of *B. adusta*. The results will be used for selecting the most promising fungal isolates for further field experiments.

## Material and methods

### Fungal isolates

In total, 11 isolates of *B. adusta*, six isolates of *S. brinkmannii*, four isolates of *P. gigantea*, two isolates of *H. annosum*, and two isolates of *H. parviporum* were used in this study (Table 1). Isolates of *B. adusta*, *S. brinkmannii* and *P. gigantea* were collected from different substrates during the years 2005 to 2016 in different regions of Latvia. Fungal isolates were obtained from decayed Scots pine, Norway spruce and alder (*Alnus* sp.) wood. The isolation procedure was described in Arhipova et al. 2011a, b. Brief-

**Table 1.** Fungal isolates used in this study

Fungal species	Isolate ID	GenBank accession number	Coordinates of the site of origin	Host	Isolation source	
<i>Bjerkandera adusta</i>	B25	-	56°33'00N 23°01'54E	<i>Picea abies</i>	Stump	
	B50	FJ903311	57°00'00N 23°01'54E	<i>Picea abies</i>	Stump	
	E67	-	57°18'51N 22°50'42E	<i>Picea abies</i>	Stem	
	E182	-	57°21'23N 25°11'49E	<i>Picea abies</i>	Stem	
	I216	-	56°50'39N 22°22'13E	<i>Alnus incana</i>	Stem	
	I345	GU062301	56°41'43N 25°55'24E	<i>Alnus incana</i>	Stem	
	L38A	-	56°44'50N 23°55'57E	<i>Picea abies</i>	Stem	
	L38B	-	56°44'50N 23°55'57E	<i>Picea abies</i>	Stem	
	M59	JF340266	56°41'59N 25°54'26E	<i>Alnus glutinosa</i>	Stem	
	M108	-	56°41'59N 25°54'26E	<i>Alnus glutinosa</i>	Stem	
	M111	-	56°41'59N 25°54'26E	<i>Alnus glutinosa</i>	Stem	
	<i>Sistotrema brinkmannii</i>	B5	FJ903297	56°59'29N 23°10'38E	<i>Picea abies</i>	Stump
		E136	-	56°59'35N 23°10'11E	<i>Picea abies</i>	Stem
		I390	GU062313	57°00'30N 23°07'22E	<i>Alnus incana</i>	Stem
I409		-	56°41'43N 25°55'24E	<i>Alnus incana</i>	Stump	
P31		-	57°00'23N 23°13'02E	<i>Pinus sylvestris</i>	Stem	
S73		MK911637	57°06'28N 25°07'09E	<i>Picea abies</i>	Root	
<i>Phlebiopsis gigantea</i>	Rotstop®	-	-	<i>Picea abies</i>	Stem	
	PG382	-	56°50'57N 24°48'18E	<i>Picea abies</i>	Stem	
	PG182	-	56°37'30N 25°50'58E	<i>Pinus sylvestris</i>	Stem	
	PG408	-	56°40'53N 25°53'24E	<i>Pinus sylvestris</i>	Stem	
<i>Heterobasidion annosum</i>	HAZ	-	56°37'49N 24°27'24E	<i>Pinus contorta</i>	Stump	
	HAU	-	57°15'43N 22°00'35E	<i>Pinus contorta</i>	Stump	
<i>Heterobasidion parviporum</i>	HPN	-	57°06'28N 25°07'09E	<i>Picea abies</i>	Root	
	HPO	-	56°45'42N 24°45'57E	<i>Picea abies</i>	Root	

ly, wood samples were taken by using a Pressler borer. In the laboratory, small pieces of decayed wood were placed on Hagem agar in Petri plates. Fungal cultures that were obtained were identified as follows. DNA was extracted, PCR reactions were run using the primer combination ITS1F and ITS4, Sanger sequenced, and BLAST search were performed using NCBI (National Center for Biotechnology Information) fungal database (NCBI 2022) (Arhipova et al. 2011a, b, Arhipova et al. 2012, Burņeviča et al. 2021). Selected sequences were deposited in the NCBI database. For the identification of *Heterobasidion* species, intersterility tests were performed (Korhonen 1978) using homokaryotic test cultures 03013 and 03015 (*H. annosum*), and 96119 and 98036 (*H. parviporum*) (courtesy of Dr. Kari Korhonen, LUKE, Finland) (Zaļuma et al. 2019, Burņeviča et al. 2021). Latvian *P. gigantea* isolates as well as the Finnish one of the fungi contained in Rotstop® were used as controls in this study for the mycelial growth rate *in vitro*, oidia production and antagonism of *B. adusta* and *S. brinkmannii* against *H. annosum* and *H. parviporum*.

#### Assessment of fungal mycelium growth rate

For the assessment of the mycelial growth *in vitro*, circle-shaped agar blocks (Ø 0.6 cm) were cut from actively growing fungal cultures and placed at the edge of a Ø 9 cm Petri plate containing malt-agar medium (7 g malt extract, 6 g agar per 1 L water). The Petri plates were incubated in the dark at 20°C. The fungal growth was determined by measuring the radial growth along the agar surface and then tracing the leading edge of the growing colony with a marker. The growth rate was calculated every 48 hours at approximately the same time for 8 days. The average growth rate of three replicates was calculated for each fungal isolate.

#### Assessment of antagonistic activity against *H. annosum* and *H. parviporum*

For the assessment of antagonistic activity of the fungal isolates against *Heterobasidion* spp., the dual culture assay technique was used (Holdenrieder and Greig 1998). Circle-shaped agar blocs (Ø 0.6 cm) of each pure fungal culture was transferred to a sterile malt-agar Petri plate, using a flame sterilized inoculation needle. A similar agar block (Ø 0.6 cm) of either *H. annosum* or *H. parviporum* was transferred to the same Petri plate at a distance of 5–6 cm from the test cultures. All the Petri plates were incubated in the dark at 20°C and the width of the inhibition zone between the two fungal isolates was measured after 14 days. For the evaluation of fungal interaction, a modified classification system developed by Holdenrieder (1984a) was used: ++ strong antagonism, antagonist overgrows *Heterobasidion* sp.; + moderate antagonism, formation of mycelium-free inhibition zone between antagonist and *Heterobasidion* sp.; +- sporadic antagonism, antagonist creates a mycelium-free inhibition zone only with one *Heterobasidion* sp. isolate; - antagonistic activity absent, *Heterobasidion* sp. overgrows the antagonist.

#### Assessment of oidia production

Oidia suspensions were prepared from each fungal isolate of *B. adusta* and *P. gigantea* by washing the agar media surface of 12-day old pure cultures with 0.5 ml of sterile water, gently shaking the mycelium with a flame sterilized scalpel. Using a 0.3 ml Eppendorf automatic pipette, oidia suspension was transferred to a sterile 1.5 ml Eppendorf tube. A similar approach was used to collect conidia of *S. brinkmannii*.

The number of oidia/conidia in the suspensions was counted in a Goryaev chamber (a variation of a hemocytometer) (Muntyan et al. 2012). The cover glass was placed on the middle grid of the chamber and rubbed so that the Newton's rings could be observed (to ensure the depth of the chamber). Using an Eppendorf automatic pipette, 100 µl of suspension was transferred to the edge of the cover glass in the Goryaev chamber. The oidia/conidia were counted in 30 small grids of the chamber at 100× magnification using a Leica DM 4000B microscope. On the upper left corner of the grid, oidia/conidia were counted in the diagonally selected four smaller squares. Following the same principle, oidia/conidia were counted in all the squares by moving the camera to the right and finding the next large square. Oidia/conidia located at the edge with at least half of the cell inside the square were considered on the left and lower line. The average number of oidia/conidia in one square and in 1 ml of suspension was calculated, considering the area of the square.

### Data analysis

Data was organized and analyzed using Microsoft Excel spreadsheet. The concentration of oidia/conidia in 1 ml of suspension was calculated using the following formula:

$$C_{oc} = 16N_{ss} \cdot N_{oc1} \cdot 25 \cdot 10^4, \quad (1)$$

where:

$N_{ss}$  is the number of small squares;

$N_{oc1}$  is the number of oidia/conidia in one small square.

For data analysis,  $t$ -test was run at  $\alpha = 0.05$ . Correlation between mycelial growth rate and oidia production was calculated using the Pearson correlation (Liepa 1974).

## Results

### Mycelial growth rate of *B. adusta*, *S. brinkmannii* and *P. gigantea* isolates

The average mycelial growth rate of each isolate is summarized in Table 2. The results show that the average mycelial growth of the fungal isolates is uneven and differs not only between fungal species, but also between isolates of the same species. For *B. adusta*, the highest average growth rate was observed for isolates E182 (0.88 cm day<sup>-1</sup>) and I345 (0.84 cm day<sup>-1</sup>), while the lowest one was observed for L38A (0.42 cm day<sup>-1</sup>). For *S. brinkmannii*, the highest growth rate was observed for isolates B5 (0.46 cm day<sup>-1</sup>) and P31 (0.42 cm day<sup>-1</sup>), while the lowest one for E136 (0.18 cm day<sup>-1</sup>) and I409 (0.16 cm day<sup>-1</sup>). The highest average growth rate among *P. gigantea* isolates was observed for the isolate from Rotstop® (0.73 cm day<sup>-1</sup>) and for isolate PG382 (0.70 cm day<sup>-1</sup>). The average growth rate of *B. adusta* isolates (0.73 ± 0.11 cm day<sup>-1</sup>) was slightly higher than the average growth rate of *P. gigantea* isolates (0.69 ± 0.03 cm day<sup>-1</sup>) although differences were not significant ( $p = 0.31$ ). However, five *B. adusta* isolates showed higher average growth rate than

**Table 2.** Mycelial growth rate, antagonism against *Heterobasidion* spp. and oidia/conidia production of different *B. adusta*, *S. brinkmannii* and *P. gigantea* isolates


Fungal species	Fungal isolate	Average mycelial growth rate on malt agar media, cm per day (± SD)	Antagonism against <i>H. annosum</i> / <i>H. parviporum</i> *	Average number of oidia/conidia in 1 mL of suspension and SD × 10 <sup>6</sup>
<i>Bjerkandera adusta</i>	B25	0.77 ± 0.15	++/++	0.17 ± 0.08
	B50	0.72 ± 0.15	-/+	0.16 ± 0.01
	E67	0.74 ± 0.11	-/+	0.36 ± 0.10
	E182	0.88 ± 0.04	+/-	0.20 ± 0.06
	I216	0.76 ± 0.09	-/+	0.26 ± 0.06
	I345	0.84 ± 0.09	-/+	0.16 ± 0.07
	L38A	0.42 ± 0.08	+/+	0.57 ± 0.52
	L38B	0.73 ± 0.17	-/-	0.07 ± 0.01
	M59	0.69 ± 0.25	++/++	0.29 ± 0.33
	M108	0.70 ± 0.17	+/-/+	0.14 ± 0.01
	M111	0.78 ± 0.18	-/+	0.27 ± 0.08
<i>Sistotrema brinkmannii</i>	B5	0.46 ± 0.09	+/+	-
	E136	0.18 ± 0.07	+/+	0.03 ± 0.01
	I390	0.34 ± 0.11	-/-	0.06 ± 0.01
	I409	0.16 ± 0.05	-/-	-
	P31	0.42 ± 0.12	++/++	0.08 ± 0.02
<i>Phlebiopsis gigantea</i>	S73	0.34 ± 0.10	+/-/+	0.06 ± 0.08
	Rot-stop®	0.73 ± 0.21	++/++	1.44 ± 0.03
	PG408	0.64 ± 0.13	++/++	1.26 ± 0.03
	PG382	0.70 ± 0.17	++/++	1.05 ± 0.01
<i>Heterobasidion annosum</i>	PG182	0.68 ± 0.22	++/++	1.50 ± 0.02
	HAZ	0.50 ± 0.07	-	-
	HAU	0.50 ± 0.14	-	-
<i>Heterobasidion parviporum</i>	HPN	0.52 ± 0.18	-	-
	HPO	0.62 ± 0.13	-	-

Notes: \* ++ strong antagonism; + moderate antagonism; +/- sporadic antagonism; - no antagonism.


that of *P. gigantea*: B25 (0.77 ± 0.15 cm day<sup>-1</sup>), E182 (0.88 ± 0.04 cm day<sup>-1</sup>), I216 (0.76 ± 0.09 cm day<sup>-1</sup>), I345 (0.84 ± 0.09 cm day<sup>-1</sup>) and M111 (0.78 ± 0.18 cm day<sup>-1</sup>). The average growth rate of *S. brinkmannii* isolates was much lower (0.32 ± 0.11 cm day<sup>-1</sup>) compared to *B. adusta* and *P. gigantea* ( $p < 0.01$ ). The growth rate of *Heterobasidion* spp. varied between 0.3–0.7 cm day<sup>-1</sup> with an average of 0.5 cm day<sup>-1</sup> for *H. annosum*, and 0.52–0.62 cm day<sup>-1</sup> with an average of 0.6 cm day<sup>-1</sup> for *H. parviporum*. Although the mycelium of *Heterobasidion* spp. had a lower growth rate in comparison with *B. adusta* and *P. gigantea*, the difference was not significant ( $p > 0.05$ ).

### Antagonistic activity against *Heterobasidion* spp.

All *P. gigantea* isolates showed high antagonistic activity against both *Heterobasidion* species, whereas the antagonistic activity of *B. adusta* and *S. brinkmannii* varied depending on isolates (Table 2). Three *B. adusta* isolates – B25 (Figure 1), L38A and M59 – showed a distinct antagonism against both *Heterobasidion* species. Similarly, to all the analyzed *P. gigantea* isolates, including that from Rotstop®, the previously mentioned *B. adusta* isolates were able to overgrow *Heterobasidion* spp. mycelia. The other



**Figure 1.** Strong antagonistic reaction of *B. adusta* isolate B25 (BAB25) against: a) *H. parviporum* (S); b) *H. annosum* (P)



**Figure 2.** Moderate antagonistic reaction of *S. brinkmannii* isolate B5 (SBB5) against: a) *H. parviporum* (S); b) *H. annosum* (P)

investigated *B. adusta* isolates either showed weak antagonistic activity against some isolates of *Heterobasidion* spp. or did not inhibit the growth of the pathogen at all.

Three *S. brinkmannii* isolates – B5 (Figure 2), E136 and P31 – showed an antagonistic reaction against both *Heterobasidion* species. The remaining *S. brinkmannii* isolates showed either a slight inhibition of the growth of some isolates of *Heterobasidion* spp. or a lack of antagonistic activity. As for *B. adusta*, most of the *S. brinkmannii* isolates had a stronger antagonistic activity against *H. parviporum* than against *H. annosum* isolates.

### Oidia/conidia production

The average number of oidia/conidia of *B. adusta*, *S. brinkmannii* and *P. gigantea* is summarized in Table 2.

Among the *B. adusta* isolates, the highest oidia production was observed for isolate L38A ( $0.57 \times 10^6$  oidia per 1 mL suspension), which was characterized by a slow mycelial growth rate. The remaining *B. adusta* isolates had lower oidia production ( $0.07 \times 10^6$ – $0.36 \times 10^6$  oidia per 1 mL suspension). A significant negative correlation was observed between mycelium growth rate of *B. adusta* and oidia production ( $r = -0.7$ ;  $p < 0.01$ ). The highest rate of conidia production of *S. brinkmannii* was observed for

isolate P31 ( $0.08 \times 10^6$  conidia per 1 mL suspension); the other isolates had slightly lower conidia production ranging from  $0.03$  to  $0.06 \times 10^6$  oidia per 1 mL suspension. Two isolates (B5 and I409) did not produce conidia on agar media. There was no significant correlation observed between mycelium growth rate of *S. brinkmannii* and conidia production ( $r = 0.3$ ;  $p > 0.05$ ).

The number of oidia of *P. gigantea* in 1 mL of suspension was comprised between  $1.05 \times 10^6$  and  $1.50 \times 10^6$  depending on isolates. No correlation was observed between the mycelial growth rate and oidia production for *P. gigantea* ( $r = 0.1$ ;  $p > 0.05$ ).

### Discussion

The results obtained in this study highlighted that *B. adusta* and *S. brinkmannii* show high intraspecific variability of the parameters investigated under laboratory conditions. *Sistotrema brinkmannii* isolates showed great variation in mycelial growth rate while *B. adusta* displayed a relevant variability in oidia production. Oidia production of *B. adusta* isolates was lower compared to that of *P. gigantea* isolates used as controls. Our results are consistent with those of Holdenrieder (1984), who reported that only

two out of the three tested isolates of *B. adusta* produced oidia and that the oidia production of this species was scarce. However, Holdenrieder (1984) suggested that sporulation of *B. adusta* can be stimulated by changing nutrient content or using UV light.

As previously mentioned, the ability of producing oidia/conidia is an important parameter for screening and assessing fungal isolates as potential biocontrol agents (Pratt et al. 2000, Sun et al. 2009). The isolates of *B. adusta* investigated in this study differed by more than 10 times in terms of ability to produce oidia, and this points to the possibility of identifying the most promising ones for further experiments.

Interestingly, the analyzed *P. gigantea* isolates, including the Rotstop® one, showed small variability in the oidia production although other authors have reported large variability of oidia production for *P. gigantea* isolates (Sun et al. 2009, Zaļuma et al. 2019). Reduced variability among isolates of this species in our study could be related to the low number of *P. gigantea* strains tested (three Latvian isolates and the Rotstop® one).

*Bjerkandera adusta* isolates showed a growth rate similar to the *P. gigantea* isolates used as controls, on average 0.73 and 0.69 cm per day for the former and the latter species, respectively. The observed growth rate of *P. gigantea* is consistent with that reported in other studies (Zaļuma et al. 2019). A study from Finland reported a negative correlation between mycelial growth rate in wood and oidia production of *P. gigantea* as well as a positive correlation between growth rate of isolates on agar medium and in wood (Sun et al. 2009). While we could not confirm the above correlations for *P. gigantea*, possibly because of the limited number of isolates included in the experiments, our results showed that the growth rate on agar medium is negatively correlated with oidia production for *B. adusta* isolates.

However, the most important parameter to evaluate *in vitro* for the selection of potential BCA against *Heterobasidion* spp. is the performance of candidates in dual culture tests (Pratt et al. 2000). For this purpose, malt agar media and mixtures with sawdust as well as conifer wood blocks or discs are used (Rishbeth 1970, Aufsess 1976a, Lundborg and Unestam 1980, Woods et al. 2005, Drenkhan et al. 2008, Pellicciaro et al. 2021a).

Our results show that all the analyzed *P. gigantea* isolates are strong antagonists against *Heterobasidion* spp. under laboratory conditions, supporting the results of previous experiments (Negrutskii 1986, Vasiliauskas 1989, Hanso and Hanso 1992, Annesi et al. 2005, Drenkhan et al. 2008, Sun et al. 2009, Mgbeahuruike et al. 2011, Zaļuma et al. 2019). Two of the tested *B. adusta* isolates and one *S. brinkmannii* isolate also showed explicit antagonism against both *H. annosum* and *H. parviporum*. In a similar study from Italy evaluating the antagonism of various fungal species against *Heterobasidion* spp. *in vitro* including *P. gigantea* used as a control, the authors concluded that

*P. gigantea* performs better than *S. brinkmannii* (Capretti and Mugnai 1989a, b). Antagonism in our study was assessed after 14 days although it was previously reported that fungal antagonism can differ significantly if evaluated after 5 or 30 days (Negrutskii 1986). The temperature of incubation of dual culture plates could also influence the outcomes of the interaction, as noted in previous studies (Capretti and Mugnai 1989a, Hanso and Hanso 1992, Drenkhan et al. 2008, Mgbeahuruike et al. 2011). In our study, plates were incubated at 20°C. *In vitro* studies (Holdenrieder 1984, Capretti and Mugnai 1989a) revealed that some *P. gigantea* isolates in dual cultures display greater antagonistic activity against *Heterobasidion* spp. at 11–12°C than at 20–21°C. This aspect is very important since in Fennoscandia and the Baltic countries active sporulation of *Heterobasidion* spp. has been reported in September–October (Brandtberg 1996, Bruna et al. 2021) but sporulation of *P. gigantea* decreases starting from late summer (Kallio and Hallaksela 1979, Vasiliauskas et al. 2005). Therefore, the possible impact of temperature should be considered when assessing potential BCAs. In contrast to *P. gigantea*, *S. brinkmannii* antagonism against *Heterobasidion* spp. did not differ at 12 and 20°C (Capretti and Mugnai 1989a). However, it is worth noting that several studies have emphasized that results obtained under laboratory conditions need to be confirmed in field experiments (Holdenrieder and Greig 1998, Mgbeahuruike et al. 2011).

Our findings confirm that *B. adusta* is a very promising antagonist of *Heterobasidion* spp. Some isolates of this species originating from Latvia, i.e. B25 and M59, ranked similarly to *P. gigantea* in terms of oidia production, growth rate and ability to overgrow the mycelium of both *H. annosum* and *H. parviporum*. These isolates, along with *S. brinkmannii* isolate P31 showing the ability of producing conidia and high antagonistic activity against *Heterobasidion* spp. in dual cultures, will be further tested in conifer wood alone but also in combination with *P. gigantea*. In fact, it was previously demonstrated that mixed suspensions of *P. gigantea* with other wood decay fungi like *Chondrostereum purpureum* may improve the performances of *P. gigantea* against *Heterobasidion* spp. (Hamberg et al. 2012).

### Acknowledgements

The study was financially supported by European Regional Development fund project (No 1.1.1.1/20/A/095) and Latvian Forest Research Institute “Silava”.

### References

- Annesi, T., Curcio, G., D’Amico, L. and Motta, E. 2005. Biological control of *Heterobasidion annosum* on *Pinus pinea* by *Phlebiopsis gigantea*. *Forest Pathology* 35(2): 127–134. <https://doi.org/10.1111/j.1439-0329.2004.00394.x>.
- Anselmi, N. and Nicolotti, G. 1997. Biological control of *Heterobasidion annosum* in the forest by non-pathogenic wood-destroying fungi. In: Root and butt rots of forest

- trees: 9th International Conference on Root and Butt Rots, Carcans-Maubuisson, (France), 1–7 September 1997, p. 421–428.
- Arhipova, N., Gaitnieks, T., Donis, J., Stenlid, J., and Vasaitis, R.** 2011a. Butt rot incidence, causal fungi and related losses in *Picea abies* stands of Latvia. *Canadian Journal of Forest Research* 41: 2337–2345. <https://doi.org/10.1139/x11-141>.
- Arhipova, N., Gaitnieks, T., Donis, J., Stenlid, J. and Vasaitis, R.** 2011b. Decay, yield loss and associated fungi in stands of grey alder (*Alnus incana*) in Latvia. *Forestry* 84(4): 337–348. <https://doi.org/10.1093/forestry/cpr018>.
- Arhipova, N., Gaitnieks, T., Donis, J., Stenlid, J. and Vasaitis, R.** 2012. Heart-rot and associated fungi in *Alnus glutinosa* stands in Latvia. *Scandinavian Journal of Forest Research* 27: 227–336. <https://doi.org/10.1080/02827581.2012.670727>.
- Aufsess, von H.** 1976a. Über die Wirkung verschiedener Antagonisten auf das Mycelwachstum von einigen Stamfäulepilzen [Effect of different antagonist on mycelial growth of some stem rot fungi]. *Material und Organismen* 11: 183–196 (in German).
- Aufsess, von H.** 1976b. Der Einfluß verschiedener Mikroorganismen auf das Abbauverhalten einiger Stammfäuleerreger [The influence of different microorganisms on the decomposition behavior of some stem rot pathogens]. *Material und Organismen* 3: 239–250 (in German).
- Berglund, M. and Rönnerberg, J.** 2004. Effectiveness of treatment of Norway spruce stumps with *Phlebiopsis gigantea* at different rates of coverage for the control of *Heterobasidion*. *Forest Pathology* 34(4): 233–243. <https://doi.org/10.1111/j.1439-0329.2004.00363.x>.
- Brandtberg, P.-O., Johansson, M. and Seger, P.** 1996. Effects of season and urea treatment on infection of stumps of *Picea abies* by *Heterobasidion annosum* in stands on former arable land. *Scandinavian Journal of Forest Research* 11: 261–268. <https://doi.org/10.1080/02827589609382935>.
- Brūna, L., Lione, G., Kenigvalde, K., Burneviča, N., Zaļuma, A., Kļaviņa, A., Gaitnieks, T. and Gonthier, P.** 2021. Inferences on susceptibility of wood of different tree species to *Heterobasidion annosum sensu lato* primary infections and the range of pathogen spores dispersal. *Forests* 12: 854. <https://doi.org/10.3390/f12070854>.
- Burneviča, N., Zaļuma, A., Kļaviņa, D., Brūna, L., Legzdina, L. and Gaitnieks, T.** 2021. Initial and long-term fungal diversity and occurrence of *Heterobasidion* spp. in Norway spruce root fragments remaining in soil after stump extraction. *Scandinavian Journal of Forest Research* 36(2-3): 117–125. <https://doi.org/10.1080/02827581.2021.1890814>.
- Capretti, P., and Mugnai, L.** 1989a. *In vitro* tests of antagonism against *Heterobasidion annosum* (Fr.) Bref. / Prove di antagonismo in vitro contro *Heterobasidion annosum* (Fr.) Bref. *Phytopathologia Mediterranea* 28(2): 150–154. Available online at: <http://www.jstor.org/stable/42685693>.
- Capretti, P. and Mugnai, L.** 1989b. Biological control of *Heterobasidion annosum* in silver fir (*Abies alba* Mill.) stands. In: Morrison, D.J. (Ed.) Proceedings of the 7<sup>th</sup> IUFRO Conference on Root and Butt Rots, Vernon and Victoria, British Columbia, Canada, 9–16 August 1988, p. 277–287.
- Carris, L.M. and Glawe, D.A.** 1989. Fungi colonizing cysts of *Heterodera glycines*. Bulletin 786, University of Illinois at Urbana-Champaign, College of Agriculture, Agricultural Experimental Station, U.S. Department of Agriculture, 93 pp.
- Drenkhan, T., Hanso, S. and Hanso, M.** 2008. Effect of stump treatment with *Phlebiopsis gigantea* against *Heterobasidion* root rot in Estonia. *Baltic Forestry* 14(1): 16–25.
- Gaitnieks, T., Bruna, L., Zaļuma, A., Burneviča, N., Klaviņa, D., Legzdina, L., Jansons, J. and Piri, T.** 2021. Development of *Heterobasidion* spp. fruit bodies on decayed *Picea abies*. *Forest Ecology and Management* 482: 118835. <https://doi.org/10.1016/j.foreco.2020.118835>.
- Garbelotto, M. and Gonthier, P.** 2013. Biology, epidemiology and control of *Heterobasidion* species worldwide. *Annual Review of Phytopathology* 51: 39–59. <https://doi.org/10.1146/annurev-phyto-082712-102225>.
- Gonthier, P. and Thor, M.** 2013. Annosus root and butt rot. In: Gonthier, P. and Nicolotti G. (Eds.). Infectious Forest Diseases. Wallingford: CAB International, p. 128–158.
- Hamberg, L., Korhonen, K. and Hantula, J.** 2012. Interaction between *Chondrostereum purpureum* and *Phlebiopsis gigantea* in mixed suspensions and their biocontrol efficacy in spruce logs and birch stumps. *Biocontrol Science and Technology* 22(10): 1239–1243. <https://doi.org/10.1080/09583157.2012.720673>.
- Hanso, M. and Hanso, S.** 1985. Poisk antagonistov kornevoi gubki I [Search for antagonists to *Heterobasidion annosum* (Fr.) Bref. I]. *Metsanduslikud uurimused / Forestry Studies* 20: 122–152 (in Russian with English and Estonian summaries).
- Hanso, M. and Hanso, S.** 1992. Poisk antagonistov kornevoi gubki II [Search for antagonists to *Heterobasidion annosum* (Fr.) Bref. II]. *Metsanduslikud uurimused / Forestry Studies* 25: 124–142 (in Russian with English and Estonian summaries).
- Holdenrieder, O.** 1982. Kristallbildung bei *Heterobasidion annosum* (Fr.) Bref. (*Fomes annosus* P. Karst.) und anderen holzbewohnenden Pilzen [Crystal formation in *Heterobasidion annosum* (Fr.) Bref. (*Fomes annosus* P. Karst.) and other wood-dwelling fungi.] *European Journal of Forest Pathology* 12: 41–58 (in German with English abstract). <https://doi.org/10.1111/j.1439-0329.1982.tb01371.x>.
- Holdenrieder, O.** 1984. Investigations on biological control of *Heterobasidion annosum* in Norway spruce with antagonistic fungi. I. Production of inoculum and interaction experiments *in vitro*. *European Journal of Plant Pathology* 141: 17–32 (in German with English summary).
- Holdenrieder, O. and Greig, B.J.W.** 1998. Biological methods of control. In: Woodward, S., Stenlid, J., Karjalainen, R. and Hüttermann, A. (Eds.) *Heterobasidion*: Biology, Ecology, Impact and Control. Wallingford / Cambridge, UK: CAB International / University Press, p. 235–258.
- Holmer, L. and Stenlid, J.** 1993. Biological control of *Heterobasidion annosum* by cord-forming basidiomycetes. In: Johansson, M. and Stenlid, J. (Eds.) 8<sup>th</sup> International Conference on Root and Butt Rots, Conference proceedings, 9–16 August 1993. Wik, Sweden and Haikko, Finland: p. 686–695.
- Holmer, L. and Stenlid, J.** 1996. Diffuse competition for heterogeneous substrate in soil among six species of wood-decomposing basidiomycetes. *Oecologia* 106: 531–538. <https://doi.org/10.1007/BF00329712>.
- Hypel, A.** 1968. Antagonistic effects of some soil fungi on *Fomes annosus* in laboratory experiments. *Studia forestalia Suecica* 64: 1–18.
- Kallio, T. and Hallaksela, A.-M.** 1979. Biological control of *Heterobasidion annosum* (Fr.) Bref. (*Fomes annosus*) in Finland. *European Journal of Forest Pathology* 9(5): 298–308.
- Kenigvalde, K., Brauners, I., Korhonen, K., Zaļuma, A., Mihailova, A. and Gaitnieks, T.** 2016. Evaluation of biological control agent Rotstop in controlling the infection of spruce and pine stumps by *Heterobasidion* in Latvia. *Scandinavian Journal of Forest Research* 31(3): 254–261. <https://doi.org/10.1080/02827581.2015.1085081>.

- Korhonen, K.** 1978. Intersterility groups of *Heterobasidion annosum*. *Communicationes Instituti Forestalis Fenniae* 94: 1–25.
- Korhonen, K.** 2003. Simulated stump treatment experiments for monitoring the efficacy of *Phlebiopsis gigantea* against *Heterobasidion annosum*. In: Laflamme, G., Bérubé, J.A. and Bussièrès, G. (Eds.) Root and butt rots of forest trees, Proceedings of the 10<sup>th</sup> International Conference on Root and Butt Rots, IUFRO Working Party 7.02.01, 16–22 September 2001, Quebec City. Canadian Forest Service, Information Report LAU-X-126, p. 206–210.
- Korhonen, K., Bobko, I., Hanso, S., Piri, T. and Vasiliuskas, A.** 1992. Intersterility groups of *Heterobasidion annosum* in some spruce and pine stands in Byelorussia, Lithuania and Estonia. *European Journal of Forest Pathology* 22(6–7): 384–391. <https://doi.org/10.1111/j.1439-0329.1992.tb00311.x>.
- Korhonen, K., Liponen, K., Bendz, M., Johansson, M., Ryen, I., Venn, K., Seiskari, P. and Niemi, M.** 1994. Control of *Heterobasidion annosum* by stump treatment with 'Rotstop', a new commercial formulation of *Phlebiopsis gigantea*. In: Johansson, M. and Stenlid, J. (Eds.) Root and butt rots. Proceedings of the IUFRO Working Party S2.06.01, Wik, Sweden and Haikko, Finland, 9–16 August 1993. Uppsala, Sweden: Swedish University of Agricultural Sciences, p. 675–685.
- Korhonen, K. and Piri, T.** 1994. The main hosts and distribution of the S and P groups of *Heterobasidion annosum* in Finland. In: Johansson, M. and Stenlid, J. (Eds.) Proceedings of the 8<sup>th</sup> International Conference on Root and Butt Rots, Wik, Sweden and Haikko, Finland, 9–16 August 1993. Uppsala, Sweden: SLU, Part 1, p. 260–267.
- Korhonen, K. and Stenlid, S.** 1998. Biology of *Heterobasidion annosum*. In: Woodward, S., Stenlid, J., Karjalainen, R. and Hüttermann, A. (Eds.) *Heterobasidion annosum*: Biology, ecology, impact and control. Wallingford: CAB International, p. 43–70.
- Liepa, I.** 1974. Biometrija [Biometrics]. Riga: Zvaigzne, 335 pp. (in Latvian).
- Lundborg, A. and Unestam, T.** 1980. Antagonism against *Fomes annosus*. Comparison between different test methods *in vitro* and *in vivo*. *Mycopathologia* 70: 107–115. <https://doi.org/10.1007/BF00443076>.
- Mgbeahuruike, A., Sun, H., Fransson, P., Kasanen, R., Daniel, G., Karlsson, M. and Asiegbu, F.** (2011). Screening of *Phlebiopsis gigantea* isolates for traits associated with biocontrol of the conifer pathogen *Heterobasidion annosum*. *Biological Control* 57(2): 118–129. <https://doi.org/10.1016/j.biocontrol.2011.01.007>.
- Morrison, D.J. and Johnson, A.L.S.** 1978. Stump colonization and spread of *Fomes annosus* 5 years after thinning. *Canadian Journal of Forest Research* 8(2): 177–180. <https://doi.org/10.1139/x78-028>.
- Muntyan, M.S., Morozov, D.A., Klishin, S.S., Khitrin, N.V. and Kolomijtseva, G.Ya.** 2012. Evaluation of the electrical potential on the membrane of the extremely alkaliphilic bacterium *Thioalkalivibrio*. *Biochemistry* 77(8): 917–924. <https://doi.org/10.1134/S0006297912080135>.
- NCBI 2022. The National Center for Biotechnology Information advances science and health by providing access to biomedical and genomic information. URL: <https://www.ncbi.nlm.nih.gov/>.
- Negrutskii, S.F.** 1986. Kornevaia gubka [Annosus but rot, *Heterobasidion annosum*]. 2<sup>nd</sup> ed. Moscow: AgroPromIzdat, 196 pp. (in Russian).
- Negrutskii, S.F., Zaporozhchenko, E.V. and Sukhomlin, M.N.** 1993. Taksonomicheskie issledovaniia *Heterobasidion annosum* (Fr.:Fr.) Bref. III. Tozhdestvo dvukh neperesekaiushchikhsia grupp s P- i S-gruppami [Taxonomical studies of *Heterobasidion annosum* (Fr.:Fr.) Bref. III. Identity of two non-crossing groups with P- and S-groups]. *Mikologiya i Fitopatologiya / Mycology and Phytopathology* 27(4): 30–33 (in Russian with English summary); ISSN: 0026-3648.
- Nicolotti, G., Gangemi, D., Lanata, F. and Anselmi, N.** 1993. Antagonistic activity of wood decay basidiomycetes against European *Armillaria* species. In: Johansson, M. and Stenlid, J. (Eds.) 8<sup>th</sup> International Conference on Root and Butt Rots, Conference proceedings, 9–16 August 1993. Wik, Sweden and Haikko, Finland: p. 725–735.
- Nicolotti, G. and Gonthier, P.** 2005. Stump treatment against *Heterobasidion* with *Phlebiopsis gigantea* and some chemicals in *Picea abies* stands in the western Alps. *Forest Pathology* 35: 365–374. <https://doi.org/10.1111/j.1439-0329.2005.00419.x>.
- Nicolotti, G., Gonthier, P. and Varese, G.C.** 1999. Effectiveness of some biocontrol and chemical treatments against *Heterobasidion annosum* on Norway spruce stumps. *European Journal of Forest Pathology* 29: 339–346. <https://doi.org/10.1046/j.1439-0329.1999.00159.x>.
- Pechmann, H. and Aufsess, H.** 1971. Study of the causes of stem decays in spruce stands. *Forstwissenschaftliches Centralblatt* 90: 259–284 (in German with English summary).
- Pellicciaro, M., Lione, G., Giordano, L. and Gonthier, P.** 2021a. Biocontrol potential of *Pseudomonas protegens* against *Heterobasidion* species attacking conifers in Europe. *Biological Control* 157: 104583. <https://doi.org/10.1016/j.biocontrol.2021.104583>.
- Pellicciaro, M., Lione, G., Ongaro, S. and Gonthier, P.** 2021b. Comparative efficacy of state-of-the-art and new biological stump treatments in forests infested by the native and the alien invasive *Heterobasidion* species present in Europe. *Pathogens* 10: 1272. <https://doi.org/10.3390/pathogens10101272>.
- Pratt, J.E.** 1998. Economical appraisal of the benefits of control treatments. In: Woodward, S., Stenlid, J., Karjalainen, R. and Hüttermann, A. (Eds.) *Heterobasidion annosum*: biology, ecology, impact and control. Wallingford: CAB International, p. 315–331.
- Pratt, J., Niemi, M. and Sierota, Z.** 2000. Comparison of three products based on *Phlebiopsis gigantea* for the control of *Heterobasidion annosum* in Europe. *Biocontrol Science and Technology* 10(4): 467–477. <https://doi.org/10.1080/09583150050115052>.
- Rishbeth, J.** 1970. The possibility of stump inoculations for conifers other than pines. In: Hodges, C.S., Rishbeth, J. and Yde-Anderson, A. (Eds.) Proceedings of the 3<sup>rd</sup> IUFRO Conference on *Fomes annosus*, July–August 1968, Denmark, Washington DC: USDA Forest Service, p. 110–120.
- Rönnerberg, J., Sidorov, E. and Petrylaite, E.** 2006. Efficacy of different concentrations of Rotstop® and Rotstop®S and imperfect coverage of Rotstop®S against *Heterobasidion* spp. spore infections on Norway spruce stumps. *Forest Pathology* 36(6): 422–433. <https://doi.org/10.1111/j.1439-0329.2006.00476.x>.
- Roy, G., Laflamme, G., Bussièrès, G. and Dessureault, M.** 2003. Field tests on biological control of *Heterobasidion annosum* by *Phaeothea dimorphospora* in comparison with *Phlebiopsis gigantea*. *Forest Pathology* 33: 127–140. <https://doi.org/10.1046/j.1439-0329.2003.00319.x>.
- Samils, N., Olson, Å. and Stenlid, J.** 2008. The capacity in *Heterobasidion annosum* s.l. to resist overgrowth by the biocontrol agent *Phlebiopsis gigantea* is a heritable trait. *Biological Control* 45: 419–426. <https://doi.org/10.1016/j.biocontrol.2008.03.010>.



- Stokland, J.N. and Larsson, K.-H.** 2011. Legacies from natural forest dynamics: Different effects of forest management on wood-inhabiting fungi in pine and spruce forests. *Forest Ecology and Management* 261(11): 1707–1721. <https://doi.org/10.1016/j.foreco.2011.01.003>.
- Sun, H., Korhonen, K., Hantula, J. and Kasanen, R.** 2009. Variation in properties of *Phlebiopsis gigantea* related to biocontrol against infection by *Heterobasidion* spp. in Norway spruce stumps. *Forest Pathology* 39(2): 133–144. <https://doi.org/10.1111/j.1439-0329.2008.00574.x>.
- Thor, M.** 2003. Operational stump treatment against *Heterobasidion annosum* in European forestry – current situation. In: Laflamme, G., Bérubé, J.A. and Bussièrès, G. (Eds.) Proceedings of the 10<sup>th</sup> IUFRO conference on root and butt rots of forest trees, 16–22 September 2001. Quebec (Canada): Laurentian Forestry Centre, p. 170–175.
- Thor, M. and Stenlid, J.** 2005. *Heterobasidion annosum* infection of *Picea abies* following manual or mechanized stump treatment. *Scandinavian Journal of Forest Research* 20(2): 154–164. <https://doi.org/10.1080/02827580510008338>.
- Vainio, E.J., Lipponen, K. and Hantula, J.** 2001. Persistence of a biocontrol strain of *Phlebiopsis gigantea* in conifer stumps and its effects on within-species genetic diversity. *Forest Pathology* 31(5): 285–295. <https://doi.org/10.1080/02827580510008338>.
- Varese, G.C., Gonthier, P. and Nicolotti, G.** 2003. Impact of biological and chemical treatments against *Heterobasidion annosum* on non-target micro-organisms. In: Laflamme, G., Bérubé, J.A. and Bussièrès, G. (Eds.) Proceedings of the 10<sup>th</sup> International Conference on Root and Butt Rots of Forest Trees, Working Party 7.02.01, Québec City, Canada, 16–22 September 2001. Natural Resources Canada, Canadian Forest Service, Laurentian Forestry Centre, p. 145–154.
- Vasaitis, R., Stenlid, J., Thomsen, I.M., Barklund, P. and Dahlberg, A.** 2008. Stump removal to control root rot in forest stands. A literature study. *Silva Fennica* 42(3): 457–483. <https://doi.org/10.14214/sf.249>.
- Vasiliauskas, A.** 1989. Kornevaia gubka i ustoichivost ekosistem khvoinykh lesov [Root and butt rot (*Heterobasidion annosum*) and sustainability of coniferous forest ecosystems]. Vilnius: Mokslas, 175 pp. (in Russian).
- Vasiliauskas, R., Jusbreveka, E., Vasiliauskas, A. and Stenlid, J.** 2002. Community of Aphyllophorales and root rot is stumps of *Picea abies* on clear-felled forest sites in Lithuania. *Scandinavian Journal of Forest Research* 17(5): 398–407.
- Vasiliauskas, R., Lygis, V., Larsson, K.-H. and Stenlid, J.** 2005. Airborne fungal colonisation of coarse woody debris in North Temperate *Picea abies* forest: impact of season and local spatial scale. *Mycological Research* 109(4): 487–496. <https://doi.org/10.1017/S0953756204002084>.
- Vasiliauskas, R., Lygis, V., Thor, M. and Stenlid, J.** 2004. Impact of biological (Rotstop) and chemical (urea) treatments on fungal community structure in freshly cut *Picea abies* stumps. *Biological Control* 31(3): 405–413. <https://doi.org/10.1016/j.biocontrol.2004.05.006>.
- Webber, J. and Thorpe, K.** 2003. Potential for biological control of *Heterobasidion annosum* in the UK using Rotstop® – In: Laflamme, G., Bérubé, J.A. and Bussièrès, G. (Eds.) Root and butt rots of forest trees. Proceedings of the 10<sup>th</sup> International Conference on Root and Butt Rots, IUFRO Working Party 7.02.01, 16–22 September 2001, Quebec City. Canadian Forest Service, Information Report LAU-X-126, p. 221–225.
- Woodward, S., Stenlid, J., Karjalainen, R. and Hüttermann, A.** 1998. Preface. In: Woodward, S., Stenlid, J., Karjalainen, R. and Hüttermann, A. (Eds.) *Heterobasidion annosum: Biology, Ecology, Impact and Control*. Wallingford-New York: CAB International, p. xi–xii.
- Woods, C.M., Woodward, S. and Redfern, D.B.** 2005. *In vitro* interactions in artificial and wood-based media between fungi colonizing stumps of Sitka spruce. *Forest Pathology* 35: 213–229. <https://doi.org/10.1111/j.1439-0329.2005.00403.x>.
- Zaļuma, A., Muižnieks, I., Gaitnieks, T., Burņeviča, N., Jansons, Ā., Jansons, J., Stenlid, J. and Vasaitis, R.** 2019. Infection and spread of root rot caused by *Heterobasidion* spp. in *Pinus contorta* plantations in Northern Europe: three case studies. *Canadian Journal of Forest Research* 49(8): 969–977.