

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Re-discovery of *Trichophyton bullosum* in North Africa as a cause of severe dermatophytosis in donkeys

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1852716> since 2022-04-08T09:50:05Z

Published version:

DOI:10.1007/s12223-021-00930-9

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 **Re-discovery of *Trichophyton bullosum* in North Africa as a cause of severe**
2 **dermatophytosis in donkeys**

3

4 **Andrea Peano**¹ (ORCID: 0000-0002-6783-4742), **Sara Arnoldi**¹, **Adéla Čmoková**^{2,3}
5 (ORCID: 0000-0002-6783-4742), **Vit Hubka**^{2,3} (ORCID: 0000-0003-4583-6496)

6

7 ¹Department of Veterinary Sciences, , University of Turin, Largo Paolo Braccini 2, 10095
8 Grugliasco (Turin), Italy

9 ²Department of Botany, Faculty of Science, Charles University, Benátská 2, 128 01 Prague 2,
10 Czech Republic

11 ³Laboratory of Fungal Genetics and Metabolism, Institute of Microbiology of the Academy of
12 Sciences of the Czech Republic, v. v. i., Vídeňská 1083, 142 20 Prague 4, Czech Republic

13

14 **ABSTRACT**

15 This article reports the first verified cases of infection by *Trichophyton bullosum* in Africa since
16 the description of the fungus, isolated in 1933 from the coat of horses in Tunisia and Mali. We
17 re-discovered the fungus in cutaneous samples obtained from donkeys suffering for severe
18 dermatitis with areas of alopecia and scaling in the surroundings of Cairo (Egypt). Fungal
19 elements (arthroconidia and hyphae) were seen at the microscopy of material collected by skin
20 scraping and digested in NaOH. Fungal colonies grown on various culture media were
21 identified through PCR and sequencing of the ITS rDNA region. Since the original report in
22 Africa and the Middle East, only a few cases have been reported thus far in humans in France
23 and two cases in horses in the Czech Republic and Japan. *Trichophyton bullosum* seems thus
24 an infrequent cause of dermatophytosis. However, the actual prevalence of this pathogen may
25 be underestimated due to the similarity with *T. verrucosum*, a diffuse cause of infection in cattle

1 and humans, occasionally found on horses and donkeys too. Indeed, the two fungi can be
2 distinguished only via molecular methods, which are poorly employed in epidemiological
3 studies on equine and bovine dermatophytosis. The results of the present study add to our
4 knowledge on the ecology of this poorly explored dermatophyte, reinforcing the idea that
5 equines are the main hosts of *T. bullosum*, and confirming the presence of this pathogen in
6 Africa.

7

8 **Keywords**

9 *Equidae*, molecular identification, animal skin diseases, *Trichophyton benhamiae* complex,
10 zoophilic dermatophytes, zoonotic mycoses

11

1 **INTRODUCTION**

2 *Trichophyton bullosum* is a poorly known zoophilic dermatophyte that was originally
3 described from the coat of horses in Africa (Tunisia and Mali) and the Middle East (Syria)
4 (Lebasque 1933). The species was almost forgotten for decades, but recently, the use of the
5 species name was justified using the sequence data and phylogeny (Heidemann et al. 2010).
6 The morphology of this slow-growing pathogen is very similar to that of *Trichophyton*
7 *verrucosum*, and reliable identification can only be achieved through molecular methods
8 (Sitterle et al. 2012; Lysková et al. 2015).

9 Horses and donkeys are supposed to be the main hosts of *T. bullosum*, in contrast to *T.*
10 *verrucosum*, with cattle as the primary host. However, infections in horses and donkeys can be
11 caused by both mentioned species (Lysková et al. 2015). The distribution of *T. bullosum* is
12 almost unknown due to the meagre number of reported cases in both animals and humans and
13 the lack of comprehensive studies on the epidemiology of dermatophytosis in cattle, horses,
14 and donkeys using molecular methods.

15 Eighty years after discovering *T. bullosum* in Africa and the Middle East, the first
16 cases were described in Europe and Asia. Two cases of human dermatophytosis were reported
17 in France (Sitterle et al. 2012; Sabou et al. 2018), but no sample was collected from the source
18 animals. The only two cases of verified animal dermatophytosis were reported in horses from
19 the Czech Republic (Lysková et al. 2015) and Japan (Watanabe et al. 2021).

20 This article reports some cases of severe dermatophytosis caused by *T. bullosum* in
21 donkeys from Egypt. These cases are the first verified infections in Africa since the
22 description of *T. bullosum* in 1933. They add to our knowledge of this poorly explored
23 dermatophyte ecology and confirm its continuing occurrence in Northern Africa.

24

25

1 MATERIAL AND METHODS

2 *Animals and sampling site*

3 The findings presented in this article take origin from experience made by one of the authors
4 (SA) within a training program on the medicine of donkeys under the guidance of local
5 Veterinarians operating for “The Donkey Sanctuary” (www.thedonkeysanctuary.org.uk).

6 Donkeys in Egypt are a common means of transport for goods and people. Though these
7 animals represent a fundamental resource for many families, they frequently receive poor or
8 null veterinary care due to the lack of financial resources and availability of veterinary services.
9 To address this criticism, since 2002, the Donkey Sanctuary serves through mobile clinics in
10 villages, offering free veterinary counselling and interventions to donkey owners. SA noticed
11 many animals with skin lesions during her practical activities and collected biological samples
12 to assess a possible fungal aetiology. The animals covered in this study were visited in August
13 2015 and April 2016 during different working sessions in villages in the surroundings of Cairo.

15 *Microbiological methods*

16 Hair and scales from the donkeys presenting dermatological lesions were collected using sterile
17 scalpel blades through superficial skin scrapings. The samples were digested in 10% NaOH for
18 3 hours at room temperature. The material was then transferred to slides and observed under
19 the microscope for the presence of fungal elements. Primary cultures were performed on a
20 commercial medium selective for dermatophytes (Mycobios Selective Agar, Biolife, Milan,
21 Italy), on Sabouraud Dextrose agar enriched with thiamine and inositol and on a medium with
22 4% casein and 0.5% yeast extract (Kane and Smitka 1978). All the media included antibacterial
23 agents and cycloheximide to reduce the growth of contaminant organisms.

24 Subcultures were grown on Sabourad’s dextrose agar (SGA, Himedia, Milano, Italy),
25 malt extract agar (MEA; malt extract from Oxoid, Basingstoke, UK) and potato dextrose agar

1 (PDA; Roth, Karlsruhe, Germany) at 25 °C in the dark; growth at 30 °C and 37 °C was tested
2 on SGA. Lactic acid with cotton blue was used as a mounting medium for observation of the
3 micromorphology. Preparations for light microscopy were made from 14–21 days old cultures
4 grown on MEA. Lactic acid (60 %) was used as a mounting medium. Microphotographs were
5 taken by Olympus BX51 microscope with an Olympus DP72 camera. Colony details were
6 observed and photographed on Olympus SZX2-ILLT (Tokyo, Japan) equipped with
7 microscope digital camera Olympus DP27 (Tokyo, Japan). The isolates were deposited into the
8 Culture Collection of Fungi (CCF) housed at the Department of Botany, Charles University,
9 Czech Republic.

10

11 *Molecular studies*

12 Genomic DNA was extracted from 14-day-old colonies grown on MEA (Oxoid Ltd.,
13 Basingstoke, UK) using the ArchivePure DNA yeast and Gram2 + kit (5PRIME Inc.,
14 Gaithersburg, MD, USA) according to the updated manufacturer's instructions (Hubka et al.
15 2018b). The ITS rDNA region was amplified with the primers ITS1 (5'-
16 TCCGTAGGTGAACCTGCGG) and ITS4 (5'-TCCTCCGCTTATTGATATGC) (White et al.
17 1990). The reaction mixture and PCR protocol were described previously (Hubka et al. 2018a).
18 The PCR product purification followed the protocol described by Sklenář et al. (Sklenář et al.
19 2021); sequencing was performed in Seqlab Sequencing Service (Charles University, Czech
20 Republic) using both terminal primers. The obtained sequences were deposited into the ENA
21 (European Nucleotide Archive) database.

22 The primer pair MF3 and MF6 was used for the detection of the MAT1-1-1 idiomorph
23 of the mating type gene (MAT) encoding protein with an alpha-domain motif, and primers MF1
24 and MF5 were used for the detection of the MAT1-2-1 idiomorph encoding a regulatory protein

1 with a HMG (high-mobility group) DNA-binding motif (Kano et al. 2012; Symoens et al.
2 2013).

3 Alignments of the ITS was performed using the FFT-NS-i option implemented in the
4 MAFFT online (Kato et al. 2019). The final dataset contained 33 taxa and a total of 667
5 characters of which 90 were variable and 51 parsimony-informative. Suitable partitioning
6 schemes and substitution models (Bayesian information criterion) were selected using
7 PartitionFinder 2 (Lanfear et al. 2017) with settings allowing ITS1, 5.8S and ITS2 regions to
8 be independent datasets. Suitable partitioning scheme and substitution models were as follows:
9 the TrNef model was proposed for the ITS1 and ITS2 regions; and K80 model for the 5.8S
10 region. The maximum likelihood tree was constructed with IQ-TREE version 1.4.4 (Nguyen et
11 al. 2015) with nodal support determined by nonparametric bootstrapping with 1000 replicates.

12

13 **RESULTS**

14 *Animals*

15 Out of 187 animals examined, 35 (18,7%) presented dermatological problems.

16 *Direct examination and primary cultures*

17 Samples from three donkeys were positive for fungal elements at microscopy. The animals were
18 adults (Table 1) belonging to different owners and visited on different days in Abu Ghaleb,
19 close to Cairo. The lesions were represented by multiple areas of alopecia localized in various
20 body sites (Table 1, Fig. 1, Supplementary Fig. S1).

21 The fungal structures observed were chains/groups of arthroconidia inside or surrounding the
22 hair shafts (Fig. 2). In the cultures of three donkeys (Nos. 1, 2, and 3) it was possible to observe
23 pale, waxy, slow-growing colonies, which were subcultured for identification. As regards
24 donkey No. 4, despite numerous attempts, the fungus seen at direct examination failed to grow,
25 with the culture plates rapidly invaded by contaminant moulds.

1 *Mycological studies*

2 In total, six isolates from three donkeys were further characterized. One isolate from each
3 culture-positive animal was deposited into the CCF collection with accession codes CCF 5730
4 (donkey No.1), CCF 5731 (donkey No. 2) and CCF 5678 (donkey No. 3). The description of
5 subcultures cultivated at 25 °C in the dark is reported below and shown in Fig. 3.

6 Colonies on MEA attained 8–12 mm diam after 7 days (10–13 mm after 14 days and
7 15–17 mm after 21 days), white to pale yellow (#F3E5AB) or vivid orange-yellow (#F6A600),
8 waxy, slightly elevated, margins entire, reverse light yellow (#F8DE7E). Colonies on SGA
9 attained 11–12 mm diam after 7 days (20–21 mm after 14 days and 28–33 mm after 21 days),
10 white to pale yellowish pink (#ECD5C5) or pale orange-yellow (#FAD6A5), submerged,
11 filamentous, reverse light yellow (#F8DE7E). Colonies on PDA attained 7–9 mm diam after 7
12 days (14–16 mm after 14 days and 25–29 mm after 21 days), white to pale yellowish pink
13 (#ECD5C5) or pale orange yellow (#FAD6A5), waxy, lanuginous in some areas, elevated in
14 the centre, margins entire to delicately filamentous, reverse light yellow (#F8DE7E). Colonies
15 on SGA with cycloheximide and chloramphenicol (Trios, Prague, Czech Republic) were
16 submerged with dendritic growth. Colonies on SGA after 7 d at 30 °C attained 11–14 mm diam
17 ($\varnothing = 13$ mm); at 37 °C 8–9 mm diam ($\varnothing = 8$ mm).

18 *Vegetative hyphae* were smooth, septate, frequently inflated, hyaline, 1.5–4 μ m diam
19 (Mean \pm sd; 2.7 \pm 0.7). Chlamydospores abundantly present, spherical, ovate or irregularly
20 shaped, 4–9(–20) μ m diam, frequently forming chains. Microconidia, macroconidia and spiral
21 hyphae were not observed.

22

23 *Molecular studies*

24 The MAT1-1-1 idiomorph of the mating type gene was amplified in all six *T. bullosum* strains
25 with primers MF3 and MF6, while the idiomorph MAT1-2-1 was not detected. The ITS

1 rDNA sequences obtained from all strains were identical. Three sequences representing
2 isolates preserved in the CCF collection are available under accession numbers LR595961–
3 LR595963 (Table 1). Using the BLAST similarity search, these sequences were identical to
4 those derived from strains of *T. bullosum* isolated from human tinea infections in France
5 (strains IHEM 24321 and 0805m150877; GenBank accession numbers: MK298921 and
6 KY885205), horse in the Czech Republic (isolate CCF 4831; LR794144), ex-type strain of *T.*
7 *bullosum* (CBS 363.35; FM992675), and strain CBS 557.50 (FM992675) from an unknown
8 source. The strain NUBS 20002 from a horse in Japan (LC592175) showed one substitution
9 in the ITS2 region compared to our strains.

10 Some sequences deposited in GenBank under the name *T. bullosum* represent in fact a
11 recently described species, namely *T. africanum*, especially some generated in the study of
12 Baert et al. (2020). In the phylogeny based on the ITS rDNA sequences (Fig. 4), this species is
13 resolved as a sister species to *T. bullosum*. Although the majority of known *T. africanum* records
14 originate from human dermatophytoses in Mozambique (IHEM 4032), South Africa (IHEM
15 4033) or Belgium (IHEM 19628), it is probably a zoophilic dermatophytes as evidenced by
16 strains IHEM 19638 and IHEM 17561 isolated from horse in South Africa and guinea pig in
17 France, respectively.

18

19 **DISCUSSION**

20 *Trichophyton bullosum* is a member of the *Trichophyton benhamiae* complex along with other
21 ten predominantly zoophilic pathogens; the only exception is an anthropophilic species, *T.*
22 *concentricum* (Čmoková et al. 2021). *Trichophyton verrucosum*, a dominant causal agent of
23 dermatophytosis in cattle and other ruminants, belongs to the most well-known members of
24 this complex (Chermette et al. 2008). Some other representatives, especially *T. benhamiae*, *T.*
25 *europaeum*, *T. japonicum* and *T. erinacei*, have also received considerable attention in recent

1 years due to the increasing incidence of infections in pets and pet owners, most noticeably in
2 the European countries (Nenoff et al. 2014; Abarca et al. 2017; Hubka et al. 2018c; Le Barzic
3 et al. 2021). Compared to these extensively studied pathogens, little is known about the host
4 spectrum of the remaining zoophilic species, which are usually known only from several case
5 reports in humans and animals.

6 *Trichophyton bullosum* is phylogenetically relatively distant from all other species in
7 the *T. benhamiae* complex. It is most closely related to *T. africanum* (Fig. 4), which is
8 strikingly different by morphology (Čmoková et al. 2020). It is also phylogenetically distant
9 from *T. verrucosum*, which is superficially and microscopically very similar, and it has
10 similar nutritional requirements and growth pattern *in vitro* (Lysková et al. 2015; Watanabe et
11 al. 2021). Both species show a low level of intraspecific genetic variability, and their
12 populations most likely consist only of strains displaying one mating type gene (MAT)
13 idiomorph. The MAT idiomorph, however, differs between these pathogens. Only *MAT1-1-1*
14 idiomorph has been detected in *T. bullosum* strains, while *MAT1-2-1* was detected in all
15 hitherto examined *T. verrucosum* strains (Kano et al. 2014; Kosanke et al. 2018; Čmoková et
16 al. 2020). These facts suggest that clonal reproduction is dominant or the only mode of
17 dissemination of these species. Because there is no reliable phenotypic character
18 distinguishing these species, molecular methods are necessary to for the identification.

19 Apart from molecular and phylogenetic differences, the main difference between *T.*
20 *bullosum* and *T. verrucosum*, therefore, lies in their host spectrum, which can, however, partly
21 overlap. Based on the known data, *T. bullosum* is specific for horses and donkeys, while *T.*
22 *verrucosum* is mainly reported from ruminants but occasionally from other animals, including
23 horses and donkeys. Lysková et al. (2015) hypothesized that a significant part of
24 dermatophytosis cases in horses and donkeys due to *T. verrucosum* could be caused by *T.*
25 *bullosum* due to the overall lack of large-scale studies examining agents of dermatophytosis in

1 these animals using molecular methods. Consequently, this is why we do not have exact data
2 on geographical distribution, host spectrum and prevalence of *T. bullosum*.

3 Probable endemic areas of *T. bullosum* are Africa and the Middle East, where the
4 species was initially found. Our findings confirm its occurrence at present in Africa.

5 Lysková et al. (2015) suggested that this species could be introduced to Europe with
6 the animal trade between France and former French colonies and then spread to other
7 countries, including the Czech Republic. In this regard, the case discovered in a horse in
8 Japan could represent a significant turning point in this thinking (Watanabe et al. 2021). This
9 case may indicate the introduction of a pathogen to remote areas along with animal transport
10 and trade. Still, on the other hand, it can also show the worldwide distribution of an
11 underdiagnosed pathogen.

12 Clinical manifestation of infection due to *T. bullosum* in donkeys has not been
13 described except in the present study. Thus the only well-documented animal cases due to this
14 pathogen come from horses and, in general, manifested as circular erythematous lesions with
15 hair loss in the saddle area, shoulders, chest, withers, and less commonly on the head, neck,
16 buttocks and limbs (Lebasque 1933; Lebasque 1934; Lysková et al. 2015; Watanabe et al.
17 2021). There are few well-documented cases of dermatophytosis in donkeys due to *T.*
18 *verrucosum* (Suleiman et al. 2017) or *T. verrucosum*-like isolates (Abdalla et al. 2005), which
19 could in fact represent *T. bullosum*, while the remaining records lack a detailed clinical
20 description. In agreement with the cases described here, these infections predominantly
21 manifested as generalized superficial infections with non-itchy, slightly elevated, well-
22 demarcated lesions accompanied by alopecia. The lesions were mainly distributed on the
23 head, neck, limbs and flank.

24 Dermatophytosis in equids is usually a self-limiting disease that resolves
25 spontaneously within several months (Rochette et al. 2003; Knottenbelt 2005; Cafarchia et al.

1 2013). However, the treatment is mandatory as it shortens disease duration and reduces the
2 possibility of spread to other animals or humans. The decontamination of the environment
3 using hypochlorite bleach and enilconazole sprays can further prevent re-infection and spread
4 of the disease (Rochette et al. 2003). Enilconazole and natamycin are registered in most
5 European countries for the treatment of dermatophytosis in Equids. These drugs are usually
6 employed in topical washes and sprays with satisfactory results (Rochette et al. 2003).
7 Although the data about efficacy are mostly based on the therapy of horse dermatophytosis,
8 enilconazole washes also show good results in treating dermatophytosis in donkeys (White et
9 al. 2019). Dermatophyte vaccine can be used as an alternative treatment option in donkeys,
10 leading to resolution (White et al. 2019).

11 Available data on treatment of verified *T. bullosum* infections include only three cases,
12 two in horses and one in man. The first horse was successfully treated with topical
13 enilconazole (Lysková et al. 2015) and the second with terbinafine ointment (Watanabe et al.
14 2021). Tinea corporis infection in a 21-year-old male was successfully treated by a
15 combination of topical ketoconazole and systemic griseofulvin therapy (Sitterle et al. 2012).
16 Regarding the donkeys of the present study, it was not possible to set up a therapeutic
17 protocol. It is essential to remind that the study context was that of a rural area characterized
18 by extreme poverty. Facing the commitment and expense of antifungal treatment in a situation
19 of subsistence rearing is difficult to accept. There are also cultural reasons behind this way of
20 thinking. The role of donkeys in human activities in many African countries, although
21 fundamental, is frequently overshadowed in the collective imagination. At the same time,
22 more attention and resources are dedicated to animals destined for food production. Popular
23 beliefs have often built a distorted vision of donkeys, considered stubborn and able to live and
24 work without any care.

1 The work context was problematic also concerning the communication with the donkeys'
2 owners, who generally provided only basic data regarding animal history. Unfortunately, this
3 made it impossible to investigate a crucial aspect: whether and to what extent in the area of
4 study the contact with donkeys causes infection by *T. bullosum* in people.

5 In conclusion, this study allowed demonstrating *T. bullosum* as an agent of dermatomycosis in
6 Northern Africa 90 years after its first report. Further studies will be necessary to ascertain the
7 involvement of animal species other than donkeys and the role of this fungus as a human
8 pathogen.

9

10 **ACKNOWLEDGEMENTS**

11 The contribution of Vit Hubka was supported by the Charles University Research Centre
12 program no. 204069.

1 **FIGURE LEGENDS**

2 **Fig. 1** Clinical manifestation of dermatophytosis due to *Trichophyton bullosum* in donkeys. a-
3 d: donkey No. 1; e-h: donkey No. 2; i-l: donkey No. 3. A detailed description of the lesions is
4 provided in Table 1.

5
6 **Fig. 2** a-c: hair infected with arthroconidia (microscopic examination after NaOH digestion,
7 250x magnification). a: donkey No. 1; b: donkey No. 2; c: donkey No. 4; d-e macromorphology
8 of primary cultures; d: donkey No. 1, culture on Sabouraud Dextrose agar with thiamine and
9 inositol (red arrows = contaminant moulds; blue arrows = *T. bullosum*); e: donkey No. 1 (culture
10 on casein medium); f: donkey No. 2 (culture on casein medium); g: donkey No. 3 (culture on
11 Mycobios selective agar).

12
13 **Fig. 3** Macromorphology and micromorphology of *Trichophyton bullosum*. a. Colonies after
14 14 days at 25 °C on MEA, SGA and PDA (from left to right). b. detail of colony on MEA. c.
15 detail of colony on SGA. d. detail of colony on PDA. e-f. colony on SGA with cycloheximide
16 and chloramphenicol (Trios, Prague, Czech Republic) after 3 months of incubation at 25 °C.
17 g-k. thick-walled vegetative hyphae disintegrating into arthroconidia; numerous thick-walled
18 irregular intercalary, terminal or free chlamydospores. Scale bars = 10 µm.

19
20 **Fig. 4** A best scoring maximum likelihood tree based on ITS rDNA sequences showing
21 relationships of *Trichophyton bullosum* to other members of the the *T. benhamiae* complex.
22 The main primary host(s) of particular species are shown as icons; uncommon occasional
23 hosts are omitted; a question mark means that host spectrum is little known and only based
24 one single or few isolations from animals. Ex-type isolates are designated by a superscript T;

1 only support values exceeding bootstrap values 70% are shown on branches; *Trichophyton*
2 *rubrum* CBS 202.88 was used as outgroup.

3

4 **Supplementary Fig. S1** Donkey No. 4. One large, irregular, alopecic patch with scaling on
5 the head. Another smaller area on the face (close to the nares). Some little alopecic areas on
6 the pinna. Large confluent alopecic areas on the limbs.

7

8 REFERENCES

- 9 Abarca M, Castellá G, Martorell J, Cabañes F (2017) *Trichophyton erinacei* in pet hedgehogs
10 in Spain: occurrence and revision of its taxonomic status. *Med Mycol* 55:164-172
- 11 Abdalla W, Suliman E, Gabbar A (2005) A report on *Trichophyton verrucosum* in donkeys in
12 the Sudan. *Sudan J Vet Res* 20:83-85
- 13 Baert F, Stubbe D, D'hooge E, Packeu A, Hendrickx M (2020) Updating the taxonomy of
14 dermatophytes of the BCCM/IHEM Collection according to the new standard: a
15 phylogenetic approach. *Mycopathologia* 185:161-168
- 16 Cafarchia C, Figueredo LA, Otranto D (2013) Fungal diseases of horses. *Vet Microbiol*
17 167:215-234
- 18 Čmoková A, Kolařík M, Dobiáš R, Hoyer LL, Janouškovcová H, Kano R, Kuklová I,
19 Lysková P, Machová L, Maier T, Mallátová N, Man M, Mencl K, Nenoff P, Peano A,
20 Prausová H, Stubbe D, Uhrlaß S, Větrovský T, Wiegand C, Hubka V (2020)
21 Resolving the taxonomy of emerging zoonotic pathogens in the *Trichophyton*
22 *benhamiae* complex. *Fungal Divers* 104:333-387
- 23 Čmoková A, Rezaei-Matehkolaei A, Kuklová I, Kolařík M, Shamsizadeh F, Ansari S,
24 Gharaghani M, Miňovská V, Najafzadeh MJ, Nouripour-Sisakht S, Yaguchi T,
25 Zomorodian K, Zarrinfar H, Hubka V (2021) Discovery of new *Trichophyton*
26 members, *T. persicum* and *T. spiraliforme* spp. nov., as a cause of highly
27 inflammatory tinea cases in Iran and Czechia. *Microbiol Spectr*: accepted
- 28 Heidemann S, Monod M, Gräser Y (2010) Signature polymorphisms in the internal
29 transcribed spacer region relevant for the differentiation of zoophilic and
30 anthropophilic strains of *Trichophyton interdigitale* and other species of *T.*
31 *mentagrophytes* sensu lato. *Brit J Dermatol* 162:282-295
- 32 Hubka V, Barrs V, Dudová Z, Sklenář F, Kubátová A, Matsuzawa T, Yaguchi T, Horie Y,
33 Nováková A, Frisvad J (2018a) Unravelling species boundaries in the *Aspergillus*
34 *viridinutans* complex (section *Fumigati*): opportunistic human and animal pathogens
35 capable of interspecific hybridization. *Persoonia* 41:142-174

- 1 Hubka V, Nováková A, Jurjević Ž, Sklenář F, Frisvad JC, Houbraken J, Arendrup MC,
2 Jørgensen KM, Siqueira JP, Gené J, Kolařík M (2018b) Polyphasic data support the
3 splitting of *Aspergillus candidus* into two species; proposal of *Aspergillus dobrogensis*
4 sp. nov. Int J Syst Evol Microbiol 68:995–1011
- 5 Hubka V, Peano A, Cmokova A, Guillot J (2018c) Common and emerging dermatophytoses
6 in animals: well-known and new threats. In: Seyedmousavi S, de Hoog GS, Guillot J,
7 Verweij PE (eds) Emerging and Epizootic Fungal Infections in Animals. Springer,
8 Cham, pp 31-79
- 9 Chermette R, Ferreira L, Guillot J (2008) Dermatophytoses in animals. Mycopathologia
10 166:385–405
- 11 Kane J, Smitka C (1978) Early detection and identification of *Trichophyton verrucosum*. J
12 Clin Microbiol 8:740-747
- 13 Kano R, Kawasaki M, Mochizuki T, Hiruma M, Hasegawa A (2012) Mating genes of the
14 *Trichophyton mentagrophytes* complex. Mycopathologia 173:103-112
- 15 Kano R, Yoshida E, Yaguchi T, Hubka V, Anzawa K, Mochizuki T, Hasegawa A, Kamata H
16 (2014) Mating type gene (MAT1-2) of *Trichophyton verrucosum*. Mycopathologia
17 177:103–112
- 18 Katoh K, Rozewicki J, Yamada KD (2019) MAFFT online service: multiple sequence
19 alignment, interactive sequence choice and visualization. Brief Bioinform 20:1160-
20 1166
- 21 Knottenbelt D (2005) Skin disorders of donkeys. In: Matthews N, Taylor T (eds) Veterinary
22 Care of Donkeys. International Veterinary Information Service, Ithaca, p A2918.0605
- 23 Kosanke S, Hamann L, Kupsch C, Garcia SM, Chopra A, Gräser Y (2018) Unequal
24 distribution of the mating type (MAT) locus idiomorphs in dermatophyte species.
25 Fungal Genet Biol 118:45-53
- 26 Lanfear R, Frandsen PB, Wright AM, Senfeld T, Calcott B (2017) PartitionFinder 2: new
27 methods for selecting partitioned models of evolution for molecular and
28 morphological phylogenetic analyses. Mol Biol Evol 34:772–773
- 29 Le Barzic C, Cmokova A, Denaes C, Arné P, Hubka V, Guillot J, Risco-Castillo V (2021)
30 Detection and control of dermatophytosis in wild European hedgehogs (*Erinaceus*
31 *europaeus*) admitted to a french wildlife rehabilitation centre. J Fungi 7:74
- 32 Lebasque J (1933) Les champignons des teignes du cheval et des bovidés. Dissertation,
33 Faculté des Sciences de Paris
- 34 Lebasque J (1934) Recherches morphologiques et biologiques sur les *Trichophyton*
35 mégasporés du cheval et du boeuf. Annales de Parasitologie 12:418–444
- 36 Lysková P, Hubka V, Petříčáková A, Dobiáš R, Čmoková A, Kolařík M (2015) Equine
37 dermatophytosis due to *Trichophyton bullosum*, a poorly known zoophilic
38 dermatophyte masquerading as *T. verrucosum*. Mycopathologia 180:407–419

- 1 Nenoff P, Uhrlaß S, Krüger C, Erhard M, Hipler UC, Seyfarth F, Herrmann J, Wetzig T,
2 Schroedl W, Gräser Y (2014) *Trichophyton* species von *Arthroderma benhamiae* – a
3 new infectious agent in dermatology. J Dtsch Dermatol Ges 12:571-582
- 4 Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ (2015) IQ-TREE: A fast and effective
5 stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol
6 32:268–274
- 7 Rochette F, Engelen M, Vanden Bossche H (2003) Antifungal agents of use in animal health–
8 practical applications. J Vet Pharmacol Ther 26:31–53
- 9 Sabou M, Denis J, Boulanger N, Forouzanfar F, Glatz I, Lipsker D, Poirier P, Candolfi E,
10 Letscher-Bru V (2018) Molecular identification of *Trichophyton benhamiae* in
11 Strasbourg, France: a 9-year retrospective study. Med Mycol 56:723–734
- 12 Sitterle E, Frealle E, Foulet F, Cabaret O, Cremer G, Guillot J, Delhaes L, Botterel F (2012)
13 *Trichophyton bullosum*: a new zoonotic dermatophyte species. Med Mycol 50:305–
14 309
- 15 Sklenář F, Jurjević Ž, Houbraken J, Kolařík M, Arendrup MC, Jørgensen KM, Siqueira JPZ,
16 Gené J, Yaguchi T, Ezekiel CN, Pereira CS, Hubka V (2021) Re-examination of
17 species limits in *Aspergillus* section *Flavipedes* using advanced species delimitation
18 methods and proposal of four new species. Stud Mycol: accepted
- 19 Suleiman EA, Abdalla WG, Ahmed AH, Abdelgabar MA (2017) Isolation and molecular
20 characterization of dermatophytes in donkeys. International Journal of Tropical
21 Disease & Health 26:1-8
- 22 Symoens F, Jousson O, Packeu A, Fratti M, Staib P, Mignon B, Monod M (2013) The
23 dermatophyte species *Arthroderma benhamiae*: intraspecies variability and mating
24 behaviour. J Med Microbiol 62:377-385
- 25 Watanabe R, Furuta H, Ueno Y, Nukada T, Niwa H, Shinyashiki N, Kano R (2021) First
26 isolation of *Trichophyton bullosum* from a horse with dermatophytosis in Japan. Med
27 Mycol Case Rep 32:81-83
- 28 White SD, Bourdeau PJ, Brément T, Vandenabeele SI, Haspeslagh M, Bruet V, Sloet van
29 Oldruitenborgh-Oosterbaan MM (2019) Skin disease in donkeys (*Equus asinus*): a
30 retrospective study from four veterinary schools. Vet Dermatol 30:247-e276
- 31 White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal
32 ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, J. SJ, White TJ
33 (eds) PCR protocols: a guide to methods and applications. Academic Press, San
34 Diego, pp 315–322
- 35
- 36