

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Overview of potential determinants of radical prostatectomy versus radiation therapy in management of clinically localized prostate cancer: results from an Italian, prospective, observational study (the Pros-IT CNR study)

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1799981> since 2021-09-08T13:44:21Z

Published version:

DOI:10.23736/S0393-2249.19.03637-3

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

ARTICLE ONLINE FIRST

This provisional PDF corresponds to the article as it appeared upon acceptance.

A copyedited and fully formatted version will be made available soon.

The final version may contain major or minor changes.

Overview of potential determinants of radical prostatectomy versus radiation therapy in management of clinically localized prostate cancer: results from an Italian, prospective, observational study (the Pros-IT CNR study)

ALESSANDRO ANTONELLI, Carlotta PALUMBO, Marianna NOALE, Walter ARTIBANI, PierFrancesco BASSI, Filippo BERTONI, Sergio BRACARDA, Alessio BRUNI, Renzo CORVò, Mauro GACCI, STEFANO MARIA MAGRINI, Rodolfo MONTIRONI, Angelo PORRECA, Andrea TUBARO, Vittorina ZAGONEL, Stefania MAGGI, and the Pros-IT CNR study group

Minerva Urologica e Nefrologica 2020 Jan 07

DOI: 10.23736/S0393-2249.19.03637-3

Article type: Original article

© 2019 EDIZIONI MINERVA MEDICA

Article first published online: January 7, 2020

Manuscript accepted: December 12, 2019

Manuscript revised: December 5, 2019

Manuscript received: September 5, 2019

Subscription: Information about subscribing to Minerva Medica journals is online at:

<http://www.minervamedica.it/en/how-to-order-journals.php>

Reprints and permissions: For information about reprints and permissions send an email to:

journals.dept@minervamedica.it - journals2.dept@minervamedica.it - journals6.dept@minervamedica.it

Overview of potential determinants of radical prostatectomy versus radiation therapy in management of clinically localized prostate cancer: results from an Italian, prospective, observational study (the Pros-IT CNR study)

Authors

Alessandro Antonelli^{1*}, Carlotta Palumbo¹, Marianna Noale², Walter Artibani³, Pierfrancesco Bassi⁴, Filippo Bertoni⁵, Sergio Bracarda⁶, Alessio Bruni⁷, Renzo Corvò⁸, Mauro Gacci⁹, Stefano Maria Magrini¹⁰, Rodolfo Montironi¹¹, Angelo Porreca¹², Andrea Tubaro¹³, Vittorina Zagonel¹⁴, Stefania Maggi² and the Pros-IT CNR study group[‡]

[‡]Members are listed at the end of the paper

Affiliations

1. Urology Unit, ASST Spedali Civili of Brescia. Department of Medical and Surgical Specialties, Radiological Science and Public Health, University of Brescia, Italy
2. Neuroscience Institute, Section of Padua, Aging Branch, National Research Council of Italy, Padua, Italy
3. Urology Unit, Azienda Ospedaliera Universitaria integrata di Verona, Verona, Italy
4. Catholic University of Rome, Policlinico Gemelli, Urology Department, Rome, Italy
5. Prostate Group of AIRO – Italian Association for Radiation Oncology, Milan, Italy
6. Medical Oncology Department, Azienda USL Toscana Sud-Est, Istituto Toscano Tumori, Arezzo, Italy
7. Radiotherapy Unit, University Hospital of Modena, Modena, Italy
8. Radiation Oncology Unit, IRCCS San Martino, Genoa, Italy
9. Urology Unit, Careggi Hospital, University of Florence, Florence, Italy
10. Radiation Oncology Unit, Spedali Civili Hospital of Brescia, University of Brescia, Italy

11. Section of Pathological Anatomy, Polytechnic University of the Marche Region, School of Medicine, United Hospitals, Ancona, Italy
12. Urology Unit, Policlinico di Abano Terme, Padua, Italy
13. Urology Unit, Sant'Andrea Hospital, University of Rome "La Sapienza", Italy
14. Istituto Oncologico Veneto IOV-IRCCS, Padua, Italy

* **Corresponding author:** Alessandro Antonelli, MD

Urology Unit, ASST Spedali Civili of Brescia. Department of Medical and Surgical Specialties, Radiological Science and Public Health, University of Brescia, Italy

Piazzale Spedali Civili 1, 25123, Brescia, Italy

Phone number: +390303995215 - Fax number: +390303995055

E-mail: alessandro_antonelli@me.com

Word count

Abstract: 275 words

Text: 2,620 words

Abstract

BACKGROUND: We assessed patients and tumor characteristics, as well as health-related quality of life (HRQoL) items, associated with curative intent treatment decision-making in clinically localized prostate cancer (PCa) patients.

METHODS: Clinically localized PCa treated with either radical prostatectomy (RP) or radiation therapy (RT) within 12 months from diagnosis were abstracted from The PROState cancer monitoring in ITaly, from the National Research Council (Pros-IT CNR) database. Multivariable logistic regression (MLR) models predicting RT vs. RP were fitted, after adjustment for HRQoL items, patients and tumor characteristics.

RESULTS: Of 1 041 patients, 631 (60.2%) were treated with RP and 410 (39.8%) with RT. Relative to RT, RP patients were younger age (mean age 64.5 ± 6.6 vs. 71.4 ± 4.9 , $p < 0.001$) and had higher rates of D'Amico low-intermediate risk groups (31.8 vs. 21.9% low, 46.3 vs. 43.5% intermediate and 21.9 vs. 34.6% high risk, $p < 0.001$). Overall, 93.2% of RP patients were enrolled by Urologists and 82.7% of RT patients by Radiation Oncologists. RP patients had generally higher means values of HRQoL items. In MLR models, higher RT rates were independently associated with more advanced age (odds ratio [OR] 6.14, $p < 0.001$) and BMI ≥ 30 kg/m² (OR 1.78, $p < 0.001$). Conversely, lower rates of RT were independently associated with married (OR 0.55, $p = 0.01$) and worker status (OR 0.52, $p = 0.004$), enrollment in academic centers (OR 0.59, $p = 0.005$) and higher physical composite score (OR 0.88, $p = 0.03$) and baseline sexual function (OR 0.92, $p < 0.001$) items.

CONCLUSIONS: Most patients with clinically localized prostate cancer undergoing definitive treatment at Italian institutions receive RP instead of RT. Moreover, those who are younger,

married, working, as well as those with better physical and sexual function are more likely to undergo surgery.

Keywords localized prostate cancer; radical prostatectomy; radiation therapy; decision-making; health related quality of life

1 Introduction

Different treatment modalities may be offered to patients diagnosed with clinically localized prostate cancer (PCa)¹⁻³. Among these, treatments with curative intent include radical prostatectomy ([RP] open, laparoscopic or robotic) and radiation therapy ([RT] external beam radiation therapy [EBRT] or brachytherapy) with or without androgen-deprivation therapy (ADT). To date, none of these two treatment modalities have been definitely proven to be superior in term of oncological benefit⁴⁻⁶. Nonetheless, each treatment is associated with specific risks and side effects which may invariably impact on patients' health-related quality of life (HRQoL)^{7,8}.

As result of lack of specific evidences supporting the preferential use of either RP or RT among the same risk categories, decision-making process for men diagnosed with clinically localized PCa is often conditioned by patients' preferences^{9,10} and features other than oncological characteristics distinctive of PCa. Indeed, several studies showed how sociodemographic characteristics¹¹⁻¹³, such as age, comorbidity, educational level, as well as provider facilities¹⁴, specialist seen, as well as multimodal discussion^{15,16}, and social support¹⁷, strongly influence this decision-making process.

In this population-based study, we assessed determinants of receipt of treatment with curative intent (i.e. RP or RT) in clinically localized PCa. We accounted for clinical, sociodemographic and tumor features, as well as HRQoL items, assessed through validated patient-reported outcomes measures (PROMs), prospectively collected within the PROState cancer monitoring in Italy project from the National Research Council (Pros-IT CNR)¹⁸.

2. Materials and Methods

2.1 Study design and population

The Pros-IT CNR project is a prospective observational longitudinal study which monitors an Italian cohort of 1 705 PCa men¹⁸⁻²⁰ through validated tools. Ninety-seven Italian public or private facilities, including 51 Urology, 39 Radiation Oncology and 7 Medical Oncology units, were involved in the Pros-IT CNR project. All the units were identified on the basis of project Steering Committee recommendations. Patients eligible for recruitment were males aged 18 years or more willing to participate into the study and that were newly diagnosed with biopsy-confirmed PCa between September 2014 and September 2015. Each participating center was required to enroll consecutive patients. Patients were evaluated every six-months for the first year, then yearly up to five years from enrollment. The approval of the Ethics Committee of coordinating center (Sant'Anna Hospital, Como, Italy; register number 45/2014), as well as of all local Committees, were obtained. Informed consent was signed by each patient.

The aim of the present study was to examine the use of curative treatment in patients with clinically localized PCa and to identify health-related quality of life (HRQoL) items, as well as patients- and physicians-associated characteristics, to drive treatment type selection. For the present analysis, only patients with clinically localized PCa treated with curative intent were included. Specifically, clinically localized stage definition, i.e. cT1 and cT2 tumors, as well as staging work-up, followed the current guidelines^{1,2}. Treatment options were categorized into two groups: (a) RP, alone or as a first step of multimodal treatment (RP plus EBRT plus/minus ADT), and (b) RT, alone or as part of multimodal treatment (EBRT plus ADT or brachytherapy plus/minus ADT). Only patients with a residual life expectancy of at least 10 years were included, according to guidelines recommendation of offering RP to patients with life expectancy of more

than 10 years. Life expectancy was calculated based on the Italian National Institute of Statistics (ISTAT) estimates for 2016. Specifically, residual life expectancy was calculated as the average number of years that a person of a specific age can expect to live, assuming that age-specific mortality levels remain constant. Further exclusion criteria consisted of lack of information of received treatment, patients treated with focal therapy, androgen deprivation therapy alone, monitoring (i.e. active surveillance or watchful waiting) and patients that did not return at least one questionnaire. These selection criteria yielded a cohort of 1 041 men with clinically localized PCa (Figure 1).

2.2 Covariates

Among patients' covariates, age at diagnosis, body-mass index (BMI), smoking status, comorbidities - assessed by the Cumulative Illness Rating Scale (CIRS) - family history of PCa, marital status, working status and area of residence were included. PROMs were assessed by validated questionnaires delivered, self-administered and returned during in-hospital visits, at diagnosis and then at 6 and 12 months after enrollment. Physical (PCS) and mental composite scores (MCS) were evaluated by the Italian versions of the Short-Form Health Survey (SF-12)²¹. Urinary, bowel and sexual functions and bothers (UF/B, BF/B, SF/B, respectively) were evaluated by the University of California Los Angeles-Prostate Cancer Index (Italian UCLA-PCI)²². Scores of both questionnaires ranged from 0 to 100, with higher scores indicating better outcomes. Using Gleason grade, PSA level and clinical stage, tumors were categorized according to European Association of Urology risk group as low, intermediate, or high risks. Finally, characteristics of institution where the initial prostate cancer was diagnosed (i.e. presence of Urology unit, Radiation Oncology Unit and/or Prostate Cancer Unit and type of institution – academic vs. non-academic), as well as physician that enrolled the patient, were collected.

2.3 Statistical Analysis

Number and percentages were presented for categorical variables, while means and standard deviation (SD) or medians and interquartile ranges (IQR) for continuous variables. No imputation of missing data was performed. Patient demographics, tumor characteristics and baseline PROMs were compared between RP vs. RT using chi-square or Fisher's Exact test for categorical variables, and mixed models or Wilcoxon rank-sum test for continuous variables. Additionally, to aid the interpretation of the results, the minimal clinically important difference (MID), which provides a measure of the smallest change in the PROMs of interest that patients perceive as important, was calculated for PROMs and defined as half standard deviation of the mean at baseline^{23,24}. Multivariable logistic regression models predicting treatment receipt (RT vs. RP) were fitted. Adjustment variables consisted of socio-demographic characteristics (age at diagnosis, education, marital status, work conditions, geographical area of residence), lifestyle and health status (smoking status, family history of prostate cancer, comorbidities, diabetes, BMI), D'Amico risk classes and PROMs. The linearity assumption for quantitative variables was evaluated on the basis of the quartiles and interactions among covariates were also evaluated. Finally, as sensitivity analysis, multivariable logistic regression models were repeated without inclusion of PROMs.

All statistical tests were two-sided with a level of significance set at $p < 0.05$. Analyses were performed using SAS software, version 9.4 (SAS Institute).

3. Results

3.1 Baseline patients' characteristics

For the purpose of the study, 1 041 patients with clinically localized PCa were considered (Figure 1). Of these, 631 (60.3%) patients were treated with RP (509 with RP alone and 122 with RP followed by RT with or without ADT) and 410 patients (39.4%) were treated with RT (242 with EBRT alone, 150 with EBRT plus ADT and 18 with brachytherapy). Baseline sociodemographic and tumors characteristics are summarized in Table 1. When compared to RT group, RP patients exhibited younger age (mean age 64.5 ± 6.6 vs. 71.4 ± 4.9 , $p < 0.001$). Additionally, RP patients were more frequently married or cohabiting (89.2 vs. 81.4%, $p < 0.001$), still working (39.6 vs. 13.0%, $p < 0.001$) and with higher educational level (79.8 vs. 66.0%, $p < 0.001$). Median PSA at diagnosis was lower in RP patients (6.6 vs. 7.3 ng/ml, $p = 0.02$). D'Amico risk classification was as follows: 31.8 vs. 21.9% low, 46.3 vs. 43.5% intermediate and 21.9 vs. 34.6% high risk, for respectively RP and RT patients ($p < 0.001$).

Additionally, 93.2% of RP patients were enrolled by Urologists, 6.5% by Radiation Oncologists and 0.3% by Medical Oncologists. Conversely, 16.6% of RT patients were enrolled by Urologists, 82.7% by Radiation Oncologists and 0.7% by Medical Oncologists (Table 2).

3.2 Baseline patients reported outcomes

There was a statistically significant difference in baseline PROMs that were higher in RP patients with the exception of MCS that was higher in RT patients (Table 3, Figure 2). However, these differences were minor. The MID did not exceed for all the PROMs with the exception of the UCLA-PCI SF. Here, the MID exceeded 0.5 SD of the baseline values (mean score of 61.4 ± 28.6 vs. 39.0 ± 29.9 , for respectively RP and RT patients, MID 15.6 $p < 0.001$).

3.3 Multivariable logistic regression models predicting treatment receipt

In multivariable logistic regression models predicting treatment receipt (Table 4), more advanced age (odds ratio [OR] 6.14, $p < 0.001$) and BMI ≥ 30 kg/m² (OR 1.78, $p < 0.001$) were independent predictors of RT treatment. Conversely, married or cohabiting patients (OR 0.55, $p = 0.01$), workers (OR 0.52, $p = 0.004$) and patients enrolled in academic centers (OR 0.59, $p = 0.005$) less frequently were treated with RT. Among HRQoL items, only PCS and SF items reached the independent predictor status. Specifically, a five-point increase in both PCS (OR 0.88, $p = 0.03$) and SF (OR 0.92, $p < 0.001$) were independent predictors of lower RT treatments.

In sensitivity analyses, after removing PROMs, more advanced age (OR 7.26, $p < 0.001$), BMI ≥ 30 kg/m² (OR 1.91 $p < 0.001$) and diabetes (OR 1.87, $p = 0.01$) were independent predictors of RT treatment. Conversely, married or cohabiting patients (OR 0.51, $p = 0.01$), workers (OR 0.45, $p = 0.004$) and patients enrolled in academic centers (OR 0.60, $p = 0.005$) less frequently were treated with RT.

4. Discussion

Men diagnosed with clinically localized PCa who desire treatment with curative intent face highly preference-sensitive choice between RP and RT. Indeed, both RP and RT are recognized as equivalent treatment options in patients with clinically localized PCa by both European and North-American guidelines^{1,2}. Therefore, selection of either RP or RT as initial treatment in clinically localized PCa may be challenging for both physicians and patients. The present study investigated patients and tumor characteristics, as well as HRQoL items, that may be associated with treatment decision-making in a contemporary nation-based cohort with clinically localized PCa and more than ten-year life-expectancy. Our analyses resulted in several noteworthy findings.

First, in this study, 60% of patients were treated with RP, while 40% were treated with RT. This finding is in agreement with previous population and nation-based analyses²⁵⁻²⁷, where rates of RP exceeded those of RT in clinically localized PCa patients. As those previous reports, RP patients were generally younger and with longer life expectancy. Additionally, RP patients were more frequently still working, more frequently married or cohabiting, and had higher educational level. Conversely, non-clinically meaningful differences were recorded according to comorbidity profile. Of these demographic patient's characteristics, only younger age, married or cohabiting status and still working condition were significantly associated with being treated with RP after multivariable adjustment.

Second, RP patients presented with lower D'Amico risk class. These findings suggest a more favorable tumor phenotype in RP patients. However, D'Amico risk class was not significantly associated with treatment receipt. These observations suggest that both RP and RT are equally offered based on tumor characteristics in clinically localized disease.

Third, the vast majority (93.2%) of RP patients were enrolled by Urologist, while the vast majority of RT patients (82.7%) were enrolled by Radiation Oncologist. Additionally, urologists enrolled 16.6% of RT, while radiation oncologists enrolled 6.5% of RP. Due to the wide difference in the proportions, it was not possible to test in multivariable models the effect of physician who enrolled the patient. Previous reports^{15,28} consistently showed that both Radiation Oncologists and Urologists mainly prescribe the treatment modalities they offer. Moreover, if we consider the time span of the previous studies^{15,28}, our findings suggested that the pattern of treatment decision-making according to physician specialty have remained unchanged for the past two decades. Finally, it should be noted that in our cohort biopsy was performed by Urologists in almost all of the cases. Therefore, the Urologist was the physician who made the upfront PCa diagnosis.

Fourth, PROMs at baseline were generally higher in RP patients. Nevertheless, this difference may be considered clinically meaningful only for SF. This observation may be further confirmed by the fact that only SF item exceed the MID. However, it should be noted that these PROMs were developed before the MID concept was widespread and that use of MID was not included in the prespecified analysis plan. Therefore, MID should be use as an aid in interpretation of the results. In the present cohort, the clinically meaningful difference in PROMs was recorded for SF item that additionally exceeded the MID. Nonetheless, this difference may be attributable to the younger age of RP patients, as previously reported²⁴. After adjustment for multiple covariates, only SF and PCS items were significantly associated with treatment receipt. Specifically, higher SF and PCS scores were associated with a lower likelihood of receiving a RP. Previous studies showed how patients' preference^{10,29}, as well as patients' bother from treatment side effects^{30,31}, may play an important role in treatment decision-making progress. However, to the best of our knowledge, this is the first study that investigated how HRQoL collected with

validated PROMs may be related to treatment choice. Indeed, choosing a treatment option that less negatively affects mental and physical health, as well as urinary, sexual or bowel function, may be prioritized by many patients.

In this contemporary nation-based cohort of clinically localized PCa patients treated with treatment with curative intent, the majority (60%) received RP. Higher RP rates were significantly associated with younger age, married status and still working conditions, as well as with higher baseline HRQoL items, with the biggest difference recorded for SF. It is worth of mention that those should be the subjects featured by the largest independence in choosing treatments, as well as those more interested in sexual life. Therefore, it could be concluded that the expectations to achieve the more radical treatment have overcome any concern on the side effects of surgery on sexual function³². Alternatively, the physicians proposing surgery undermined such effects. Of interest, neither comorbidity profile nor D'Amico risk class were significantly associated with treatment receipt in our cohort, suggesting that contemporary surgical indication is not influenced by these factors due to the improvement of peri- and intra-operative care and the widening indication to more aggressive tumors. These findings offer an insight in the present Italian scenario and are of remarkable importance to further interpret the outcomes in patients treated with RP and RT that will be the matter of next future studies. Additionally, our findings further encourage the importance of a multidisciplinary approach³³, in order to provide the best tools for decision-making. Indeed, it has been demonstrated that multimodal discussion may alter management plans in 25% to 50% of PCa cases¹⁶.

This study has several strengths and potential limitations that should be acknowledged.

First of all, a strength of the study is the population-based design of contemporary patients, which

yielded a cohort that is more representative than institutional reports. However, the findings of the current study derived from exploratory analyses of an observational study design, which could have made results susceptible to confounders. Additionally, the involvement of participating centers on a voluntary basis, with almost half from the North of Italy, could have limited the representability of all Italian scenario. Second, this study is also unique in assessing patient reported HRQoL items through prospectively and systematically collected validated questionnaires. Nonetheless, patients' involvement in treatment decision-making could not be assessed. Similarly, also data on patient's income and social network could not be assessed. Therefore, we could not test the effect for these variables on treatment choice. Third, physician recommendations that may have influenced treatment choice could not be assessed. Moreover, whether multimodal discussion was performed also could not be adequately assessed. Nonetheless, we accounted for center characteristics, such as presence of Prostate Unit and academic status. Fourth, due to the low number of patients treated with active surveillance, and due to the lack of information on active surveillance strategy, in this study was not possible to investigate also active surveillance among treatment modalities.

5. Conclusions

Most patients with clinically localized prostate cancer undergoing definitive treatment at Italian institutions receive RP instead of RT. Moreover, those who are younger, married, working, as well as those with better physical and sexual function are more likely to undergo surgery.

Notes**Conflict of Interest**

All the authors report no potential conflict of interest to disclose.

Funding

None

Authors contribution

Concept and design: Antonelli, Palumbo, Noale, Maggi

Data collection: all the authors, Pros-IT CNR Group:

Analysis and interpretation of data: all the authors

Statistical analysis: Noale

Drafting the manuscript: Antonelli, Palumbo, Noale

Revision for important intellectual content: Artibani, Bassi, Bertoni, Bracarda, Bruni, Corvò,

Gacci, Magrini, Montironi, Porreca, Tubaro, Zagonel, Maggi

Supervision: Antonelli, Maggi

The Pros-IT CNR Group:

Steering Committee: Gaetano CREPALDI (Padova); Stefania MAGGI (Padova); Marianna NOALE (Padova); Angelo PORRECA (Padova); Walter ARTIBANI (Verona); Pierfrancesco BASSI (Roma); Sergio BRACARDA (Arezzo); Giario Natale CONTI (Como); Renzo CORVÒ (Genova); Pierpaolo GRAZIOTTI (Milano); Riccardo Maurizi ENRICI (Roma); Vincenzo MIRONE (Napoli); Rodolfo MONTIRONI (Ancona).

Scientific committee: Filippo BERTONI (Brescia); Mauro GACCI (Firenze); Stefano Maria MAGRINI (Brescia); Giovanni MUTO (Roma); Stefano PECORARO (Avellino); Umberto RICARDI (Torino); Andrea TUBARO (Roma); Vittorina ZAGONEL (Padova); Filiberto ZATTONI (Padova).

Working Group: Anna Rita ALITTO (Roma); Enrica AMBROSI (Brescia); Alessandro ANTONELLI (Brescia); Cynthia ARISTEI (Perugia); Michele BARBIERI (Napoli); Franco BARDARI (Asti); Lilia BARDOSCIA (Brescia); Salvina BARRA (Genova); Sara BARTONCINI (Torino); Umberto BASSO (Padova); Carlotta BECHERINI (Firenze); Rita BELLAVITA (Perugia); Franco BERGAMASCHI (Reggio Emilia); Stefania BERLINGHERI (Brescia);

Alfredo BERRUTI (Brescia); Marco BORGHESI (Bologna); Roberto BORTOLUS (Pordenone);
 Valentina BORZILLO (Napoli); Davide BOSETTI (Milano); Giuseppe BOVE (Foggia); Pierluigi
 BOVE (Roma); Maurizio BRAUSI (Modena); Alessio BRUNI (Modena); Giorgio BRUNO
 (Ravenna); Eugenio BRUNOCILLA (Bologna); Alberto BUFFOLI (Brescia); Michela
 BUGLIONE (Brescia); Consuelo BUTTIGLIERO (Torino); Giovanni CACCIAMANI (Verona);
 Michela CALDIROLI (Varese); Giuseppe CARDO (Bari); Giorgio CARMIGNANI (Genova);
 Giuseppe CARRIERI (Foggia); Emanuele CASTELLI (Torino); Elisabetta CASTREZZATI
 (Brescia); Gianpiero CATALANO (Milano); Susanna CATTARINO (Roma); Francesco
 CATUCCI (Roma); Dario CAVALLINI FRANCOLINI (Pavia); Ofelia CECCARINI (Bergamo);
 Antonio CELIA (Vicenza); Francesco CHIANCONE (Napoli); Tommaso CHINI (Firenze);
 Claudia CIANCI (Pisa); Antonio CISTERNINO (Foggia); Devis COLLURA (Torino); Franco
 CORBELLÀ (Pavia); Matteo CORINTI (Como); Paolo CORSI (Verona); Fiorenza CORTESE
 (Alessandria); Luigi CORTI (Padova); Cosimo DE NUNZIO (Roma); Olga CRISTIANO
 (Avellino); Rolando D'ANGELILLO (Roma); Luigi DA POZZO (Bergamo); Daniele D'agostino
 (Padova); David D'ANDREA (Bolzano); Matteo DANDREA (Padova); Michele DE ANGELIS
 (Arezzo); Ottavio DE COBELLI (Milano); Bernardino DE CONCILIO (Vicenza); Antonello DE
 LISA (Cagliari); Stefano DE LUCA (Torino); Agostina DE STEFANI (Bergamo); Chiara
 Lucrezia DEANTONI (Milano); Claudio DEGLI ESPOSTI (Bologna); Anna DESTITO
 (Catanzaro); Beatrice DETTI (Firenze); Nadia DI MUZIO (Milano); Andrea DI STASIO
 (Alessandria); Calogero DI STEFANO (Ravenna); Danilo DI TRAPANI (Palermo); Giuseppe
 DIFINO (Foggia); Marco FABIANO (Napoli); Sara FALIVENE (Napoli); Giuseppe FARULLO
 (Roma); Paolo FEDELINI (Napoli); Iliaria FERRARI (Varese); Francesco FERRAU (Messina);
 Matteo FERRO (Milano); Andrei FODOR (Milano); Francesco FONTANA (Novara); Francesco
 FRANCESCA (Pisa); Giulio FRANCOLINI (Firenze); Paolo FRATA (Brescia); Giovanni
 FREZZA (Bologna); Pietro GABRIELE (Torino); Maria GALEANDRO (Reggio Emilia);
 Elisabetta GARIBALDI (Torino); Pietro Giovanni GENNARI (Arezzo); Alessandro
 GENTILUCCI (Roma); Alessandro GIACOBBE (Torino); Laura GIUSSANI (Varese); Giuseppe
 GIUSTI (Cagliari); Paolo GONTERO (Torino); Alessia GUARNERI (Torino); Cesare GUIDA
 (Avellino); Alberto GURIOLI (Torino); Dorijan HUQI (Bolzano); Ciro IMBIMBO (Napoli);
 Gianluca INGROSSO (Roma); Cinzia IOTTI (Reggio Emilia); Corrado ITALIA (Bergamo);
 Pierdaniele LA MATTINA (Milano); Enza LAMANNA (Ravenna); Luciana LASTRUCCI
 (Arezzo); Grazia LAZZARI (Taranto); Fabiola LIBERALE (Biella); Giovanni LIGUORI
 (Trieste); Roberto LISI (Roma); Frank LOHR (Modena); Riccardo LOMBARDO (Roma); Jon
 LOVISOLO (Varese); Giuseppe Mario LUDOVICO (Bari); Nicola MACCHIONE (Novara);
 Francesca MAGGIO (Imperia); Michele MALIZIA (Bologna); Gianluca MANASSE (Perugia);
 Giovanni MANDOLITI (Rovigo); Giovanna MANTINI (Roma); Luigi MARAFIOTI (Cosenza);
 Luisa MARCIELLO (Prato); Alberto Mario MARCONI (Varese); Antonietta MARTILLOTTA
 (Cosenza); Salvino MARZANO (Prato); Stefano MASCIULLO (Bergamo); Gloria MASO
 (Verbania); Adele MASSENZO (Cosenza); Ercole MAZZEO (Modena); Luigi MEARINI
 (Perugia); Serena MEDORO (Ferrara); Rosa MOLÈ (Catanzaro); Giorgio MONESI (Novara);
 Emanuele MONTANARI (Milano); Franco MONTEFIORE (Alessandria); Giampaolo
 MONTESI (Rovigo); Giuseppe MORGIA (Catania); Gregorio MORO (Biella); Giorgio
 MUSCAS (Cagliari); Daniela MUSIO (Roma); Paolo MUTO (Napoli); Giovanni MUZZONIGRO
 (Ancona); Giorgio NAPODANO (Salerno); Carlo Luigi Augusto NEGRO (Asti); Mattia NIDINI
 (Mantova); Maria NTRETA (Bologna); Marco ORSATTI (Imperia); Carmela PALAZZOLO
 (Messina); Isabella PALUMBO (Perugia); Alessandro PARISI (Bologna); Paolo PARMA

(Mantova); Nicola PAVAN (Trieste); Martina PERICOLINI (Roma); Francesco PINTO (Roma); Antonio PISTONE (Salerno); Valerio PIZZUTI (Grosseto); Angelo PLATANIA (Messina); Caterina POLLI (Prato); Giorgio POMARA (Pisa); Elisabetta PONTI (Roma); Antonio Benito PORCARO (Verona); Francesco PORPIGLIA (Torino); Dario PUGLIESE (Roma); Armin PYCHA (Bolzano); Giuseppe RAGUSO (Reggio Emilia); Andrea RAMPINI (Arezzo); Donato Franco RANDONE (Torino); Valentina ROBOLDI (Bergamo); Marco ROSCIGNO (Bergamo); Maria Paola RUGGIERI (Reggio Emilia); Giuseppe RUOPPO (Reggio Emilia); Roberto SANSEVERINO (Salerno); Anna SANTACATERINA (Messina); Michele SANTARSIERI (Pisa); Riccardo SANTONI (Roma); Giorgio Vittorio SCAGLIOTTI (Torino); Mauro SCANZI (Brescia); Marcello SCARCIA (Bari); Riccardo SCHIAVINA (Bologna); Alessandro SCIARRA (Roma); Carmine SCIORIO (Lecco); Tindaro SCOLARO (La Spezia); Salvatore SCUZZARELLA (Lecco); Oscar SELVAGGIO (Foggia); Armando SERAO (Alessandria); Sergio SERNI (Firenze); Marco Andrea SIGNOR (Udine); Mauro SILVANI (Biella); Giovanni SILVANO (Taranto); Franco SILVESTRIS (Bari); Claudio SIMEONE (Brescia); Valeria SIMONE (Bari); Girolamo SPAGNOLETTI (Foggia); Matteo Giulio SPINELLI (Milano); Luigi SQUILLACE (Pavia); Vincenzo TOMBOLINI (Roma); Mariastella TONINELLI (Brescia); Luca TRIGGIANI (Brescia); Alberto TRINCHIERI (Lecco); Luca Eolo TRODELLA (Roma); Lucio TRODELLA (Roma); Carlo TROMBETTA (Trieste); Lidia TRONNOLONE (Roma); Marcello TUCCI (Torino); Daniele URZÌ (Catania); Riccardo VALDAGNI (Milano); Maurizio VALERIANI (Roma); Maurizio VANOLI (La Spezia); Elisabetta VITALI (Bergamo); Stefano ZARAMELLA (Novara); Guglielmo ZECCOLINI (Vicenza); Giampaolo ZINI (Ferrara).

Acknowledgments

Pros-IT CNR is a non-profit observational study. Takeda Italia S.p.A. provided an unconditional grant to the CNR, to cover the costs of the preparatory meetings, the meetings of the advisory committees and of the PIs, and the cost of developing a web platform for data entry.

References

1. Mohler JL, Hurwitz M, Richey S. NCCN Guidelines Index Table of Contents Discussion. *Prostate Cancer*. 2019;163.
2. Mottet N, Bellmunt J, Bolla M, Briers E, Cumberbatch MG, De Santis M, et al. EAU-ESTRO-SIOG Guidelines on Prostate Cancer. Part 1: Screening, Diagnosis, and Local Treatment with Curative Intent. *Eur Urol*. 2017 Apr;71(4):618–29.
3. Fontanella P, Benecchi L, Grasso A, Patel V, Albala D, Abbou C, et al. Decision-making tools in prostate cancer: from risk grouping to nomograms. *Minerva Urol E Nefrol Ital J Urol Nephrol*. 2017 Dec;69(6):556–66.
4. Hamdy FC, Donovan JL, Lane JA, Mason M, Metcalfe C, Holding P, et al. 10-Year Outcomes after Monitoring, Surgery, or Radiotherapy for Localized Prostate Cancer. *N Engl J Med*. 2016 Oct 13;375(15):1415–24.
5. Kishan AU, Cook RR, Ciezki JP, Ross AE, Pomerantz MM, Nguyen PL, et al. Radical Prostatectomy, External Beam Radiotherapy, or External Beam Radiotherapy With Brachytherapy Boost and Disease Progression and Mortality in Patients With Gleason Score 9–10 Prostate Cancer. *JAMA*. 2018 Mar 6;319(9):896–905.
6. Berg S, Cole AP, Krimphove MJ, Nabi J, Marchese M, Lipsitz SR, et al. Comparative Effectiveness of Radical Prostatectomy Versus External Beam Radiation Therapy Plus Brachytherapy in Patients with High-risk Localized Prostate Cancer. *Eur Urol*. 2019 Apr;75(4):552–5.
7. Donovan JL, Hamdy FC, Lane JA, Mason M, Metcalfe C, Walsh E, et al. Patient-Reported Outcomes after Monitoring, Surgery, or Radiotherapy for Prostate Cancer. *N Engl J Med*. 2016 Oct 13;375(15):1425–37.
8. Wallis CJD, Saskin R, Choo R, Herschorn S, Kodama RT, Satkunasingam R, et al. Surgery Versus Radiotherapy for Clinically-localized Prostate Cancer: A Systematic Review and Meta-analysis. *Eur Urol*. 2016 Jul;70(1):21–30.
9. Jayadevappa R, Chhatre S, Gallo JJ, Wittink M, Morales KH, Lee DI, et al. Patient-Centered Preference Assessment to Improve Satisfaction With Care Among Patients With Localized Prostate Cancer: A Randomized Controlled Trial. *J Clin Oncol*. 2019 Apr 20;37(12):964–73.
10. Hurwitz LM, Cullen J, Elsamanoudi S, Kim DJ, Hudak J, Colston M, et al. A prospective cohort study of treatment decision-making for prostate cancer following participation in a multidisciplinary clinic. *Urol Oncol Semin Orig Investig*. 2016 May;34(5):233.e17–233.e25.
11. Lunardi P, Ploussard G, Grosclaude P, Roumiguié M, Soulié M, Beauval JB, et al. Current impact of age and comorbidity assessment on prostate cancer treatment choice and over/undertreatment risk. *World J Urol*. 2017 Apr;35(4):587–93.
12. Gray PJ, Lin CC, Cooperberg MR, Jemal A, Efsthathiou JA. Temporal Trends and the Impact of Race, Insurance, and Socioeconomic Status in the Management of Localized Prostate Cancer. *Eur Urol*. 2017 May;71(5):729–37.
13. Zattoni F, Montebelli F, Rossanese M, Crestani A, Giannarini G, Ficarra V. Should radical prostatectomy be encouraged at any age? A critical non-systematic review. *Minerva Urol E Nefrol Ital J Urol Nephrol*. 2018 Feb;70(1):42–52.
14. Satkunasingam R, Lo M, Stern M, Gill IS, Flensing S, Wu X-C, et al. The Role of Provider Characteristics in the Selection of Surgery or Radiation for Localized Prostate Cancer and Association With Quality of Care Indicators. *Am J Clin Oncol*. 2018 Apr 17;

15. Kim SP, Gross CP, Nguyen PL, Nguyen PY, Smaldone MC, Thompson RH, et al. Specialty bias in treatment recommendations and quality of life among radiation oncologists and urologists for localized prostate cancer. *Prostate Cancer Prostatic Dis.* 2014 Jun;17(2):163–9.
16. De Luca S, Fiori C, Tucci M, Poggio M, Allis S, Bollito E, et al. Prostate cancer management at an Italian tertiary referral centre: does multidisciplinary team meeting influence diagnostic and therapeutic decision-making process? A snapshot of the everyday clinical practice. *Minerva Urol E Nefrol Ital J Urol Nephrol.* 2019 Sep 4;
17. Reamer E, Yang F, Holmes-Rovner M, Liu J, Xu J. Influence of Men's Personality and Social Support on Treatment Decision-Making for Localized Prostate Cancer. *BioMed Res Int.* 2017;2017:1–8.
18. Noale M, Maggi S, Artibani W, Bassi PF, Bertoni F, Bracarda S, et al. Pros-IT CNR: an Italian prostate cancer monitoring project. *Aging Clin Exp Res.* 2017 Apr;29(2):165–72.
19. Porreca A, Noale M, Artibani W, Bassi PF, Bertoni F, Bracarda S, et al. Disease-specific and general health-related quality of life in newly diagnosed prostate cancer patients: the Pros-IT CNR study. *Health Qual Life Outcomes [Internet].* 2018 Dec [cited 2018 Aug 9];16(1). Available from: <https://hql.o.biomedcentral.com/articles/10.1186/s12955-018-0952-5>
20. Antonelli A, Palumbo C, Noale M, Porreca A, Maggi S, Simeone C, et al. Impact of Surgical Approach on Patient-Reported Outcomes after Radical Prostatectomy: A Propensity Score-Weighted Analysis from a Multicenter, Prospective, Observational Study (The Pros-IT CNR Study). *Urol Int.* 2019 Feb 7;1–11.
21. Apolone G, Mosconi P, Quattrocioni L, Gianicolo EAL, Groth N, Ware JJ E. Questionario sullo stato di salute SF-12. Versione Italiana. Milano, Guerini e Associati Editore, 2001.
22. Gacci M, Livi L, Pajar F, Detti B, Litwin MS, Bartoletti R, et al. Quality of life after radical treatment of prostate cancer: validation of the Italian version of the University of California-Los Angeles Prostate Cancer Index. *Urology.* 2005 Aug;66(2):338–43.
23. Revicki D, Hays RD, Cella D, Sloan J. Recommended methods for determining responsiveness and minimally important differences for patient-reported outcomes. *J Clin Epidemiol.* 2008 Feb;61(2):102–9.
24. Lane A, Metcalfe C, Young GJ, Peters TJ, Blazeby J, Avery KNL, et al. Patient-reported outcomes in the ProtecT randomized trial of clinically localized prostate cancer treatments: study design, and baseline urinary, bowel and sexual function and quality of life. *BJU Int.* 2016 Dec;118(6):869–79.
25. Cooperberg MR, Broering JM, Carroll PR. Time Trends and Local Variation in Primary Treatment of Localized Prostate Cancer. *J Clin Oncol.* 2010 Mar;28(7):1117–23.
26. Weissbach L, Stuerzebecher S, Mumperow E, Klotz T, Schnell D. HAROW: the first comprehensive prospective observational study comparing treatment options in localized prostate cancer. *World J Urol.* 2016 May;34(5):641–7.
27. Tyson MD, Alvarez J, Koyama T, Hoffman KE, Resnick MJ, Wu X-C, et al. Racial Variation in Patient-Reported Outcomes Following Treatment for Localized Prostate Cancer: Results from the CEASAR Study. *Eur Urol.* 2017 Aug;72(2):307–14.
28. Jang TL. Physician Visits Prior to Treatment for Clinically Localized Prostate Cancer. *Arch Intern Med.* 2010 Mar 8;170(5):440.
29. Johnson DC, Mueller DE, Deal AM, Dunn MW, Smith AB, Woods ME, et al. Integrating Patient Preference into Treatment Decisions for Men with Prostate Cancer at the Point of Care. *J Urol.* 2016 Dec;196(6):1640–4.

30. Ross LE, Howard DL, Bowie JV, Thorpe RJ, Kinlock BL, Burt C, et al. Factors Associated with Men's Assessment of Prostate Cancer Treatment Choice. *J Cancer Educ.* 2016 Jun;31(2):301–7.
31. Broughman JR, Basak R, Nielsen ME, Reeve BB, Usinger DS, Spearman KC, et al. Prostate Cancer Patient Characteristics Associated With a Strong Preference to Preserve Sexual Function and Receipt of Active Surveillance. *JNCI J Natl Cancer Inst.* 2018 Apr 1;110(4):420–5.
32. Antonelli A, Sodano M, Peroni A, Mittino I, Palumbo C, Furlan M, et al. Positive surgical margins and early oncological outcomes of robotic vs open radical prostatectomy at a medium case-load institution. *Minerva Urol E Nefrol Ital J Urol Nephrol.* 2017;69(1):63–8.
33. Zattoni F, Morlacco A, Matrone F, Arcicasa M, Buttazzi L, Maruzzi D, et al. Multimodal treatment for high-risk locally-advanced prostate cancer following radical prostatectomy and extended lymphadenectomy. *Minerva Urol E Nefrol Ital J Urol Nephrol.* 2019 Oct;71(5):508–15.

Table 1. Baseline sociodemographic and tumors characteristics of 1 041 patients stratified according to treatment type (radical prostatectomy vs. radiation therapy) for clinically localized prostate cancer, identified within the PROState cancer monitoring in Italy, from the National Research Council (Pros-IT CNR) study.

	Radical Prostatectomy* (n=631, 60.6%)	Radiation Therapy# (n=410, 39.4%)	p
Age at diagnosis, years, mean \pm SD	64.5 \pm 6.6	71.4 \pm 4.9	<0.001
Age at diagnosis, years, median (IQR)	66 (60-69)	73 (69-75)	<0.001
Life expectancy at diagnosis, years, mean \pm SD	19.7 \pm 5.4	14.3 \pm 3.8	<0.001
Education \geq lower secondary school, n (%)	501 (79.8)	266 (66.0)	<0.001
Marital status, married or cohabiting, n (%)	563 (89.2)	333 (81.4)	<0.001
Work condition, still working, n (%)	246 (39.6)	53 (13.0)	<0.001
Geographical area of residence, n (%)			0.004
Northern regions	312 (50.7)	237 (60.8)	
Central regions	181 (29.4)	82 (21.0)	
Southern regions	122 (19.8)	71 (18.2)	
BMI, kg/m ² , mean \pm SD	26.4 \pm 3.2	27.1 \pm 4.0	0.02
Smoking status, current smoker, n (%)	102 (16.5)	48 (12.0)	0.05
Diabetes mellitus, n (%)	64 (10.2)	85 (20.8)	<0.001
CIRS Comorbidities Index, median (IQR)	1 (0-2)	1 (0-2)	0.001
CIRS Severity Index, median (IQR)	1.31 (1.15-1.46)	1.31 (1.23-1.54)	0.002
Family history of prostate cancer, n (%)	129 (20.7)	53 (13.0)	0.002
Clinical T stage at diagnosis, n (%)			<0.001
cT1	378 (59.9)	187 (45.6)	
cT2	253 (40.1)	223 (54.4)	
Gleason score at diagnosis, n (%)			0.1
\leq 6	311 (49.8)	179 (44.1)	
3+4	151 (24.2)	96 (23.6)	
4+3	81 (13.0)	58 (14.3)	
\geq 8	81 (13.0)	73 (18.0)	
PSA at diagnosis, ng/mL, median (IQR)	6.6 (5.0-9.4)	7.3 (5.1-10.1)	0.02
D'Amico risk class, n (%)			<0.001
Low	198 (31.8)	89 (21.9)	
Intermediate	288 (46.3)	177 (43.5)	
High	136 (21.9)	141 (34.6)	

* Radical prostatectomy includes 509 exclusive radical prostatectomy and 122 radical prostatectomy followed by radiation therapy \pm androgen deprivation therapy

Radiation therapy includes 242 external beam radiation therapy, 150 external beam radiation therapy + androgen deprivation therapy and 18 brachytherapy \pm androgen deprivation therapy

Abbreviations. SD=Standard Deviation; IQR =interquartile range; BMI=Body Mass Index;

CIRS=Cumulative Illness Rating Scale

Table 2. Provider and facility characteristics of 1 041 patients stratified according to treatment type (radical prostatectomy vs. radiation therapy) for clinically localized prostate cancer, identified within the PROState cancer monitoring in Italy, from the National Research Council (Pros-IT CNR) cohort.

	Radical Prostatectomy (n=631, 60.6%)	Radiation Therapy (n=410, 39.4%)	p
Presence of Urology Unit, n (%)			<0.001
Yes	627 (99.4)	375 (91.5)	
No	4 (0.6)	35 (8.5)	
Presence of Radiation Oncology Unit, n (%)			<0.001
Yes	475 (75.3)	383 (93.4)	
No	156 (24.7)	27 (6.8)	
Presence of Medical Oncology Unit, n (%)			
Yes	551 (87.3)	399 (97.3)	<0.001
No	80 (12.7)	11 (2.7)	
Presence of Prostate Unit, n (%)			<0.001
Yes	118 (18.7)	47 (11.5)	
No	513 (81.3)	363 (88.5)	
Hospital teaching status, n (%)			
Teaching	348 (55.2)	176 (42.9)	0.02
Non-teaching	283 (44.9)	234 (57.1)	
Patient enrolled by, n (%)			<0.001
Urologist	588 (93.2)	68 (16.6)	
Radiation Oncologist	41 (6.5)	339 (82.7)	
Medical Oncologist	2 (0.3)	3 (0.7)	

Table 3. Baseline health related quality of life items of 1,041 patients stratified according to treatment type (radical prostatectomy vs. radiation therapy) for clinically localized prostate cancer, identified within the PROState cancer monitoring in ITaly, from the National Research Council (Pros-IT CNR) study.

Scores at diagnosis	Radical Prostatectomy (n=631, 60.6%)	Radiation Therapy (n=410, 39.4%)	p value	MID
UCLA-PCI scores (mean±SD)				
Urinary function	94.9±13.7	92.0±16.7	<0.001	7.5 [§]
Urinary bother	91.1±21.5	85±25.7	<0.001	11.5 [§]
Bowel function	95.7±10.9	92.3±15.0	<0.001	6.4 [§]
Bowel bother	95.5±14.2	91.2±21	<0.001	8.7 [§]
Sexual function	61.4±28.6	39±29.9	<0.001	15.6 [^]
Sexual bother	67.4±33.6	58.9±35.6	<0.001	17.3 [§]
SF-12 scores (mean±SD)				
Physical Component Subscale	53.2±6.1	50.3±8.1	<0.001	3.5 [§]
Mental Component Subscale	49±9.4	50.1±9.6	0.01	4.7 [§]

Abbreviations. SD=Standard Deviation; MID=minimal clinically important difference; SF-12=Short-Form Health Survey; UCLA-PCI=University of California Los Angeles-Prostate Cancer Index. All scores ranges from 0 to 100, with higher scores representing better quality of life.

[§] the MID did exceed between groups (p>0.05)

[^] the MID exceeded between groups (p=0.0001)

Table 4. Multivariable logistic regression models predicting use of radiation therapy vs. radical prostatectomy as treatment with curative intent in a cohort of 1 041 patients with clinically localized prostate cancer, identified within the PROState cancer monitoring in ITaly, from the National Research Council (Pros-IT CNR) study.

	Odds ratio	95% Confidence Interval	p-value
Age at diagnosis ≥ 70 years	6.14	4.27-8.82	<0.001
Education \geq lower secondary school	1.25	0.86-1.81	0.2
Marital status, married or cohabiting	0.55	0.34-0.88	0.01
Work condition, still working	0.52	0.34-0.81	0.004
Geographical area of residence			
Central vs. Northern regions	0.99	0.59-1.65	0.9
Southern vs. Northern regions	0.69	0.44-1.08	0.1
BMI ≥ 30 kg/m ²	1.78	1.13-2.81	0.01
Smoking status			
Past vs. never smoker	1.08	0.75-1.55	0.7
Current vs. never smoker	1.58	0.95-2.64	0.07
Family history of prostate cancer, yes vs. no	0.77	0.48-1.21	0.2
CIRS Comorbidity Index ≥ 2	0.72	0.48-1.06	0.1
CIRS Severity Index ≥ 1.3	0.87	0.56-1.34	0.5
Diabetes mellitus	1.46	0.91-2.35	0.1
D'Amico risk class, n (%)			
Intermediate vs. low	1.02	0.67-1.56	0.9
High vs. low	1.41	0.89-2.25	0.1
SF-12 PCS (5 points increment)	0.88	0.78-0.99	0.03
SF-12 MCS (5 points increment)	1.09	0.99-1.19	0.07
UCLA-PCI UF (5 points increment)	1.06	0.99-1.14	0.08
UCLA-PCI UB (5 points increment)	0.95	0.91-1.00	0.05
UCLA-PCI BF (5 points increment)	0.99	0.91-1.07	0.8
UCLA-PCI BB (5 points increment)	0.96	0.90-1.02	0.2
UCLA-PCI SF (5 points increment)	0.92	0.89-0.95	<0.001
UCLA-PCI SB (5 points increment)	1.00	0.98-1.03	0.8
Presence of Prostate Unit, yes vs. no	0.67	0.37-1.20	0.2
Academic centre, yes vs. no	0.59	0.41-0.86	0.005

Abbreviations. BMI=Body Mass Index; CIRS=Cumulative Illness Rating Scale; SF-12=Short-Form Health Survey; PCS=Physical Component Subscale; MCS=Mental Component Subscale. Scores ranges from 0 to 100, with higher scores representing better quality of life. UCLA-PCI=University of California Los Angeles-Prostate Cancer Index; UF=Urinary Function; UB=Urinary Bother; BF=Bowel Function; BB=Bowel Bother; SF=Sexual Function; SB=Sexual Bother. Scores ranges from 0 to 100, with higher scores representing better quality of life in relation to functions or symptoms

Figure legends

Figure 1. CONSORT flow diagram illustrating inclusion and exclusion criteria

Figure 2. Baseline health-related quality of life scores of 1 041 clinically-localized prostate cancer patients stratified according to treatment type (radical prostatectomy vs. radiation therapy), identified within the PROState cancer monitoring in ITaly from the National Research Council (Pros-IT CNR) study.

Abbreviations. RP=radical prostatectomy; RT=external beam radiation therapy; UCLA-PCI=University of California Los Angeles-Prostate Cancer Index; UF=Urinary Function; UB=Urinary Bother; BF=Bowel Function; BB=Bowel Bother; SF=Sexual Function; SB=Sexual Bother; SF-12=Short-Form Health Survey; PCS=Physical Component Subscale; MCS=Mental Component Subscale.

RP was performed either alone or as a first step of multimodal treatment (RP plus RT with/without androgen deprivation therapy). RT was performed either alone or as part of multimodal treatment (external beam RT with/without androgen deprivation therapy or brachytherapy with/without androgen deprivation therapy).

The bars represent mean values. All scores ranges from 0 to 100, with higher scores representing better quality of life.



