

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Anxiety, depression, and posttraumatic stress in nurses during the COVID-19 outbreak

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1837422> since 2022-01-31T16:14:15Z

Published version:

DOI:10.1016/j.iccn.2021.103014

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 **Anxiety, depression, and posttraumatic stress in nurses during the COVID-19 outbreak**

2 **Funding Source**

3 This research received no specific grant from any funding agency in the public, commercial, or not-
4 for-profit sectors.

5 **Conflict of Interest**

6 The authors declare that they have no conflicts of interest.

7 **Acknowledgements**

8 The authors would like to thank the participants involved in the study.

9 During the COVID-19 pandemic, nurses and physicians faced exhausting work conditions, making
10 ethically and morally difficult decisions, and taking health risks to themselves and their loved ones
11 (Maben and Bridges, 2020). This situation also affected their mental health, with high levels of
12 trauma- and stress-related symptoms that have been reported by nurses and physicians since the
13 early months of pandemic (Benfante et al., 2020).

14 The present study mainly aimed to assess the psychological impact of the COVID-19
15 outbreak on nurses and physicians working in XXX.

16 The responses of 73 nurses and 72 physicians were collected through an anonymized online
17 survey. The current data were acquired as part of a larger project investigating the psychological
18 impact of COVID-19 in XXX population. Participants were asked to indicate sociodemographic,
19 clinical and work-related information, and to complete three self-report questionnaires to assess
20 anxiety (State-Trait Anxiety Inventory-Form Y1, STAI Y1), depressive (Beck Depression
21 Inventory, BDI-II), and posttraumatic (PTSD Checklist for DSM-5, PCL-5) symptoms. The study
22 was approved by the University of XXX ethics committee (Prot. n. 142069) and was conducted in
23 accordance with the Declaration of Helsinki; all the participants gave their informed consent.

24 In order to investigate the psychological impact of the COVID-19 outbreak on nurses and
25 physicians, independent *t*-tests were performed to evaluate possible differences between those two
26 groups on sociodemographic, clinical, and work-related variables, health-related questions, and
27 psychological variables. Results of comparisons are presented in **Table 1**. Significant differences
28 between nurses and physicians were found on both Visual Analogue Scales (VAS) we used for the
29 assessment of health-related aspects. Indeed, nurses rated lower their health and reported to be more
30 worried about contracting COVID-19 with respect to physicians. Similarly, significant differences
31 emerged between nurses and physicians on anxiety symptoms and PTSS, with the former reporting
32 higher scores on the STAI Y1 and PCL-5, respectively.

33 These results are not surprising considering that nurses are usually one of the healthcare
34 professional groups dealing with patients at the frontline across healthcare settings and for this

35 reason they are continually exposed to stressful events. Particularly, during the COVID-19
36 pandemic, being more in contact with risky patients than physicians makes nurses more vulnerable
37 and at risk to develop psychological distress. In line with our findings, previous studies showed that
38 PTSS (Şahin et al., 2020; Johnson et al., 2020) and anxiety symptoms (Johnson et al., 2020) were
39 significantly higher among nurses than physicians.

40 Spending much more time with patients may also explain nurses' greater fear of contracting
41 infection. A previous study confirmed our results showing that nurses reported an increased fear of
42 being infected, and consequently of infecting their loved ones, leading to high levels of
43 psychological distress (Sampaio et al., 2020).

44 Despite this study being limited by its cross-sectional nature and its reduced sample, our
45 results highlight that nurses experience higher levels of anxiety symptoms and PTSS, and are more
46 worried about contracting COVID-19 compared to physicians. In conclusion, the present findings
47 underline the importance of paying attention to the psychological health of this healthcare
48 professional group.

49 **References**

- 50 1. Benfante, A., Di Tella, M., Romeo, A., Castelli, L., 2020. Traumatic stress in healthcare
51 workers during COVID-19 pandemic: A review of the immediate impact. *Front. Psychol.*, *11*,
52 569935. doi: 10.3389/fpsyg.2020.569935.
- 53 2. Johnson, S. U., Ebrahimi, O. V., Hoffart, A., 2020. PTSD symptoms among health workers
54 and public service providers during the COVID-19 outbreak. *PloS one*, *15*(10), e0241032.
55 doi: 10.1371/journal.pone.0241032.
- 56 3. Maben, J., Bridges, J., 2020. Covid-19: Supporting nurses' psychological and mental health.
57 *J. Clin. Nurs.*, *29*(15-16), 2742–2750. doi: 10.1111/jocn.15307. Epub 2020 Jun 2.
- 58 4. Şahin, M. K., Aker, S., Şahin, G., Karabekiroğlu, A., 2020. Prevalence of depression, anxiety,
59 distress and insomnia and related factors in healthcare workers during COVID-19 pandemic
60 in Turkey. *J. Community Health*, *45*(6), 1168–1177. doi: 10.1007/s10900-020-00921-w.
- 61 5. Sampaio, F., Sequeira, C., Teixeira, L., 2020. Nurses' mental health during the Covid-19
62 outbreak: A cross-sectional study. *J. Occup. Environ Med.*, *62*(10), 783–787. doi:
63 10.1097/JOM.0000000000001987.

64 **Table 1.** Sociodemographic, clinical, and work-related characteristics of nurse and physician
 65 groups. Mean (SD), percentage, *t*-test, chi-square test, and Cohen's *d* are listed.

	Nurses (N = 73)	Physicians (N = 72)	Test (df)	<i>p</i>	Effect size
Age (years)	44.3 (10.6)	41.5 (11.7)	t(143) = -1.51	.134	<i>d</i> = 0.25
Gender			$\chi^2(1) = 0.63$.427	
Male	18 (24.7%)	22 (30.6%)			
Female	55 (75.3%)	50 (69.4%)			
Marital status			$\chi^2(1) = 0.61$.433	
Not in a relationship	31 (42.5%)	26 (36.1 %)			
In a relationship	42 (57.5%)	46 (63.9%)			
Type of ward			$\chi^2(1) = 0.01$.925	
COVID-19 unit	32 (43.8%)	31 (43.1%)			
Other unit	41 (56.2%)	41 (56.9%)			
Medical condition			$\chi^2(1) = 2.14$.144	
Yes	23 (31.5%)	15 (20.8%)			
No	50 (68.5%)	57 (79.2%)			
Psychological aspects					
Health evaluation (VAS)*	7.11 (2.0)	8.4 (1.4)	t(126.64) = 4.62	<.001	<i>d</i> = 0.77
Health concern (VAS)#	7.04 (2.5)	6.0 (2.6)	t(143) = -2.42	.017	<i>d</i> = 0.40
STAI Y1	53.8 (14.5)	48.8 (12.9)	t(143) = -2.21	.029	<i>d</i> = 0.37
Scored above the STAI Y1 cut-off point (≥ 41)	55 (75.3%)	48 (66.7%)	$\chi^2(1) = 1.33$.250	
BDI-II	12.3 (10.5)	10.7 (8.4)	t(143) = -0.96	.340	<i>d</i> = 0.20
Scored above the BDI-II cut-off point (> 13)	21 (28.8%)	24 (33.3%)	$\chi^2(1) = 0.35$.552	
PCL-5	29.1 (18.6)	20.3 (14.2)	t(134.52) = -3.19	.002	<i>d</i> = 0.53
Scored above the PCL-5 cut-off point (≥ 33)	24 (32.9%)	14 (19.4%)	$\chi^2(1) = 3.38$.066	

66 SD = Standard Deviation; VAS = Visual Analogue Scale; BDI-II = Beck Depression Inventory;

67 STAI Y1 = State-Trait Anxiety Inventory Form Y1; PCL-5 = PTSD Checklist for DSM-5.

68 * Health evaluation question = 'How do you currently rate your health?'

69 # Health concern question = 'How concerned are you about contracting COVID-19?'