

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

A fountain of positive bubbles on a Coron's problem for a competitive weakly coupled gradient system

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1895228> since 2023-03-13T09:45:09Z

Published version:

DOI:10.1016/j.matpur.2019.09.004

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

A FOUNTAIN OF POSITIVE BUBBLES ON A CORON'S PROBLEM FOR A COMPETITIVE WEAKLY COUPLED GRADIENT SYSTEM

ANGELA PISTOIA, NICOLA SOAVE, AND HUGO TAVARES

ABSTRACT. We consider the following critical elliptic system:

$$\begin{cases} -\Delta u_i = \mu_i u_i^3 + \beta u_i \sum_{j \neq i} u_j^2 & \text{in } \Omega_\varepsilon \\ u_i = 0 \text{ on } \partial\Omega_\varepsilon, \quad u_i > 0 \text{ in } \Omega_\varepsilon \end{cases} \quad i = 1, \dots, m,$$

in a domain $\Omega_\varepsilon \subset \mathbb{R}^4$ with a small shrinking hole $B_\varepsilon(\xi_0)$. For $\mu_i > 0$, $\beta < 0$, and $\varepsilon > 0$ small, we prove the existence of a non-synchronized solution which looks like a fountain of positive bubbles, i.e. each component u_i exhibits a towering blow-up around ξ_0 as $\varepsilon \rightarrow 0$. The proof is based on the Ljapunov-Schmidt reduction method, and the velocity of concentration of each layer within a given tower is chosen in such a way that the interaction between bubbles of different components balance the interaction of the first bubble of each component with the boundary of the domain, and in addition is dominant when compared with the interaction of two consecutive bubbles of the same component.

1. INTRODUCTION

This paper deals with the existence of solutions to the elliptic critical system

$$\begin{cases} -\Delta u_i = \mu_i u_i^p + \beta u_i \sum_{j \neq i} u_j^{\frac{p+1}{2}} & \text{in } \Omega \\ u_i = 0 \text{ on } \partial\Omega, \quad u_i > 0 \text{ in } \Omega \end{cases} \quad i = 1, \dots, m, \quad (1.1)$$

when Ω is a bounded smooth domain in \mathbb{R}^N , and $p = \frac{N+2}{N-2} = 2^* - 1$, with 2^* critical Sobolev exponent. Thinking at u_i as a density function (which is natural since (1.1) is studied in connection with problems in nonlinear optics and Bose-Einstein condensation), the sign of the real parameters μ_i describes the self-interaction between particles of the same density u_i , and will always be positive: that is, we have attractive self-interaction. On the contrary, the coupling parameter β , which describes the interaction between particles of different densities, will always be negative: that is, we have repulsive mutual interaction.

Date: December 12, 2018.

2010 Mathematics Subject Classification. 35B09; 35B33; 35B44; 35J20; 35J50.

Key words and phrases. Competitive systems, Concentration Phenomena, Coron's Problem, Critical Exponent, Elliptic Systems, Fountain of Bubbles, Ljapunov-Schmidt Reduction, Positive solutions, Weakly coupled gradient systems.

Acknowledgments. A. Pistoia is partially supported by Sapienza research grant "Nonlinear PDE's in geometry and physics". N. Soave is partially supported by the PRIN-2015KB9WPT_010 Grant: "Variational methods, with applications to problems in mathematical physics and geometry". H. Tavares is partially supported by the Portuguese government through FCT - Fundação para a Ciência e a Tecnologia, I.P., both under the project PTDC/MAT-PUR/28686/2017 and through the grant UID/MAT/04561/2013. The second and third author are also supported by the ERC Advanced Grant 2013 n. 339958 "Complex Patterns for Strongly Interacting Dynamical Systems - COMPAT".

The system (1.1) has the trivial solution, i.e. all the components u_i vanish. It can also have a semi-trivial solution, i.e. only $\ell < m$ components vanish. It is clear that in this case (1.1) reduces to a system with $m - \ell$ nontrivial components, so we are naturally lead to find *fully nontrivial solutions*, namely solutions where all the components are nontrivial. In fact, we will be concerned with *positive solutions*, namely fully nontrivial solutions with $u_i > 0$ for every i .

It is useful to point out that (1.1) can have solutions with synchronized components, i.e. all the components satisfy $u_i = s_i u$ for some $s_i \in \mathbb{R}$ and u solves the single equation

$$-\Delta u = u^p \text{ in } \Omega, \quad u = 0 \text{ on } \partial\Omega, \quad u > 0 \text{ in } \Omega. \quad (1.2)$$

For instance, if the number of components is $m = 2$, the space dimension is $N = 4$ (so that $p = 3$), and

$$-\sqrt{\mu_1 \mu_2} < \beta < \min\{\mu_1, \mu_2\} \quad \text{or} \quad \beta > \max\{\mu_1, \mu_2\},$$

then a solution of (1.2) gives rise to a synchronized solution. In this way, results available for the single equation can be translated in terms of (1.1): for instance, if Ω has nontrivial \mathbb{Z}_2 -homology, then the celebrated Bahri-Coron's result [2] claims the existence of a positive solution for (1.2), and in turn this gives existence of a synchronized solution for (1.1). It is worthwhile to recall also the Coron's result [11], where the case of a domain with a small hole has been considered, namely Ω is replaced by $\Omega_\varepsilon := \Omega \setminus \overline{B_\varepsilon(\xi_0)}$, and problem (1.2) has a solution which blows-up at ξ_0 as $\varepsilon \rightarrow 0$ (see also [16, 22]). Again, this family of solutions can be used to construct an associated family of synchronized solutions for (1.1).

The assumptions on the domain are natural, since, exactly as in the scalar case, a Pohozaev-type identity shows that there is no solution if Ω is starshaped (see for instance [7, p. 519] or [8]).

The above discussion induced the first two authors to investigate the following problem: does (1.1) have non-synchronized solutions? An affirmative answer is given in [20], where (1.1) is posed in a domain $\Omega_\varepsilon \subset \mathbb{R}^N$, with $N = 3, 4$, having κ distinct holes; that is, $\Omega_\varepsilon := \Omega \setminus \bigcup_{i=1}^{\kappa} \overline{B_\varepsilon(\xi_i)}$, with $2 \leq \kappa \leq m$; for a quite general choice of interaction terms β_{ij} (which can be both of cooperative type, and of competitive type), Pistoia and Soave proved existence and concentration results of solutions whose components are splitted in several groups G_1, \dots, G_κ , in such a way that each component within a given group G_i concentrates around a point ξ_i in a somehow synchronized fashion (in the sense that the velocity of concentration of different components belonging the same group is the same), while the different groups concentrate around different points. In particular, the main results in [20] regard the case when at least two components concentrate around different points, and hence cannot be synchronized.

In view of the above discussion, it is natural to ask the following question: *if the domain has only one small hole, is it still possible to find a non-synchronized solution?* The main purpose of this paper is to give a positive answer for $\beta < 0$ and $N = 4$ - so that $p = 3$ (for a discussion of the cases $N = 3$ or other dimensions, see Remark 1.9 below). More precisely, we take

$$\Omega \subset \mathbb{R}^4 \text{ bounded domain, symmetric with respect to one of its points } \xi_0 \in \Omega, \quad (1.3)$$

i.e. $x \in \Omega$ if and only if $2\xi_0 - x \in \Omega$, and consider the following elliptic problem with $m \in \mathbb{N}$ equations:

$$\begin{cases} -\Delta u_i = \mu_i u_i^3 + \beta u_i \sum_{j \neq i} u_j^2 & \text{in } \Omega_\varepsilon \\ u_i = 0 \text{ on } \partial\Omega_\varepsilon, \quad u_i > 0 & \text{in } \Omega_\varepsilon \end{cases} \quad i = 1, \dots, m, \quad (1.4)$$

where Ω_ε is a domain with one hole, $\Omega_\varepsilon := \Omega \setminus \overline{B_\varepsilon(\xi_0)} \subset \mathbb{R}^4$, and $B_\varepsilon(\xi_0)$ denotes the open ball of \mathbb{R}^4 centered at ξ_0 with radius ε . Throughout this paper we take $\mu_i > 0$, the so called focusing case, and $\beta < 0$, which means that the coupling terms in (1.4) are of competitive type.

We find solutions of (1.4) which look like a fountain of bubbles, namely their components are a superposition of bubbles centered at ξ_0 with different rates of concentration. In particular, all the components have a towering blow-up point at ξ_0 . This new phenomena is quite surprising, since it is in sharp contrast with the case of the single equation for which positive solutions cannot have neither clustering or towering blow-up points, i.e. at every blow-up point there is at most one bubble concentrating there (see Schoen [23]). We also mention that it is somehow unexpected that in a competitive regime (with a possibly large $|\beta|$) we find solutions whose components concentrate at the same point; this is only possible because the concentration rates are different, and in particular such solutions are not synchronized.

In order to state our results we need to introduce some notations. We define

$$U_{\delta,\xi} = \alpha_4 \frac{\delta}{\delta^2 + |x - \xi|^2}, \quad \delta > 0, \quad x, \xi \in \mathbb{R}^N \quad (1.5)$$

(a *bubble*) with $\alpha_4 = 2\sqrt{2}$: these functions are all the positive solutions of the problem

$$-\Delta U = U^3, \quad U \in \mathcal{D}^{1,2}(\mathbb{R}^4).$$

(see [1, 4, 24]). Also, we denote by $P_\varepsilon : \mathcal{D}^{1,2}(\mathbb{R}^4) \rightarrow H_0^1(\Omega_\varepsilon)$ the projection map and we define the projection of the bubble defined in (1.5) as $W := P_\varepsilon U_{\delta,\xi} \in H_0^1(\Omega_\varepsilon)$, which is the unique solution of

$$-\Delta W = -\Delta U_{\delta,\xi} = U_{\delta,\xi}^3 \text{ in } \Omega_\varepsilon, \quad W = 0 \text{ on } \partial\Omega_\varepsilon. \quad (1.6)$$

Take $k \in \mathbb{N}$ (the total number of bubbles) larger than or equal to m . Consider $I_1, \dots, I_m \subset \{1, \dots, k\}$ satisfying the following properties:

- (1) $1 \in I_1$;
- (2) $I_i \neq \emptyset$ for every $i = 1, \dots, m$;
- (3) $I_i \cap I_j = \emptyset$ whenever $i \neq j$;
- (4) $I_1 \cup \dots \cup I_m = \{1, \dots, k\}$;
- (5) for every $j \in \{1, \dots, k\}$ and $i \in \{1, \dots, m\}$, if $j \in I_i$ then $j - 1, j + 1 \notin I_i$.

Observe that one considers condition (1) without loss of generality, simply to fix ideas and simplify some statements. Conditions (2)-(3)-(4) imply that I_1, \dots, I_m form a partition of $\{1, \dots, k\}$, while condition (5) means that each set I_i does not contain two consecutive integers. Our main result is the following

Theorem 1.1. *Take Ω satisfying (1.3) and let $\mu_i > 0$, $\beta < 0$. For any integer $k \geq m$ and for every partition I_1, \dots, I_m of $\{1, \dots, k\}$ satisfying (1)–(5), there exists $\varepsilon_0 > 0$ such that for any $\varepsilon \in (0, \varepsilon_0)$ problem (1.4) has a solution (symmetric with respect to ξ_0) of the form*

$$u_{i,\varepsilon} = \mu_i^{-\frac{1}{p-1}} \sum_{j \in I_i} P_\varepsilon U_{\delta_j^\varepsilon, \xi_0} + \phi_i^\varepsilon, \quad i = 1, \dots, m$$

with

$$\delta_j^\varepsilon = d_j^\varepsilon \varepsilon^{\frac{j}{k+1}} \left(\log \frac{1}{\varepsilon} \right)^{\frac{1}{2} - \frac{j}{k+1}} \text{ for some } d_j^\varepsilon \rightarrow d_j^*, \quad j = 1, \dots, k \quad (1.7)$$

for

$$d_j^* = \Gamma^{\frac{i}{2(k+1)}} (A^2 \tau(0))^{\frac{i}{2(k+1)} - \frac{1}{2}} \left(\frac{|\beta| \alpha_4^4 |\mathbb{S}^3|}{k+1} \right)^{\frac{1}{2} - \frac{i}{k+1}}$$

(see the upcoming (1.16) and (1.17) for the expressions of the constants A, Γ) and

$$\|\phi_i^\varepsilon\|_{H_0^1(\Omega_\varepsilon)} \rightarrow 0 \text{ as } \varepsilon \rightarrow 0, \quad i = 1, \dots, m.$$

Remark 1.2. As stated in the theorem, each component of the solution, $u_{i,\varepsilon}$, belongs to the space

$$H_{\varepsilon,\xi_0} = \{u \in H_0^1(\Omega_\varepsilon) : u(x) = u(2\xi_0 - x) \forall x \in \Omega_\varepsilon\}.$$

Since also $U_{\delta,\xi}$ is symmetry with respect to ξ_0 , then $P_\varepsilon U_{\delta_j^\varepsilon,\xi_0} \in H_{\varepsilon,\xi_0}$, as well as the remainder terms ϕ_i^ε .

In order to better explain our result, let us take a particular case of (1.4) and Theorem 1.1:

$$m = 2, \quad k \geq 2, \quad \text{and } N = 4 \quad (\text{so that } p = 3). \quad (1.8)$$

and the following partition of $\{1, \dots, k\}$:

$$I_1 = \{\text{odd numbers between 1 and } k\}, \quad I_2 = \{\text{even numbers between 1 and } k\}. \quad (1.9)$$

Clearly, I_1, I_2 satisfies conditions (1)–(5), and it is actually the only admissible partition for $m = 2$. Problem (1.4) now reads as

$$\begin{cases} -\Delta u_1 = \mu_1 u_1^3 + \beta u_1 u_2^2 \text{ in } \Omega_\varepsilon, \\ -\Delta u_2 = \mu_2 u_2^3 + \beta u_1^2 u_2 \text{ in } \Omega_\varepsilon \\ u_1 = u_2 = 0 \text{ on } \partial\Omega_\varepsilon, \quad u_1, u_2 > 0 \text{ in } \Omega_\varepsilon. \end{cases} \quad (1.10)$$

In this particular situation, Theorem 1.1 can be stated in the following way.

Theorem 1.3. *Take Ω satisfying (1.3) and let $\mu_1, \mu_2 > 0, \beta < 0$. For any integer $k \geq 2$, let I_1, I_2 be respectively the set of all odd and even numbers between 1 and k , as in (1.9). Then there exists $\varepsilon_0 > 0$ such that for any $\varepsilon \in (0, \varepsilon_0)$ problem (1.10) has a solution (symmetric with respect to ξ_0) of the form*

$$u_{1,\varepsilon} = \mu_1^{-\frac{1}{2}} \sum_{j \in I_1} P_\varepsilon U_{\delta_j^\varepsilon,\xi_0} + \phi_1^\varepsilon \quad \text{and} \quad u_{2,\varepsilon} = \mu_2^{-\frac{1}{2}} \sum_{j \in I_2} P_\varepsilon U_{\delta_j^\varepsilon,\xi_0} + \phi_2^\varepsilon$$

with

$$\delta_j^\varepsilon = d_j^\varepsilon \varepsilon^{\frac{j}{k+1}} \left(\log \frac{1}{\varepsilon} \right)^{\frac{1}{2} - \frac{j}{k+1}} \quad \text{for some } d_j^\varepsilon \rightarrow d_j^*, \quad j = 1, \dots, k \quad (1.11)$$

for

$$d_j^* = \Gamma^{\frac{j}{2(k+1)}} (A^2 \tau(0))^{\frac{j}{2(k+1)} - \frac{1}{2}} \left(\frac{|\beta| \alpha_4^4 |\mathbb{S}^3|}{k+1} \right)^{\frac{1}{2} - \frac{j}{k+1}}$$

and

$$\|\phi_1^\varepsilon\|_{H_0^1(\Omega_\varepsilon)} \rightarrow 0, \quad \|\phi_2^\varepsilon\|_{H_0^1(\Omega_\varepsilon)} \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0.$$

In order to avoid insignificant technicalities that would make the presentation harder to follow, we will simply prove Theorem 1.3; in order to convince the reader that the proof of Theorem 1.4 follows precisely in the same way we will make some remarks along the paper (see Remarks 2.2, 3.7, 4.8 and 5.2).

Our result is inspired by the construction performed by Musso and Pistoia in [18] and Ge, Musso and Pistoia in [12], where the authors built sign-changing solutions to Coron's problem whose shape resembles a superposition of bubbles centered at the point ξ_0 with alternating sign and with different rate of concentration. The proof here also follows the same scheme which is based on a Ljapunov-Schmidt procedure: we find a good first order approximation term (see (3.2)), we perform a linear theory for the linearized system around the ansatz (see Proposition 3.2), we reduce the problem to a finite dimensional one (see Proposition 3.1) and finally we study the reduced problem (see Section 4). However, the main steps of our proof require rather delicate and careful estimates, see for instance the estimates involving the interacting term in the study of the

linear part in Subsection 3.1, the asymptotic expansion of the interaction energy (Lemma 4.4), and the estimate of the remainder term in Lemma 4.6. Indeed, the interaction between bubbles of different components has to balance the interaction of the first bubble of each component with the boundary of the domain, and most of all it has to be dominant compared with the interaction of two consecutive bubbles of the same component. Actually, this is possible because of the presence of an $|\log \varepsilon|$ -order term which turns out to be crucial in our construction (see estimate (4.30)).

Remark 1.4. We prove the existence of solutions which look like fountains of positive bubble all centered at the point ξ_0 when Ω is symmetric with respect ξ_0 . It is clear that using the same arguments of Ge, Musso and Pistoia [12] we can remove the symmetry assumption, just centering all the bubbles U_{δ_i, ξ_i} at suitable points $\xi_i = \xi_i(\varepsilon)$ which approach ξ_0 with a suitable rate as $\varepsilon \rightarrow 0$.

Remark 1.5. For the sake of completeness, we also mention some recent results concerning the existence of solutions to system (1.1) when Ω is the whole space \mathbb{R}^N . As far as we know, all the results deal with systems with only two components. Guo, Li and Wei in [15] established the existence of infinitely many positive nonradial solutions of (1.1), only when $N = 3$, in the competitive case. Peng, Peng and Wang discussed in [19] uniqueness of the least energy solution for $\beta > 0$, and the non-degeneracy of the manifold of the synchronized positive solutions. Clapp and Pistoia in [10] proved that system (1.1) in any dimension has infinitely many fully nontrivial solutions, which are not conformally equivalent. Gladiali, Grossi and Troestler in [13, 14] obtained radial and nonradial solutions to some critical systems like (1.1) using bifurcation methods.

Remark 1.6. A Brezis-Nirenberg type problem has been studied for systems, see for instance [6, 7, 8] for existence results, while for concentration and blow-up type results see [5, 21].

Remark 1.7. As already mentioned, appropriate assumptions on β allows to obtain a synchronized solution to (1.1) if Ω has nontrivial \mathbb{Z}_2 -homology. We conjecture that system (1.1) has at least one (actually we would say infinitely many) positive non-synchronized solution if Ω has nontrivial \mathbb{Z}_2 -homology (as in Bahri-Coron's result for the single equation (1.2)) and $\beta < 0$ is arbitrary. A first attempt in this direction is due to Clapp and Faya [9], who establish the existence of a prescribed number of fully nontrivial solutions to the system with only two components under suitable symmetry assumptions on the topologically nontrivial domain Ω .

We would like to remark that the difficulty in finding positive solutions to system (1.1), even with only two components, is similar to the difficulty in finding sign-changing solutions for the single equation (1.2). One key point is the blow-up analysis of solutions: in the case of positive solutions the blow-up, whenever it occurs, is isolated and simple, while in the case of sign-changing solution multiple bubbling naturally appears.

Without loss of generality, we will work from now on with

$$\mu_1 = \mu_2 = 1, \quad \text{and take} \quad \xi_0 = 0 \in \Omega, \tag{1.12}$$

assuming that Ω is symmetric with respect to the origin. Observe that we are conduced to such situation by eventually replacing u_i with $\mu_i^{-\frac{1}{2}} u_i(x + \xi_0)$.

Remark 1.8. Solutions of (1.4) correspond to critical points with nontrivial components of the C^1 -energy functional $J_\varepsilon : H_0^1(\Omega; \mathbb{R}^m) \rightarrow \mathbb{R}$ defined by

$$J_\varepsilon(u_1, \dots, u_m) = \sum_{i=1}^m \int_{\Omega_\varepsilon} \left(\frac{|\nabla u_i|^2}{2} - \frac{\mu_i (u_i^+)^{p+1}}{p+1} \right) - \frac{2\beta}{p+1} \sum_{\substack{i,j=1 \\ i < j}}^m \int_{\Omega_\varepsilon} |u_i|^{\frac{p+1}{2}} |u_j|^{\frac{p+1}{2}}.$$

Indeed, if (u_1, \dots, u_m) is a critical point of J_ε , then it satisfies

$$-\Delta u_i = \mu_i (u_i^+)^p + \beta \sum_{j \neq i} u_i |u_i|^{\frac{p-3}{2}} |u_j|^{\frac{p+1}{2}}, \quad i = 1, \dots, m.$$

Multiplying this equation by u_i^- and integrating by parts yields (since $\beta < 0$)

$$0 \geq - \int_{\Omega_\varepsilon} |\nabla u_i^-|^2 = -\beta \sum_{j \neq i} \int_{\Omega_\varepsilon} |u_i^-|^{\frac{p+1}{2}} |u_j|^{\frac{p+1}{2}} \geq 0.$$

If $u_i \not\equiv 0$, then by the maximum principle we deduce that $u_i > 0$.

Remark 1.9. The Sobolev critical exponent is defined only for $N \geq 3$. On the other hand, for p defined as before, the right hand sides of (1.1) are C^1 nonlinearities if and only if we have $\frac{p-1}{2} \geq 1$, if and only if $N \leq 4$. Therefore, it is reasonable to work in dimension $N = 3$ or $N = 4$. Here we chose to deal with the case $N = 4$ only since it requires less technicalities: all the exponents are positive integers, which makes some expansions explicit. Using Taylor expansions we could have tackled the case $N = 3$. We conjecture that in this case the main results (and in particular the rates) would be the same.

Remark 1.10. A similar approach could also be used to find solutions for critical systems in pierced domains when the interaction term is more in general like (e.g. Lotka-Volterra systems)

$$\begin{cases} -\Delta u_i = \mu_i u_i^p + \beta_i u_i^{q_i} \sum_{j \neq i} u_j^{q_j} & \text{in } \Omega_\varepsilon \\ u_i = 0 & \text{on } \partial\Omega_\varepsilon, \quad u_i > 0 & \text{in } \Omega_\varepsilon \end{cases} \quad i = 1, \dots, m, \quad (1.13)$$

when $\mu_i > 0$, $\beta_i < 0$ and $q_i, q_j > 1$. In the non-variational cases, one has to replace the asymptotic estimates on the energy of Section 4 with an argument that simply uses the system like in [17, Section 2].

Notations. Working with dimension $N = 4$, we deal with the following bubbles concentrated at the origin

$$U_{\delta,0}(x) = \alpha_4 \frac{\delta}{\delta^2 + |x|^2}$$

(where $\alpha_4 = 2\sqrt{2}$), which we denote also by U_δ ; in many cases we deal with different concentration parameters δ_i , $i = 1, \dots, k$, and we shall simply write $U_{\delta_i} = U_i$. These correspond to all positive solutions of $-\Delta U = U^3$ in \mathbb{R}^4 which are symmetric with respect to the origin. It is well known (see [3]) that the space of solutions of the linearized equation

$$-\Delta V = 3U_\delta^2 V \quad (1.14)$$

has dimension $4 + 1 = 5$ in $\mathcal{D}^{1,2}(\mathbb{R}^5)$, being spanned by

$$\frac{\partial U_\delta}{\partial \delta}(x) = \alpha_4 \frac{|x|^2 - \delta^2}{(\delta^2 + |x|^2)^2}, \quad \frac{\partial U_\delta}{\partial \xi_i}(x) = 2\alpha_4 \frac{\delta x_i}{(\delta^2 + |x|^2)^2}, \quad i = 1, \dots, 4.$$

Therefore, the space of solutions to (1.14) which belong to

$$\mathcal{D}_s^{1,2}(\mathbb{R}^4) := \{\psi \in \mathcal{D}^{1,2}(\mathbb{R}^4) : \psi(-x) = \psi(x) \forall x \in \mathbb{R}^4\}$$

has dimension 1, being spanned by $\frac{\partial U_\delta}{\partial \delta}$. For future convenience, we observe that

$$\left| \frac{\partial U_\delta}{\partial \delta}(x) \right| \leq \frac{U_\delta(x)}{\delta}. \quad (1.15)$$

We take the following inner product and norm in $H_0^1(\Omega_\varepsilon)$:

$$\langle u, v \rangle_{H_0^1} := \int_{\Omega_\varepsilon} \nabla u \cdot \nabla v, \quad \|u\|_{H_0^1}^2 = \int_{\Omega_\varepsilon} |\nabla u|^2$$

and the standard L^p norm by $\|\cdot\|_p$ (we omit the dependence on ε for simplicity).

The Green function of the Laplace operator in Ω with Dirichlet boundary conditions is denoted by $G(x, y)$, and can be decomposed as

$$G(x, y) = \frac{\gamma_4}{|x - y|^2} - H(x, y),$$

where $\gamma_4 := (2|\partial B_1|)^{-1}$, and H is the regular part of G which, for every $x \in \Omega$, satisfies

$$\begin{cases} -\Delta_y H(x, y) = 0 & \text{for } y \in \Omega, \\ H(x, y) = \frac{\gamma_4}{|x - y|^2} & \text{for } y \in \partial\Omega. \end{cases}$$

The Robin function of Ω is defined as $\tau(x) := H(x, x)$, and satisfies $\tau(x) \rightarrow +\infty$ as $\text{dist}(x, \partial\Omega) \rightarrow 0$. Throughout the paper, we will always label the following constants:

$$A := \int_{\mathbb{R}^4} U_{1,0}^3 = \int_{\mathbb{R}^4} \frac{\alpha_4^3}{(1 + |y|^2)^3} dy, \quad B := \int_{\mathbb{R}^4} U_{1,0}^4 = \int_{\mathbb{R}^4} \frac{\alpha_4^4}{(1 + |y|^2)^4} dy, \quad (1.16)$$

$$\Gamma := \int_{\mathbb{R}^N} \frac{\alpha_4^4}{|y|^2(1 + |y|^2)^3} dy, \quad (1.17)$$

and use $B_\varepsilon, \partial B_\varepsilon$ instead of $B_\varepsilon(0), \partial B_\varepsilon(0)$ respectively. We will denote the $L^p(\Omega_\varepsilon)$ norms by $\|\cdot\|_{L^p}$, while $\|u\|_{H_0^1}^2 := \int_{\Omega_\varepsilon} |\nabla u|^2$ for every $u \in H_0^1(\Omega_\varepsilon)$.

2. THE ANSATZ AND REDUCTION SCHEME

Recall that, without loss of generality, we assume (1.12); due to the symmetry, by the principle of symmetric criticality we can work in the space

$$H_\varepsilon := H_{\varepsilon,0} = \{u \in H_0^1(\Omega_\varepsilon) : u(-x) = u(x) \ \forall x \in \Omega_\varepsilon\}. \quad (2.1)$$

We deal with solutions of

$$\begin{cases} -\Delta u_1 = f(u_1) + \beta u_1 u_2^2 \\ -\Delta u_2 = f(u_2) + \beta u_2 u_1^2 \\ u_1, u_2 \in H_0^1(\Omega_\varepsilon), \end{cases} \quad (2.2)$$

where $f : \mathbb{R} \rightarrow \mathbb{R}$, $f(s) := (s^+)^3$. Denote by $\mathcal{I}^* : L^{\frac{4}{3}}(\Omega_\varepsilon) \rightarrow H_0^1(\Omega_\varepsilon)$ the adjoint operator of the canonical Sobolev embedding $\mathcal{I} : H_0^1(\Omega_\varepsilon) \rightarrow L^4(\Omega_\varepsilon)$. This means that $v := \mathcal{I}^* u$ can be defined as the (unique) weak solution of

$$-\Delta v = u \text{ in } \Omega_\varepsilon, \quad v = 0 \text{ on } \partial\Omega_\varepsilon.$$

Observe that, if u is symmetric with respect to the origin, so is $\mathcal{I}^* u$. The operator \mathcal{I}^* is continuous: there exists $C > 0$, independent of ε , such that

$$\|\mathcal{I}^* u\|_{H_0^1} \leq C \|u\|_{L^{\frac{4}{3}}} \quad \forall u \in L^{\frac{4}{3}}(\Omega_\varepsilon).$$

Using this operator, we can rewrite (2.2) as

$$u_1 = \mathcal{I}^* (f(u_1) + \beta u_1 u_2^2), \quad u_2 = \mathcal{I}^* (f(u_2) + \beta u_2 u_1^2). \quad (2.3)$$

Denote $U_j := U_{\delta_j}$ for $j = 1, \dots, k$. Our ansatz is the following: for any integer $k \geq 2$, we look for a solution of (2.2) in H_ε of the form

$$u_1 = \sum_{j \in I_1} P_\varepsilon U_j + \phi_1 \quad \text{and} \quad u_2 = \sum_{j \in I_2} P_\varepsilon U_j + \phi_2, \quad (2.4)$$

where

$$\delta_j = d_j \varepsilon^{\frac{j}{k+1}} \left(\log \frac{1}{\varepsilon} \right)^{\frac{1}{2} - \frac{j}{k+1}}, \quad j = 1, \dots, k, \quad (2.5)$$

$\mathbf{d} = (d_1, \dots, d_k)$ belongs to the set

$$X_\eta = \{ \mathbf{d} \in \mathbb{R}^k : \eta < d_1, \dots, d_k < 1/\eta \} \quad \text{for some } \eta \ll 1, \quad (2.6)$$

and $\phi_1, \phi_2 \in H_\varepsilon$.

Remark 2.1. For future reference, we collect in this remark several important relations between the different rates δ_j . Given $\eta > 0$, we have

$$\frac{\varepsilon}{\delta_j} = \frac{1}{d_j} \varepsilon^{\frac{k+1-j}{k+1}} \left(\log \frac{1}{\varepsilon} \right)^{\frac{j}{k+1} - \frac{1}{2}} \rightarrow 0 \quad \text{and} \quad \frac{\delta_{j+1}}{\delta_j} = \frac{d_{j+1}}{d_j} \varepsilon^{\frac{1}{k+1}} \left(\log \frac{1}{\varepsilon} \right)^{-\frac{1}{k+1}} \rightarrow 0 \quad (2.7)$$

as $\varepsilon \rightarrow 0$, uniformly for $\mathbf{d} \in X_\eta$.

For each $\varepsilon > 0$ small, our aim is to find $\eta > 0$, $\mathbf{d} \in X_\eta$ and $\phi_1, \phi_2 \in H_\varepsilon$ such that, for $i, j = 1, 2$, $i \neq j$,

$$\sum_{l \in I_i} P_\varepsilon U_l + \phi_i = \mathcal{I}^* \left(f \left(\sum_{l \in I_i} P_\varepsilon U_l + \phi_i \right) + \beta \left(\sum_{l \in I_i} P_\varepsilon U_l + \phi_i \right) \left(\sum_{l \in I_j} P_\varepsilon U_l + \phi_j \right)^2 \right). \quad (2.8)$$

Given $\varepsilon > 0$ and $d_1, \dots, d_k > 0$, for δ_i defined as before define

$$\psi_i(x) := \frac{\partial U_i}{\partial \delta_i}(x) = \alpha_4 \frac{|x|^2 - \delta_i^2}{(\delta_i^2 + |x|^2)^2}$$

(recall the Notation section) and

$$K_1 = K_{1, \mathbf{d}, \varepsilon} := \text{span} \{ P_\varepsilon \psi_j : j \in I_1 \}, \quad K_2 = K_{2, \mathbf{d}, \varepsilon} := \text{span} \{ P_\varepsilon \psi_j : j \in I_2 \}, \quad \mathbf{K}_{\mathbf{d}, \varepsilon} := K_1 \times K_2.$$

Observe that $\mathbf{K}_{\mathbf{d}, \varepsilon}^\perp = K_1^\perp \times K_2^\perp$. Moreover, consider the projection maps

$$\Pi_i : H_\varepsilon \rightarrow K_i, \quad \Pi_i^\perp : H_\varepsilon \rightarrow K_i^\perp, \quad i = 1, 2.$$

We can rewrite (2.11) as a system of 4 equations: for $i, j = 1, 2$, $j \neq i$,

$$\Pi_i \left(\sum_{l \in I_i} P_\varepsilon U_l + \phi_i \right) = \Pi_i \circ \mathcal{I}^* \left(f \left(\sum_{l \in I_i} P_\varepsilon U_l + \phi_i \right) + \beta \left(\sum_{l \in I_i} P_\varepsilon U_l(x) + \phi_i \right) \left(\sum_{l \in I_j} P_\varepsilon U_l + \phi_j \right)^2 \right), \quad (2.9)$$

$$\Pi_i^\perp \left(\sum_{l \in I_i} P_\varepsilon U_l + \phi_i \right) = \Pi_i^\perp \circ \mathcal{I}^* \left(f \left(\sum_{l \in I_i} P_\varepsilon U_l + \phi_i \right) + \beta \left(\sum_{l \in I_i} P_\varepsilon U_l + \phi_i \right) \left(\sum_{l \in I_j} P_\varepsilon U_l(x) + \phi_j \right)^2 \right). \quad (2.10)$$

In the next section, given $\varepsilon, \eta > 0$ sufficiently small and $\mathbf{d} \in X_\eta$, we find a unique $(\phi_1, \phi_2) = (\phi_1^{\mathbf{d}, \varepsilon}, \phi_2^{\mathbf{d}, \varepsilon}) \in K_{\mathbf{d}, \varepsilon}^\perp$ solution to (2.10). By plugging this result in (2.9), we end up having a problem with unknown $\mathbf{d} \in \mathbb{R}^k$ (thus a finite dimensional problem), which can be stated in terms of a *reduced energy*. We analyse this reduced energy in Section 4.

Remark 2.2. For the general system (1.4) and given a partition I_1, \dots, I_m of $\{1, \dots, k\}$, the ansatz is exactly the same: $u_i = \sum_{j \in I_i} U_j + \phi_i$, for $i = 1, \dots, m$, where $\phi_i \in K_i$. We denote in this case $\mathbf{K}_{\mathbf{d}, \varepsilon} = K_1^\perp \times \dots \times K_m^\perp$, and split the system of m equations:

$$\sum_{l \in I_i} P_\varepsilon U_l + \phi_i = \mathcal{I}^* \left(f \left(\sum_{l \in I_i} P_\varepsilon U_l + \phi_i \right) + \beta \left(\sum_{l \in I_i} P_\varepsilon U_l + \phi_i \right) \sum_{\substack{j=1 \\ j \neq i}}^m \left(\sum_{l \in I_j} P_\varepsilon U_l + \phi_j \right)^2 \right) \quad (2.11)$$

($i = 1, \dots, m$) in $2m$ equations using the projection maps Π_i and Π_i^\perp .

3. REDUCTION TO A FINITE DIMENSIONAL PROBLEM

In this section we study the solvability of (2.10). We rewrite (2.10) as

$$L_{\mathbf{d}, \varepsilon}^i(\phi) = N_{\mathbf{d}, \varepsilon}^i(\phi) + R_{\mathbf{d}, \varepsilon}^i, \quad (3.1)$$

where L stays for the linear part

$$L_{\mathbf{d}, \varepsilon}^1(\phi) = \Pi_1^\perp \left\{ \phi_1 - \mathcal{I}^* \left[f' \left(\sum_{j \in I_1} P_\varepsilon U_j \right) \phi_1 + \beta \left(\sum_{j \in I_2} P_\varepsilon U_j \right)^2 \phi_1 + 2\beta \left(\sum_{j \in I_1} P_\varepsilon U_j \right) \left(\sum_{j \in I_2} P_\varepsilon U_j \right) \phi_2 \right] \right\}, \quad (3.2)$$

N stays for the nonlinear part

$$\begin{aligned} N_{\mathbf{d}, \varepsilon}^1(\phi) &= \Pi_1^\perp \circ \mathcal{I}^* \left[f \left(\sum_{j \in I_1} P_\varepsilon U_j + \phi_1 \right) - f \left(\sum_{j \in I_1} P_\varepsilon U_j \right) - f' \left(\sum_{j \in I_1} P_\varepsilon U_j \right) \phi_1 \right. \\ &\quad + \beta \left(\sum_{j \in I_1} P_\varepsilon U_j + \phi_1 \right) \left(\sum_{j \in I_2} P_\varepsilon U_j + \phi_2 \right)^2 - \beta \left(\sum_{j \in I_1} P_\varepsilon U_j \right) \left(\sum_{j \in I_2} P_\varepsilon U_j \right)^2 \\ &\quad \left. - \beta \left(\sum_{j \in I_2} P_\varepsilon U_j \right)^2 \phi_1 - 2\beta \left(\sum_{j \in I_1} P_\varepsilon U_j \right) \left(\sum_{j \in I_2} P_\varepsilon U_j \right) \phi_2 \right] \\ &= \Pi_1^\perp \circ \mathcal{I}^* \left[f \left(\sum_{j \in I_1} P_\varepsilon U_j + \phi_1 \right) - f \left(\sum_{j \in I_1} P_\varepsilon U_j \right) - f' \left(\sum_{j \in I_1} P_\varepsilon U_j \right) \phi_1 \right. \\ &\quad \left. + \beta \left(\sum_{j \in I_1} P_\varepsilon U_j \right) \phi_2^2 + 2\beta \left(\sum_{j \in I_2} P_\varepsilon U_j \right) \phi_1 \phi_2 + \beta \phi_1 \phi_2^2 \right], \end{aligned} \quad (3.3)$$

and R is the remainder term

$$\begin{aligned} R_{\mathbf{d}, \varepsilon}^1 &= \Pi_1^\perp \left\{ - \sum_{j \in I_1} P_\varepsilon U_j + \mathcal{I}^* \left[f \left(\sum_{j \in I_1} P_\varepsilon U_j \right) + \beta \left(\sum_{j \in I_1} P_\varepsilon U_j \right) \left(\sum_{j \in I_2} P_\varepsilon U_j \right)^2 \right] \right\} \\ &= \Pi_1^\perp \circ \mathcal{I}^* \left[f \left(\sum_{j \in I_1} P_\varepsilon U_j \right) - \sum_{j \in I_1} f(U_j) + \beta \left(\sum_{j \in I_1} P_\varepsilon U_j \right) \left(\sum_{j \in I_2} P_\varepsilon U_j \right)^2 \right] \end{aligned} \quad (3.4)$$

where the last equality is a consequence of the definitions of \mathcal{I}^* and of f (analogue expressions hold for $L_{\mathbf{d}, \varepsilon}^2$, $N_{\mathbf{d}, \varepsilon}^2$ and $R_{\mathbf{d}, \varepsilon}^2$).

We also define

$$\mathbf{L}_{\mathbf{d},\varepsilon} := (L_{\mathbf{d},\varepsilon}^1, L_{\mathbf{d},\varepsilon}^2) : \mathbf{K}_{\mathbf{d},\varepsilon}^\perp \rightarrow \mathbf{K}_{\mathbf{d},\varepsilon}^\perp,$$

and $\mathbf{R}_{\mathbf{d},\varepsilon}$ and $\mathbf{N}_{\mathbf{d},\varepsilon}$ in an analogue way.

Proposition 3.1. *Let $\beta < 0$. Then for every $\eta > 0$ sufficiently small there exists $\varepsilon_0 > 0$ and $C > 0$ such that, whenever $\varepsilon \in (0, \varepsilon_0)$ and $\mathbf{d} \in X_\eta$, there exists a unique function $\phi = \phi^{\mathbf{d},\varepsilon} \in \mathbf{K}_{\mathbf{d},\varepsilon}^\perp$ solving the equation*

$$\mathbf{L}_{\mathbf{d},\varepsilon}(\phi) = \mathbf{R}_{\mathbf{d},\varepsilon} + \mathbf{N}_{\mathbf{d},\varepsilon}(\phi).$$

and satisfying

$$\|\phi^{\mathbf{d},\varepsilon}\|_{H_0^1(\Omega_\varepsilon)} \leq C\varepsilon^{\frac{1}{k+1}} \left(\log \left(\frac{1}{\varepsilon} \right) \right)^{-\frac{1}{k+1}} = o(\delta_1)$$

Moreover, the map $X_\eta \rightarrow \mathbf{K}_{\mathbf{d},\varepsilon}^\perp$, $\mathbf{d} \mapsto \phi^{\mathbf{d},\varepsilon}$ is of class \mathcal{C}^1 .

The proof of the proposition takes the rest of this section, and is divided into several intermediate lemmas.

3.1. Study of the linear part. As a first step, it is important to understand the solvability of the linear problem associated with (3.1), i.e.

$$L_{\mathbf{d},\varepsilon}^i(\phi) = f_i, \quad \text{with } f_i \in K_i^\perp. \quad (3.5)$$

Proposition 3.2. *For every $\eta > 0$ small enough there exists $\varepsilon_0 > 0$ small, and $C > 0$, such that if $\varepsilon \in (0, \varepsilon_0)$ then*

$$\|\mathbf{L}_{\mathbf{d},\tau,\varepsilon}(\phi)\|_{H_0^1(\Omega_\varepsilon)} \geq C\|\phi\|_{H_0^1(\Omega_\varepsilon)} \quad \forall \phi \in H_0^1(\Omega_\varepsilon, \mathbb{R}^2) \quad (3.6)$$

for every $\mathbf{d} \in X_\eta$. Moreover, $\mathbf{L}_{\mathbf{d},\varepsilon}$ is invertible in $\mathbf{K}_{\mathbf{d},\varepsilon}^\perp$, with continuous inverse.

The long proof proceeds by contradiction. For a fixed $\eta > 0$ small, let us suppose that there exist sequences

$$\{\varepsilon_n\} \subset \mathbb{R}^+, \quad \varepsilon_n \rightarrow 0, \quad \{\mathbf{d}_n\} \subset X_\eta, \quad \{\phi_n\} \subset K_{1,n}^\perp \times K_{2,n}^\perp$$

such that

$$\|\phi_n\|_{H_0^1(\Omega_{\varepsilon_n})} = 1 \quad \text{and} \quad \|\mathbf{L}_n(\phi_n)\|_{H_0^1(\Omega_{\varepsilon_n})} \rightarrow 0$$

as $n \rightarrow \infty$, where we wrote $K_{i,n} := K_{i,\mathbf{d}_n,\varepsilon_n}$ and $\mathbf{L}_n := \mathbf{L}_{\mathbf{d}_n,\varepsilon_n}$ for short. In the same spirit, in this proof we write $P_n := P_{\varepsilon_n}$, $U_{i,n} := U_{\delta_{i,n},0}$, $\psi_{i,n} := \psi_{\delta_{i,n},0}$, and $\Omega_n := \Omega_{\varepsilon_n}$.

Let $\mathbf{h}_n := \mathbf{L}_n(\phi_n)$. Then, by definition of \mathbf{L}_n ,

$$\begin{aligned} \phi_{1,n} &= h_{1,n} + w_{1,n} \\ &+ \mathcal{I}^* \left[3 \left(\sum_{j \in I_1} P_n U_{j,n} \right)^2 \phi_{1,n} + \beta \left(\sum_{j \in I_2} P_n U_{j,n} \right)^2 \phi_{1,n} + 2\beta \left(\sum_{j \in I_1} P_n U_{j,n} \right) \left(\sum_{j \in I_2} P_n U_{j,n} \right) \phi_{2,n} \right] \end{aligned} \quad (3.7)$$

(an analogue equation holds for $\phi_{2,n}$) for some $w_{i,n} \in K_{i,n}$.

Lemma 3.3. $\|w_{i,n}\|_{H_0^1(\Omega_n)} \rightarrow 0$ as $n \rightarrow \infty$.

Proof. We focus on $w_{1,n}$, the proof for $w_{2,n}$ is analogue. As $w_{1,n} \in K_{1,n} = \text{span}\{P_n \psi_{j,n} : j \in I_1\}$, there exist constants $c_{j,n}$ such that

$$w_{1,n} = \sum_{j \in I_1} c_{j,n} \delta_{j,n} P_n \psi_{j,n}.$$

Now we consider the scalar product in $H_0^1(\Omega_\varepsilon)$ of both sides in (3.7) with $\delta_{i,n}P_n\psi_{i,n}$, with $i \in I_1$: as $h_{1,n}, \phi_{1,n} \in K_{1,n}^\perp$, we obtain

$$\begin{aligned} \delta_{i,n} \int_{\Omega_n} \nabla w_{1,n} \cdot \nabla (P_n \psi_{i,n}) &= 3\delta_{i,n} \int_{\Omega_n} \left(\sum_{j \in I_1} P_n U_{j,n} \right)^2 \phi_{1,n} (P_n \psi_{i,n}) \\ &+ \delta_{i,n} \beta \int_{\Omega_n} \left(\sum_{j \in I_2} P_n U_{j,n} \right)^2 \phi_{1,n} (P_n \psi_{i,n}) \\ &+ 2\beta \delta_{i,n} \int_{\Omega_n} \left(\sum_{j \in I_1} P_n U_{j,n} \right) \left(\sum_{j \in I_2} P_n U_{j,n} \right) \phi_{2,n} (P_n \psi_{i,n}). \end{aligned} \quad (3.8)$$

The left hand side can be estimated using [12, Remark 5.2] and (1.15) (see also [21, p. 417], noting that therein $\psi_{i,n}$ corresponds to $\delta_{i,n}\psi_{i,n}$ in the present paper) and obtaining

$$\int_{\Omega_n} \nabla w_{1,n} \cdot \nabla (\delta_{i,n} P_n \psi_{i,n}) = c_{i,n}(\sigma_0 + o(1)) + o(1) \sum_{\substack{j \in I_1 \\ j \neq i}} c_{j,n} \quad (3.9)$$

as $n \rightarrow \infty$, where

$$\sigma_0 = 3\alpha_4^4 \int_{\mathbb{R}^4} \frac{(|y|^2 - 1)^2}{(1 + |y|^2)^6} dy.$$

The first integral on the right hand side in (3.8) can be estimated as in [12, Formula (5.7)]:

$$3\delta_{i,n} \int_{\Omega_n} \left(\sum_{j \in I_1} P_n U_{j,n} \right)^2 \phi_{1,n} (P_n \psi_{i,n}) = o(1) \quad (3.10)$$

as $n \rightarrow \infty$. We have now to estimate the interaction terms. To this purpose, we observe that by Hölder and Sobolev inequality, and by (1.15),

$$\begin{aligned} \left| \int_{\Omega_n} \left(\sum_{j \in I_2} P_n U_{j,n} \right)^2 \phi_{1,n} (P_n \psi_{i,n}) \right| &\leq \left(\int_{\Omega_n} \left(\sum_{j \in I_2} P_n U_{j,n} \right)^{\frac{8}{3}} |P_n \psi_{i,n}|^{\frac{4}{3}} \right)^{\frac{3}{4}} \|\phi_{1,n}\|_{L^4} \\ &\leq C \left(\int_{\Omega_n} \left(\sum_{j \in I_2} U_{j,n} \right)^{\frac{8}{3}} |\psi_{i,n}|^{\frac{4}{3}} \right)^{\frac{3}{4}} \|\phi_{1,n}\|_{H_0^1} + h.o.t. \\ &\leq \frac{C}{\delta_{i,n}} \sum_{j \in I_2} \left(\int_{\Omega_n} U_{j,n}^{\frac{8}{3}} U_{i,n}^{\frac{4}{3}} \right)^{\frac{3}{4}} + h.o.t. \end{aligned} \quad (3.11)$$

as $n \rightarrow \infty$. The precise rate of the higher order terms (*h.o.t.*) does not play any role, and in any case can be derived using Lemmas A.1 and A.2. Moreover, the leading integral on the right hand side can be estimated using Lemma A.4, obtaining

$$\int_{\Omega_n} U_{j,n}^{\frac{8}{3}} U_{i,n}^{\frac{4}{3}} = \begin{cases} O\left(\left(\frac{\delta_{i,n}}{\delta_{j,n}}\right)^{\frac{4}{3}}\right) & \text{if } i > j \\ O\left(\left(\frac{\delta_{j,n}}{\delta_{i,n}}\right)^{\frac{4}{3}}\right) & \text{if } j > i. \end{cases}$$

Coming back to (3.11), we have

$$\delta_{i,n} \left| \int_{\Omega_n} \left(\sum_{j \in I_2} P_n U_{j,n} \right)^2 \phi_{1,n} (P_n \psi_{i,n}) \right| = \begin{cases} O\left(\frac{\delta_{i,n}}{\delta_{j,n}}\right) = o(1) & \text{if } i > j \\ O\left(\frac{\delta_{j,n}}{\delta_{i,n}}\right) = o(1) & \text{if } i < j \end{cases} \quad (3.12)$$

as $n \rightarrow \infty$, which proves that the second integral on the right hand side in (3.8) is of order $o(\delta_{i,n})$. As far as the third integral is concerned, we note that

$$\begin{aligned}
& \left| \int_{\Omega_n} \left(\sum_{j \in I_1} P_n U_{j,n} \right) \left(\sum_{j \in I_2} P_n U_{j,n} \right) \phi_{2,n}(P_n \psi_{i,n}) \right| \\
& \leq \left(\int_{\Omega_n} \left(\sum_{j \in I_1} P_n U_{j,n} \right)^{\frac{4}{3}} \left(\sum_{j \in I_2} P_n U_{j,n} \right)^{\frac{4}{3}} |P_n \psi_{i,n}|^{\frac{4}{3}} \right)^{\frac{3}{4}} \|\phi_{2,n}\|_{L^4} \\
& \leq C \left(\int_{\Omega_n} \left(\sum_{j \in I_1} U_{j,n} \right)^{\frac{4}{3}} \left(\sum_{j \in I_2} U_{j,n} \right)^{\frac{4}{3}} |\psi_{i,n}|^{\frac{4}{3}} \right)^{\frac{3}{4}} \|\phi_{2,n}\|_{H_0^1} + h.o.t. \quad (3.13) \\
& \leq \frac{C}{\delta_{i,n}} \sum_{h \in I_1} \sum_{j \in I_2} \left(\int_{\Omega_n} U_{h,n}^{\frac{4}{3}} U_{j,n}^{\frac{4}{3}} U_{i,n}^{\frac{4}{3}} \right)^{\frac{3}{4}} + h.o.t. \\
& = \frac{C}{\delta_{i,n}} o(1)
\end{aligned}$$

as $n \rightarrow \infty$. The last inequality follows by Lemma A.6 if $h \neq i$, and by Lemma A.4 if $h = i$. In any case

$$\delta_{i,n} \left| \int_{\Omega_n} \left(\sum_{j \in I_1} P_n U_{j,n} \right) \left(\sum_{j \in I_2} P_n U_{j,n} \right) \phi_{2,n}(P_n \psi_{i,n}) \right| = o(1)$$

as $n \rightarrow \infty$. To sum up, by expanding (3.8), we proved that for every index $i \in I_1$ it results that

$$c_{i,n}(\sigma_0 + o(1)) + o(1) \sum_{\substack{j \in I_1 \\ j \neq i}} c_{j,n} = o(1) \quad (3.14)$$

as $n \rightarrow \infty$. From this and by Cramer's rule, we deduce that $c_{i,n} \rightarrow 0$ for every $i \in I_1$. From this, the conclusion $\|w_{1,n}\| \rightarrow 0$ follows. \square

Let us set now $z_{i,n} := \phi_{i,n} - h_{i,n} - w_{i,n}$. Notice that, since $\|h_{i,n}\|_{H_0^1(\Omega_n)}, \|w_{i,n}\|_{H_0^1(\Omega_n)} \rightarrow 0$, we have $\|z_{1,n}\|_{H_0^1(\Omega_n)}^2 + \|z_{2,n}\|_{H_0^1(\Omega_n)}^2 \rightarrow 1$. In terms of $z_{i,n}$, equation (3.7) can be rewritten as

$$\begin{aligned}
z_{1,n} = \mathcal{I}^* \left\{ \right. & \left[3 \left(\sum_{j \in I_1} P_n U_{j,n} \right)^2 + \beta \left(\sum_{j \in I_2} P_n U_{j,n} \right)^2 \right] (z_{1,n} + h_{1,n} + w_{1,n}) \\
& \left. + 2\beta \left(\sum_{j \in I_1} P_n U_{j,n} \right) \left(\sum_{j \in I_2} P_n U_{j,n} \right) (z_{2,n} + h_{2,n} + w_{2,n}) \right\}. \quad (3.15)
\end{aligned}$$

Of course, a similar equation holds for $z_{2,n}$.

Lemma 3.4. *It results that at least one of the following lower estimates holds:*

$$\liminf_{n \rightarrow \infty} \left\{ \int_{\Omega_n} \left[3 \left(\sum_{j \in I_1} P_n U_{j,n} \right)^2 + \beta \left(\sum_{j \in I_2} P_n U_{j,n} \right)^2 \right] z_{1,n}^2 \right. \\
\left. + 2\beta \int_{\Omega_n} \left(\sum_{j \in I_1} P_n U_{j,n} \right) \left(\sum_{j \in I_2} P_n U_{j,n} \right) z_{1,n} z_{2,n} \right\} > 0,$$

or

$$\liminf_{n \rightarrow \infty} \left\{ \int_{\Omega_n} \left[3 \left(\sum_{j \in I_2} P_n U_{j,n} \right)^2 + \beta \left(\sum_{j \in I_1} P_n U_{j,n} \right)^2 \right] z_{2,n}^2 + 2\beta \int_{\Omega_n} \left(\sum_{j \in I_1} P_n U_{j,n} \right) \left(\sum_{j \in I_2} P_n U_{j,n} \right) z_{1,n} z_{2,n} \right\} > 0.$$

Proof. Since $\|z_{1,n}\|_{H_0^1(\Omega_n)}^2 + \|z_{2,n}\|_{H_0^1(\Omega_n)}^2 \rightarrow 1$, we can suppose that up to a subsequence $\{\|z_{1,n}\|_{H_0^1(\Omega_n)}^2\}_n$ or $\{\|z_{2,n}\|_{H_0^1(\Omega_n)}^2\}_n$ is uniformly bounded from below by $1/2$. Suppose for instance that $\{\|z_{1,n}\|_{H_0^1(\Omega_n)}^2\}_n$ is bounded from below. Then we test equation (3.15) with $z_{1,n}$, obtaining

$$\begin{aligned} \|z_{1,n}\|_{H_0^1(\Omega_n)}^2 &= \int_{\Omega_n} \left[3 \left(\sum_{j \in I_1} P_n U_{j,n} \right)^2 + \beta \left(\sum_{j \in I_2} P_n U_{j,n} \right)^2 \right] z_{1,n}^2 \\ &\quad + 2\beta \int_{\Omega_n} \left(\sum_{j \in I_1} P_n U_{j,n} \right) \left(\sum_{j \in I_2} P_n U_{j,n} \right) z_{1,n} z_{2,n} \\ &\quad + 3 \int_{\Omega_n} \left(\sum_{j \in I_1} P_n U_{j,n} \right)^2 (h_{1,n} + w_{1,n}) z_{1,n} + \beta \left(\sum_{j \in I_2} P_n U_{j,n} \right)^2 (h_{1,n} + w_{1,n}) z_{1,n} \\ &\quad + 2\beta \int_{\Omega_n} \left(\sum_{j \in I_1} P_n U_{j,n} \right) \left(\sum_{j \in I_2} P_n U_{j,n} \right) (h_{2,n} + w_{2,n}) z_{1,n}. \end{aligned} \tag{3.16}$$

Arguing as in [12, Formula (5.12)], we can easily check that the last two integrals are 0. Therefore, in this case the first lim inf in the thesis is positive. If $\{\|z_{2,n}\|_{H_0^1(\Omega_n)}^2\}_n$ is bounded from below, in the same way we find that the second lim inf is positive. \square

We aim to obtain a contradiction with Lemma 3.4. To this end, we fix $\rho > 0$ so that $B_\rho \subset \subset \Omega$, and we decompose $B_\rho \setminus \overline{B_{\varepsilon_n}}$ into the union of disjoint annuli as follows:

$$B_\rho \setminus \overline{B_{\varepsilon_n}} = \bigcup_{\ell=1}^k \mathcal{A}_{\ell,n}, \quad \text{where } \mathcal{A}_{\ell,n} = B_{\sqrt{\delta_{\ell,n} \delta_{\ell-1,n}}} \setminus \overline{B_{\sqrt{\delta_{\ell,n} \delta_{\ell+1,n}}}} \quad \text{for } \ell = 1, \dots, k,$$

with the convention $\delta_{0,n} = \delta_{1,n}^{-1} \rho^2$ and $\delta_{k+1,n} = \delta_{k,n}^{-1} \varepsilon_n^2$. Recall from Remark 2.1 that $\delta_{l+1,n} / \delta_{l,n} \rightarrow 0$ as $n \rightarrow \infty$. We also set

$$\mathcal{B}_{\ell,n} = B_{2\sqrt{\delta_{\ell,n} \delta_{\ell-1,n}}} \setminus \overline{B_{\sqrt{\delta_{\ell,n} \delta_{\ell+1,n}/2}}},$$

and, for every $\ell = 1, \dots, k$, we define a cut-off function $\chi_{\ell,n} \in C_c^\infty(\mathbb{R}^N)$ with the properties that

$$\begin{cases} \chi_{\ell,n} = 1 & \text{in } \mathcal{A}_{\ell,n}, \quad \chi_{\ell,n} = 0 & \text{in } \mathbb{R}^4 \setminus \mathcal{B}_{\ell,n}, \\ |\nabla \chi_{\ell,n}| \leq \frac{C}{\sqrt{\delta_{\ell,n} \delta_{\ell-1,n}}}, & |D^2 \chi_{\ell,n}| \leq \frac{C}{\delta_{\ell,n} \delta_{\ell-1,n}} & \text{in } \mathbb{R}^4, \end{cases} \tag{3.17}$$

for a positive universal constant C . Finally, we define for $\ell = 1, \dots, k$ and $i = 1, 2$ the $\mathcal{D}^{1,2}(\mathbb{R}^4)$ functions

$$\hat{z}_{i,n}^\ell(x) := \delta_{\ell,n} z_{i,n}(\delta_{\ell,n} x) \chi_{\ell,n}(\delta_{\ell,n} x) \quad \text{for } x \in \frac{\mathcal{B}_{\ell,n}}{\delta_{\ell,n}} =: \hat{\mathcal{B}}_{\ell,n},$$

naturally extended by 0 in $\mathbb{R}^4 \setminus \hat{\mathcal{B}}_{\ell,n}$. We have $\hat{z}_{i,n}^\ell(x) = \delta_{\ell,n} z_{i,n}(\delta_{\ell,n} x)$ if $x \in \tilde{\mathcal{A}}_{\ell,n} := \mathcal{A}_{\ell,n} / \delta_{\ell,n}$.

Lemma 3.5. *It results that $\hat{z}_{i,n}^\ell \rightarrow 0$ weakly in $\mathcal{D}^{1,2}(\mathbb{R}^4)$, and strongly in $L_{\text{loc}}^q(\mathbb{R}^4)$, for every $q \in [2, 2^*)$, for every $i = 1, 2$, $\ell = 1, \dots, k$.*

Proof. We have

$$\nabla \hat{z}_{i,n}^\ell(x) = \delta_{\ell,n}^2 [\chi_{\ell,n}(\delta_{\ell,n}x) \nabla z_{i,n}(\delta_{\ell,n}x) + z_{i,n}(\delta_{\ell,n}x) \nabla \chi_{\ell,n}(\delta_{\ell,n}x)]$$

and

$$\begin{aligned} \Delta \hat{z}_{i,n}^\ell(x) &= \delta_{\ell,n}^3 [\chi_{\ell,n}(\delta_{\ell,n}x) \Delta z_{i,n}(\delta_{\ell,n}x) + 2 \nabla z_{i,n}(\delta_{\ell,n}x) \cdot \nabla \chi_{\ell,n}(\delta_{\ell,n}x) \\ &\quad + z_{i,n}(\delta_{\ell,n}x) \Delta \chi_{\ell,n}(\delta_{\ell,n}x)] \end{aligned} \quad (3.18)$$

for $x \in \tilde{\mathcal{B}}_{\ell,n}$, that is,

$$\frac{1}{2} \sqrt{\frac{\delta_{\ell+1,n}}{\delta_{\ell,n}}} < |x| < 2 \sqrt{\frac{\delta_{\ell-1,n}}{\delta_{\ell,n}}}.$$

Notice that $\tilde{\mathcal{B}}_{\ell,n}$ exhausts \mathbb{R}^N as $n \rightarrow \infty$, by Remark 2.1. Now

$$\begin{aligned} \int_{\mathbb{R}^4} |\nabla \hat{z}_{i,n}^\ell|^2 &\leq 2 \delta_{\ell,n}^4 \int_{\tilde{\mathcal{B}}_{\ell,n}} (|\nabla z_{i,n}(\delta_{\ell,n}x)|^2 + z_{i,n}^2(\delta_{\ell,n}x) |\nabla \chi_{\ell,n}(\delta_{\ell,n}x)|^2) dx \\ &= 2 \int_{\mathcal{B}_{\ell,n}} (|\nabla z_{i,n}(y)|^2 + z_{i,n}^2(y) |\nabla \chi_{\ell,n}(y)|^2) dy. \end{aligned}$$

The integral of $|\nabla z_{i,n}|^2$ is clearly bounded, since $\|z_{i,n}\|_{H_0^1(\Omega_n)} \leq 1$. Also, by (3.17),

$$\begin{aligned} \int_{\mathcal{B}_{\ell,n}} z_{i,n}^2 |\nabla \chi_{\ell,n}|^2 &\leq \frac{C}{\delta_{\ell,n} \delta_{\ell-1,n}} \int_{\mathcal{B}_{\ell,n}} z_{i,n}^2 \leq \frac{C}{\delta_{\ell,n} \delta_{\ell-1,n}} |\mathcal{B}_{\ell,n}|^{\frac{1}{2}} \|z_{i,n}\|_{H_0^1(\Omega_n)}^2 \\ &\leq C \delta_{\ell,n} \delta_{\ell-1,n} \|z_{i,n}\|_{H_0^1(\Omega_n)}^2 \leq C, \end{aligned}$$

and we infer that $\|\hat{z}_{i,n}^\ell\|_{\mathcal{D}^{1,2}(\mathbb{R}^4)} \leq C$. Then, up to a subsequence, we have that $\hat{z}_{i,n}^\ell \rightharpoonup \hat{z}_i^\ell$ weakly in $\mathcal{D}^{1,2}$, and $\hat{z}_{i,n}^\ell \rightarrow \hat{z}_i^\ell$ strongly in $L_{\text{loc}}^q(\mathbb{R}^4)$ for $q \in [2, 2^*)$. The equation satisfied by the weak limit can be determined using (3.15) and (3.18): for every $\varphi \in C_c^\infty(\mathbb{R}^4 \setminus \{0\})$, by combining (3.15) with (3.18) we have that

$$\begin{aligned} &\int_{\mathbb{R}^N} \nabla \hat{z}_{1,n}^\ell \cdot \nabla \varphi \\ &= \delta_{\ell,n}^3 \int_{\tilde{\mathcal{B}}_{\ell,n}} \chi_{\ell,n}(\delta_{\ell,n}x) \left[3 \left(\sum_{i \in I_1} P_n U_{j,n}(\delta_{\ell,n}x) \right)^2 + \beta \left(\sum_{j \in I_2} P_n U_{j,n}(\delta_{\ell,n}x) \right)^2 \right] \\ &\quad \cdot (z_{1,n}(\delta_{\ell,n}x) + h_{1,n}(\delta_{\ell,n}x) + w_{1,n}(\delta_{\ell,n}x)) \varphi(x) dx \\ &+ 2\beta \delta_{\ell,n}^3 \int_{\tilde{\mathcal{B}}_{\ell,n}} \chi_{\ell,n}(\delta_{\ell,n}x) \left(\sum_{i \in I_1} P_n U_{j,n}(\delta_{\ell,n}x) \right) \left(\sum_{i \in I_2} P_n U_{j,n}(\delta_{\ell,n}x) \right) \\ &\quad \cdot (z_{2,n}(\delta_{\ell,n}x) + h_{2,n}(\delta_{\ell,n}x) + w_{2,n}(\delta_{\ell,n}x)) \varphi(x) dx \\ &- \delta_{\ell,n}^3 \int_{\tilde{\mathcal{B}}_{\ell,n}} (2 \nabla \chi_{\ell,n}(\delta_{\ell,n}x) \cdot \nabla z_{1,n}(\delta_{\ell,n}x) + z_{1,n}(\delta_{\ell,n}x) \Delta \chi_{\ell,n}(\delta_{\ell,n}x)) \varphi(x) dx. \end{aligned}$$

The last integral and all the terms involving $h_{i,n}$ and $w_{i,n}$ tend to 0 as $n \rightarrow \infty$, exactly as in [12, Formula (5.20)]. Therefore,

$$\begin{aligned}
 & \int_{\mathbb{R}^N} \nabla \hat{z}_{1,n}^\ell \cdot \nabla \varphi = o(1) \\
 & + \delta_{\ell,n}^3 \int_{\tilde{\mathcal{B}}_{\ell,n}} \chi_{\ell,n}(\delta_{\ell,n}x) \left[3 \left(\sum_{i \in I_1} P_n U_{j,n}(\delta_{\ell,n}x) \right)^2 + \beta \left(\sum_{j \in I_2} P_n U_{j,n}(\delta_{\ell,n}x) \right)^2 \right] z_{1,n}(\delta_{\ell,n}x) \varphi(x) dx \\
 & + 2\beta \delta_{\ell,n}^3 \int_{\tilde{\mathcal{B}}_{\ell,n}} \chi_{\ell,n}(\delta_{\ell,n}x) \left(\sum_{i \in I_1} P_n U_{j,n}(\delta_{\ell,n}x) \right) \left(\sum_{i \in I_2} P_n U_{j,n}(\delta_{\ell,n}x) \right) z_{2,n}(\delta_{\ell,n}x) \varphi(x) dx \\
 & = o(1) + \delta_{\ell,n}^2 \int_{\tilde{\mathcal{B}}_{\ell,n}} 3 \left(\sum_{i \in I_1} P_n U_{j,n}(\delta_{\ell,n}x) \right)^2 \hat{z}_{1,n}^\ell(x) \varphi(x) dx \\
 & + \beta \delta_{\ell,n}^2 \int_{\tilde{\mathcal{B}}_{\ell,n}} \left(\sum_{j \in I_2} P_n U_{j,n}(\delta_{\ell,n}x) \right)^2 \hat{z}_{1,n}(x) \varphi(x) dx \\
 & + 2\beta \delta_{\ell,n}^2 \int_{\tilde{\mathcal{B}}_{\ell,n}} \left(\sum_{i \in I_1} P_n U_{j,n}(\delta_{\ell,n}x) \right) \left(\sum_{i \in I_2} P_n U_{j,n}(\delta_{\ell,n}x) \right) \hat{z}_{2,n}^\ell(x) \varphi(x) dx \\
 & =: o(1) + A_1 + A_2 + A_3.
 \end{aligned}$$

In order to study the behavior of the integrals as $n \rightarrow \infty$, it is convenient to observe (see Lemma A.1) that, if $\ell \in I_1$, then

$$\begin{aligned}
 \sum_{j \in I_1} P_n U_{j,n}(\delta_{\ell,n}x) &= \sum_{j \in I_1} U_{j,n}(\delta_{\ell,n}x) + h.o.t. = \sum_{j \in I_1} \alpha_4 \frac{\delta_{j,n}}{\delta_{j,n}^2 + \delta_{\ell,n}^2 |x|^2} + h.o.t. \\
 &= \frac{1}{\delta_{\ell,n}} U_{1,0}(x) + \sum_{\substack{j \in I_1 \\ j \neq \ell}} \alpha_4 \frac{\delta_{j,n}}{\delta_{j,n}^2 + \delta_{\ell,n}^2 |x|^2} + h.o.t. \\
 &= \frac{1}{\delta_{\ell,n}} U_{1,0}(x) + \sum_{\substack{j \in I_1 \\ j < \ell}} O\left(\frac{1}{\delta_{j,n}}\right) + \sum_{\substack{j \in I_1 \\ j > \ell}} O\left(\frac{\delta_{j,n}}{\delta_{\ell,n}^2 |x|^2}\right) + h.o.t.,
 \end{aligned} \tag{3.19}$$

as $n \rightarrow \infty$. If instead $\ell \notin I_1$, then we have a similar expansion, but without the term $U_{1,0}(x)/\delta_{\ell,n}$. We focus at first on the first possibility. We have,

$$\begin{aligned}
A_1 &= 3\delta_{\ell,n}^2 \int_{\tilde{B}_{\ell,n}} \left(\frac{U_{1,0}(x)}{\delta_{\ell,n}} \right)^2 \hat{z}_{1,n}^\ell(x) \varphi(x) + \left[\sum_{\substack{j \in I_1 \\ j < \ell}} O\left(\frac{1}{\delta_{j,n}^2}\right) + \sum_{\substack{j \in I_1 \\ j > \ell}} O\left(\frac{\delta_{j,n}^2}{\delta_{\ell,n}^4 |x|^4}\right) \right] \hat{z}_{1,n}^\ell(x) \varphi(x) dx \\
&+ 3\delta_{\ell,n}^2 \int_{\tilde{B}_{\ell,n}} \frac{2}{\delta_{\ell,n}} U_{1,0}(x) \left[\sum_{\substack{j \in I_1 \\ j < \ell}} O\left(\frac{1}{\delta_{j,n}}\right) + \sum_{\substack{j \in I_1 \\ j > \ell}} O\left(\frac{\delta_{j,n}}{\delta_{\ell,n}^2 |x|^2}\right) \right] \hat{z}_{1,n}^\ell(x) \varphi(x) dx \\
&+ 3\delta_{\ell,n}^2 \int_{\tilde{B}_{\ell,n}} 2 \sum_{\substack{i \in I_1 \\ i < \ell}} \sum_{\substack{j \in I_1 \\ j > \ell}} O\left(\frac{\delta_{j,n}}{\delta_{i,n} \delta_{\ell,n}^2 |x|^2}\right) \hat{z}_{1,n}^\ell(x) \varphi(x) dx + h.o.t.
\end{aligned} \tag{3.20}$$

Now, for every $j < \ell$

$$\frac{\delta_{\ell,n}^2}{\delta_{j,n}^2} \left| \int_{\tilde{B}_{\ell,n}} \hat{z}_{1,n}^\ell \varphi \right| \leq o(1) \|\hat{z}_{1,n}^\ell\|_{L^4} \rightarrow 0,$$

and, by Lemma A.3,

$$\left| \frac{\delta_{\ell,n}}{\delta_{j,n}} \int_{\tilde{B}_{\ell,n}} U_{1,0} \hat{z}_{1,n}^\ell \varphi \right| \leq \frac{\delta_{\ell,n}}{\delta_{j,n}} \|U_{1,0}\|_{L^4} \|\hat{z}_{1,n}^\ell\|_{L^4} \|\varphi\|_{L^2} \rightarrow 0.$$

Moreover, for every $j > \ell$, using the fact that $\text{supp } \varphi \subset B_R \setminus B_\rho$ for suitable $0 < \rho < R$, we have that

$$\frac{\delta_{j,n}^2}{\delta_{\ell,n}^2} \left| \int_{\tilde{B}_{\ell,n}} \hat{z}_{1,n}^\ell(x) \frac{\varphi(x)}{|x|^4} dx \right| \leq C \frac{\delta_{j,n}^2}{\delta_{\ell,n}^2} \|\hat{z}_{1,n}^\ell\|_{L^4} \left(\int_{B_R \setminus B_\rho} \frac{dx}{|x|^{\frac{16}{3}}} \right)^{\frac{3}{4}} \leq C \frac{\delta_{j,n}^2}{\delta_{\ell,n}^2} \rightarrow 0,$$

and that

$$\frac{\delta_{j,n}}{\delta_{\ell,n}} \left| \int_{\tilde{B}_{\ell,n}} U_{1,0}(x) \hat{z}_{1,n}^\ell(x) \frac{\varphi(x)}{|x|^2} dx \right| \leq C \frac{\delta_{j,n}}{\delta_{\ell,n}} \|\hat{z}_{1,n}^\ell\|_{L^4} \left(\int_{B_R \setminus B_\rho} \frac{U_{1,0}^{\frac{4}{3}}(x)}{|x|^{\frac{8}{3}}} dx \right)^{\frac{3}{4}} \leq C \frac{\delta_{j,n}}{\delta_{\ell,n}} \rightarrow 0.$$

The previous estimates yield

$$A_1 = 3 \int_{\tilde{B}_{\ell,n}} U_{1,0}^2 \hat{z}_{1,n}^\ell \varphi + o(1) \rightarrow 3 \int_{\mathbb{R}^4} U_{1,0}^2 \hat{z}_1^\ell \varphi \tag{3.21}$$

as $n \rightarrow \infty$, for every $\varphi \in C_c^\infty(\mathbb{R}^4 \setminus \{0\})$.

Notice that, in the above computations, we never used the fact that the indexes j were in I_1 . Therefore, we directly deduce that

$$A_2 \rightarrow 0 \quad \text{as } n \rightarrow \infty. \tag{3.22}$$

Finally, in an analogue way

$$\begin{aligned}
 A_3 = 2\beta\delta_{\ell,n}^2 \int_{\tilde{B}_{\ell,n}} & \left[\frac{1}{\delta_{\ell,n}} U_{1,0}(x) + \sum_{\substack{j \in I_1 \\ j < \ell}} O\left(\frac{1}{\delta_{j,n}}\right) + \sum_{\substack{j \in I_1 \\ j > \ell}} O\left(\frac{\delta_{j,n}}{\delta_{\ell,n}^2 |x|^2}\right) \right] \\
 & \cdot \left[\sum_{\substack{j \in I_2 \\ j < \ell}} O\left(\frac{1}{\delta_{j,n}}\right) + \sum_{\substack{j \in I_2 \\ j > \ell}} O\left(\frac{\delta_{j,n}}{\delta_{\ell,n}^2 |x|^2}\right) \right] \hat{z}_{1,n}^\ell(x) \varphi(x) dx + h.o.t. \rightarrow 0
 \end{aligned} \tag{3.23}$$

as $n \rightarrow \infty$.

Collecting together (3.21), (3.22) and (3.23), and coming back to (3.20), we finally obtain that the weak limit of $\hat{z}_{1,n}^\ell$ satisfies

$$-\Delta \hat{z}_1^\ell = 3U_{1,0}^2 \hat{z}_1^\ell \quad \text{in } \mathbb{R}^4 \setminus \{0\}.$$

Let now $\theta_\rho \in C^\infty(\mathbb{R}^N)$ be such that $\theta_\rho \equiv 1$ in $B_{2\rho}^c$, $\theta_\rho \equiv 0$ in B_ρ , and $|\nabla \theta_\rho| \leq C/\rho$; and let $\varphi \in C_c^\infty(\mathbb{R}^4)$; testing the above equation with $\theta_\rho \varphi$, and passing to the limit as $\rho \rightarrow 0^+$, using the fact that $\hat{z}_1^\ell \in \mathcal{D}^{1,2}(\mathbb{R}^4)$ (since it is the weak limit of $\mathcal{D}^{1,2}$ functions), we easily deduce that

$$-\Delta \hat{z}_1^\ell = 3U_{1,0}^2 \hat{z}_1^\ell \quad \text{in the whole space } \mathbb{R}^4.$$

In order to show that $\hat{z}_1^\ell \equiv 0$, recalling that it is symmetric with respect to 0, it is sufficient to verify that $\hat{z}_1^\ell \perp \psi_{1,0}$. This can be done exactly as in [12, Formula (5.19)], and completes the proof.

It still remains to analyze the case $\ell \notin I_1$. In such a situation we can proceed exactly as before, but this time we end up with

$$-\Delta \hat{z}_1^\ell = \beta U_{1,0}^2 \hat{z}_1^\ell \quad \text{in the whole space } \mathbb{R}^4.$$

Since $\beta < 0$ and $\hat{z}_1^\ell \in \mathcal{D}^{1,2}(\mathbb{R}^4)$, we infer that

$$0 \leq \int_{\mathbb{R}^N} |\nabla \hat{z}_1^\ell|^2 = \beta \int_{\mathbb{R}^N} (U_{1,0} \hat{z}_1^\ell)^2 \leq 0,$$

and the conclusion follows also in this case. \square

Conclusion of the proof of Proposition 3.2. Using Lemma 3.5, we will obtain a contradiction with Lemma 3.4. Let us consider

$$\int_{\Omega_n} \left(\sum_{j \in I_1} P_n U_{j,n} \right)^2 z_{1,n}^2 \leq C \sum_{i \in I_1} \int_{\Omega_n} U_{i,n}^2 z_{1,n}^2 = C \sum_{i \in I_1} \left(\int_{\Omega_n \setminus B_\rho} U_{i,n}^2 z_{1,n}^2 + \sum_{\ell=1}^k \int_{\mathcal{A}_{\ell,n}} U_{i,n}^2 z_{1,n}^2 \right). \tag{3.24}$$

We show that the right hand side tends to 0 as $n \rightarrow \infty$. At first, we have

$$\int_{\Omega_n \setminus B_\rho} U_{i,n}^2 z_{1,n}^2 \leq C \delta_{i,n}^2 \int_{\Omega_n \setminus B_\rho} z_{1,n}^2 \leq C \delta_{i,n}^2 \|z_{i,n}\|_{H_0^1(\Omega_n)}^2 \rightarrow 0. \tag{3.25}$$

Now, let $i \neq \ell$. Then we have

$$\int_{\mathcal{A}_{\ell,n}} U_{i,n}^2 z_{1,n}^2 \leq C \|z_{i,n}\|_{H_0^1(\Omega_n)}^2 \left(\int_{\mathcal{A}_{\ell,n}} U_{i,n}^4 \right)^{\frac{1}{2}} = C \|z_{i,n}\|_{H_0^1(\Omega_n)}^2 \left(\int_{\mathcal{A}_{\ell,n}/\delta_{i,n}} U_{1,0}^4 \right)^{\frac{1}{2}}.$$

Since $i \neq \ell$, we have that

$$\frac{\mathcal{A}_{\ell,n}}{\delta_{i,n}} \subset \begin{cases} B\left(0, \frac{\sqrt{\delta_{\ell-1,n}\delta_{\ell,n}}}{\delta_{i,n}}\right) & \text{if } i < \ell \\ \mathbb{R}^N \setminus B\left(0, \frac{\sqrt{\delta_{\ell+1,n}\delta_{\ell,n}}}{\delta_{i,n}}\right) & \text{if } i > \ell, \end{cases}$$

and

$$\frac{\sqrt{\delta_{\ell-1,n}\delta_{\ell,n}}}{\delta_{i,n}} \rightarrow 0 \quad \text{if } i < \ell, \quad \text{and} \quad \frac{\sqrt{\delta_{\ell+1,n}\delta_{\ell,n}}}{\delta_{i,n}} \rightarrow +\infty \quad \text{if } i > \ell.$$

Therefore, the fact that

$$\int_{\mathcal{A}_{\ell,n}} U_{i,n}^2 z_{1,n}^2 \rightarrow 0 \quad \text{as } n \rightarrow \infty, \quad \forall i \neq \ell \quad (3.26)$$

follows from the integrability of $U_{1,0}^4$ on \mathbb{R}^N . If moreover $i = \ell$, recalling that $\tilde{\mathcal{A}}_{\ell,n} = \mathcal{A}_{\ell,n}/\delta_{\ell,n}$ we have

$$\int_{\mathcal{A}_{\ell,n}} U_{\ell,n}^2 z_{1,n}^2 = \delta_{\ell,n}^2 \int_{\tilde{\mathcal{A}}_{\ell,n}} U_{\ell,n}^2 (\delta_{\ell,n} x) (\hat{z}_{1,n}^\ell)^2(x) dx = \alpha_4^2 \int_{\mathbb{R}^N} \left(\frac{1}{1+|x|^2}\right)^2 (\hat{z}_{1,n}^\ell)^2(x) dx + o(1) \rightarrow 0 \quad (3.27)$$

as $n \rightarrow \infty$, since $U_{1,0}^2 \in L^2(\mathbb{R}^N)$ and $(\hat{z}_{1,n}^\ell)^2 \rightharpoonup 0$ weakly in $L^2(\mathbb{R}^4)$ by Lemma 3.5. By (3.25) and (3.26), we obtain in (3.24) that

$$\int_{\Omega_n} \left(\sum_{j \in I_1} P_n U_{j,n}\right)^2 z_{1,n}^2 \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (3.28)$$

In a completely analogue way, we also have

$$\int_{\Omega_n} \left(\sum_{j \in I_2} P_n U_{j,n}\right)^2 z_{1,n}^2 \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (3.29)$$

Finally,

$$\left| \int_{\Omega_n} \left(\sum_{j \in I_1} P_n U_{j,n}\right) \left(\sum_{j \in I_2} P_n U_{j,n}\right) z_{1,n} z_{2,n} \right| \leq \left(\sum_{j \in I_1} \int_{\Omega_n} U_{j,n}^2 z_{1,n}^2\right) \left(\sum_{j \in I_2} \int_{\Omega_n} U_{j,n}^2 z_{2,n}^2\right) \rightarrow 0 \quad (3.30)$$

as $n \rightarrow \infty$. But (3.24), estimates (3.28), (3.29) and (3.30) imply that

$$\liminf_{n \rightarrow \infty} \left\{ \int_{\Omega_n} \left[3 \left(\sum_{j \in I_1} P_n U_{j,n}\right)^2 + \beta \left(\sum_{j \in I_2} P_n U_{j,n}\right)^2 \right] z_{1,n}^2 + 2\beta \int_{\Omega_n} \left(\sum_{j \in I_1} P_n U_{j,n}\right) \left(\sum_{j \in I_2} P_n U_{j,n}\right) z_{1,n} z_{2,n} \right\} = 0,$$

in contradiction with Lemma 3.4. \square

3.2. Estimates on the reminder term. In this subsection we prove the following

Proposition 3.6. *Let $\eta > 0$. There exists $\varepsilon_0 > 0$ and $C > 0$ such that*

$$\|R_{\mathbf{d},\varepsilon}^i\| \leq C \varepsilon^{\frac{1}{k+1}} \left(\log\left(\frac{1}{\varepsilon}\right)\right)^{-\frac{1}{k+1}},$$

for $i = 1, 2$, for every $\mathbf{d} \in X_\eta$ and for every $\varepsilon \in (0, \varepsilon_0)$.

Proof. We focus on $i = 1$. By continuity of Π_1^1 and of \mathcal{I}^* , there exists $C > 0$ such that

$$\begin{aligned} \|R_{\mathbf{d},\varepsilon}^1\| &\leq C \left\| f\left(\sum_{j \in I_1} P_\varepsilon U_j\right) - \sum_{j \in I_1} f(U_j) + \beta \left(\sum_{j \in I_1} P_\varepsilon U_j\right) \left(\sum_{j \in I_2} P_\varepsilon U_j\right)^2 \right\|_{L^{\frac{4}{3}}} \\ &\leq C \left\| f\left(\sum_{j \in I_1} P_\varepsilon U_j\right) - \sum_{j \in I_1} f(U_j) \right\|_{L^{\frac{4}{3}}} + C \left\| \left(\sum_{j \in I_1} P_\varepsilon U_j\right) \left(\sum_{j \in I_2} P_\varepsilon U_j\right)^2 \right\|_{L^{\frac{4}{3}}} \\ &=: C(\mathcal{A} + \mathcal{B}). \end{aligned} \quad (3.31)$$

We estimate separately \mathcal{A} and \mathcal{B} . At first we note that

$$\mathcal{A} \leq \left\| f\left(\sum_{j \in I_1} P_\varepsilon U_j\right) - \sum_{j \in I_1} f(P_\varepsilon U_j) \right\|_{L^{\frac{4}{3}}} + \left\| \sum_{j \in I_1} (f(P_\varepsilon U_j) - f(U_j)) \right\|_{L^{\frac{4}{3}}} =: A_1 + A_2. \quad (3.32)$$

Recalling that $f(s) = (s^+)^3$, and using the fact that

$$(a_1 + \dots + a_n)^3 \leq (a_1^3 + \dots + a_n^3) + C_n \sum_{1 \leq j \neq h \leq n} a_j^2 a_h$$

for a positive constant C_n depending only on n , we obtain

$$A_1^{\frac{4}{3}} \leq C \int_{\Omega_\varepsilon} \left| \sum_{\substack{j \neq h \\ j, h \in I_1}} (P_\varepsilon U_j)^2 (P_\varepsilon U_h) \right|^{\frac{4}{3}} \leq C \sum_{\substack{j \neq h \\ j, h \in I_1}} \int_{\Omega_\varepsilon} (U_j^2 U_h)^{\frac{4}{3}}. \quad (3.33)$$

Let us fix $j \neq h$. Then, by Lemma A.4,

$$\int_{\Omega_\varepsilon} (U_j^2 U_h)^{\frac{4}{3}} \leq \begin{cases} C \left(\frac{\delta_h}{\delta_j}\right)^{\frac{4}{3}} & \text{if } h > j \\ C \left(\frac{\delta_j}{\delta_h}\right)^{\frac{4}{3}} & \text{if } h < j. \end{cases} \quad (3.34)$$

Recalling (2.5), we see that if $h > j$ and $\mathbf{d} \in X_\eta$

$$\frac{\delta_h}{\delta_j} \leq C \left(\varepsilon^{\frac{h-j}{k+1}} \left(\log \left(\frac{1}{\varepsilon} \right) \right)^{-\frac{h-j}{k+1}} \right) \leq C \varepsilon^{\frac{2}{k+1}} \left(\log \frac{1}{\varepsilon} \right)^{-\frac{2}{k+1}},$$

where C denotes a positive constant depending on η (but not on \mathbf{d}) and we used the fact that $h - j \geq 2$ since $j, h \in I_1$ with $j \neq h$. The same estimate holds in case $h < j$. Plugging this into (3.34), and coming back to (3.33), we finally conclude that

$$A_1 \leq C \varepsilon^{\frac{2}{k+1}} \left(\log \frac{1}{\varepsilon} \right)^{-\frac{2}{k+1}} \leq C \varepsilon^{\frac{1}{k+1}} \left(\log \frac{1}{\varepsilon} \right)^{-\frac{1}{k+1}}. \quad (3.35)$$

Regarding A_2 , we have

$$A_2 \leq \sum_{j \in I_1} \left\| (P_\varepsilon U_j)^3 - U_j^3 \right\|_{L^{\frac{4}{3}}} \leq \sum_{j \in I_1} \left(\left\| (U_j - P_\varepsilon U_j)^3 \right\|_{L^{\frac{4}{3}}} + C \left\| U_j^2 (U_j - P_\varepsilon U_j) \right\|_{L^{\frac{4}{3}}} \right). \quad (3.36)$$

Using the estimate for $U_i - P_\varepsilon U_i$ contained in Lemma A.1, we deduce that

$$\begin{aligned} \left\| (U_j - P_\varepsilon U_j)^3 \right\|_{L^{\frac{4}{3}}}^{\frac{4}{3}} &= \int_{\Omega_\varepsilon} |U_j - P_\varepsilon U_j|^4 \leq C \int_{\Omega_\varepsilon} \left(\delta_j^4 + \left(\frac{\varepsilon}{\delta_j} \right)^8 \frac{\delta_j^4}{|x|^8} \right) \\ &\leq C \delta_j^4 + C \frac{\varepsilon^8}{\delta_j^4} \int_\varepsilon^R r^{-5} dr \leq C \left(\delta_j^4 + \left(\frac{\varepsilon}{\delta_j} \right)^4 \right). \end{aligned} \quad (3.37)$$

In a similar way

$$\begin{aligned} \|U_j^2(U_j - P_\varepsilon U_j)\|_{L^{\frac{4}{3}}} &\leq C \int_{\Omega_\varepsilon} \left(\delta_j^{\frac{4}{3}} U_j^{\frac{8}{3}}(x) + \delta_j^{\frac{4}{3}} \left(\frac{\varepsilon U_j(x)}{\delta_j |x|} \right)^{\frac{8}{3}} \right) dx \\ &\leq C \delta_j^{\frac{8}{3}} \int_{\Omega_\varepsilon/\delta_j} \frac{dx}{(1+|x|^2)^{\frac{8}{3}}} + C \left(\frac{\varepsilon}{\delta_j} \right)^{\frac{8}{3}} \int_{\Omega_\varepsilon/\delta_j} \frac{dx}{(1+|x|^2)^{\frac{8}{3}} |x|^{\frac{8}{3}}} \leq C \left(\delta_j^{\frac{8}{3}} + \left(\frac{\varepsilon}{\delta_j} \right)^{\frac{8}{3}} \right). \end{aligned} \quad (3.38)$$

Plugging (3.37) and (3.38) into (3.36), and recalling again Remark 2.1, we obtain

$$A_2 \leq C \sum_{j \in I_1} \left(\delta_j^2 + \left(\frac{\varepsilon}{\delta_j} \right)^2 \right) \leq C \varepsilon^{\frac{1}{k+1}} \left(\log \left(\frac{1}{\varepsilon} \right) \right)^{-\frac{1}{k+1}}. \quad (3.39)$$

Therefore, (3.32), (3.35) and (3.39) give

$$\mathcal{A} \leq C \varepsilon^{\frac{1}{k+1}} \left(\log \left(\frac{1}{\varepsilon} \right) \right)^{-\frac{1}{k+1}}, \quad (3.40)$$

and it remains to estimate \mathcal{B} . By Lemma A.4

$$\begin{aligned} \mathcal{B} &\leq C \sum_{(j,h) \in I_1 \times I_2} \left\| (P_\varepsilon U_j)(P_\varepsilon U_h)^2 \right\|_{L^{\frac{4}{3}}} \leq C \sum_{(j,h) \in I_1 \times I_2} \left\| U_j U_h^2 \right\|_{L^{\frac{4}{3}}} \\ &= C \sum_{(j,h) \in I_1 \times I_2} \left(\int_{\Omega_\varepsilon} U_j^{\frac{4}{3}} U_h^{\frac{8}{3}} \right)^{\frac{3}{4}} = \begin{cases} O \left(\frac{\delta_j}{\delta_h} \right) & \text{if } j > h \\ O \left(\frac{\delta_h}{\delta_j} \right) & \text{if } j < h, \end{cases} \end{aligned} \quad (3.41)$$

and for any $h > j$ we have

$$\frac{\delta_h}{\delta_j} \leq C \varepsilon^{\frac{h-j}{k+1}} \left(\log \left(\frac{1}{\varepsilon} \right) \right)^{-\frac{h-j}{k+1}} \leq C \varepsilon^{\frac{1}{k+1}} \left(\log \left(\frac{1}{\varepsilon} \right) \right)^{-\frac{1}{k+1}},$$

The same estimate holds for δ_j/δ_h in case $j > h$. Therefore, gathering (3.31), (3.40) and (3.41), we obtain the desired result. \square

3.3. The nonlinear part: end of the proof of Proposition 3.1. In virtue of Proposition 3.2, solving the equation

$$\mathbf{L}_{\mathbf{d},\varepsilon}(\phi) = \mathbf{R}_{\mathbf{d},\varepsilon} + \mathbf{N}_{\mathbf{d},\varepsilon}(\phi).$$

reduces to finding a fixed point of the operator

$$\mathbf{T}_{\mathbf{d},\varepsilon}(\phi) := (\mathbf{L}_{\mathbf{d},\varepsilon})^{-1} (\mathbf{R}_{\mathbf{d},\varepsilon} + \mathbf{N}_{\mathbf{d},\varepsilon}(\phi)).$$

in the ball

$$\mathcal{B}_\rho := \left\{ \phi \in \mathbf{K}_{\mathbf{d},\varepsilon}^\perp : \|\phi\|_{H_0^1} \leq \rho \varepsilon^{\frac{1}{k+1}} \left(\log \left(\frac{1}{\varepsilon} \right) \right)^{-\frac{1}{k+1}} \right\}$$

for some $\rho > 0$. It is quite standard to show that $\mathbf{T}_{\mathbf{d},\varepsilon} : \mathcal{B}_\rho \rightarrow \mathcal{B}_\rho$ is a contraction mapping for ε small enough. Indeed, Proposition 3.2 together with straightforward computations lead to

$$\|\mathbf{T}_{\mathbf{d},\varepsilon}(\phi)\|_{H_0^1} \leq C \left(\|\mathbf{R}_{\mathbf{d},\varepsilon}\|_{H_0^1} + \|\mathbf{N}_{\mathbf{d},\varepsilon}(\phi)\|_{H_0^1} \right) \leq C \left(\|\mathbf{R}_{\mathbf{d},\varepsilon}\|_{H_0^1} + \|\phi\|_{H_0^1}^2 \right)$$

and

$$\|\mathbf{T}_{\mathbf{d},\varepsilon}(\phi_1) - \mathbf{T}_{\mathbf{d},\varepsilon}(\phi_2)\|_{H_0^1} \leq C \left(\|\mathbf{N}_{\mathbf{d},\varepsilon}(\phi_1) - \mathbf{N}_{\mathbf{d},\varepsilon}(\phi_2)\|_{H_0^1} \right) \leq \ell \|\phi_1 - \phi_2\|_{H_0^1} \quad \text{for some } \ell \in (0, 1).$$

A standard argument also shows that the map $\mathbf{d} \rightarrow \phi^{\mathbf{d},\varepsilon}$ is of class C^1 .

Remark 3.7. Suppose that, instead of dealing with the set of odd and even numbers of $\{1, \dots, k\}$ in the two equation case, we are dealing with system (1.4) with m equations and with a general partition I_1, \dots, I_m satisfying (1)–(5). Having already splitted the original problem into $2m$ equations (see Remark 2.2), we can repeat the argument used for $m = 2$ without substantial changes, using the fact that each set I_j does not contain consecutive integers.

4. EXPANSION OF THE REDUCED ENERGY

Recall that the energy functional is given by

$$J_\varepsilon(u_1, u_2) = \sum_{i=1}^2 \int_{\Omega_\varepsilon} \left(\frac{|\nabla u_i|^2}{2} - F(u_i) \right) - \frac{\beta}{2} \int_{\Omega_\varepsilon} u_1^2 u_2^2.$$

where $F(s) = (s^+)^4/4$. Recall that we denote $f(s) := F'(s) = (s^+)^3$. For every $\eta > 0$ small fixed, we introduce the reduced functional $\tilde{J}_\varepsilon : X_\eta \rightarrow \mathbb{R}$ as being

$$\tilde{J}_\varepsilon(\mathbf{d}) = J_\varepsilon \left(\sum_{j \in I_1} P_\varepsilon U_{\delta_j} + \phi_1^{\mathbf{d}, \varepsilon}, \sum_{j \in I_2} P_\varepsilon U_{\delta_j} + \phi_2^{\mathbf{d}, \varepsilon} \right)$$

This is a C^1 functional due to Proposition 3.1 and since δ_i depends on d_i via (2.5). Finding critical point of \tilde{J}_ε corresponds to find solutions of our original system, as we prove next.

Lemma 4.1. *Given $\varepsilon \in (0, \varepsilon_0)$ and $\eta > 0$ small, let $\mathbf{d} \in X_\eta$. We have that*

$$\left(\sum_{j \in I_1} P_\varepsilon U_{\delta_j, 0} + \phi_1^{\mathbf{d}, \varepsilon}, \sum_{j \in I_2} P_\varepsilon U_{\delta_j, 0} + \phi_2^{\mathbf{d}, \varepsilon} \right) \text{ is a solution of (2.2)}$$

if, and only if,

\mathbf{d} is a critical point of \tilde{J}_ε .

Proof. To simplify notations, define $V_i^{\mathbf{d}, \varepsilon} := \sum_{j \in I_i} P_\varepsilon U_{\delta_j} + \phi_i^{\mathbf{d}, \varepsilon}$ for $i = 1, 2$. From (2.5) we see that

$$\partial_{d_i} \tilde{J}_\varepsilon(\mathbf{d}) = \varepsilon^{\frac{l}{\kappa+1}} \left(\log \frac{1}{\varepsilon} \right)^{\frac{1}{2} - \frac{l}{\kappa+1}} J'_\varepsilon(V_1^{\mathbf{d}, \varepsilon}, V_2^{\mathbf{d}, \varepsilon}) [\partial_{\delta_i} V_1^{\mathbf{d}, \varepsilon}, \partial_{\delta_i} V_2^{\mathbf{d}, \varepsilon}] \quad (4.1)$$

Hence, if $(V_1^{\mathbf{d}, \varepsilon}, V_2^{\mathbf{d}, \varepsilon})$ solves (2.2) then $J'_\varepsilon(V_1^{\mathbf{d}, \varepsilon}, V_2^{\mathbf{d}, \varepsilon}) = 0$ and so $\tilde{J}'_\varepsilon(\mathbf{d}) = 0$. Conversely, assume $\mathbf{d} \in X_\eta$ is a solution of $\tilde{J}'_\varepsilon(\mathbf{d}) = 0$. For $i \in \{1, 2\}$ and $l \in I_i$, recalling that $\psi_l := \partial_{\delta_l} U_{\delta_l}$, we have from (4.1) that

$$\begin{aligned} 0 &= J'_\varepsilon(V_1^{\mathbf{d}, \varepsilon}, V_2^{\mathbf{d}, \varepsilon}) [\partial_{\delta_l} V_1^{\mathbf{d}, \varepsilon}, \partial_{\delta_l} V_2^{\mathbf{d}, \varepsilon}] \\ &= J'_\varepsilon(V_1^{\mathbf{d}, \varepsilon}, V_2^{\mathbf{d}, \varepsilon}) \left[\sum_{j \in I_1} P_\varepsilon \partial_{\delta_l} U_{\delta_j} + \partial_{\delta_l} \phi_1^{\mathbf{d}, \varepsilon}, \sum_{j \in I_2} P_\varepsilon \partial_{\delta_l} U_{\delta_j} + \partial_{\delta_l} \phi_2^{\mathbf{d}, \varepsilon} \right] \\ &= \partial_{\delta_l} J_\varepsilon(V_1^{\mathbf{d}, \varepsilon}, V_2^{\mathbf{d}, \varepsilon}) [P_\varepsilon \psi_l] + J'_\varepsilon(V_1^{\mathbf{d}, \varepsilon}, V_2^{\mathbf{d}, \varepsilon}) [\partial_{\delta_l} \phi_1^{\mathbf{d}, \varepsilon}, \partial_{\delta_l} \phi_2^{\mathbf{d}, \varepsilon}] \\ &= \langle V_i^{\mathbf{d}, \varepsilon} - \mathcal{I}^*(f(V_i^{\mathbf{d}, \varepsilon}) + \beta V_i^{\mathbf{d}, \varepsilon} \sum_{j \neq i} (V_j^{\mathbf{d}, \varepsilon})^2), P_\varepsilon \psi_l \rangle \\ &\quad + \sum_{n=1}^2 \langle V_n^{\mathbf{d}, \varepsilon} - \mathcal{I}^*(f(V_n^{\mathbf{d}, \varepsilon}) + \beta V_n^{\mathbf{d}, \varepsilon} \sum_{m \neq n} (V_m^{\mathbf{d}, \varepsilon})^2), \partial_{\delta_l} \phi_n^{\mathbf{d}, \varepsilon} \rangle \end{aligned}$$

From (2.10), Proposition 3.1 and recalling that K_i is spanned by $P_\varepsilon\psi_j$ for $j \in I_i$, we deduce the existence of coefficients $c_i^j = c_i^j(\varepsilon, \mathbf{d})$, $j \in I_i$ such that

$$V_i^{\mathbf{d},\varepsilon} - \mathcal{I}^*(f(V_i^{\mathbf{d},\varepsilon}) + \beta V_i^{\mathbf{d},\varepsilon} \sum_{j \neq i} (V_j^{\mathbf{d},\varepsilon})^2) = \sum_{j \in I_i} c_i^j \delta_j P_\varepsilon \psi_j, \quad i = 1, 2. \quad (4.2)$$

In conclusion, for $i \in \{1, 2\}$ and $l \in I_i$,

$$\sum_{j \in I_i} c_i^j \langle \delta_j P_\varepsilon \psi_j, \delta_l P_\varepsilon \psi_l \rangle + \sum_{n=1}^2 \sum_{j \in I_n} c_n^j \langle \delta_j P_\varepsilon \psi_j, \delta_l \partial_{\delta_l} \phi_n^{\mathbf{d},\varepsilon} \rangle = 0$$

A straightforward computation shows that

$$\langle \delta_j P_\varepsilon \psi_j, \delta_l P_\varepsilon \psi_l \rangle = o(1) \text{ for } l \neq j, \quad \langle \delta_l P_\varepsilon \psi_l, \delta_l P_\varepsilon \psi_l \rangle = \|\delta_l P_\varepsilon \psi_l\|^2 = \sigma_{ll} + o(1) \quad \text{as } \varepsilon \rightarrow 0 \quad (4.3)$$

for some constant $\sigma_{ll} > 0$ (see for instance [21, p. 417]). On the other hand, we have $\langle P_\varepsilon \psi_j, \partial_{\delta_l} \phi_n^{\mathbf{d},\varepsilon} \rangle = o(1)$. Indeed, since $\phi_n^{\mathbf{d},\varepsilon} \in K_n^\perp$ ($n = 1, 2$), then $\langle P_\varepsilon \psi_j, \phi_n^{\mathbf{d},\varepsilon} \rangle = 0$ for every \mathbf{d} . Therefore, taking the derivative of the previous identity with respect to δ_l ($l \in I_n$), we get $\langle P_\varepsilon \psi_j, \partial_{\delta_l} \phi_n^{\mathbf{d},\varepsilon} \rangle = -\langle \partial_{\delta_l} P_\varepsilon \psi_j, \phi_n^{\mathbf{d},\varepsilon} \rangle$. Combining (1.15) with Lemma A.3 we have $\|\delta_l \partial_{\delta_l} P_\varepsilon \psi_j\| = O(1)$, while Proposition 3.1 yields $\|\phi_n^{\mathbf{d},\varepsilon}\| = o(\delta_1)$. Therefore, $\langle P_\varepsilon \psi_j, \delta_l \partial_{\delta_l} \phi_n^{\mathbf{d},\varepsilon} \rangle = o(\delta_1) = o(1)$, as claimed. In conclusion, we end up with a linear system of the form

$$c_i^l \sigma_{ll} + \sum_{j \in I_i \setminus \{l\}} c_i^j o(1) + \sum_{n=1}^2 \sum_{j \in I_n} c_n^j o(1) = 0, \quad i = 1, 2, l \in I_i$$

which, as $\varepsilon \rightarrow 0$, has the unique solution $c_i^j = 0$ for every $i = 1, 2, j \in I_i$. Looking back at (4.2) we see that $(V_1^{\mathbf{d},\varepsilon}, V_2^{\mathbf{d},\varepsilon})$ solves (2.2), as we wanted. \square

We now compute the leading term of the reduced energy. For simplicity, and when there is no risk of confusion, we denote $\phi_i = \phi_i^{\mathbf{d},\varepsilon}$ and $U_i = U_{\delta_i}$. We have

$$\tilde{J}_\varepsilon(\mathbf{d}) = J_\varepsilon \left(\sum_{j \in I_1} P_\varepsilon U_j + \phi_1, \sum_{j \in I_2} P_\varepsilon U_j + \phi_2 \right) \quad (4.4)$$

$$= \sum_{i=1}^2 \int_{\Omega_\varepsilon} \left(\frac{1}{2} |\nabla(\sum_{j \in I_i} P_\varepsilon U_j + \phi_i)|^2 - F(\sum_{j \in I_i} P_\varepsilon U_j + \phi_i) \right) - \frac{\beta}{2} \int_{\Omega_\varepsilon} (\sum_{j \in I_1} P_\varepsilon U_j + \phi_1)^2 (\sum_{j \in I_2} P_\varepsilon U_j + \phi_2)^2 \quad (4.5)$$

$$= \sum_{i=1}^2 \sum_{j \in I_i} \int_{\Omega_\varepsilon} \left(\frac{1}{2} |\nabla P_\varepsilon U_j|^2 - F(P_\varepsilon U_j) \right) - \frac{\beta}{2} \sum_{i \in I_1, j \in I_2} \int_{\Omega_\varepsilon} (P_\varepsilon U_i)^2 (P_\varepsilon U_j)^2 + R(\mathbf{d}, \varepsilon) \quad (4.6)$$

$$= \sum_{i=1}^k \int_{\Omega_\varepsilon} \left(\frac{1}{2} |\nabla P_\varepsilon U_i|^2 - \frac{1}{4} (P_\varepsilon U_i)^4 \right) - \frac{\beta}{2} \sum_{i \in I_1, j \in I_2} \int_{\Omega_\varepsilon} (P_\varepsilon U_i)^2 (P_\varepsilon U_j)^2 + R(\mathbf{d}, \varepsilon), \quad (4.7)$$

where

$$R(\mathbf{d}, \varepsilon) = J_\varepsilon \left(\sum_{j \in I_1} P_\varepsilon U_j + \phi_1, \sum_{j \in I_2} P_\varepsilon U_j + \phi_2 \right) - \sum_{i \in I_1, j \in I_2} J_\varepsilon(P_\varepsilon U_i, P_\varepsilon U_j)$$

will be an higher order term.

In what follows we show that the reduced energy reads as

$$\tilde{J}_\varepsilon(\mathbf{d}) = c_1 + c_2\delta_1^2 + c_3 \left(\frac{\varepsilon}{\delta_k}\right)^2 - \beta c_4 \sum_{i=1}^{k-1} \left(\frac{\delta_{i+1}}{\delta_i}\right)^2 \log \frac{\delta_i}{\delta_{i+1}} + h.o.t., \quad (4.8)$$

for some constants $c_1, c_2, c_3, c_4 > 0$. This yields the choice of parameters (2.5) (which for convenience of the reader we recall)

$$\delta_j := d_j \varepsilon^{\frac{j}{k+1}} \left(\log \frac{1}{\varepsilon}\right)^{\frac{1}{2} - \frac{j}{k+1}} \quad \text{with } d_j > 0 \text{ for } j = 1, \dots, k.$$

and the existence of towers of bubbles as we want. Observe that, as $\varepsilon \rightarrow 0$,

$$\delta_1^2 \sim \left(\frac{\varepsilon}{\delta_k}\right)^2 \sim \left(\frac{\delta_{i+1}}{\delta_i}\right)^2 \log \frac{\delta_i}{\delta_{i+1}} \sim \varepsilon^{\frac{2}{k+1}} \left(\log \frac{1}{\varepsilon}\right)^{\frac{k-1}{k+1}}$$

(see ahead for the details).

Lemma 4.2. *Given $i = 1, \dots, k$ we have*

$$\int_{\Omega_\varepsilon} \left(\frac{1}{2} |\nabla P_\varepsilon U_i|^2 - \frac{1}{4} (P_\varepsilon U_i)^4 \right) = \frac{B}{4} + \frac{A^2}{2} \tau(0) \delta_i^2 + \frac{\Gamma}{2} \left(\frac{\varepsilon}{\delta_i}\right)^2 + o(\delta_i^2) + o\left(\left(\frac{\varepsilon}{\delta_i}\right)^2\right)$$

as $\varepsilon \rightarrow 0$, uniformly for every $\mathbf{d} \in X_\eta$. We recall that A, B and Γ are defined in (1.16)–(1.17), while τ is the Robin function (see the notation section).

Proof. We reason similarly to [20, Lemma 4.3], to which we refer for more details.

First of all, using (1.6), we have that

$$\begin{aligned} \frac{1}{2} \int_{\Omega_\varepsilon} |\nabla P_\varepsilon U_i|^2 - \frac{1}{4} \int_{\Omega_\varepsilon} (P_\varepsilon U_i)^4 &= \frac{1}{2} \int_{\Omega_\varepsilon} (P_\varepsilon U_i) U_i^3 - \frac{1}{4} \int_{\Omega_\varepsilon} (P_\varepsilon U_i)^4 \\ &= \frac{1}{4} \int_{\Omega_\varepsilon} U_i^4 + \frac{1}{2} \int_{\Omega_\varepsilon} (P_\varepsilon U_i - U_i) U_i^3 - \frac{1}{4} \int_{\Omega_\varepsilon} ((P_\varepsilon U_i)^4 - U_i^4) \end{aligned} \quad (4.9)$$

Using a Taylor expansion up to second order, we have that

$$(P_\varepsilon U_i)^4 - U_i^4 = 4U_i^3(P_\varepsilon U_i - U_i) + 6(U_i^2 + \xi(P_\varepsilon U_i - U_i))^2(P_\varepsilon U_i - U_i)^2,$$

for some function $\xi(x) \in [0, 1]$. Therefore, we can rewrite (4.9) as

$$\frac{1}{4} \int_{\Omega_\varepsilon} U_i^4 - \frac{1}{2} \int_{\Omega_\varepsilon} (P_\varepsilon U_i - U_i) U_i^3 - \frac{3}{2} \int_{\Omega_\varepsilon} (U_i + \xi(P_\varepsilon U_i - U_i))^2 (P_\varepsilon U_i - U_i)^2. \quad (4.10)$$

We now estimate each one of the three terms separately. The first term in (4.10) is, after a change of variables $x = \delta_i y$ and recalling that $\Omega_\varepsilon = \Omega \setminus B_\varepsilon$ and $\varepsilon/\delta_i \rightarrow 0$,

$$\int_{\Omega_\varepsilon} U_i^4 = \int_{\Omega_\varepsilon} \frac{\alpha_4^4 \delta_i^4}{(\delta_i^2 + |x|^2)^4} dx = \int_{\Omega_\varepsilon \setminus \delta_i} \frac{\alpha_4^4}{(1 + |y|^2)^4} dy \quad (4.11)$$

$$= B + \int_{\mathbb{R}^N \setminus (\Omega/\delta_i)} \frac{\alpha_4^4}{(1 + |y|^2)^4} dy + \int_{B_{\varepsilon/\delta_i}} \frac{\alpha_4^4}{(1 + |y|^2)^4} dy = B + O(\delta_i^4) + O\left(\left(\frac{\varepsilon}{\delta_i}\right)^4\right) \quad (4.12)$$

$$= B + o(\delta_i^2) + o\left(\left(\frac{\varepsilon}{\delta_i}\right)^2\right) \quad (4.13)$$

As for the second term, we use the fact that

$$P_\varepsilon U_i - U_i = -A\delta_i H(x, 0) - \frac{\alpha_4}{\delta_i} \frac{\varepsilon^2}{|x|^2} + R(x)$$

(by Lemma A.1, which we can apply since $\varepsilon/\delta_i \rightarrow 0$ as $\varepsilon \rightarrow 0$). We have

$$\int_{\Omega_\varepsilon} (P_\varepsilon U_i - U_i) U_i^3 = \int_{\Omega_\varepsilon} \left(-A\delta_i H(x, 0) - \frac{\alpha_4 \varepsilon^2}{\delta_i |x|^2}\right) U_i^3 + \int_{\Omega_\varepsilon} R(x) U_i^3 \quad (4.14)$$

$$= \int_{\Omega_\varepsilon} \frac{\alpha_4^3 \delta_i^3}{(\delta_i^2 + |x|^2)^3} \left(-A\delta_i H(x, 0) - \frac{\alpha_4 \varepsilon^2}{\delta_i |x|^2}\right) + \int_{\Omega_\varepsilon} R(x) U_i^3 \quad (4.15)$$

$$= -A \int_{\Omega_\varepsilon/\delta_i} \frac{\alpha_4^3 \delta_i^2}{(1 + |y|^2)^3} H(\delta_i y, 0) - \int_{\Omega_\varepsilon/\delta_i} \left(\frac{\varepsilon}{\delta_i}\right)^2 \frac{\alpha_4^4}{|y|^2 (1 + |y|^2)^3} + \int_{\Omega_\varepsilon} R(x) U_i^3 \quad (4.16)$$

$$= -A^2 \tau(0) \delta_i^2 - \Gamma \left(\frac{\varepsilon}{\delta_i}\right)^2 + o(\delta_i^2) + o\left(\left(\frac{\varepsilon}{\delta_i}\right)^2\right) \quad (4.17)$$

where we have used the estimates for the remainder term R contained in Lemma A.1.

As for the last term in (4.10), since $0 \leq P_\varepsilon U_i \leq U_i$ (by the maximum principle) and $\xi(x) \in [0, 1]$, we have $0 \leq U_i + \xi(PU_i - U_i) \leq U_i$. Combining this with Lemma A.2 and since $\varepsilon/\delta_i \rightarrow 0$,

$$\left| \int_{\Omega_\varepsilon} (U_i + \xi(PU_i - U_i))^2 (PU_i - U_i)^2 \right| \leq \int_{\Omega_\varepsilon} U_i^2 (PU_i - U_i)^2 = O\left(\delta_i^4 |\log \delta_i| + \left(\frac{\varepsilon}{\delta_i}\right)^4 \left| \log \left(\frac{\varepsilon}{\delta_i}\right) \right|\right) \quad (4.18)$$

$$= o(\delta_i^2) + o\left(\left(\frac{\varepsilon}{\delta_i}\right)^2\right). \quad (4.19)$$

The result follows combining (4.10) with (4.13)–(4.17)–(4.19). \square

Corollary 4.3. *The following estimate holds*

$$\begin{aligned} \sum_{i=1}^k \int_{\Omega_\varepsilon} \frac{1}{2} |\nabla P_\varepsilon U_i|^2 - \frac{1}{4} (P_\varepsilon U_i)^4 dx &= k \frac{B}{4} + \frac{A^2}{2} \tau(0) \delta_1^2 + \frac{\Gamma}{2} \left(\frac{\varepsilon}{\delta_k}\right)^2 + o(\delta_1^2) + o\left(\left(\frac{\varepsilon}{\delta_k}\right)^2\right) \\ &= k \frac{B}{4} + \left(\frac{A^2}{2} \tau(0) d_1^2 + \frac{\Gamma}{2} \left(\frac{1}{d_k}\right)^2\right) \varepsilon^{\frac{2}{k+1}} \left(\log \frac{1}{\varepsilon}\right)^{\frac{k-1}{k+1}} + o\left(\varepsilon^{\frac{2}{k+1}} \left(\log \frac{1}{\varepsilon}\right)^{\frac{k-1}{k+1}}\right) \end{aligned} \quad (4.20)$$

as $\varepsilon \rightarrow 0$, uniformly for every $\mathbf{d} \in X_\eta$.

Proof. From the previous lemma we see that

$$\sum_{i=1}^k \int_{\Omega_\varepsilon} \frac{1}{2} |\nabla P_\varepsilon U_i|^2 - \frac{1}{4} (P_\varepsilon U_i)^4 dx = k \frac{B}{4} + \frac{A^2}{2} \tau(0) \sum_{i=1}^k \delta_i^2 + \frac{\Gamma}{2} \sum_{i=1}^k \left(\frac{\varepsilon}{\delta_i}\right)^2 + \sum_{i=1}^k \left(o(\delta_i^2) + o\left(\left(\frac{\varepsilon}{\delta_i}\right)^2\right)\right)$$

and the first identity of the lemma follows because $d \in N_\eta$ and $\delta_i = o(\delta_1)$ for every $i \in \{2, \dots, k\}$ (recall Remark 2.1), which implies that $\varepsilon^2/\delta_i^2 = o(\varepsilon^2/\delta_k^2)$ for $i \in \{1, \dots, k-1\}$ as $\varepsilon \rightarrow 0$. The second identity follows directly from the definition of d_i (see (2.5)). \square

Lemma 4.4. *Given $i, j \in \{1, \dots, k\}$ with $i > j$, we have*

$$\int_{\Omega_\varepsilon} (P_\varepsilon U_i)^2 (P_\varepsilon U_j)^2 = \alpha_4^4 |\mathbb{S}^3| \left(\frac{\delta_i}{\delta_j} \right)^2 \log \frac{\delta_j}{\delta_i} + o \left(\left(\frac{\delta_i}{\delta_j} \right)^2 \log \frac{\delta_j}{\delta_i} \right)$$

as $\varepsilon \rightarrow 0$, uniformly for every $\mathbf{d} \in X_\eta$.

Proof. First, we rewrite

$$\int_{\Omega_\varepsilon} (P_\varepsilon U_i)^2 (P_\varepsilon U_j)^2 = \int_{\Omega_\varepsilon} U_i^2 U_j^2 + \int_{\Omega_\varepsilon} ((P_\varepsilon U_i)^2 (P_\varepsilon U_j)^2 - (U_i)^2 (U_j)^2) = \int_{\Omega_\varepsilon} U_i^2 U_j^2 + h.o.t. \quad (4.21)$$

(by Lemma A.1). We estimate the leading term as follows: for $\varepsilon > 0$ small and $r > 0$ such that $B_\varepsilon \subset B_r \subset \Omega$ and $\varepsilon < \sqrt{\delta_i \delta_j}$,

$$\int_{\Omega_\varepsilon} U_i^2 U_j^2 = \underbrace{\int_{\{\varepsilon \leq |x| \leq \sqrt{\delta_i \delta_j}\}} U_i^2 U_j^2}_{(I)} + \underbrace{\int_{\{\sqrt{\delta_i \delta_j} \leq |x| \leq r\}} U_i^2 U_j^2}_{(II)} + \underbrace{\int_{\Omega \setminus B_r} U_i^2 U_j^2}_{(III)}.$$

Asymptotic estimate of (I): scaling $x = \delta_i y$,

$$(I) = \int_{\{\varepsilon \leq |x| \leq \sqrt{\delta_i \delta_j}\}} U_i^2 U_j^2 = \alpha_4^4 \delta_i^2 \delta_j^2 \int_{\{\varepsilon \leq |x| \leq \sqrt{\delta_i \delta_j}\}} \frac{1}{(\delta_i^2 + |x|^2)^2} \frac{1}{(\delta_j^2 + |x|^2)^2} dx \quad (4.22)$$

$$= \alpha_4^4 \delta_i^2 \delta_j^2 \int_{\left\{ \frac{\varepsilon}{\delta_i} \leq |y| \leq \sqrt{\frac{\delta_j}{\delta_i}} \right\}} \frac{1}{(1 + |y|^2)^2} \frac{1}{(\delta_j^2 + \delta_i^2 |y|^2)^2} dy \quad (4.23)$$

$$= \alpha_4^4 \left(\frac{\delta_i}{\delta_j} \right)^2 \int_{\left\{ \frac{\varepsilon}{\delta_i} \leq |y| \leq \sqrt{\frac{\delta_j}{\delta_i}} \right\}} \frac{1}{(1 + |y|^2)^2} \frac{1}{(1 + (\delta_i/\delta_j)^2 |y|^2)^2} dy \quad (4.24)$$

We have:

$$\begin{aligned} & \int_{\left\{ \frac{\varepsilon}{\delta_i} \leq |y| \leq \sqrt{\frac{\delta_j}{\delta_i}} \right\}} \frac{1}{(1 + |y|^2)^2} \frac{1}{(1 + (\delta_i/\delta_j)^2 |y|^2)^2} dy \\ &= \underbrace{\int_{\left\{ \frac{\varepsilon}{\delta_i} \leq |y| \leq \sqrt{\frac{\delta_j}{\delta_i}} \right\}} \frac{1}{(1 + |y|^2)^2} dy}_{(I.a)} + \underbrace{\int_{\left\{ \frac{\varepsilon}{\delta_i} \leq |y| \leq \sqrt{\frac{\delta_j}{\delta_i}} \right\}} \frac{1}{(1 + |y|^2)^2} \left(\frac{1}{(1 + (\delta_i/\delta_j)^2 |y|^2)^2} - 1 \right) dy}_{(I.b)}. \end{aligned}$$

The first term can be estimated as follows:

$$\begin{aligned}
(I.a) &= |\mathbb{S}^3| \int_{\frac{\varepsilon}{\delta_i}}^{\sqrt{\frac{\delta_j}{\delta_i}}} \frac{r^3}{(1+r^2)^2} dr = \frac{1}{2} |\mathbb{S}^3| \left[\frac{1}{1+r^2} + \log(1+r^2) \right]_{r=\frac{\varepsilon}{\delta_i}}^{r=\sqrt{\frac{\delta_j}{\delta_i}}} \\
&= \frac{1}{2} |\mathbb{S}^3| \left(\frac{1}{1+\delta_j/\delta_i} - \frac{1}{1+\varepsilon^2/\delta_i^2} + \log \left(1 + \frac{\delta_j}{\delta_i} \right) - \log \left(1 + \frac{\varepsilon^2}{\delta_i^2} \right) \right) \\
&= \frac{1}{2} |\mathbb{S}^3| \log \frac{\delta_j}{\delta_i} + o \left(\log \frac{\delta_j}{\delta_i} \right),
\end{aligned} \tag{4.25}$$

since, as $\delta_j/\delta_i \rightarrow \infty$ ($i > j$) and $\varepsilon/\delta_i \rightarrow 0$ (recall Remark 2.1):

$$\frac{1}{1+\delta_j/\delta_i} - \frac{1}{1+\varepsilon^2/\delta_i^2} = -1 + o(1) = o \left(\log \frac{\delta_j}{\delta_i} \right), \quad \log \left(1 + \delta_j/\delta_i \right) = \log \frac{\delta_j}{\delta_i} + o \left(\log \frac{\delta_j}{\delta_i} \right)$$

and

$$\log \left(1 + \varepsilon^2/\delta_i^2 \right) = o(1) = o \left(\log \frac{\delta_j}{\delta_i} \right).$$

As for the second term, because $(\delta_i/\delta_j)|y| \leq \sqrt{\delta_i/\delta_j} \leq c$ for $|y| \leq \sqrt{\delta_j/\delta_i}$, as $\varepsilon \rightarrow 0$, and recalling the computation done for (I.a), we have

$$\begin{aligned}
|(I.b)| &= \int_{\left\{ \frac{\varepsilon}{\delta_i} \leq |y| \leq \sqrt{\frac{\delta_j}{\delta_i}} \right\}} \frac{1}{(1+|y|^2)^2} \left(\frac{|(\delta_i/\delta_j)^4|y|^4 + 2(\delta_i/\delta_j)^2|y|^2}{(1+(\delta_i/\delta_j)^2|y|^2)^2} \right) dy \\
&\leq c' \frac{\delta_i}{\delta_j} \int_{\left\{ \frac{\varepsilon}{\delta_i} \leq |y| \leq \sqrt{\frac{\delta_j}{\delta_i}} \right\}} \frac{1}{(1+|y|^2)^2} dy \leq c'' \frac{\delta_i}{\delta_j} \log \frac{\delta_j}{\delta_i} = o \left(\log \frac{\delta_j}{\delta_i} \right).
\end{aligned}$$

Combining the expansions of (I.a) and (I.b) with (4.24) yields, in conclusion, that

$$(I) = \frac{\alpha_4^4}{2} |\mathbb{S}^3| \left(\frac{\delta_i}{\delta_j} \right)^2 \log \frac{\delta_j}{\delta_i} + o \left(\left(\frac{\delta_i}{\delta_j} \right)^2 \log \frac{\delta_j}{\delta_i} \right).$$

Asymptotic estimate of (II): by using this time the scaling $x = \delta_j y$ and the fact that

$$\int \frac{1}{r(1+r^2)} dr = \log r - \frac{1}{2} \log(1+r^2) + \frac{1}{2(1+r^2)},$$

we have

$$\begin{aligned}
 (II) &= \int_{\{\sqrt{\delta_i \delta_j} \leq |x| \leq r\}} U_i^2 U_j^2 = \alpha_4^4 \delta_i^2 \delta_j^2 \int_{\{\sqrt{\delta_i \delta_j} \leq |x| \leq r\}} \frac{1}{(\delta_i^2 + |x|^2)^2} \frac{1}{(\delta_j^2 + |x|^2)^2} dx \\
 &= \alpha_4^4 \delta_i^2 \delta_j^2 \int_{\left\{\sqrt{\frac{\delta_i}{\delta_j}} \leq |y| \leq \frac{r}{\delta_j}\right\}} \frac{1}{(\delta_i^2 + \delta_j^2 |y|^2)^2} \frac{1}{(1 + |y|^2)^2} dy \\
 &= \alpha_4^4 \left(\frac{\delta_i}{\delta_j}\right)^2 \int_{\left\{\sqrt{\frac{\delta_i}{\delta_j}} \leq |y| \leq \frac{r}{\delta_j}\right\}} \frac{1}{((\delta_i/\delta_j)^2 + |y|^2)^2} \frac{1}{(1 + |y|^2)^2} dy \\
 &= \alpha_4^4 \left(\frac{\delta_i}{\delta_j}\right)^2 \int_{\left\{\sqrt{\frac{\delta_i}{\delta_j}} \leq |y| \leq \frac{r}{\delta_j}\right\}} \frac{1}{|y|^4 (1 + |y|^2)^2} dy + o\left(\left(\frac{\delta_i}{\delta_j}\right)^2 \log \frac{\delta_j}{\delta_i}\right) \\
 &= \alpha_4^4 \left(\frac{\delta_i}{\delta_j}\right)^2 |\mathbb{S}^3| \int_{\frac{\sqrt{\frac{\delta_i}{\delta_j}}}{\delta_j}}^{\frac{r}{\delta_j}} \frac{1}{r(1+r^2)^2} dr + o\left(\left(\frac{\delta_i}{\delta_j}\right)^2 \log \frac{\delta_j}{\delta_i}\right) \\
 &= \frac{\alpha_4^4}{2} |\mathbb{S}^3| \left(\frac{\delta_i}{\delta_j}\right)^2 \log \frac{\delta_j}{\delta_i} + o\left(\left(\frac{\delta_i}{\delta_j}\right)^2 \log \frac{\delta_j}{\delta_i}\right).
 \end{aligned}$$

Asymptotic estimate of (III):

$$\begin{aligned}
 0 \leq \int_{\Omega \setminus B_r} U_i^2 U_j^2 &= \alpha_4^4 \delta_i^2 \delta_j^2 \int_{\Omega \setminus B_r} \frac{1}{(\delta_i^2 + |y|^2)^2} \frac{1}{(\delta_j^2 + |y|^2)^2} = \alpha_4^4 \left(\frac{\delta_i}{\delta_j}\right)^2 \int_{\Omega \setminus B_r} \frac{1}{(\delta_i^2 + |y|^2)^2} \frac{1}{(1 + \delta_j^{-2} |y|^2)^2} \\
 &\leq \alpha_4^4 \left(\frac{\delta_i}{\delta_j}\right)^2 \int_{\Omega \setminus B_r} \frac{1}{|y|^4} \leq c \left(\frac{\delta_i}{\delta_j}\right)^2 = o\left(\left(\frac{\delta_i}{\delta_j}\right)^2 \log \frac{\delta_j}{\delta_i}\right).
 \end{aligned}$$

By combining the estimates of (I), (II) and (III) we deduce that

$$\int_{\Omega_\varepsilon} U_i^2 U_j^2 = \alpha_4^4 |\mathbb{S}^3| \left(\frac{\delta_i}{\delta_j}\right)^2 \log \frac{\delta_j}{\delta_i} + o\left(\left(\frac{\delta_i}{\delta_j}\right)^2 \log \frac{\delta_j}{\delta_i}\right)$$

which yields the desired conclusion. □

Corollary 4.5. *We have, as $\varepsilon \rightarrow 0$, uniformly for every $\mathbf{d} \in X_\eta$,*

$$\begin{aligned}
 \sum_{i \in I_1, j \in I_2} \int_{\Omega_\varepsilon} (P_\varepsilon U_i)^2 (P_\varepsilon U_j)^2 &= \alpha_4^4 |\mathbb{S}^3| \sum_{i=1}^{k-1} \left(\frac{\delta_{i+1}}{\delta_i}\right)^2 \log \frac{\delta_i}{\delta_{i+1}} + \sum_{i=1}^{k-1} o\left(\left(\frac{\delta_{i+1}}{\delta_i}\right)^2 \log \frac{\delta_i}{\delta_{i+1}}\right) \\
 &= \frac{\alpha_4^4}{k+1} |\mathbb{S}^3| \sum_{i=1}^{k-1} \left(\frac{d_{i+1}}{d_i}\right)^2 \varepsilon^{\frac{2}{k+1}} \left(\log \frac{1}{\varepsilon}\right)^{\frac{k-1}{k+1}} + o\left(\varepsilon^{\frac{1}{k+1}} \left(\log \frac{1}{\varepsilon}\right)^{\frac{k-1}{k+1}}\right)
 \end{aligned} \tag{4.26}$$

Proof. The first identity is a simple consequence of the previous lemma together with the fact that $\delta_l = o(\delta_i)$ as $\varepsilon \rightarrow 0$, for $l > i$. In fact, since each one of the sets I_1 and I_2 do not contain two consecutive integers, and that given $i > j$ with $|i - j| > 1$ it holds

$$\left(\frac{\delta_i}{\delta_j}\right)^2 \log \frac{\delta_j}{\delta_i} = o\left(\left(\frac{\delta_{j+1}}{\delta_j}\right)^2 \log \frac{\delta_j}{\delta_{j+1}}\right),$$

then

$$\sum_{i \in I_1, j \in I_2} \int_{\Omega_\varepsilon} (P_\varepsilon U_i)^2 (P_\varepsilon U_j)^2 = \sum_{i \in I_1, j \in I_2} \left(\alpha_4^4 |\mathbb{S}^3| \left(\frac{\delta_i}{\delta_j}\right)^2 \log \frac{\delta_j}{\delta_i} + o\left(\left(\frac{\delta_i}{\delta_j}\right)^2 \log \frac{\delta_j}{\delta_i}\right) \right) \quad (4.27)$$

$$= \alpha_4^4 |\mathbb{S}^3| \sum_{i=1}^{k-1} \left(\frac{\delta_{i+1}}{\delta_i}\right)^2 \log \frac{\delta_i}{\delta_{i+1}} + \sum_{i=1}^{k-1} o\left(\left(\frac{\delta_{i+1}}{\delta_i}\right)^2 \log \frac{\delta_i}{\delta_{i+1}}\right). \quad (4.28)$$

The last identity of the statement is a consequence of the definition of δ_i and the fact that

$$\begin{aligned} \left(\frac{\delta_{i+1}}{\delta_i}\right)^2 \log \frac{\delta_i}{\delta_{i+1}} &= \left(\frac{d_{i+1}}{d_i}\right)^2 \varepsilon^{\frac{2}{k+1}} \left(\log \frac{1}{\varepsilon}\right)^{-\frac{2}{k+1}} \log \left(\frac{d_i}{d_{i+1}} \left(\frac{1}{\varepsilon}\right)^{\frac{1}{k+1}} \left(\log \frac{1}{\varepsilon}\right)^{\frac{1}{k+1}} \right) \\ &= \frac{1}{k+1} \left(\frac{d_{i+1}}{d_i}\right)^2 \varepsilon^{\frac{2}{k+1}} \left(\log \frac{1}{\varepsilon}\right)^{\frac{k-1}{k+1}} + o\left(\varepsilon^{\frac{2}{k+1}} \left(\log \frac{1}{\varepsilon}\right)^{\frac{k-1}{k+1}}\right) \quad \square \end{aligned}$$

Lemma 4.6. *We have*

$$R(\mathbf{d}, \varepsilon) = o(\delta_1^2) = o\left(\varepsilon^{\frac{2}{k+1}} \left(\log \frac{1}{\varepsilon}\right)^{\frac{k-1}{k+1}}\right) \quad (4.29)$$

as $\varepsilon \rightarrow 0$, uniformly for every $\mathbf{d} \in X_\eta$.

Proof. Recall that $F(s) = (s^+)^4/4$, and we denote $f(s) := F'(s) = (s^+)^3$. We have

$$\begin{aligned} R(\mathbf{d}, \varepsilon) &= J_\varepsilon \left(\sum_{j \in I_1} P_\varepsilon U_j + \phi_1, \sum_{j \in I_2} P_\varepsilon U_j + \phi_2 \right) - \sum_{i \in I_1, j \in I_2} J_\varepsilon(P_\varepsilon U_i, P_\varepsilon U_j) \\ &= \frac{1}{2} \sum_{i=1}^2 \sum_{\substack{j, k \in I_i \\ j \neq k}} \int_{\Omega_\varepsilon} \nabla P_\varepsilon U_j \cdot \nabla P_\varepsilon U_k + \frac{1}{2} \sum_{i=1}^2 \int_{\Omega_\varepsilon} |\nabla \phi_i|^2 + \sum_{i=1}^2 \sum_{j \in I_i} \int_{\Omega_\varepsilon} \nabla P_\varepsilon U_j \cdot \nabla \phi_i \\ &\quad + \sum_{i=1}^2 \int_{\Omega_\varepsilon} \left(\sum_{j \in I_i} F(P_\varepsilon U_j) - F\left(\sum_{j \in I_i} P_\varepsilon U_j + \phi_i\right) \right) \\ &\quad + \frac{\beta}{2} \sum_{i \in I_1, j \in I_2} \int_{\Omega_\varepsilon} (P_\varepsilon U_i)^2 (P_\varepsilon U_j)^2 - \frac{\beta}{2} \int_{\Omega_\varepsilon} \left(\sum_{j \in I_1} P_\varepsilon U_j + \phi_1 \right)^2 \left(\sum_{j \in I_2} P_\varepsilon U_j + \phi_2 \right)^2 \end{aligned}$$

Recalling the definition of P_ε from (1.6) and adding and subtracting terms of type $F(\sum_{j \in I_i} P_\varepsilon U_j)$ and $f(\sum_{j \in I_i} P_\varepsilon U_j)\phi_i$, we have

$$\begin{aligned}
 & \frac{1}{2} \sum_{i=1}^2 \sum_{\substack{j,k \in I_i \\ j \neq k}} \int_{\Omega_\varepsilon} \nabla P_\varepsilon U_j \cdot \nabla P_\varepsilon U_k + \frac{1}{2} \sum_{i=1}^2 \int_{\Omega_\varepsilon} |\nabla \phi_i|^2 + \sum_{i=1}^2 \sum_{j \in I_i} \int_{\Omega_\varepsilon} \nabla P_\varepsilon U_j \cdot \nabla \phi_i \\
 & + \sum_{i=1}^2 \int_{\Omega_\varepsilon} \left(\sum_{j \in I_i} F(P_\varepsilon U_j) - F(\sum_{j \in I_i} P_\varepsilon U_j + \phi_i) \right) \\
 & = \frac{1}{2} \sum_{i=1}^2 \sum_{\substack{j,k \in I_i \\ j \neq k}} \int_{\Omega_\varepsilon} U_j^3 P_\varepsilon U_k \, dx + \sum_{i=1}^2 \int_{\Omega_\varepsilon} \left(\sum_{j \in I_i} F(P_\varepsilon U_j) - F(\sum_{j \in I_i} P_\varepsilon U_j) \right) \\
 & + \frac{1}{2} \sum_{i=1}^2 \int_{\Omega_\varepsilon} |\nabla \phi_i|^2 - \sum_{i=1}^2 \int_{\Omega_\varepsilon} \left(F(\sum_{j \in I_i} P_\varepsilon U_j + \phi_i) - F(\sum_{j \in I_i} P_\varepsilon U_j) - f(\sum_{j \in I_i} P_\varepsilon U_j)\phi_i \right) \\
 & + \sum_{i=1}^2 \int_{\Omega_\varepsilon} \left(\sum_{j \in I_i} f(U_j) - f(\sum_{j \in I_i} P_\varepsilon U_j) \right) \phi_i.
 \end{aligned}$$

Moreover,

$$\begin{aligned}
 & \frac{\beta}{2} \sum_{i \in I_1, j \in I_2} \int_{\Omega_\varepsilon} (P_\varepsilon U_i)^2 (P_\varepsilon U_j)^2 - \frac{\beta}{2} \int_{\Omega_\varepsilon} (\sum_{j \in I_1} P_\varepsilon U_j + \phi_1)^2 (\sum_{j \in I_2} P_\varepsilon U_j + \phi_2)^2 \\
 & = -\frac{\beta}{2} \int_{\Omega_\varepsilon} \sum_{\substack{i,j \in I_1 \\ i \neq j}} \sum_{\substack{k,l \in I_2 \\ k \neq l}} P_\varepsilon U_i P_\varepsilon U_j P_\varepsilon U_k P_\varepsilon U_l - \frac{\beta}{2} \int_{\Omega_\varepsilon} \sum_{\substack{i,j=1 \\ i \neq j}}^2 \sum_{k,l \in I_i} P_\varepsilon U_k P_\varepsilon U_l \sum_{m \in I_j} P_\varepsilon U_m \phi_j \\
 & - \frac{\beta}{2} \int_{\Omega_\varepsilon} \sum_{\substack{i,j=1 \\ i \neq j}}^2 \sum_{k,l \in I_i} P_\varepsilon U_k P_\varepsilon U_l \phi_j^2 - \frac{\beta}{2} \int_{\Omega_\varepsilon} 4 \sum_{i \in I_1, j \in I_2} P_\varepsilon U_i P_\varepsilon U_j \phi_1 \phi_2 \\
 & - \frac{\beta}{2} \int_{\Omega_\varepsilon} 2 \sum_{\substack{i,j=1 \\ i \neq j}}^2 \sum_{k \in I_i} P_\varepsilon U_k \phi_i \phi_j^2 - \frac{\beta}{2} \int_{\Omega_\varepsilon} \phi_1^2 \phi_2^2.
 \end{aligned}$$

Let us rewrite $R(\mathbf{d}, \varepsilon)$ as

$$\begin{aligned}
R(\mathbf{d}, \varepsilon) &= \underbrace{\frac{1}{2} \sum_{i=1}^2 \sum_{\substack{j, h \in I_i \\ j \neq h}} \int_{\Omega_\varepsilon} U_j^3 P_\varepsilon U_h - \frac{\beta}{2} \int_{\Omega_\varepsilon} \sum_{\substack{i, j \in I_1 \\ i \neq j}} \sum_{\substack{h, l \in I_2 \\ h \neq l}} P_\varepsilon U_i P_\varepsilon U_j P_\varepsilon U_h P_\varepsilon U_l}_{:=a_1} \\
&+ \underbrace{\int_{\Omega_\varepsilon} \sum_{i=1}^2 \int_{\Omega_\varepsilon} \left(\sum_{j \in I_i} F(P_\varepsilon U_j) - F\left(\sum_{j \in I_i} P_\varepsilon U_j\right) \right)}_{:=a_2} \\
&+ \underbrace{\sum_{i=1}^2 \int_{\Omega_\varepsilon} \left(F\left(\sum_{j \in I_i} P_\varepsilon U_j\right) - F\left(\sum_{j \in I_i} P_\varepsilon U_j + \phi_i\right) + f\left(\sum_{j \in I_i} P_\varepsilon U_j\right) \phi_i \right)}_{:=a_3} \\
&+ \underbrace{\sum_{i=1}^2 \int_{\Omega_\varepsilon} \left(\sum_{j \in I_i} f(U_j) - f\left(\sum_{j \in I_i} P_\varepsilon U_j\right) \right) \phi_i - \frac{\beta}{2} \int_{\Omega_\varepsilon} \sum_{\substack{i, j=1 \\ i \neq j}}^2 \sum_{h, l \in I_i} P_\varepsilon U_h P_\varepsilon U_l \sum_{m \in I_j} P_\varepsilon U_m \phi_j}_{:=a_4} \\
&- \underbrace{\frac{\beta}{2} \int_{\Omega_\varepsilon} \sum_{\substack{i, j=1 \\ i \neq j}}^2 \sum_{h, l \in I_i} P_\varepsilon U_h P_\varepsilon U_l \phi_j^2 - \frac{\beta}{2} \int_{\Omega_\varepsilon} 4 \sum_{i \in I_1, j \in I_2} P_\varepsilon U_i P_\varepsilon U_j \phi_1 \phi_2}_{:=a_5} \\
&- \underbrace{\frac{\beta}{2} \int_{\Omega_\varepsilon} 2 \sum_{\substack{i, j=1 \\ i \neq j}}^2 \sum_{h \in I_i} P_\varepsilon U_h \phi_i \phi_j^2 + \frac{1}{2} \sum_{i=1}^2 \int_{\Omega_\varepsilon} |\nabla \phi_i|^2 - \frac{\beta}{2} \int_{\Omega_\varepsilon} \phi_1^2 \phi_2^2}_{:=a_6}.
\end{aligned}$$

Estimates for a_1, a_2 . First of all, we check that the first two terms satisfy $a_1, a_2 = o(\delta_1^2)$. Indeed, since $0 \leq P_\varepsilon U_i \leq U_i$ (by the maximum principle), $a_1 + a_2$ is controlled by a sum of terms of the form $\int_{\Omega_\varepsilon} P_\varepsilon U_i P_\varepsilon U_j P_\varepsilon U_h P_\varepsilon U_l$ for indices j, l, h, i not all equal at the same time; each term is of higher order with respect to the leading term δ_1^2 , as we will now check. Indeed, if $i \neq j$ we have by Lemma A.4 that

$$\begin{aligned}
\int_{\Omega_\varepsilon} U_i^3 U_j &= \begin{cases} O\left(\frac{\delta_i}{\delta_j}\right) \int_{\Omega_\varepsilon/\delta_i} \frac{1}{(1+|y|^2)^3} = O\left(\frac{\delta_i}{\delta_j}\right) & \text{if } i > j \\ O\left(\frac{\delta_j}{\delta_i}\right) \int_{\Omega_\varepsilon/\delta_i} \frac{1}{(1+|y|^2)^3 |y|^2} = O\left(\frac{\delta_j}{\delta_i}\right) & \text{if } j > i. \end{cases} \\
&= o(\delta_1^2)
\end{aligned}$$

because we are always in a situation that i, j belong to the same set I_h , thus $|i - j| > 1$, and then by the choices we did in (2.5),

$$\frac{\delta_i}{\delta_j} = O\left(\varepsilon^{\frac{i-j}{k+1}} \left(\ln \frac{1}{\varepsilon}\right)^{-\frac{i-j}{k+1}}\right) = o(\delta_1^2) \quad \text{whenever } i - j \geq 2.$$

Moreover, if $i \neq j$, then assuming without loss of generality that $i > j$ with $|i - j| > 1$ we have by Lemma A.5

$$\int_{\Omega_\varepsilon} U_i^2 U_j^2 = O\left(\frac{\delta_i^2}{\delta_j^2} |\log \delta_i|\right) = o\left(\frac{\delta_i}{\delta_j}\right) = o(\delta_1^2).$$

(note that this term only appears in a_2). In a similar way, if $i, j, l \in I_h$ for some $h \in \{1, 2\}$, then $|i - j|, |i - l|, |j - l| > 1$ and

$$\begin{aligned} \int_{\Omega_\varepsilon} U_i^2 U_j U_l dx &= \int_{\Omega_\varepsilon} \frac{\alpha_4^2 \delta_i^2}{(\delta_i^2 + |x|^2)^2} \frac{\alpha_4 \delta_j}{\delta_j^2 + |x|^2} \frac{\alpha_4 \delta_l}{\delta_l^2 + |x|^2} dx = \int_{\Omega_\varepsilon / \delta_i} \frac{\alpha_4^4 \delta_i^2 \delta_j \delta_l}{(1 + |y|^2)^2 (\delta_j^2 + \delta_i^2 |y|^2) (\delta_l^2 + \delta_i^2 |y|^2)} dy \\ &= \begin{cases} O\left(\frac{\delta_i \delta_l}{\delta_j^2} |\log \delta_i|\right) & \text{if } i < j < l \\ O\left(\frac{\delta_l}{\delta_j}\right) & \text{if } j < i < l \\ O\left(\frac{\delta_i^2}{\delta_j \delta_l} |\log \delta_i|\right) & \text{if } j < l < i \end{cases} \\ &= o(\delta_1^2) \end{aligned}$$

(note that this term only appears in a_2). Finally, if all the indices are different, then assuming without loss of generality that $i > j > h > l$,

$$\begin{aligned} \int_{\Omega_\varepsilon} U_i U_j U_h U_l dx &= \int_{\Omega_\varepsilon} \frac{\alpha_4 \delta_i}{\delta_i^2 + |x|^2} \frac{\alpha_4 \delta_j}{\delta_j^2 + |x|^2} \frac{\alpha_4 \delta_h}{\delta_h^2 + |x|^2} \frac{\alpha_4 \delta_l}{\delta_l^2 + |x|^2} dx \\ &\leq \int_{\Omega_\varepsilon / \delta_i} \frac{\alpha_4^4 \delta_i^3 \delta_j \delta_h \delta_l}{(1 + |y|^2) \delta_h^2 \delta_l^2 |y|^2} dy = O\left(\delta_1^2 \frac{\delta_i}{\delta_h} \frac{\delta_j}{\delta_l} |\log \delta_i|\right) = o(\delta_1^2). \end{aligned}$$

Estimates for a_3 . Arguing as in the proof of Lemma 7.2 in [12] (see equation (7.6) therein), the term a_3 is quadratic in ϕ_1 and ϕ_2 , and so by Proposition 3.1 it satisfies $a_2 = o(\delta_1^2)$.

Estimates for a_4 . The first term in a_4 can be estimated as

$$\begin{aligned} \left| \int_{\Omega_\varepsilon} \left(\sum_{j \in I_i} f(U_j) - f\left(\sum_{j \in I_i} P_\varepsilon U_j\right) \right) \phi_i dx \right| &\leq \left\| \sum_{j \in I_i} f(U_j) - f\left(\sum_{j \in I_i} P_\varepsilon U_j\right) \right\|_{L^{\frac{4}{3}}(\Omega_\varepsilon)} \|\phi_i\|_{L^4(\Omega_\varepsilon)} \\ &= O(\delta_1^2) \|\phi_i\|_{H_0^1(\Omega_\varepsilon)} = o(\delta_1^2), \end{aligned}$$

because, by (3.40),

$$\left\| \sum_{j \in I_i} f(U_j) - f\left(\sum_{j \in I_i} P_\varepsilon U_j\right) \right\|_{L^{\frac{4}{3}}(\Omega_\varepsilon)} = O(\delta_1^2).$$

(this term corresponds to the quantity A defined in the proof of Proposition 3.6). As for the second term in a_4 , given $i, j \in \{1, 2\}$ with $i \neq j$ and $h, l \in I_i$ with $h < l$, $m \in I_j$, by Lemma A.6 we have

$$\begin{aligned} \left| \int_{\Omega_\varepsilon} P_\varepsilon U_h P_\varepsilon U_l P_\varepsilon U_m \phi_j \right| &\leq \|P_\varepsilon U_h P_\varepsilon U_l P_\varepsilon U_m\|_{L^{4/3}} \|\phi_j\|_{L^4} \leq \|U_h U_l U_m\|_{L^{4/3}} \|\phi_j\|_{L^4} \\ &= \begin{cases} O\left(\frac{\delta_l}{\delta_h}\right) \|\phi_j\|_{H_0^1} & \text{if } h < l < m \\ O\left(\frac{\delta_m}{\delta_k}\right) \|\phi_j\|_{H_0^1} & \text{if } h < m < l \\ O\left(\frac{\delta_h}{\delta_m}\right) \|\phi_j\|_{H_0^1} & \text{if } m < h < l \end{cases} \\ &= o(\delta_1^2) \end{aligned}$$

since, for instance when $h < l < m$,

$$O\left(\frac{\delta_l}{\delta_h}\right) \|\phi_j\|_{H_0^1} \delta_1^{-2} = O(1) \varepsilon^{l-h-1} \left(\ln \frac{1}{\varepsilon}\right)^{-\frac{k+l-h}{k+1}} \rightarrow 0. \quad (4.30)$$

Estimates for a_5 . Starting from the first term, by Proposition 3.1 and Lemma A.5 we see that, given $i, j \in \{1, 2\}$ with $i \neq j$, and $h, l \in I_i$ with $h < l$,

$$\left| \int_{\Omega_\varepsilon} P_\varepsilon U_h P_\varepsilon U_l \phi_j^2 \right| \leq C \|U_h U_l\|_{L^2} \|\phi_j\|_{H_0^1}^2 = o\left(\frac{\delta_l \delta_1^2}{\delta_h}\right) = o(\delta_1^2).$$

Similarly, the second term in a_5 is also an $o(\delta_1^2)$.

Estimates for a_6 . We have, by Proposition 3.1 and Lemma (A.3),

$$\left| \int_{\Omega_\varepsilon} P_\varepsilon U_h \phi_i \phi_j^2 \right| \leq C \|U_h\|_{L^4} \|\phi\|_{H_0^1}^3 = o(\delta_1^2)$$

while the second and third terms in a_6 are respectively of second and fourth order in ϕ , thus an $o(\delta_1^2)$. This ends the proof. \square

As a direct consequence of (4.7), Corollary 4.3, Corollary 4.5 and Lemma 4.6, we have the following result, which gives us the leading term of the expansion of the reduced energy.

Proposition 4.7. *We have*

$$\begin{aligned} \tilde{J}_\varepsilon(\mathbf{d}) &= k \frac{B}{4} + \left(\frac{A^2}{2} \tau(0) d_1^2 + \frac{\Gamma}{2} \left(\frac{1}{d_k}\right)^2 - \frac{\beta}{2} \frac{\alpha_4^4}{k+1} |\mathbb{S}^3| \sum_{i=1}^{k-1} \left(\frac{d_{i+1}}{d_i}\right)^2 \right) \varepsilon^{\frac{2}{k+1}} \left(\log \frac{1}{\varepsilon}\right)^{\frac{k-1}{k+1}} \\ &\quad + o\left(\varepsilon^{\frac{2}{k+1}} \left(\log \frac{1}{\varepsilon}\right)^{\frac{k-1}{k+1}}\right) \end{aligned} \quad (4.31)$$

as $\varepsilon \rightarrow 0$, uniformly in $\mathbf{d} \in X_\eta$.

Remark 4.8. Suppose that, instead of dealing with the set of odd and even numbers of $\{1, \dots, k\}$ in the two equation case, we are dealing with system (1.4) with m equations and with a general

partition I_1, \dots, I_m satisfying (1)–(5). Then the reduced energy reads as

$$\begin{aligned} \tilde{J}_\varepsilon(\mathbf{d}) &= J_\varepsilon \left(\sum_{j \in I_1} P_\varepsilon U_{\delta_j} + \phi_1^{\mathbf{d}, \varepsilon}, \dots, \sum_{j \in I_m} P_\varepsilon U_{\delta_j} + \phi_m^{\mathbf{d}, \varepsilon} \right) \\ &= \sum_{i=1}^k \int_{\Omega_\varepsilon} \left(\frac{1}{2} |\nabla P_\varepsilon U_i|^2 - \frac{1}{4} (P_\varepsilon U_i)^4 \right) - \frac{\beta}{2} \sum_{\substack{h_1, h_2=1 \\ h_1 \neq h_2}}^k \sum_{\substack{i \in I_{h_1} \\ j \in I_{h_2}}} \int_{\Omega_\varepsilon} (P_\varepsilon U_i)^2 (P_\varepsilon U_j)^2 + R(\mathbf{d}, \varepsilon) \end{aligned}$$

with

$$R(\mathbf{d}, \varepsilon) = J_\varepsilon \left(\sum_{j \in I_1} P_\varepsilon U_{\delta_j} + \phi_1^{\mathbf{d}, \varepsilon}, \dots, \sum_{j \in I_m} P_\varepsilon U_{\delta_j} + \phi_m^{\mathbf{d}, \varepsilon} \right) - \sum_{\substack{h_1, h_2=1 \\ h_1 \neq h_2}}^k \sum_{\substack{i \in I_{h_1} \\ j \in I_{h_2}}} J_\varepsilon (P_\varepsilon U_i, P_\varepsilon U_j)$$

With an analogous proof of the one of Lemma 4.1, we can show that critical points of this functional correspond to solutions of (1.4). The choice of rates is still (2.5) in the general case. As pointed out in the proofs of Corollary 4.5 and Lemma 4.6, besides the exact shape of the rates, the other crucial step is that each set I_h does not contain two consecutive integers. Since this property is valid for a general partition (it corresponds to property (5)), it is straightforward to adapt the proofs of these results and show that the quantity

$$\sum_{\substack{h_1, h_2=1 \\ h_1 \neq h_2}}^k \sum_{\substack{i \in I_{h_1} \\ j \in I_{h_2}}} \int_{\Omega_\varepsilon} (P_\varepsilon U_i)^2 (P_\varepsilon U_j)^2$$

has the asymptotic expansion (4.26), and that R satisfies (4.29). Combining this with Corollary 4.3 yields that, in the general case, the reduced expression has the exact same expansion, namely (4.31).

5. PROOF OF THE MAIN RESULT

In this section we conclude the proof of Theorem 1.3. Define $\Psi : (\mathbb{R}^+)^k \rightarrow \mathbb{R}$ as

$$\Psi(x_1, \dots, x_k) = a_1 x_1 + \frac{a_2}{x_k} + a_3 \sum_{i=1}^{k-1} \frac{x_{i+1}}{x_i}$$

where, since $\beta < 0$,

$$a_1 = \frac{A^2}{2} \tau(0) > 0, \quad a_2 = \frac{\Gamma}{2} > 0, \quad a_3 = -\frac{\beta}{2} \frac{\alpha_4^4}{k+1} |\mathbb{S}^3| > 0.$$

Lemma 5.1. *The function Ψ achieves a unique global minimum at (x_1^*, \dots, x_k^*) , with*

$$x_i^* := \left(\frac{a_2}{a_3} \right)^{\frac{i}{k+1}} \left(\frac{a_3}{a_1} \right)^{\frac{k+1-i}{k+1}} = \Gamma^{\frac{i}{k+1}} (A^2 \tau(0))^{\frac{i-1-k}{k+1}} \left(\frac{|\beta| \alpha_4^4 |\mathbb{S}^3|}{k+1} \right)^{\frac{k+1-2i}{k+1}} > 0$$

for $i = 1, \dots, k$. In particular, the conclusion of Theorem 1.3 holds true.

Proof. First of all, observe that

$$\Psi(\mathbf{x}) \rightarrow \infty \quad \text{as } |\mathbf{x}| \rightarrow \infty,$$

since $a_1, a_2, a_3 > 0$ and $a_1 x_1 \rightarrow \infty$ if $x_1 \rightarrow \infty$, $x_{i+1}/x_i \rightarrow \infty$ if $x_{i+1} \rightarrow \infty$ and x_i is bounded. Moreover, given $\bar{\mathbf{x}} \in (\mathbb{R}_0^+)^k$ with $\bar{x}_i = 0$,

$$\Psi(\mathbf{x}) \rightarrow \infty \quad \text{as } \mathbf{x} \rightarrow \bar{\mathbf{x}},$$

since in this case at least one of the $k+1$ terms in the expression of Ψ diverges to ∞ . In conclusion, Ψ admits a global minimum. Let us see next that it is unique, and deduce its expression.

We have

$$\frac{\partial \Psi}{\partial x_1} = a_1 - a_3 \frac{x_2}{x_1^2}, \quad \frac{\partial \Psi}{\partial x_k} = -\frac{a_2}{x_k^2} + \frac{a_3}{x_{k-1}}$$

and

$$\frac{\partial \Psi}{\partial x_j} = a_3 \left(\frac{1}{x_{j-1}} - \frac{x_{j+1}}{x_j^2} \right). \quad (j = 2, \dots, k-1)$$

Hence, at a critical point,

$$x_1^2 a_1 = a_3 x_2, \quad x_j^2 = x_{j-1} x_{j+1} \quad (j = 2, \dots, k-1), \quad a_3 x_k^2 = a_2 x_{k-1}$$

which yields, by direct substitution,

$$x_j = \left(\frac{a_1}{a_3} \right)^{j-1} x_1^j \quad (j = 2, \dots, k), \quad a_3 \left(\frac{a_1}{a_3} \right)^{2k-2} x_1^{2k} = a_2 \left(\frac{a_1}{a_3} \right)^{k-2} x_1^{k-1}.$$

The last identity gives

$$x_1 = \left(\frac{a_2}{a_3} \right)^{\frac{1}{k+1}} \left(\frac{a_3}{a_1} \right)^{\frac{k}{k+1}},$$

and the rest of the proof follows. \square

End of the proof of Theorem 1.3. From the definition of Ψ and by Proposition 4.7, we have

$$J(\mathbf{d}) = k \frac{B}{4} + \varepsilon^{\frac{2}{k+1}} \left(\log \frac{1}{\varepsilon} \right)^{\frac{k-1}{k+1}} (\Psi(d_1^2, \dots, d_k^2) + o(1))$$

where $o(1) \rightarrow 0$ as $\varepsilon \rightarrow 0$, uniformly in $\mathbf{d} \in X_\eta$. Let $d_i^* := \sqrt{x_i^*}$ (cf. Lemma 5.1), and take $\eta > 0$ small enough so that $(d_1^*, \dots, d_k^*) \in X_\eta$. Let $K \Subset X_\eta$ be a compact set such that $((d_1^*)^2, \dots, (d_k^*)^2) \in \text{int} K$ and

$$\Psi((d_1^*)^2, \dots, (d_k^*)^2) = \min_K \Psi < \min_{\partial K} \Psi.$$

Then

$$\min_K J_\varepsilon \leq J_\varepsilon(\mathbf{d}^*) < \min_{\partial K} J_\varepsilon$$

Therefore $J_\varepsilon|_K$ has a minimizer \mathbf{d}_ε , which converges to \mathbf{d}^* (by the uniqueness stated in Lemma 5.1). Thus $J'_\varepsilon(\mathbf{d}^*) = 0$. By invoking Lemma 4.1, the proof is finished. \square

Remark 5.2. The proof of the general case, Theorem 1.1, follows exactly in the same way since, as we commented on Remark 4.8, the reduced functional is the same as in the two equation case.

APPENDIX A. ASYMPTOTIC ESTIMATES

In this appendix we collect several important asymptotic estimates which are used in the paper. We assume in every statement that $N = 4$, that is, $\Omega \subset \mathbb{R}^4$.

The following two results are taken from [12], see Lemmas 3.1 and 3.2 therein.

Lemma A.1. *Let $a \in \Omega$, $r > 0$, and $\tau \in \mathbb{R}^4$. Assume that $\xi = a + \delta\tau$, with $\delta = \delta(\varepsilon) \rightarrow 0$ and $\varepsilon/\delta \rightarrow 0$ as $\varepsilon \rightarrow 0^+$. Then, for $A = \int_{\mathbb{R}^N} U_{1,0}^3 = \frac{\alpha_4}{\gamma_4}$ and R defined by*

$$P_\varepsilon U_{\delta,\xi} = U_{\delta,\xi} - A\delta H(x, \xi) - \frac{\alpha_4}{\delta(1 + |\tau|^2)} \left(\frac{r\varepsilon}{|x - a|} \right)^2 + R(x)$$

there exists $C = C(\tau, \text{dist}(a, \partial\Omega)) > 0$ such that, for any $x \in \Omega \setminus B_{r\varepsilon}(a)$,

$$\begin{aligned} |R(x)| &\leq C\delta \left[\frac{\varepsilon^2(1 + \varepsilon\delta^{-3})}{|x - a|^2} + \delta^2 + \left(\frac{\varepsilon}{\delta}\right)^2 \right] \\ |\partial_\delta R(x)| &\leq C \left[\frac{\varepsilon^2(1 + \varepsilon\delta^{-3})}{|x - a|^2} + \delta^2 + \left(\frac{\varepsilon}{\delta}\right)^2 \right] \\ |\partial_{\tau_i} R(x)| &\leq C\delta^2 \left[\frac{\varepsilon^2(1 + \varepsilon\delta^{-4})}{|x - a|^2} + \delta^2 + \frac{\varepsilon^2}{\delta^3} \right] \end{aligned}$$

Lemma A.2. *Under the assumptions and notations of the previous lemma, we have the following estimate:*

$$\int_{\Omega_\varepsilon} U_{\delta,\xi}^2 (P_\varepsilon U_{\delta,\xi} - U_{\delta,\xi})^2 = O\left(\delta^4 |\log \delta| + \left(\frac{\varepsilon}{\delta}\right)^4 \left|\log \frac{\varepsilon}{\delta}\right|\right).$$

The following concerns the asymptotic study of L^q norms of the bubble. For the proof, see for instance [20, Lemma A.3] or [21, Lemma A.2].

Lemma A.3. *We have, as $\delta \rightarrow 0$,*

$$\int_{\Omega} U_{\delta}^q = \begin{cases} O(\delta^q) & \text{if } 0 < q < 2, \\ O(\delta^2 |\log \delta|) & \text{if } q = 2, \\ O(\delta^{4-q}) & \text{if } 2 < q < \infty, q \neq 4, \\ O(1) & \text{if } q = 4. \end{cases}$$

The following lemmas will be used many times in order to estimate interaction integrals.

Lemma A.4. *Let $1 < q < 2 < p$ be such that $p + q = 4$. Let $\rho_1 = \rho_1(\varepsilon) > 0$, $\rho_2 = \rho_2(\varepsilon) > 0$, be such that*

$$\frac{\rho_2}{\rho_1} \rightarrow 0, \quad \frac{\varepsilon}{\rho_1} \rightarrow 0, \quad \frac{\varepsilon}{\rho_2} \rightarrow 0$$

as $\varepsilon \rightarrow 0^+$. Then

$$\int_{\Omega_\varepsilon} U_{\rho_1}^p U_{\rho_2}^q = O\left(\left(\frac{\rho_2}{\rho_1}\right)^q\right), \quad \text{and} \quad \int_{\Omega_\varepsilon} U_{\rho_2}^p U_{\rho_1}^q = O\left(\left(\frac{\rho_2}{\rho_1}\right)^q\right).$$

Proof. We proceed by direct computations:

$$\begin{aligned}
\int_{\Omega_\varepsilon} U_{\rho_1}^p U_{\rho_2}^q &= \alpha_4^4 \int_{\Omega_\varepsilon} \left(\frac{\rho_1}{\rho_1^2 + |x|^2} \right)^p \left(\frac{\rho_2}{\rho_2^2 + |x|^2} \right)^q dx \\
&= \alpha_4^4 \rho_1^{4-p} \int_{\Omega_\varepsilon/\rho_1} \left(\frac{1}{1 + |y|^2} \right)^p \left(\frac{\rho_2}{\rho_2^2 + \rho_1^2 |y|^2} \right)^q dy \\
&\leq \alpha_4^4 \left(\frac{\rho_2}{\rho_1} \right)^q \int_{\Omega_\varepsilon/\rho_1} \left(\frac{1}{1 + |y|^2} \right)^p \frac{1}{|y|^{2q}} dy \\
&\leq \alpha_4^4 \left(\frac{\rho_2}{\rho_1} \right)^q \int_{\mathbb{R}^4} \left(\frac{1}{1 + |y|^2} \right)^p \frac{1}{|y|^{2q}} dy = O\left(\left(\frac{\rho_2}{\rho_1} \right)^q \right),
\end{aligned}$$

as desired. Analogously

$$\begin{aligned}
\int_{\Omega_\varepsilon} U_{\rho_2}^p U_{\rho_1}^q &= \alpha_4^4 \int_{\Omega_\varepsilon} \left(\frac{\rho_2}{\rho_2^2 + |x|^2} \right)^p \left(\frac{\rho_1}{\rho_1^2 + |x|^2} \right)^q dx \\
&= \alpha_4^4 \rho_2^{4-p} \int_{\Omega_\varepsilon/\rho_2} \left(\frac{1}{1 + |y|^2} \right)^p \left(\frac{\rho_1}{\rho_1^2 + \rho_2^2 |y|^2} \right)^q dy \\
&\leq \alpha_4^4 \left(\frac{\rho_2}{\rho_1} \right)^q \int_{\Omega_\varepsilon/\rho_2} \left(\frac{1}{1 + |y|^2} \right)^p dy \\
&\leq \alpha_4^4 \left(\frac{\rho_2}{\rho_1} \right)^q \int_{\mathbb{R}^4} \left(\frac{1}{1 + |y|^2} \right)^p dy = O\left(\left(\frac{\rho_2}{\rho_1} \right)^q \right). \quad \square
\end{aligned}$$

Lemma A.5. *Let $\rho_1 = \rho_1(\varepsilon) > 0$, $\rho_2 = \rho_2(\varepsilon) > 0$, be such that*

$$\frac{\rho_2}{\rho_1} \rightarrow 0, \quad \frac{\varepsilon}{\rho_1} \rightarrow 0, \quad \frac{\varepsilon}{\rho_2} \rightarrow 0$$

as $\varepsilon \rightarrow 0^+$. Then

$$\int_{\Omega_\varepsilon} U_{\rho_1}^2 U_{\rho_2}^2 = O\left(\frac{\rho_2}{\rho_1} |\log \rho_2| \right).$$

Proof. We have

$$\begin{aligned}
\int_{\Omega_\varepsilon} U_{\rho_1}^2 U_{\rho_2}^2 &= \alpha_4^4 \int_{\Omega_\varepsilon} \left(\frac{\rho_1}{\rho_1^2 + |x|^2} \right)^2 \left(\frac{\rho_2}{\rho_2^2 + |x|^2} \right)^2 dx \\
&= \alpha_4^4 \rho_1^2 \rho_2^2 \int_{\Omega_\varepsilon/\rho_2} \left(\frac{1}{\rho_1^2 + \rho_2^2 |y|^2} \right)^2 \left(\frac{1}{1 + |y|^2} \right)^2 dx \\
&\leq \alpha_4^4 \left(\frac{\rho_2}{\rho_1} \right)^2 \int_{\Omega_\varepsilon/\rho_2} \frac{dy}{(1 + |y|^2)^2}.
\end{aligned}$$

Since Ω is bounded, there exists a sufficiently large radius $R > 0$ such that

$$\int_{\Omega_\varepsilon/\rho_2} \frac{dy}{(1 + |y|^2)^2} \leq \int_{B_{R/\rho_2}} \frac{dy}{(1 + |y|^2)^2} \leq C + C \int_1^{R/\rho_2} r^{-1} dr = C + C \log \frac{R}{\rho_2}.$$

To sum up

$$\int_{\Omega_\varepsilon} U_{\rho_1}^2 U_{\rho_2}^2 \leq \alpha_4^4 \left(\frac{\rho_2}{\rho_1} \right)^2 \left(C + C \log \frac{R}{\rho_2} \right),$$

and the thesis follows. \square

Lemma A.6. *Let $\rho_1 = \rho_1(\varepsilon) > 0$, $\rho_2 = \rho_2(\varepsilon) > 0$, $\rho_3 = \rho_3(\varepsilon) > 0$ be such that*

$$\frac{\rho_j}{\rho_i} \rightarrow 0 \quad \text{if } 1 \leq i < j \leq 3, \quad \frac{\varepsilon}{\rho_h} \rightarrow 0 \quad \text{for every } h,$$

as $\varepsilon \rightarrow 0^+$. Then

$$\int_{\Omega_\varepsilon} (U_{\rho_1} U_{\rho_2} U_{\rho_3})^{\frac{4}{3}} = O\left(\left(\frac{\rho_2}{\rho_1}\right)^{\frac{4}{3}}\right)$$

as $\varepsilon \rightarrow 0^+$.

Proof. Again, by direct computations

$$\begin{aligned} \int_{\Omega_\varepsilon} (U_{\rho_1} U_{\rho_2} U_{\rho_3})^{\frac{4}{3}} &= \alpha_4^4 \int_{\Omega_\varepsilon} \left(\frac{\rho_1 \rho_2 \rho_3}{(\rho_1^2 + |x|^2)^2 (\rho_2^2 + |x|^2)^2 (\rho_3^2 + |x|^2)} \right)^{\frac{4}{3}} dx \\ &= \alpha_4^4 (\rho_3^2 \rho_1 \rho_2)^{\frac{4}{3}} \int_{\Omega_\varepsilon / \rho_3} \frac{dy}{(1 + |y|^2)^{\frac{4}{3}} (\rho_1^2 + \rho_3^2 |y|^2)^{\frac{4}{3}} (\rho_2^2 + \rho_3^2 |y|^2)^{\frac{4}{3}}} \\ &\leq \alpha_4^4 \left(\frac{\rho_2}{\rho_1}\right)^{\frac{4}{3}} \int_{\mathbb{R}^4} \frac{dy}{(1 + |y|^2)^{\frac{4}{3}} |y|^{\frac{8}{3}}}. \quad \square \end{aligned}$$

REFERENCES

- [1] Thierry Aubin. Problèmes isopérimétriques et espaces de Sobolev. *J. Differential Geometry*, 11(4):573–598, 1976.
- [2] A. Bahri and J.-M. Coron. On a nonlinear elliptic equation involving the critical Sobolev exponent: the effect of the topology of the domain. *Comm. Pure Appl. Math.*, 41(3):253–294, 1988.
- [3] Gabriele Bianchi and Henrik Egnell. A note on the Sobolev inequality. *J. Funct. Anal.*, 100(1):18–24, 1991.
- [4] Luis A. Caffarelli, Basilis Gidas, and Joel Spruck. Asymptotic symmetry and local behavior of semilinear elliptic equations with critical sobolev growth. *Communications on Pure and Applied Mathematics*, 42(3):271–297, 1989.
- [5] Zhijie Chen and Chang-Shou Lin. Asymptotic behavior of least energy solutions for a critical elliptic system. *International Mathematics Research Notices*, 21:11045–11082, 2015.
- [6] Zhijie Chen, Chang-Shou Lin, and Wenming Zou. Sign-changing solutions and phase separation for an elliptic system with critical exponent. *Comm. Partial Differential Equations*, 39(10):1827–1859, 2014.
- [7] Zhijie Chen and Wenming Zou. Positive least energy solutions and phase separation for coupled Schrödinger equations with critical exponent. *Arch. Ration. Mech. Anal.*, 205(2):515–551, 2012.
- [8] Zhijie Chen and Wenming Zou. Positive least energy solutions and phase separation for coupled Schrödinger equations with critical exponent: higher dimensional case. *Calculus of Variations and Partial Differential Equations*, 52(1):423–467, 2015.
- [9] Mónica Clapp and Jorge Faya. Multiple solutions to a weakly coupled purely critical elliptic system in bounded domains. arXiv: 1805.10304.
- [10] Mónica Clapp and Angela Pistoia. Existence and phase separation of entire solutions to a pure critical competitive elliptic system. *Calc. Var. Partial Differential Equations*, 57(1):Art. 23, 20, 2018.
- [11] Jean-Michel Coron. Topologie et cas limite des injections de Sobolev. *C. R. Acad. Sci. Paris Sér. I Math.*, 299(7):209–212, 1984.
- [12] Yuxin Ge, Monica Musso, and Angela Pistoia. Sign changing tower of bubbles for an elliptic problem at the critical exponent in pierced non-symmetric domains. *Communications in Partial Differential Equations*, 35(8):1419–1457, 2010.
- [13] Francesca Gladiali, Massimo Grossi, and Christophe Troestler. Entire radial and nonradial solutions for systems with critical growth. *Calc. Var. Partial Differential Equations*, 57(2):Art. 53, 26, 2018.
- [14] Francesca Gladiali, Massimo Grossi, and Christophe Troestler. A non-variational system involving the critical sobolev exponent. the radial case. arXiv: 1603.05641.
- [15] Yuxia Guo, Bo Li, and Juncheng Wei. Entire nonradial solutions for non-cooperative coupled elliptic system with critical exponents in \mathbb{R}^3 . *J. Differential Equations*, 256(10):3463–3495, 2014.
- [16] Roger Lewandowski. Little holes and convergence of solutions of $-\Delta u = u^{(N+2)/(N-2)}$. *Nonlinear Anal.*, 14(10):873–888, 1990.

- [17] Monica Musso and Angela Pistoia. Multispikes solutions for a nonlinear elliptic problem involving the critical Sobolev exponent. *Indiana Univ. Math. J.*, 51(3):541–579, 2002.
- [18] Monica Musso and Angela Pistoia. Sign changing solutions to a nonlinear elliptic problem involving the critical Sobolev exponent in pierced domains. *J. Math. Pures Appl. (9)*, 86(6):510–528, 2006.
- [19] Shuangjie Peng, Yan-fang Peng, and Zhi-Qiang Wang. On elliptic systems with Sobolev critical growth. *Calc. Var. Partial Differential Equations*, 55(6):Art. 142, 30, 2016.
- [20] Angela Pistoia and Nicola Soave. On coron’s problem for weakly coupled elliptic systems. *Proceedings of the London Mathematical Society*, 116(1):33–67, 2018.
- [21] Angela Pistoia and Hugo Tavares. Spiked solutions for Schrödinger systems with Sobolev critical exponent: the cases of competitive and weakly cooperative interactions. *J. Fixed Point Theory Appl.*, 19(1):407–446, 2017.
- [22] Olivier Rey. Sur un problème variationnel non compact: l’effet de petits trous dans le domaine. *C. R. Acad. Sci. Paris Sér. I Math.*, 308(12):349–352, 1989.
- [23] Richard Schoen. Topics in differential geometry. Notes from graduate lectures in Stanford University, <http://www.math.washington.edu/pollack/research/Schoen-1988-notes.html>, 1988.
- [24] Giorgio Talenti. Best constant in Sobolev inequality. *Ann. Mat. Pura Appl. (4)*, 110:353–372, 1976.

ANGELA PISTOIA

DIPARTIMENTO DI METODI E MODELLI MATEMATICI, UNIVERSITÀ DI ROMA “LA SAPIENZA”
VIA ANTONIO SCARPA 16, 00161 ROMA (ITALY)
E-mail address: `angela.pistoia@uniroma1.it`

NICOLA SOAVE

DIPARTIMENTO DI MATEMATICA, POLITECNICO DI MILANO,
VIA EDOARDO BONARDI 9, 20133 MILANO (ITALY)
E-mail address: `nicola.soave@gmail.com`; `nicola.soave@polimi.it`

HUGO TAVARES

CMAFCIO & DEPARTAMENTO DE MATEMÁTICA
FACULDADE DE CIÊNCIAS DA UNIVERSIDADE DE LISBOA
EDIFÍCIO C6, PISO 1, CAMPO GRANDE 1749-016 LISBOA (PORTUGAL)
E-mail address: `hrtavares@ciencias.ulisboa.pt`