



UNIVERSITÀ DEGLI STUDI DI TORINO

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

## Bunch placement effects on dehydration kinetics and physico-chemical composition of Nebbiolo grapes

This is the author's manuscript		
Original Citation:		
Availability:		
This version is available http://hdl.handle.net/2318/1868858	since	2022-07-09T09:00:54Z
Publisher:		
International Viticulture and Enology Society		
Terms of use:		
Open Access		
Anyone can freely access the full text of works made available as under a Creative Commons license can be used according to the t of all other works requires consent of the right holder (author or p protection by the applicable law.	"Open A erms an ublisher	ccess". Works made available d conditions of said license. Use ) if not exempted from copyright

(Article begins on next page)

# Bunch placement effects on dehydration kinetics and physico-chemical composition of Nebbiolo grapes

Sforzato di Valtellina DOCG is a special reinforced red wine produced using withered Nebbiolo grapes. The withering process takes place in traditional rooms under natural environmental conditions; it starts immediately after the harvest and ends not before the 1st December of the same year. The process can be performed with different bunch placements that can influence the grapes features. The purpose of the study is to compare the effect on grape physico-chemical parameters for four withering bunch placement systems: hanged clusters (HC), plastic crates (CT), breathable mesh fabric on wooden frames panels (MF), and reed mats (RM). For all the systems studied, the withering length was two months at a temperature between 6 and 19 °C and a relative humidity of 41-88%. The grapes were sampled at the beginning, at half time, and at the end of withering. For each sampling point, weight loss rate, skins mechanical properties (i.e. hardness and thickness), must technological parameters, and skins and seeds extractable polyphenols, flavonoids, anthocyanins, and condensed tannins were studied. At the end of withering, the berry weight loss resulted very different among the systems, ranging from 18.79 to 12.73%. HC showed the fastest weight loss, followed by MF, CT, and RM. Interestingly, the dehydration kinetics showed different trends over the process: for HC the rate of weight loss (WLR, %/day) resulted higher in the first half of the process and then decreased; on the contrary, the kinetics of CT and MF were slower at the beginning of withering compared with the second phase; for RM, instead, the WLR remained fairly constant throughout the entire period. These differences yielded different consequences on the complex balance between concentration-synthesis and loss of compounds during withering. HC led to a significantly higher sugar content than the others after two months of withering. Instead, no significant differences were found among the systems for total acidity, pH, acetic acid, and glycerol (markers of microbial development), and mechanical properties. As regards phenolics, RM led to a significant reduction in skin extractable polyphenols and flavonoids when expressed as mg/kg berries, possibly because the lower concentration effect did not exceed the greater loss of these compounds compared to the other placements. No significant differences were found among systems in seeds polyphenols. In conclusion, under the same environmental conditions bunch placement influenced weight loss, dehydration kinetics, skins polyphenols, and to a lesser extent also the sugar content. The best compromise between weight loss and grape features seems to be the use of plastic crates, whereas hanged clusters placement allowed to achieve the same weight loss faster, although resulting in a higher sugar content. Therefore, these results can provide knowledge to choose the withering system with awareness according to the established oenological objective.

**Authors**: Scalzini Giulia<sup>1</sup>, Giacosa Simone<sup>1</sup>, Paissoni Maria Alessandra<sup>1</sup>, Río Segade Susana<sup>1</sup>, Rolle Luca<sup>1</sup> and Gerbi Vincenzo<sup>1</sup>

<sup>1</sup>University of Turin, Department of Agricultural, Forest and Food Sciences

#### \*corresponding author: giulia.scalzini@unito.it (mailto:giulia.scalzini@unito.it)

Keywords: bunch placement, grape dehydration, weight loss rate, physico-chemical parameters, special wines





(https://ives-openscience.eu/ives-conferenceseries/)

TAGS: (IVAS 2022 (Https://ives-openscience.eu/tag/ivas-2022/)

IVES Conference Series (Https://ives-openscience.eu/tag/ives-conference-series/)

< PREVIOUS POST (HTTPS://IVES-OPENSCIENCE.EU/14338/)

NEXT POST (HTTPS://IVES-OPENSCIENCE.EU/14345/) >

Related Posts			
(https://ives-openscience. /13067/)	eu 🛱 31 May 2022 Influence of weather and climatic conditions on the viticultural production in Croatia (https://ives-openscience.eu/13067/)		
(https://ives- openscience.eu/5169/)	口 31 July 2020 Spatio-temporal analysis of grapevine water behaviour in hillslope vineyards. the example of corton hill, Burgundy (https://ives-openscience.eu/5169/)		
(https://ives- openscience.eu/13192/)	31 May 2022 Making sense of available information for climate change adaptation and building resilience into wine production systems across the world (https://ives-openscience.eu/13192/)		
(https://ives-openscience. /14432/)	eu 🗂 24 June 2022 Cell Walls Of Grape Mesocarp Possible Fining Agents For Red And White Wine (https://ives- openscience.eu/14432/)		
(https://ives-openscience. /10915/)	EU To February 2022 Study and valorization of vineyards "terroirs" in the Val de Loire (https://ives- openscience.eu/10915/)		
(https://ives- openscience.eu/6741/)	23 March 2021 Prospects of thermal imaging as a non-invasive tool to assess water status for irrigation scheduling in commercial vineyards (https://ives-openscience.eu/6741/)		

Search

#### Social links

- (https://www.linkedin.com/company/international-viticulture-and-enology-society-ives/)

#### Subscribe our newsletter

\* your@email.com Subscribe

### **Contact information**

Email: jdumercq[@]ives-openscience.eu Siret 839 985 843 00011 / RNA W332022472

Copyright© 2018 - IVES open science (https://ives-openscience.eu/)