

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

## The context of available options affect healthcare decisions: A generalization study

### This is the author's manuscript

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/104145> since 2016-04-01T14:25:08Z

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

# The Context of Available Options Affects Health Care Decisions: A Generalization Study

Florence Dumas, PhD, Michel Gonzalez, PhD, Vittorio Girotto, PhD, Christophe Pascal, PhD, Jean-François Botton, MD, Vincenzo Crupi, PhD

**Background.** When a given option is presented along with 2 alternatives, similar to each other, health care professionals choose it more often than when it is presented with just one of the alternatives. This inconsistent decision pattern may depend on the conflict generated from choosing between 2 highly similar options. **Objective.** To generalize the effect by using realistic scenarios that involve 2 alternatives displaying various degrees of similarity. **Methods.** One hundred fifty-five psychiatrists, 149 gynecologists, and 89 nurse managers had to indicate the treatment they would recommend in clinical scenarios containing either 3 options or just 2 of them. The similarity between the 2 alternatives varied across scenarios, ranging from a very high (psychiatric scenario) to an only moderately high (nursing management scenario) to a limited level (gynecological

scenario). **Results.** Professionals chose the focal option more often when both alternatives were available. The paradoxical effect occurred for all scenarios—namely, when the alternatives were medication variants (psychiatric scenario), when most of the features they shared produced their effect at a different extent in the 2 cases (nursing management scenario), and some of their consequences were at variance (gynecological problem). **Conclusions.** The context of available options affects professionals' choices when the alternatives are similar but also when they present diverging features. Professionals need to be aware of such a source of practice variability and are encouraged to consider each option per se before they compare the available options. **Keywords:** clinical decision making; context effect; similarity (*Med Decis Making* 2012;32:815–819)

The regularity principle of rational decision making states that if an agent prefers A among

options A, B, and C (e.g., a patient prefers to consult doctor A among 3 physicians working in town) and is informed that option C is not available anymore (e.g., doctor C left the town), he or she should continue to prefer A (e.g., doctor A). In other words, the patient's preferences should be independent from the absence or presence of an option that he or she would not pursue. Despite the soundness of this principle, individuals, including health care professionals, sometimes violate it.<sup>1–9</sup> In particular, Redelmeier and Shafir<sup>7</sup> asked a group of physicians to consider a patient who was treated with anti-inflammatory medications without success. In one

Received 2 August 2011 from LPC UMR CNRS 7290, Aix-Marseille University, Marseille, France (FD, MG, VG); University IUAV of Venice, Italy (VG); Jean Moulin Lyon 3 University, IFROSS, France (CP, FD); Department of Medical Information, Medical Center of Villefranche, France (J-FB); and University of Turin, Italy, and Ludwig Maximilian University, Munich, Germany (VC). The research was supported by grants from the Italian Ministry of Scientific Research (PRIN grant 20083NAH2L\_001) and by the Alexander von Humboldt Foundation to Vincenzo Crupi and from Swiss Global-Fondazione Ca' Foscari to Vittorio Girotto. The funding agreement ensured the authors' independence in designing the study, interpreting the data, and writing and publishing the report. The authors thank Don Redelmeier and 2 anonymous reviewers for their comments on a previous version of the paper, as well as 2 groups of nurse managers enrolled in continuing nursing education at Lyon III University (France) for their support in our study. They also thank a group of 33 nurse managers for their help in the construction of the material and another group of 38 nurse managers for their assistance in collecting data. Revision accepted for publication 28 January 2012.

DOI: 10.1177/0272989X12445285

Supplementary material for this article is available on the *Medical Decision Making* Web site at <http://mdm.sagepub.com/supplemental>.

Address correspondence to Florence Dumas, PhD, Université d'Aix-Marseille, Fédération de Recherche 3 C, LPC UMR CNRS 7290, Centre Saint Charles, Case D, 13331 Marseille Cedex 3, France; telephone: 133 413 550 978; fax: 133 478 587 644; e-mail: [florence.dumas@univ-amu.fr](mailto:florence.dumas@univ-amu.fr).