



UNIVERSITÀ DEGLI STUDI DI TORINO

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Application of a real-time gPCR method to measure the methanogen concentration during anaerobic digestion as an indicator of biogas production capacity.

This is the author's manuscript

Original Citation:

Availability:

This version is available http://hdl.handle.net/2318/118197 since 2015-09-07T14:51:37Z

Published version:

DOI:10.1016/j.jenvman.2012.07.021

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This is an author version of the contribution published on: Questa è la versione dell'autore dell'opera: Journal of Environmental Management 111 (2012) 173e177

10.1016/j.jenvman.2012.07.021

The definitive version is available at: La versione definitiva è disponibile alla URL: <u>http://www.sciencedirect.com/science/article/pii/S0301479712003854</u>

| 1 TITLE PAGE | : |
|--------------|---|
|--------------|---|

- 2 Application of a Real-time qPCR method to evaluate methanogen load as a high-efficiency process bio-
- 3 indicator for wet anaerobic co-digestion of organic wastes after different pre-treatments.

4

5 Author names:

- 6 D. Traversi¹, S.Villa¹, E. Lorenzi², R. Degan¹, G. Gilli¹
- 7 ¹ Department of Public Health and Microbiology,
- 8 University of the Study of Turin, via Santena 5 bis, 10126 Turin, Italy
- 9 phone +390116705810
- 10 fax +390112605810
- 11 e-mail: <u>deborah.traversi@unito.it</u>
- 12 ² SMAT S.p.A., corso XI Febbraio 14, 10152 Torino, Italy
- 13 ***Corresponding author:** e-mail: <u>deborah.traversi@unito.it</u>
- 14
- 15 **KEY WORDS**:
- 16 methanogen, anaerobic digestion, biogas production, renewable energy, Archea communities

- 18
- 19

20 ABSTRACT:

21 Biogas is a biofuel with a large energy value and primarily consists of methane. It is a renewable energy 22 source and produced via anaerobic digestion of various organic materials, including waste water sludge 23 and organic urban wastes. In reactors, anaerobic microorganisms can degrade the waste organic matter 24 into two different products: digestate and biogas. If the microbial community is optimised, then the 25 methane production rate could be improved, and the digestate could be used as a soil fertiliser. 26 Methanogen is the major microbiologic group responsible for methane production. We detected the 27 methanogen load during two wet digestion processes fed with pre-treated urban organic waste and 28 waste water sludge. Two different pre-treatments were involved in the experimental digestions: 29 pressure-extrusion and turbo-mixing. The applied methodology is based on Real Time quantitative PCR 30 (RT-qPCR) using an mcrA target. First, we evaluated the validity of the analyses, including standard 31 calibration, specificity, accuracy and precision. Next, we applied this method to 50 digestate samples. A 32 positive and significant correlation between the biogas production rate and methanogen abundance was 33 observed (r=0.579, p<0.001). The difference in the methanogen load between the two digestions is 34 significant (F=41.190, p<0.01). Moreover, considering all the collected data and using 0.6 m³/Kg TVS as 35 the 'cut-off', representing an optimal production, a higher methanogen load mean was observed with 36 the optimal production rates (F= 7.053; p<0.05). This suggests a role for methanogen load as diagnostic 37 tool in digestion pre-treatment and optimisation.

38

40 **1. INTRODUCTION**

41 Two serious environmental and public health problems our society faces today include, first, reduction 42 and treatment of waste, especially in high-density areas, and, second, finding an answer to the energetic 43 request to limit use of conventional fuels (das Neves et al., 2009). In urban communities, these goals 44 have no clear resolution, though renewable energy sources are likely a key strategy (Balat and Balat, 45 2009). The anaerobic digestion process for organic waste combines the removal of organic pollutants, 46 the reduction of organic waste volume, and energy conservation via biogas production (Rozzi and 47 Remigi, 2004). Numerous organic wastes are subject to anaerobic digestion, such as wastewater sludge, 48 pre-treated organic household waste (Bouallagui et al., 2005; Schievano et al., 2009). Biogas production 49 is the consequence of a series of metabolic interactions among bacterial and Archeal micro-organisms 50 (Ward et al., 2008). A particular ecosystem is present in an anaerobic reactor where several groups of 51 microorganisms work interactively in the conversion of complex organic matter into biogas. The 52 following four stages compose the digestion process: hydrolysis, acidogenesis, acetogenesis and 53 methanogenesis (Lozano et al., 2009). In each step, various microorganisms are involved. During the first 54 stage a group of micro-organisms secrete enzymes which hydrolyse polymers to monomers to convert 55 particulate materials into dissolved materials (Whitman et al., 2006). They remove the small amounts of 56 O₂ present and create anaerobic conditions. Subsequently, the acidogenic phase includes the action of a 57 large and diverse group of fermentative bacteria, usually belonging to the Clostridia class and the 58 Bacteroidaceaea family. These bacteria hydrolyse and ferment the organic materials and they produce 59 organic acids, CO₂ and H₂. Next, in the third phase, the acetogenic bacteria convert these monomers to 60 H₂ and volatile fatty acids. Aceticlastic methanogens perform the final step in biogas production, 61 primarily Methanosarcina in a high-acetate environment (>10⁻³M) and Methanosaeta which grows only 62 by the aceticlastic reaction, as well as hydrogenotrophic methanogens. Methanogenesis is considered 63 the rate-limiting step. Moreover, this phase is most vulnerable to either temperature or pH variations 64 and toxic chemicals (Liu and Whitman, 2008b). Low methanogen activity produces an accumulation of H_2

65 and short chain fatty acids with a consequent decrease in pH. Therefore, enhancing methanogenesis is a 66 promising method for improving anaerobic reactor performance. There are three primary useful 67 substrates for methanogens. Acetate is the predominant intermediate in the anaerobic food chain, and 68 as much as two-thirds of biologically generated methane is produced from this molecule (Liu and 69 Whitman, 2008a). Although every pathway begins differently, they all end with the same step, the 70 reaction of methyl-coenzyme M (HS-CoM) with a second thiol coenzyme B) that forms 71 methane and a mixed disulphide between coenzyme M and coenzyme B. This reaction is catalysed by 72 methyl-coenzyme M reductase (Mcr), thus, Mcr is the key enzyme in methanogenesis (Friedrich, 2005). 73 In its active site, this enzyme contains a unique prosthetic group, which is a nickel (Ni) porphinoid 74 referred to as coenzyme F430 (Hedderich and Whitman, 2006). As HS-CoM has been found in all 75 methanogens examined, it has been proposed as a sensitive biomarker for quantitative and qualitative 76 methanogen identification in different anaerobic environments. Mcr is exclusive to methanogens, with 77 the exception of methane-oxidising Archaea (Knittel and Boetius, 2009; Whitman et al., 2006), and 78 specific primers have been developed for the Mcr α -subunit gene sequence (mcrA) (Franke-Whittle et 79 al., 2009; Luton et al., 2002; Steinberg and Regan, 2008). McrA analysis can be used either in conjunction 80 with or independently of the 16S rRNA gene (Bergmann et al., 2010a; Bergmann et al., 2010b; Cardinali-81 Rezende et al., 2009; Nettmann et al., 2008); this gene is absent in all non-methanogens, with the 82 exception of the anaerobic methane-oxidizing Archea (Steinberg and Regan, 2008). The aim for the work 83 herein was to apply a recently proposed methodology for determining methanogen abundance using 84 Real-Time quantitative PCR (RT-qPCR) targeting the mcrA gene to evaluate methanogen load during the 85 digestion process and identify its correlation with the biogas production efficiency. The purpose is to 86 develop a useful Archea bio-indicator for process performance. The analyses involved two different 87 digestion sessions that differed only in pre-treatment of the organic material input. We observed 88 methanogen modulation during the digestions referring to the recorded methane production rate.

90 2. METHODS AND MATERIALS

91 **2.1** Digestion processes

92 Two pilot reactors were fed with a pre-treated organic fraction of municipal solid waste (OFMSW) and 93 waste-water sludge. The pre-treatment methods used in this study included a pressure-extrusion (A) 94 and a turbo-mixing (B) system. In method A, a specially designed extruder press separates the input 95 waste into two fractions: dry, which is sent to thermal conversion, and semi-solid. The municipal waste, 96 fed into the extruder press, is exposed to high pressure (280 bar) in a perforated extrusion chamber that 97 results in a fluid organic portion (food residues, various putrescible fractions). The organic fluid is 98 pushed out by the difference in pressure between the interior and exterior of the chamber, yielding a 99 more homogeneous sample. This is a consequence from separating the more mechanically resistant 100 portions (paper residues, cellulose tissues, wood and lignin parts, etc.). The pressure-extrused OFMSW, 101 obtained using the previously described procedure, was then diluted with waste-water sludge (1:3). In 102 contrast, method B (the turbo-mixing system) is a wet process that operates with a TS content lower 103 than 8%. Mixing and treating occur via a rotating plate, with hummers at the bottom of the turbo-mixing 104 chamber that, rotating at high velocity, induces the suspension to shear and crush. After 20 minutes of 105 mixing, the turbo-mixer was empty and a vessel was filled with the mixture, where density separated 106 the contents. The particles that weigh more than water precipitate to the bottom, where they are taken 107 up using a screw and collected in an external vessel. The same procedure was applied to the particles 108 that weigh less than water and float on the surface. The remaining suspension is the organic fraction, 109 which is pumped to a storage basin after a passage through a shredding pump. In this case, the OFMSW 110 was directly turbo-mixed with waste water sludge. Full control of the final TS content suspension is 111 impossible in the turbo-mixing system. The primary physical-chemical characteristics for each type of 112 feed used herein, immediately prior to the reactor, are shown in Table 1.

We compared the analytical results obtained from feed samples pre-treated with methods A (pressureextrusion) and B (turbo-mixing). OFMSW pre-treated with press-extrusion has a higher percentage of volatile solids that may possibly transform into biogas in feed pre-treated with method B where the organic matter value is lower. Further, the C/N ratio for the feed mixtures was evaluated. It has been shown that optimum values for the carbon-to-nitrogen (C/N) ratio are within the 20 to 70 range for the anaerobic digestion process (Burton and Turner, 2003), but even lower values (12 to 16) have also been reported (Mshandete et al., 2004) and, recently, more widely studied (Mata-Alvarez et al., 2011). The C/N values herein were better after pre-treatment A than pre-treatment B.

121 Anaerobic co-digestion was examined using a reactor, illustrated in Figure 1, with a 15 L volume capacity 122 and a 10 L working volume. The temperature was mesophilic and maintained at 38±2°C using a water 123 recirculation system connected to a thermostatic valve. The biogas produced was collected and 124 measured in a calibrated gasometer, and a mixing system with the re-circulated biogas from the 125 anaerobic digestion process was used. The reactors were equipped with two openings, one at the top 126 for feeding and one below to collect effluent discharge. Every day, 500 ml of digestate was removed 127 from each reactor before an additional 500 ml of fresh feed was added. The reactors were operated at a constant organic load rate of 4.5±0.3 kg TVS /m³*day for OFMSW pressure-extrusion and 1.7±0.5 kg TVS 128 129 /m³*day, as a mean value, for the OFMSW pulper pre-treatment. The pre-treatments were examined 130 over two 20-day consecutive hydraulic retention time periods for each organic loading rate. The first 131 period would ensure the highest replacement parts of the waste material inside the reactors, and in the 132 second 20 days, we analysed the process under stable conditions, as all the feed had replaced the 133 inoculum content.

Every 2 days, representative samples from the anaerobic reactor effluent were analysed. The following parameters were analysed three times per week in accordance with Standard Methods (APHA, 1995): pH, total solids (TS), total volatile solids (TVS), alkalinity, acidity, N, and total carbon. Daily biogas production was measured using a liquid displacement system connected to the digester. The biogas volume was corrected using standard temperature and pressure conditions. The biogas composition (methane and carbon dioxide percentages) was analysed once per week using a portable analyser and 140 confirmed via gas chromatography. For methanogen analysis, the samples were collected three times 141 per week in a 50 ml sterile tube and frozen at -20°C until extraction. More than 80 digest effluent 142 samples were collected from more than a year of digestion operations beginning March 2009.

143

144 **2.2 DNA extraction and purification**

145 The digestate aliquots were thawed at 4°C overnight and centrifuged at 3000 g for 20 minutes; the 146 supernatant was then removed, and the semi-dry aliquots were used for the following steps. The total 147 DNA was extracted from 0.25 g of particulate matter (residue humidity was 31±5%) using the PowerSoil 148 DNA Isolation Kit followed by UltraClean Soil DNA Kit (MoBio Laboratories). The DNA quantity extracted 149 was 3.51 ± 1.53 ng/µl (mean), and its quality was evaluated using gel electrophoresis before the chain 150 reaction. Only the samples over 1 ng/ μ l and of sufficient DNA quality, tested by gel electrophoresis, 151 were used for the following step. Lower quantity or a bed quality of the DNA makes the successive 152 quantification difficult so this kind of DNA quality cut-off was applied. 50 samples were sufficient in 153 distinct DNA quantity and quality.

154

155 **2.3 RT-qPCR analysis**

156 After DNA extraction and purification, total methanogen was quantified using methanogen-specific 157 short primers for an mcrA sequence described by Steinberg and Regan (Steinberg and Regan, 2008) and 158 synthesised by ThermoBiopolymer. For the reactions, we used a standard super-mix (Bio-Rad SsoFast[™] 159 EvaGreen SuperMix), the RT-PCR Chromo4 (Bio-Rad) and the Opticon Monitor 3 Software. The reaction 160 conditions were previously described (Steinberg and Regan, 2008). The standard reference was a 161 Methanosarcina acetivorans mcrA sequence of approximately 470 base pairs (Lueders et al., 162 2001)included in a pCR21 vector (Invitrogen) supplied by L.M. Steinberg and J.M. Regan of the 163 Pennsylvania State University. This plasmid was amplified by transforming Escherichia coli Top10 cells 164 according to the manufacturer's instruction. The transformed cells were selected on LB agar in presence 165 of ampicillin, and the plasmid was extracted using a plasmid DNA purification kit (NucleoSpin Plasmid, 166 Macherey-Nagel). The standard curve had six points (figure 2) and was calculated using the threshold cycle method; for the highest yield, 2,3 ng of plasmid was amplified (~4.5*10⁸ plasmid copies). Between 167 168 each standard curve point there is a 1:10 dilution. The standards and samples were tested in triplicate. 169 The triplicate value was accepted only if the coefficient of variation was below 20%. The correlation 170 coefficient was considered sufficient if above 0.990. PCR efficiency was 97%, and the resolution limit for 171 the method was $4.5*10^3$ copies of *mcrA*. We used in the sample amplification 2 μ l of a 1:10 dilution for 172 each DNA extract. This quantity was evaluated as the best among various tested quantities for a 173 standard curve and with an acceptable PCR efficiency. The 1:10 dilution limited the effect of inhibiting 174 substances present in this type of sample. Figure 3 shows the quantification precision obtained, 175 beginning with the same two samples re-extracted 10-fold. This test illustrated that the sampling 176 procedure does not affect the final analysis. The variation coefficient was below 10% for sample 1 and 177 below 20% for sample 2. The 50 samples were sufficient in DNA quantity and quality.

178

179 **2.4 Statistics**

Statistical analyses were performed using the SPSS Package, version 17.0 for Windows. A Pearson correlation coefficient was used to assess the relationship between the variables. A T-test of independent variables was used in the mean evaluations. The differences and correlations were considered significant at p<0.05 and highly significant at p<0.01.

184

3. RESULTS AND DISCUSSION

Table 2 shows the primary process-control factors that were monitored, such as biogas production parameters, solid reduction data and acidity values. Pre-treatment A not only yields more relevant quantities of biogas, but the biogas production was also double the value observed in treatment B. Thus, treatment A provides better input preparation likely due to a reduction in particle size, increasing the available specific surface and enhancing the contact surface between bacteria. This was further
 confirmed via a %TVS reduction. The digestion process fed with pre-treatment A material performs
 better with respect to pre-treatment B, despite the unfavourable input pH (table 1).

The correlation between methanogen abundance and biogas production is positive and statistically significant (p<0.01), as shown in figure 4. This correlation holds both considering all the collected data (first and second digestions) and separating the two data sets. As a measure of the correlation, Pearson's rho is 0.579 (p<0.01). In figure 4, the correlations are expressed for each collected data set. Further, the single digestion showed a positive and significant correlation between methanogen quantification and biogas production rate.

199 Even if biomolecular methodology for methanogen analyses during anaerobic processes became 200 nowadays diffused, this is one of first studies to analyse and correlate methanogen load with the biogas 201 production during full digestion sessions. Further, this are very few studies that monitored by such 202 biological method different process and that observed a significant correlation using a mcrA-based PCR 203 method. Although, anaerobic methanotrophs that are phylogenetically related to methanogens have an 204 Mcr-like protein that catalyses methane oxidation (Nunoura et al., 2008), the abundance of this 205 microbial population is likely negligible with respect to the methanogen community, as deduced from 206 the scarcity of methanotroph products such as N_2 and HS^- (Balat and Balat, 2009).

207 Figure 5 shows a marked and statistically significant difference between the methanogen load present in 208 the digester feed from pre-treated A and pre-treated B organics. This trend is similar to biogas 209 production (table 2), even if only the methanogen load difference is statistically significant (p<0.01). 210 Setting a 'cut-off' for optimal production generated an additional measure of correlated activity. As also 211 observed in the literature, production rate ranged between 0.3 to 0.8 m^3 biogas/kg TVS in base of 212 various parameters such as input characteristics, process temperature, organic load, and so on. Optimal 213 production can be defined as 0.6 m³ biogas/kg TVS (Cuetos et al., 2009; Gomez et al., 2006). Shown in 214 figure 5, the methanogen load is lower in the samples with sub-optimal production rates compared with

215 optimal samples (T-test, p<0.05).

216 However, we would also discuss in this paper whether, overall, the data support methanogen 217 abundance, determined with the previously described RT-qPCR method, decrease as a biomarker for 218 methanogenesis efficiency. The stability parameters typically measured when monitoring digestions, 219 such as pH, alkalinity and acidity of the mixture as well as split volatile fatty acid concentration in the 220 reactor, were not sufficient early predictors during this process. A variation of these parameters 221 correlates with a contemporaneous decrease in production. In this instance, corrective steps to support 222 a productive balance in the reactor are ineffective. As shown in figure 6, prediction may only be relevant at certain times during the process, such as between the 3rd and 17th of April, 8th and 15th of May and 223 22nd and 29th of May. On certain days, we observed a decrease in the *mcrA* copies just before a decrease 224 225 in the specific biogas production rate; however, this event was not regular during the digestion process.

Recent studies show that methanogen population dynamics and community structures can vary during the digestion processes (Lee et al., 2009a; Lee et al., 2009b; Yu et al., 2006), as can volatile fatty acid concentrations (Wang et al., 2009). All of these studies discussed the microbial communities' capacity for responding to changes in anaerobic environments, such as altered feeding (Kovacik et al., 2010) and different temperature (Sasaki et al., 2011), among others.

231

232

4. CONCLUSIONS

Despite the environmental complexities inherent in the work herein, we conducted a representative sampling procedure and a valid DNA extraction. The results show proper quantitation for the majority of the collected samples (63%). The experiments conducted during the digestion process in the reactors demonstrate a good method for determining methanogen concentration with respect to the biogas production. The hypothesis, that methanogen community abundance and composition are strictly related to the methane production, is confirmed. In the last few years, the research community has enhanced our knowledge regarding Archea microorganisms from both an ecological standpoint (Leigh et al., 2011; Liu and Whitman, 2008b) and a human health perspective (Johnston et al., 2010). The economic costs are really onerous for these analyses (Kalia and Purohit, 2008) moreover the proposed PCR method herein is a useful tool for studying methanogen populations and their modulation related to biogas production rate (Lee et al., 2009a; Patil et al., 2010), even if the results herein cannot yet provide a clear activity prediction.

246 McrA gene-based methods could be even more diffused methods (Zhu et al., 2011), also relating with 247 digestion performance (Bialek et al., 2011) and feeding matrices, which currently pose a relevant 248 concern for waste management and public health (Cardinali-Rezende et al., 2009). A fundamental step 249 includes analysis of, at least, the different families of methanogens to identify a better early bio-250 indicator among the total methanogen population (Steinberg and Regan, 2009; Vavilin et al., 2008). 251 From these studies, an alarm threshold for micro-organism levels could be a useful, fundamental 252 process-control parameter but the introduction of this analysis must be economically sustainable. 253 Prediction ability with respect to biogas loss over several days might also be interesting from an 254 economic standpoint. The problems related with arrested methanogenesis are one of the largest 255 obstacles to anaerobic digestion diffusion. In the approach proposed herein, it is essential to examine 256 the community composition and the genus contribution to optimise the digestion process and maximise 257 CH₄ yield.

258

259 Acknowledgements

The Authors wish to thank the Piedmont Region for funding; this work is a portion of a larger project referred to as DigestedEnergy, and it was founded for the 2006 call for Pre-competitive Development and Industrial Research. This funding scheme was promoted by the Piedmont Region with European Community resources. Special acknowledgments are due to L. Steinberg and J. Regan for the plasmid standard supply. Finally, the Authors thank all of the numerous collaborators employed in each of the

- 265 involved institutions: Università degli Studi del Piemonte Orientale "A. Avogadro", Politecnico di Torino,
- 266 SMAT S.p.A., Amiat S.p.A., Ansaldo FC S.p.A., Acsel Susa S.p.A., VM-press s.r.l., Federsviluppo, E.R.A.P.R.A
- 267 Piemonte, and Università degli Studi di Torino.

269 **FIGURE CHAPTERS**

270

271 Figure 1 – The pilot hardware description is illustrated. The same reactor, in different six-month 272 fermentation sessions, was used with two different feed pre-treatments.

273

274 Figure 2 – Example of a standard curve obtained during experiment. The same curve is included in all 275 analysis sessions, and the samples are placed within the standard points.

276

277 Figure 3 - RT-PCR quantification of 10 different, consecutive extractions for the same two samples. 278 Different samples collected on different days were used. Cross-shaped points represent each different 279 extract quantification. For each set, the last point (score) represents the mean from all the analyses. For 280 each data point, the standard deviation is shown.

281

282 Figure 4 - Linear regression model for RT-PCR quantification samples collected from reactors and the 283 logarithm of the target gene copies for each μ I of DNA sample. (A) circles represent pressure-extrusion 284 pre-treatment data, y=1.0303x+6.5118 (correlation=0.418, p<0.05); (B) rhombus represent the turbo-285 mixing pre-treatment data, y=2.5629x+4.3782 (correlation=0.622, p<0.05).

286

287 Figure 5 - McrA abundance evaluated during the digestion process, subdivided by input material pre-288 treatment (T-test p<0.01) and optimal biogas production (0.6 m^3/kg SV) (T-test p<0.05).

289

290 Figure 6 - Temporal trends of mcrA abundance (squares) and biogas production (rhombus) during 291 digestion of the pressure-extruded input for the reactor.

292

294295 TABLES

296

| | Pre-treatment A | Pre-treatment B |
|---------------------------------|-----------------|-----------------|
| рН | 4.4 ± 0.3 | 6.0 ± 0.7 |
| Total Solids (TS) (%) | 9.9 ± 0.7 | 4.6 ± 1.1 |
| Total volatile solids (TVS) (%) | 8.7 ± 0.7 | 3.3 ± 1.1 |
| TVS/ST (%) | 86.8 ± 0.2 | 70.6 ± 4.9 |
| Carbon (C) (%ST) | 46.0 ± 0.9 | 37.0 ± 3.4 |
| Nitrogen (N) (%ST) | 3.1 ± 0.2 | 3.5 ± 0.3 |
| C/N ratio | 15.2 ± 1.1 | 10.4 ± 1.5 |

297

Table 1: Characteristics of the pre-treated inputs from the two different methods used in the anaerobic

299 co-digestion processes.

300

| Parameters | Pre-treatment A | Pre-treatment B |
|--|-----------------|-----------------|
| Daily biogas production (L/die) | 27.08 ± 3.01 | 4.87 ± 2.46 |
| Specific biogas production (m ³ /kg VS _{added}) | 0.64 ±0.07 | 0.30 ± 0.13 |
| TS reduction (%) | 64.44 ± 7.57 | 31.67 ± 6.23 |
| TVS reduction (%) | 73.84 ± 5.87 | 38.13 ± 6.70 |
| рН | 7.36 ± 0.34 | 6.82 ± 0.52 |
| Ac./Alc. ratio | 0.37 ± 0.18 | 2.47± 2.41 |
| CH ₄ (%) | 60.60 ± 2.90 | 57.50 ± 6.10 |
| CO ₂ (%) | 37.70 ± 3.20 | 41.00 ± 6.44 |

301

302 **Table 2:** Evaluation of relevant parameters for the co-digestion processes subdivided by pre-treated

303 method.

- 305 **BIBLIOGRAFY**
- 306APHA, Standard methods for the examination of water and wastewater. American Water307Works Association, Water Environment Federation, Washington, DC, USA,, 1995.
- Balat, M., Balat, H., 2009. Biogas as a Renewable Energy SourceA Review. Energy Sources Part
 a-Recovery Utilization and Environmental Effects. 31, 1280-1293.
- Bergmann, I., et al., 2010a. Influence of DNA isolation on Q-PCR-based quantification of
 methanogenic Archaea in biogas fermenters. Systematic and Applied Microbiology.
 33, 78-84.
- 313Bergmann, I., et al., 2010b. Determination of methanogenic Archaea abundance in a314mesophilic biogas plant based on 16S rRNA gene sequence analysis. Canadian Journal315of Microbiology. 56, 440-444.
- 316Bialek, K., et al., 2011. Quantitative and qualitative analyses of methanogenic community317development in high-rate anaerobic bioreactors. Water Research. 45, 1298-1308.
- 318Bouallagui, H., et al., 2005. Bioreactor performance in anaerobic digestion of fruit and319vegetable wastes. Process Biochemistry. 40, 989-995.
- 320Burton, C. H., Turner, C., 2003. Anaerobic treatment options for animal manures. UK: Silsoe321Research Institute, Silsoe, Bedford.
- Cardinali-Rezende, J., et al., 2009. Molecular identification and dynamics of microbial
 communities in reactor treating organic household waste. Applied Microbiology and
 Biotechnology. 84, 777-789.
- Cuetos, M. J., et al., 2009. Anaerobic co-digestion of poultry blood with OFMSW: FTIR and TG DTG study of process stabilization. Environmental Technology. 30, 571-582.
- das Neves, L. C. M., et al., 2009. Biogas Production: New Trends for Alternative Energy
 Sources in Rural and Urban Zones. Chemical Engineering & Technology. 32, 1147-1153.
- 329Franke-Whittle, I. H., et al., 2009. Design and testing of real-time PCR primers for the330quantification of Methanoculleus, Methanosarcina, Methanothermobacter, and a331group of uncultured methanogens. Canadian Journal of Microbiology. 55, 611-616.
- Friedrich, M. W., 2005. Methyl-Coenzyme M reductase genes: unique functional markers for
 methanogenic and anaerobic methane-oxidizing Archea. Methods in Enzymology, 428 442.
- 335Gomez, X., et al., 2006. Anaerobic co-digestion of primary sludge and the fruit and vegetable336fraction of the municipal solid wastes Conditions for mixing and evaluation of the337organic loading rate. Renewable Energy. 31, 2017-2024.

- Hedderich, R., Whitman, W. B., 2006. Physiology and Biochemistry of the mathane-producing
 Archeae. Prokaryotes Volume 2, 1050-1079.
- 340Johnston, C., et al., 2010. A real-time qPCR assay for the detection of the nifH gene of341Methanobrevibacter smithii, a potential indicator of sewage pollution. J Appl342Microbiol. 109, 1946-56.
- Kalia, V. C., Purohit, H. J., 2008. Microbial diversity and genomics in aid of bioenergy. J Ind
 Microbiol Biotechnol. 35, 403-19.
- Knittel, K., Boetius, A., 2009. Anaerobic Oxidation of Methane: Progress with an Unknown
 Process. Annual Review of Microbiology. 63, 311-334.
- Kovacik, W. P., et al., 2010. Microbial dynamics in upflow anaerobic sludge blanket (UASB)
 bioreactor granules in response to short-term changes in substrate feed.
 Microbiology-Sgm. 156, 2418-2427.
- Lee, C., et al., 2009a. Quantitative analysis of methanogenic community dynamics in three
 anaerobic batch digesters treating different wastewaters. Water Research. 43, 157 165.
- Lee, C., et al., 2009a. Quantitative and qualitative transitions of methanogen community
 structure during the batch anaerobic digestion of cheese-processing wastewater.
 Applied Microbiology and Biotechnology. 87, 1963-1973.
- Lee, M., et al., 2009b. Comparative performance and microbial diversity of hyperthermophilic
 and thermophilic co-digestion of kitchen garbage and excess sludge. Bioresour
 Technol. 100, 578-85.
- Leigh, J. A., et al., 2011. Model organisms for genetics in the domain Archaea: methanogens,
 halophiles, Thermococcales and Sulfolobales. FEMS Microbiol Rev.
- Liu, Y. C., Whitman, W. B., 2008a. Metabolic, phylogenetic, and ecological diversity of the
 methanogenic archaea. Incredible Anaerobes: From Physiology to Genomics to Fuels.
 1125, 171-189.
- 364Liu, Y. C., Whitman, W. B., Metabolic, phylogenetic, and ecological diversity of the365methanogenic archaea. In: J. Wiegel, et al., Eds.), Incredible Anaerobes: From366Physiology to Genomics to Fuels. Blackwell Publishing, Oxford, 2008b, pp. 171-189.
- Lozano, C. J. S., et al., 2009. Microbiological characterization and specific methanogenic
 activity of anaerobe sludges used in urban solid waste treatment. Waste
 Management. 29, 704-711.
- 370Lueders, T., et al., 2001. Molecular analyses of methyl-coenzyme M reductase alfa-subunit371(mcrA) genes in rice field soil and enrichment cultures reveal the methanogenic372phenotype of a novel archaeal lineage. Environmental Microbiology. 3, 194-204.

- Luton, P. E., et al., 2002. The mcrA gene as an alternative to 16S rRNA in the phylogenetic analysis of methanogen populations in landfill. Microbiology-Sgm. 148, 3521-3530.
- 375Mata-Alvarez, J., et al., 2011. Codigestion of solid wastes: A review of its uses and376perspectives including modeling. Critical Reviews in Biotechnology. 31, 99-111.
- 377Mshandete, A., et al., 2004. Anaerobic batch co-digestion of sisal pulp and fish wastes.378Bioresource Technology. 95, 19-24.
- Nettmann, E., et al., 2008. Archaea diversity within a commercial biogas plant utilizing herbal
 biomass determined by 16S rDNA and mcrA analysis. Journal of Applied Microbiology.
 105, 1835-1850.
- 382Nunoura, T., et al., 2008. Quantification of mcrA by fluorescent PCR in methanogenic and383methanotrophic microbial communities. Fems Microbiology Ecology. 64, 240-247.
- Patil, S. S., et al., 2010. Microbial community dynamics in anaerobic bioreactors and algal
 tanks treating piggery wastewater. Applied Microbiology and Biotechnology. 87, 353 363.
- Rozzi, A., Remigi, E., 2004. Methods of assessing microbial activity and inhibition under
 anaerobic conditions: A literature review. Reviews in Environmental Science and
 Biotechnology. 3, 93-115.
- 390Sasaki, K., et al., 2011. Bioelectrochemical system accelerates microbial growth and391degradation of filter paper. Applied Microbiology and Biotechnology. 89, 449-455.
- Schievano, A., et al., 2009. Substituting energy crops with organic wastes and agro-industrial
 residues for biogas production. Journal of Environmental Management. 90, 2537 2541.
- Steinberg, L. M., Regan, J. M., 2008. Phylogenetic Comparison of the Methanogenic
 Communities from an Acidic, Oligotrophic Fen and an Anaerobic Digester Treating
 Municipal Wastewater Sludge. Applied and Environmental Microbiology. 74, 6663 6671.
- Steinberg, L. M., Regan, J. M., 2009. mcrA-Targeted Real-Time Quantitative PCR Method To
 Examine Methanogen Communities. Applied and Environmental Microbiology. 75,
 401 4435-4442.
- 402 Vavilin, V. A., et al., 2008. Methanosarcina as the dominant aceticlastic methanogens during
 403 mesophilic anaerobic digestion of putrescible waste. Antonie Van Leeuwenhoek
 404 International Journal of General and Molecular Microbiology. 94, 593-605.
- 405Wang, Y. Y., et al., 2009. Effects of volatile fatty acid concentrations on methane yield and406methanogenic bacteria. Biomass & Bioenergy. 33, 848-853.

- Ward, A. J., et al., 2008. Optimisation of the anaerobic digestion of agricultural resources.
 Bioresource Technology. 99, 7928-7940.
- 409 Whitman, W. B., et al., 2006. The Methanogenic Bacteria. Prokaryotes. 3, 165-207.
- Yu, Y., et al., 2006. Use of real-time PCR for group-specific quantification of aceticlastic
 methanogens in anaerobic processes: Population dynamics and community structures.
 Biotechnology and Bioengineering. 93, 424-433.
- Zhu, C. G., et al., 2011. Diversity of methanogenic archaea in a biogas reactor fed with swine
 feces as the mono-substrate by mcrA analysis. Microbiological Research. 166, 27-35.
- 416