

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Integrated strategies for the control of Fusarium head blight and deoxynivalenol contamination in winter wheat.

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/124201> since 2016-07-04T19:29:06Z

Published version:

DOI:10.1016/j.fcr.2012.04.004

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in “Integrated strategies for the control of Fusarium head blight and deoxynivalenol contamination in winter wheat.”, *Field Crops Research*, 2012; 133: 139-149, DOI: [10.1016/j.fcr.2012.04.004](https://doi.org/10.1016/j.fcr.2012.04.004)

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

- (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.
- (2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.
- (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>), [+ DOI: [10.1016/j.fcr.2012.04.004](https://doi.org/10.1016/j.fcr.2012.04.004)]

Integrated strategies for the control of Fusarium head blight and deoxynivalenol contamination in winter wheat.

Massimo Blandino^{a*}, Miriam Haidukowski^b, Michelangelo Pascale^b, Luca Plizzari^c,
Diego Scudellari^d, Amedeo Reyneri^a.

Affiliation:

^a Dipartimento di Agronomia, Selvicoltura e Gestione del Territorio, Università di Torino, via Leonardo Da Vinci 44, 10095 Grugliasco (TO), Italy.

^b Institute of Sciences of Food Production (ISPA), National Research Council (CNR), via G. Amendola 122/O, 70126 Bari, Italy.

^c CRA - SCV, via Forlani 3, 26866, S. Angelo Lodigiano (LO), Italy

^d Grandi Colture Italiane S.c.a., viale Volano 195/A, 44123 Ferrara, Italy

* Corresponding author: Tel: +39-011-6708895; fax +39-011-6708798.

E-mail address: massimo.blandino@unito.it

Address to which the proofs should be sent:

Dr. Massimo Blandino, Dipartimento di Agronomia, Selvicoltura e Gestione del Territorio, Università di Torino, via Leonardo Da Vinci 44, 10095 Grugliasco (TO), Italy.

Abbreviations: FHB, Fusarium head blight; DON, deoxynivalenol; GDDs, growing degree days; GS, growth stage; MR, moderately resistant cultivar; S, susceptible cultivar.

1 **Abstract**

2 Fusarium head blight (FHB) disease and deoxynivalenol (DON) contamination of
3 wheat grains depend on multiple factors, above all climatic conditions, but also
4 agronomic factors such as crop rotation, debris management, variety susceptibility
5 and fungicide applications. Although it is generally believed that multiple strategies
6 are more successful than a single strategy, only a few studies have shown the
7 quantitative effect of combining multiple strategies.

8 Field experiments have been conducted over three growing seasons in three sites in
9 Northern Italy to evaluate the effect of previous crop residue management through
10 tillage, variety susceptibility and triazole fungicide application on common wheat,
11 according to a full factorial scheme. The following parameters were analyzed: FHB
12 severity, grain yield and DON contamination.

13 The collected data have clearly shown a close interaction between the factors
14 involved in FHB severity and DON content, while the interactions were less
15 significant for grain yield. In all nine trials, the DON contamination was significantly
16 affected by the interaction of at least two of the compared factors, while the
17 interaction between all three factors involved was significant in four trials. The most
18 favourable scenario to avoid DON contamination (ploughing, moderately resistant
19 variety, triazole application at heading) reduced the DON content by 97% compared
20 to the worst one (direct sowing, susceptible variety, no fungicide application).

21 Since the interaction between the agricultural practices have shown a synergistic
22 effect, integrated multiple strategies, in areas characterized by a high risk of FHB,
23 can be considered the very effective management means of reducing FHB and DON
24 contamination in wheat.

- 1 **Keywords:** winter wheat, residue management, variety susceptibility, fungicide
- 2 application, Fusarium head blight, deoxynivalenol.

1 **1. Introduction**

2 Fusarium Head Blight (FHB) is the most diffuse wheat ear disease throughout the
3 world and it is caused by *Microdochium nivale* and different *Fusarium* species
4 (Champeil et al., 2004a). This disease causes total or partial ear premature
5 senescence with a consequent reduction in both crop yields and grain quality
6 (Pirgozliev et al., 2003). *F. graminearum* and *F. culmorum*, the most important
7 species responsible for FHB, are also the main causes of the accumulation of
8 deoxynivalenol (DON) in wheat kernels, a mycotoxin of the trichotecenes group,
9 inhibits protein biosynthesis in eukaryotes (Bottalico and Perrone, 2002).

10 FHB infection and DON contamination of wheat grains depend on multiple factors,
11 above all climatic conditions, particularly at flowering (Xu, 2003), but also agronomic
12 factors such as crop rotation, debris management, variety susceptibility and fungicide
13 applications (Pirgozliev et al., 2003; Koch et al., 2006), which aim at reducing
14 infection or growth of toxigenic fungi (Aldred and Magan, 2004).

15 The primary reservoir of inoculum is debris from the previous crop (Xu, 2003). FHB
16 epidemics are supported by cropping systems that leave high amount of crop debris
17 on the soil surface (Pereyra and Dill-Macky, 2008; Blandino et al., 2010) and
18 pathogens survive longer on residues that do not degrade easily, such as stem
19 nodes or stalks (Sutton, 1982). Thus, FHB disease and DON contamination are more
20 severe if the preceding crops are maize or sorghum, rather than wheat or barley and
21 even less contamination is observed following other crops (Champeil et al., 2004b;
22 Smith-White et al., 2004).

23 Limited soil tillage or no-tillage increase the frequency of FHB, whereas deep tillage,
24 such as ploughing, decreases it (Miller et al., 1998). Maiorano et al. (2008) reported a

1 close relationship between DON contamination in wheat grains and the quantity of
2 maize crop residues on the soil surface at anthesis. Moreover, FHB severity and
3 DON content are clearly affected by the interaction of previous crop residues and
4 tillage practice applied (Dill-Macky and Jones, 2000).

5 As far as variety susceptibility to FHB and DON is concerned, breeding progress in
6 cereals, using conventional methods, molecular markers or through transgenic
7 approaches, have been discussed in great detail in several reviews (Hollins et al.,
8 2003; Snijders, 2004). At present, no durable, fully FHB-resistant wheat cultivars
9 exist, therefore their control relies on the use of commercial cultivars with partial
10 resistance (Mesterházy et al., 2005), although wheat varieties more resistant to FHB
11 have been shown to reduce DON production to almost nil in recent studies (Tóth et
12 al., 2008).

13 The effect of fungicide application on FHB and DON contamination control has been
14 well documented. Several studies conducted on *in vitro* experiments (Ramirez et al.,
15 2004), on field trials in which wheat was artificially inoculated (Mesterházy et al.,
16 2003; Chala et al., 2003) or under natural infection conditions (Blandino et al., 2006)
17 have demonstrated that good levels of control can be achieved with fungicides. The
18 outcome of the use of fungicides seems to depend on the fungal species that are
19 present and the effect that the particular fungicide has on these species. Fungicides
20 containing triazole, imidazole or triazolinthione active ingredients, which inhibit the
21 biosynthesis of ergosterol, were the most active against FHB infection and DON
22 contamination (Haidukowski et al., 2005; loos et al., 2005). Of the azole group,
23 metconazole and prothioconazole, which have been developed more recently, have
24 been reported to be the most effective fungicides in controlling *Fusarium* spp. and
25 reducing the DON level in wheat grain (Paul et al., 2008).

1 Previous studies show that individual control methods can decrease the impact of the
2 disease significantly, but combining control methods can be expected to be more
3 efficient, especially if the climatic conditions are favourable for FHB infection
4 (Edwards, 2004). Therefore good agricultural practice (GAP) requires an integrated
5 approach that addresses all the possible risk factors in order to prevent DON
6 contamination (Pirgozliev et al., 2003). Moreover, although information is available on
7 the basic effect of individual agricultural practices on *Fusarium* infection and DON
8 contamination in wheat, only a few studies have been conducted to quantify the
9 relative importance of each of these factors compared to the others or to verify their
10 interactions and combined effects.

11 The aim of this study was to determine the effect of residue management, variety
12 susceptibility and fungicide application on FHB infection and DON contamination in
13 wheat kernels, with a particular attention to their interactions under natural infection
14 conditions.

15

2. Materials and Methods

2.1. Experimental site and treatments

The experiments were carried out between 2005 and 2008 at 3 sites in North Italy: Imola (IM), Riva presso Chieri (RC) and Sant'Angelo Lodigiano (SL). The geographic and the main agronomic information about the experimental fields is reported in Table 1.

The compared treatments were factorial combinations, in natural conditions, of:

- The previous crop residue management through tillage: ploughing to a 30 cm depth, thus incorporating the debris in the soil, followed by disk harrowing to prepare a proper seedbed vs. direct sowing;
- variety susceptibility: a variety classified as moderately resistant (MR) to FHB infection and DON contamination vs. a susceptible (S) one;
- fungicide application: a triazole fungicide application at heading [growth stage (GS) 59] (Zadoks et al., 1974) vs. an untreated control.

The treatments were assigned to experimental units using a split-plot design, with the previous crop residue management as the main-plot treatment and the variety susceptibility and fungicide application as the sub-plot treatments. Each trial was replicated three times in the IM and SL sites and four times in the RC site. The sub-plot was 7 x 2 m.

The previous crop was grain sorghum at site A (growing seasons 2005-06 and 2006-07) and grain maize at site A (2007-08), B and C. Since maize and sorghum are the most dangerous previous crops in the context of FHB epidemics and DON contamination (Gourdain, 2008), they have been selected to verify the effect of residue management through tillage in the more risky crop rotation conditions.

1 The MR variety that was used in each year and site was cv. Bologna , while cv. Serio
2 was the S one (Mayerle et al. 2007).

3 In each trial, the triazole fungicide was metconazole (Caramba[®], Basf, Italy,
4 formulation: suspension concentrate) and it was applied at 0.06 kg active ingredient
5 (AI) ha⁻¹. The fungicide was applied with a 4 nozzle precision sprayer (T-Jeet 110/04)
6 using a fine mist at a slow walk to ensure an effective coverage. The delivery
7 pressure at the nozzle was 324 KPa.

8 Planting was conducted in 12 cm wide rows at a seeding rate of 450 seeds m⁻². The
9 weed control was conducted with isoproturon and diflufenican at wheat tillering (GS
10 31). Glyphosate was applied to the non tilled field before direct sowing. A total of 140
11 kg N ha⁻¹ was applied to plots as a granular ammonium nitrate fertilizer, and was split
12 equally between GS 31 and 39. The sowing, fungicide application and harvesting
13 date for each year and each site are reported in Table 1.

14 Grain yields were obtained by harvesting with a Walter Wintersteiger cereal plot
15 combine-harvester. The grain yield results were adjusted to a 120 g kg⁻¹ moisture
16 content. The harvested grains were accurately mixed, and 2 kg grain samples were
17 taken from each plot to analyse the DON content.

18

19 2.2. FHB symptom evaluation

20 FHB severity was recorded for each plot at the soft dough stage (GS 85) by carrying
21 out visual evaluations of the disease. FHB severity was computed as the percentage
22 of spikelets per ear with symptoms. A scale of 1 to 7 was used in which each
23 numerical value corresponds to a percentage interval of surfaces exhibiting visible
24 symptoms of the disease according to the following schedule: 1 = 0-5%, 2 = 5-15 %, 3 = 15-30 %, 4 = 30-45 %, 5 = 45-60 %, 6 = 60-75 %, 7 = 75-100 %.

1 3 = 15-30%; 4 = 30-50 %, 5 = 50-75%, 6 = 75-90%, 7 = 90-100% (Parry et al., 1995).
2 The scores were converted to percentages of the ear exhibiting symptoms and each
3 score was replaced with the mid-point of the interval.
4

5 2.3. DON analyses

6 A 2 kg representative sample of wheat kernels from each plot was finely ground
7 using a Model MLI 204 Bühler laboratory mill (Bühler S.p.A, Milan, Italy) to pass a 1
8 mm sieve. The DON concentrations were determined according to the method
9 reported by Neumann et al. (2009) on the basis of an immunoaffinity column clean-
10 up of the extracts, and mycotoxin was determined by means of HPLC/UV.

11 Briefly, 25 g of ground samples were added to polyethylene glycol (PEG-8000) and
12 extracted with water by blending. Extracts were filtered through filter paper (Whatman
13 no. 4) and glass microfibre filter (Whatman GF/A) and cleaned up by DONTest
14 immunoaffinity column (VICAM, Milford, MA, USA). The toxin was determined by
15 reversed-phase HPLC apparatus with a diode-array UV detector set at 220 nm (1100
16 Series HPLC Value System, Agilent Technologies Inc., Santa Clara, CA, USA) . The
17 column was a Synergi 4 µm Hydro RP 80A, 150 × 3 mm (Phenomenex, Torrance,
18 CA, USA). The mobile phase consisted of a mixture of acetonitrile:water (10:90)
19 eluted at a flow rate of 0.5 mL min⁻¹. Recovery experiments were performed in
20 triplicate using DON free wheat samples spiked at levels of 100, 500, 1000 and 2000
21 µg kg⁻¹. Recoveries were higher than 80% with relative standard deviations less than
22 10%. DON standard used for recovery experiments and HPLC calibration curves was
23 purchased from Sigma-Aldrich s.r.l. (Milan, Italy).

24 Appropriate dilutions of the sample extracts contaminated with higher DON levels

1 than 2000 $\mu\text{g kg}^{-1}$ were necessary before loading them into the immunoaffinity
2 columns in order to avoid saturation of the DON-antibody binding sites. The detection
3 limit of the method was 20 $\mu\text{g kg}^{-1}$ (signal-to-noise ratio of 3:1).
4

5 2.4. Statistical analysis

6 The effect of agronomic factors on FHB severity, grain yield and DON content was
7 tested by means of a repeated measure analysis of variance (RM-ANOVA) in which
8 the management of the previous crop residue was the between-subject factor, while
9 the variety susceptibility and fungicide application were the within-subject factors.
10 The residual normal distribution was verified using the Kolmogorov-Smirnov test,
11 while variance homogeneity was verified using the Levene test. When the
12 interactions between the factors were significant, the resultant means were
13 compared using the protected Fisher Least Significant Difference (LSD) adjusted for
14 multiple comparison using the Bonferroni procedure. The RM-ANOVA was
15 conducted separately for all the year and site combinations, in order to verify clearly
16 in each experiment the interactions between the involved agronomic factors. The
17 statistical package SPSS for Windows, Version 17.0 (SPSS Inc., Chicago) was used
18 for the statistical analysis.

19

1 **3. Results**

2 **3.1. Weather conditions**

3 The three growing seasons showed different meteorological trends from the
4 beginning of the stem elongation stage to the harvest (Table 2). In 2006, the
5 precipitations were not particularly elevated, but they were concentrated close to
6 anthesis (GS 65), particularly at the IM and RC sites. In 2007, frequent rainfall
7 occurred at the IM and RC sites, but only at the end of ripening (June), while the
8 rainfall was higher from anthesis to the milk stage (GS 75) at the SL site. In 2008,
9 instead, the precipitations were frequent and regular from April to June, above all
10 from the beginning of flowering to the soft dough stage at the RC and SL sites, thus
11 prolonging the harvest till the middle of July. In 2006 and 2007, the growing degree
12 days (GDDs) were particularly high in June, thus quickening the canopy senescing
13 process and leading to a reduction in the grain filling period and to an early maturity
14 of the crop.

16 **3.2. FHB severity**

17 In eight of the nine trials, the FHB severity was significantly affected by the
18 interaction of at least two of the compared factors, while the interaction between all
19 three factors involved was significant in four trials: in 2006 at site IM, in 2007 at site
20 SL and in 2008 at site RC and SL (Table 3).

21 In 2006, at site IM, when applied on their own in order to control FHB, none of the
22 factors was able to reduce FHB severity compared to the worst scenario (direct
23 sowing, S variety, no fungicide application), while a significant reduction in disease

1 symptoms was always observed with the preventive combination of all the factors
2 (ploughing, MR variety, fungicide application) (Table 4). In 2007, at site SL, the
3 ploughing or the use of an MR variety, but not the use of a fungicide, significantly
4 reduced FHB severity compared to the combination of direct sowing, S variety and
5 no fungicide application; on the other hand, all the combinations of two factors in
6 order to prevent FHB symptoms, reduced disease severity significantly more than the
7 worst scenario. In 2008, at site RC, the application of one of the considered factors
8 on its own to prevent FHB significantly reduced the disease severity compared to the
9 worst scenario (direct sowing, S variety, no fungicide application). Moreover, a
10 significant further reduction in FHB severity was obtained with a fungicide application
11 to the MR variety in direct sowing conditions and to the S variety after ploughing,
12 compared to the untreated control, or with an MR variety instead of the S one in the
13 ploughed and untreated plot. In 2008, at site SL, ploughing and fungicide
14 application, but not the adoption of the MR variety, were able to significantly reduce
15 FHB severity compared to the worst scenario (direct sowing, S variety, no fungicide
16 application). In both sites, no significant further reduction was observed, even for the
17 best combination of the three factors.

18

19 3.3. Grain yield

20 Grain yield was affected significantly by the interaction of at least two of the factors
21 compared in the trial conducted in 2006 at the SL site, in 2007 at the RC site and SL
22 site (Table 5). In the other trials, the main effect of at least one of the factors resulted
23 to be significant. Ploughing significantly increased grain yield compared to direct
24 sowing in the trial conducted at the RC site by 17% in 2006 ($P < 0.01$), by 26% in 2007

1 (P<0.01) and by 62% in 2008 (P<0.001), and by 66% in 2008 at the SL site
2 (P<0.001) (Table 6).

3 The MR variety was significantly more productive than the S one at the IM site in
4 2006 (7% more, P<0.05) and in 2007 (9% more, P<0.01) and at the RC site in 2006
5 (11% more, P<0.01).

6 The fungicide application at heading significantly increased the yield at the IM site
7 (P<0.01) and the RC site (P<0.01), by 13% and 4%, respectively in 2006, at the IM
8 site (P<0.01) and the SL site (P<0.01), by 19% and 14%, respectively in 2007, and at
9 the IM site (P<0.01), the RC site (P<0.001) and the SL site (P<0.01), by 14%, 55%
10 and 10%, respectively in 2008.

11 In 2006, at the SL site, the interaction between the three factors was significant
12 (P<0.01): when applied on their own to control FHB, none of the factors was able to
13 increase grain yield compared to the worst situation (direct sowing, S variety, no
14 fungicide application), while a significant increase in yield of 27% was observed for
15 the preventive combination of all the factors (ploughing, MR variety, fungicide
16 application). The interaction between variety and fungicide was significant in 2007 at
17 the RC site (P<0.05): when the fungicide was applied at heading, the S variety
18 showed a significantly higher grain yield than the MR one. On the other hand, no
19 differences were observed in the untreated conditions and the fungicide did not
20 significantly increase the yield in either variety (Table 6). In 2007, at the SL site, the
21 interaction between the tillage and variety was significant (P<0.01): both the
22 ploughing and the use of an MR variety significantly increased grain yield compared
23 to the combination of direct sowing and the S variety, while no significant further
24 increase was observed for the combination of ploughing and the MR variety (Table
25 6).

1 3.4. DON contamination

2 The average DON content was clearly related to the meteorological conditions,
3 particularly close to anthesis, in each year and site. The mean DON contamination
4 was low ($< 100 \mu\text{g kg}^{-1}$) at the SL site in 2006 and at the IM site and the RC site in
5 2007. However, mean DON content was extremely high in 2008 at the RC ($12995 \mu\text{g}$
6 kg^{-1}) and SL ($9310 \mu\text{g kg}^{-1}$) sites. In the other trials, the mean DON contamination
7 was between 262 and $710 \mu\text{g kg}^{-1}$ (Table 8).

8 In all the trials, the DON contamination was significantly affected by the interaction of
9 at least two of the compared factors, while the interaction between all the three
10 factors involved was significant in four trials: in 2006 at SL site, in 2007 at RC site
11 and in 2008 at IM and SL sites (Table 7). In 2006 at the SL site and in 2008 at IM
12 site, when applied on their own to control FHB, all the factors were able to reduce
13 DON contamination compared to the worst scenario (direct sowing, S variety, no
14 fungicide application). Furthermore, no significant further reductions were observed,
15 even for the best combination of the three factors (Table 8). In 2007, at the RC site, a
16 significant difference was only observed between the best (ploughing, MR variety
17 and fungicide application) and the worst scenario (direct sowing, S variety, no
18 fungicide application). In 2008, at the SL site, the ploughing and the fungicide
19 application, but not the MR variety, significantly reduced the DON content compared
20 to the combination of direct sowing, S variety and no fungicide application. On the
21 other hand, all the two-factor combinations to prevent FHB significantly led to a
22 further reduction in DON occurrence. Compared to the best scenario (ploughing, MR
23 variety, fungicide application), direct sowing or the adoption of the S variety
24 significantly increased the DON contamination.

1 The reduction in DON level by means of a factor application (variety, tillage,
2 fungicide) can be expressed by a parameter, efficacy (E), which is defined by the
3 following ratio (Blandino et al., 2011):

$$E(\%) = \left[\frac{(\text{control DON level} - \text{treatment DON level})}{\text{control DON level}} \right] \times 100$$

4
5 On average, the three investigated factors showed a different efficacy in reducing the
6 DON content, when assessed separately in the trials with low ($\text{DON} < 100 \mu\text{g kg}^{-1}$),
7 medium ($100 < \text{DON} < 1250 \mu\text{g kg}^{-1}$) or high ($\text{DON} > 1250 \mu\text{g kg}^{-1}$) FHB pressure
8 (Table 8). The efficacy of the variety for medium and high disease pressure, were
9 higher (77%) than in the trials in which the *Fusarium* infection was low (29%). The
10 efficacy of the fungicides in reducing DON on average decreased moving from low
11 (69%) to high (46%) FHB severity. The efficacy was higher for tillage (75%) for high
12 and medium FHB pressure than trials with a low DON content (63%). As far as
13 previous crop is concerned, the efficacy for tillage after sorghum was 83% (2006 and
14 2007, at site IM) and it was 68% after maize, in the other experiments.

15 The average data of the relative DON content for the different agronomic situations
16 extrapolated from the experiments are reported in Figure 1. The data are expressed
17 in relation to the worst possible case scenario (direct sowing, S variety, no fungicide)
18 in each trial. When applied on its own, the management of the previous crop residue
19 through tillage resulted to be the best agronomic practices to minimize the field
20 contamination of this mycotoxin (-68%) compared to the worst scenario, and this was
21 followed by the use of an MR variety (-61%) and the triazole application at heading (-
22 41%). The effect of the combination of the compared factors to control DON was
23 simulated: the individual efficacies (E) observed for each factor compared to the

1 worst scenario were combined in an additive way. In the new scenarios, obtained by
2 the introduction of a control factor, the DON content was calculated using the
3 following equation:

4

$$5 \quad S_{1,2} = S_1 - (S_1 * E_2 / 100)$$

6

7 where $S_{1,2}$ = DON content in the scenario which applies factors 1 and 2; S_1 = DON
8 content in the scenario which applies factors 1, with the highest efficacy; E_2 = efficacy
9 observed for factor 2 compared to the worst scenario.

10 The simulated results of the combination of each factor were compared with the
11 average effective data observed in field trials. The combinations of avoided risk
12 factors decreased the DON content in a synergistic manner, since the observed data
13 always showed a greater reduction in DON than the simulated ones. The most
14 favourable scenario for DON contamination (ploughing, MR variety, triazole
15 application at heading) reduced the DON content by 97%.

16

4. Discussion

The results of these experiments, conducted over three years characterized by extremely different meteorological trends, confirm the significant link between agronomic practices and FHB infection and DON contamination.

The collected data clearly underline that previous crop residue management through tillage, wheat variety susceptibility and triazole application are important tools that can help growers minimize DON contents in wheat grain.

As proposed by Koch et al. (2006), information on the relative effect of management options on DON contamination can be obtained through a simplified approach that calculates the severity of the relative effect of individual factors. In this study, the severity of the effect of the individual agricultural practices was calculated as follows: the mean DON value of the treatment with the highest DON concentration divided by the mean value of the lowest treatment. The data obtained from our experiments are reported in Table 9 and compared with other data available from literature. In all these studies, the effect of at least two agricultural practices on DON contamination were compared, in naturally-infected field conditions. Based on these data, the main factors that influence DON formation in wheat grain can be put in a ranking order as follows: susceptibility of wheat variety (3.8) \geq the preceding crop (3.1) > soil tillage (2.4) \geq fungicide application at anthesis of wheat (2.3). Thus, DON control in wheat should start in the field, and should first focus on the agronomic factors that influence FHB infection. Above all, conditions such as preceding host crops, especially maize and sorghum, which leave high amount of infected residues in the field, and the cultivation of a susceptible variety contribute to heavy *Fusarium* infections of wheat crops. The ranking order summarized from these first data obtained in non-inoculated

1 trials, need to be confirmed for different environmental and management conditions
2 from those here reported.

3 Our data clearly underline how the efficacy of agronomic practices on controlling
4 DON is affected to a greater extent by the climatic conditions and different
5 meteorological trends could therefore change the order of importance of the involved
6 factors. Variety susceptibility plays a more important role for low or medium disease
7 pressure, while, when high inoculums are present, as observed in our experiment in
8 2008 at the RC and SL sites, the difference between susceptible or moderately
9 resistant varieties may not be significant. A clear interaction between climatic
10 conditions and cultivar was observed concerning the composition of the FHB species
11 on wheat heads in Germany (Klix et al., 2008). Schaafsma et al. (2001) reported a
12 significant interaction between the effect of variety on DON levels and year: no
13 difference was observed among varieties in the years when the meteorological trend
14 was unfavourable for *Fusarium* infection.

15 On the other hand, the no tillage practice, which leaves crop residues unburied,
16 clearly increased the DON contamination in all the trials. The effect of the presence
17 of residues on the soil surface, increased the DON content much more when the
18 climatic conditions were favourable, but not excessive, for inoculum production and
19 spore dispersal. In a previous work (Blandino et al., 2010), it was shown that the
20 effect of maize residue density on DON content is less evident in dry conditions
21 during the susceptible stages of wheat development or with very frequent rainfall,
22 which probably greatly disadvantages or advantages inoculum production,
23 respectively. Lori et al. (2009) reported a significant difference between conventional
24 and no tillage practices, but only when the weather conditions were moderate for
25 FHB.

1 The fungicide treatment was the agricultural practice which showed the greatest
2 grain yield advantage. Since the application of a fungicide from heading to anthesis is
3 associated with yield increases, due to the maintenance of the photosynthetic life of
4 the canopy during grain filling (Ruske et al., 2003), the effect of this agricultural
5 practice on yield was also observed in the trials with low FHB pressure. Moreover,
6 the effect of triazole fungicide application at heading on FHB and DON control has
7 been pointed out in all the trials. The DON contamination was reduced by the
8 fungicide to a greater extent in the trials with low FHB pressure than in those with
9 high *Fusarium* infection. Mesterhazy et al. (2003) achieved a higher efficacy when
10 the fungicides were applied to a moderately resistant cultivar rather than to a
11 susceptible one, while McMullen et al. (2008) reported that the effect of a triazole
12 application in reducing DON doubled when the previous crop was canola, which
13 determines a clearly lower FHB infection, rather than wheat.

14 Overall, when the meteorological trend, particularly around wheat anthesis, does not
15 lead to a high *Fusarium* infection, the DON content in the grains at harvesting is only
16 significantly higher for a combination of several risk factors, while, in these
17 conditions, the presence of an individual risk factor does not increase the
18 contamination to any great extent. On the other hand, for climatic conditions that
19 promote a high production of *Fusarium* inoculum and the consequent fungal infection
20 and development on the wheat ears, the effect of different agronomic scenarios
21 shows a greater impact on the final DON content.

22 Therefore, although the knowledge of the relative importance of the individual factors
23 that influence DON formation is crucial for the development of decision support
24 systems that aim at minimizing DON concentrations in wheat grain, our data have

1 clearly shown a close interaction between the agronomic factors involved in FHB
2 severity and DON content.

3 The collected data clearly underline that a combination of two or more agricultural
4 practices in a integrated multiple management strategy can result in a better control
5 of DON contamination.

6 As far as the maize residue level, fungicide application and cultivar resistance on
7 DON concentrations in spring wheat is concerned, Nita et al. (2006) reported
8 significant positive interactions between the compared agronomic factors in several
9 cases. The highest grain yield and the lowest DON content in spring and winter
10 wheat were also achieved with multiple, rather than single, management strategies
11 by McMullen et al. (2008). Obst et al. (2000), in a 4-year study in Germany,
12 determined four risk factors: (i) maize as the previous crop, (ii) minimum tillage after
13 maize, (iii) use of a moderately or highly susceptible wheat variety and (iv)
14 application of a strobilurin product. In this experiment, an individual risk factor
15 increased the relative risk of DON contamination three-fold, while the combination of
16 four risk factors increased the relative DON risk 56-fold. In France, ARVALIS, use a
17 grid, derived from a 7-year study, to manage wheat lots during harvesting. This grid
18 has 7 DON contamination risk levels, based on 3 combined risk factors: previous
19 crop, tillage and varietal susceptibility (Gourdain, 2008). Comparing the most
20 favourable and the worst DON control scenarios, Koch et al., (2006), reported, in a 2-
21 year trial, which involved the same factors as our experiment, the same reduction in
22 DON content (97%) observed in our work.

23 As far as our data are concerned, only in one of the nine field trials, was the
24 application of the best integrated strategies not able to reduce DON contamination
25 under the present EU admissible maximum levels for common wheat (i.e. 1250 µg

1 kg⁻¹; EC 2006), but, it is important to underline that the experiment was conducted in
2 climatic conditions that were extremely favourable for FHB infection and DON
3 content.

4 Moreover, the data collected on the integrated multiple agronomic strategies clearly
5 confirm the hypothesis advanced by Edward (2004) and Beyer et al. (2006),
6 concerning the fact that the impact of combined risk factors on DON contamination is
7 synergistic rather than additive.

8 In the future, other factors which have proven to have a possible, although often
9 conflicting, effect on DON control in wheat, such as N fertilization (Lemmens et al.,
10 2004), planting date and canopy density (Champeil et al., 2004a), weed control
11 (Jenkinson and Parry, 1994), seed dressing with fungicide (Poels et al., 2006;
12 Campagna and Fusarini, 2010) and biological control (Palazzini, 2007), need to be
13 introduced into this integrated approach and tested to establish their impact on DON
14 content.

15 In short, our results, which were obtained under naturally-infected field conditions,
16 provide useful information to help measure the impact of previous crop residue
17 management practices, variety susceptibility and triazole fungicide application, some
18 of the most important practices adopted to control FHB in wheat, on grain yield and
19 on DON contamination. Since the interaction between the agricultural practices have
20 shown a synergistic effect, integrated multiple strategies, in the areas characterized
21 by a probable FHB infection, would seem to be the best management way of
22 reducing FHB and DON contamination in wheat.

23

1 **5. Acknowledgements**

2 The authors would like to thank Fabio Carnaroglio, Marzia Cattaneo, Maria Corbellini,
3 Mattia Ciro Mancini, Alessandro Peila, Angelo Sarti, and Francesca Vanara for their
4 precious help and cooperation in the laboratory and field work.

5 The research was conducted with the financial support of the Regione Lombardia, as
6 a part of the MICOCER interregional project.

6. References

- Aldred, D., Magan, N., 2004. Prevention strategies for trichothecenes. *Toxicol. Lett.* 153, 165-171.
- Beyer, M., Klix, M.B., Klink, H., Verreet J.-A., 2006. Quantifying the effect of previous crop, tillage, cultivar and triazole fungicides on the deoxynivalenol content of wheat grain – a review. *J. Plant Dis. Prot.* 113, 241-246.
- Blandino, M., Minelli, L., Reyneri, A., 2006. Strategies for the chemical control of *Fusarium* head blight: effect on yield, alveographic parameters and deoxynivalenol contamination in winter wheat grain. *Eur. J. Agron.* 25, 193-201.
- Blandino, M., Pilati, A., Reyneri, A., Scudellari, D., 2010. Effect of maize crop residue density on *Fusarium* head blight and on deoxynivalenol contamination of common wheat grains. *Cereal Res. Commun.*, 38, 550-559.
- Blandino, M., Pascale, M., Haidukowski, M., Reyneri, A., 2011. Influence of agronomic conditions on the efficacy of different fungicides applied to wheat at heading: effect on flag leaf senescence, *Fusarium* head blight attack, grain yield and deoxynivalenol contamination. *Ital. J. Agron.* 4, 204-211.
- Bottalico, A., Perrone, G., 2002. Toxigenic *Fusarium* species and mycotoxins associated with head blight in small-grain cereals in Europe. *Eur. J. Plant Pathol.* 108, 611-624.
- Campagna, C., Fusarini, L., 2010. Contribution of Celest in seed treatment to management of mycotoxins in wheat. *Proceedings of the Giornate Fitopatologiche*, 9-12 March 2010, Cervia (RA) Italy, Vol. 2, pp. 381-386.
- Chala, A., Weinert, J., Wolf, G.A., 2003. An integrated approach to the evaluation of the efficacy of fungicides against *Fusarium culmorum*, the cause of head blight of wheat. *J. Phytopathol.* 151, 673-678.
- Champeil, A., Dore, T., Fourbet, J.F., 2004a. *Fusarium* head blight: epidemiological origin of the effects of cultural practices on head blight attacks and the production of mycotoxins by *Fusarium* in wheat grains. *Plant Sci.* 166, 1389-1415.
- Champeil, A., Fourbet, J.F., Dore, T., Rossignol, L., 2004b. Influence of cropping system on *Fusarium* head blight and mycotoxin levels in winter wheat. *Crop Prot.* 23, 531-537.
- Dill-Macky, R., Jones, R.K., 2000. The effect of previous crop residues and tillage on *Fusarium* head blight of wheat. *Plant Dis.* 84, 71-76.
- EC, 2006. Commission Regulation (EC) No 1881/2006 of 19 December 2006 setting maximum levels for certain contaminants in foodstuffs. *Official Journal of the European Union*, L364, p. 5-24.

- Edwards, S.G., 2004. Influence of agricultural practices on *Fusarium* infection of cereals and subsequent contamination of grain by trichothecene mycotoxins. *Toxicol. Lett.* 153, 29-35.
- Gourdain, E., 2008. DON: Maîtriser le risque sur les cultures de blés: quels outils pour quelles utilisations? In Proc. 2^e Séminaire Mycotoxines des céréales. April 3, 2008, Paris, Pages 27-40.
- Haidukowski, M., Pascale, M., Perrone, G., Pancaldi, D., Campagna, C., Visconti, A., 2005. Effect of fungicides on the development of *Fusarium* head blight, yield and deoxynivalenol accumulation in wheat inoculated under field conditions with *Fusarium graminearum* and *Fusarium culmorum*. *J. Sci. Food Agric.* 85, 191-198.
- Hollins, T.W., Ruckenbauer, P., De Jong, H., 2003. Progress towards wheat varieties with resistance to *Fusarium* head blight. *Food Control* 14, 239-244.
- loos, R., Belhadj, A., Menez, M., Faure, A., 2005. The effect of fungicides on *Fusarium* spp. and *Microdochium nivale* and their associated trichothecene mycotoxins in French naturally-infected cereal grains. *Crop Prot.* 24, 894-902.
- Jenkinson, P., Parry, D.W., 1994. Isolation of *Fusarium* species from common broad-leaved weeds and their pathogenicity to winter wheat. *Mycol. Res.* 98, 776-780.
- Klix, M.B., Beyer M., Verreet, J.-A., 2008. Effect of cultivar, agronomic practices, geographic location, and meteorological conditions on the composition of selected *Fusarium* species on wheat heads. *Can. J. Plant Pathol.* 20, 46-57.
- Koch, H.-J., Pringas, C., Maerlaender, B., 2006. Evaluation of environmental and management effects on *Fusarium* head blight infection and deoxynivalenol concentration in the grain of winter wheat. *Eur. J. Agron.* 24, 357-366.
- Lemmens, M., Haim, K., Lew, H., Ruckenbauer, P., 2004. The effect of nitrogen fertilization on *Fusarium* head blight development and deoxynivalenol contamination in wheat. *J. Phytopathol.* 152, 1-8.
- Lori, G.A., Sisterna, M.N., Sarandon, S.J., Rizzo, I., Chidichimo, H., 2009. *Fusarium* head blight in wheat: impact of tillage and other agronomic practices under natural infection. *Crop Prot.* 28, 495-502.
- Maiorano, A., Blandino, M., Reyneri, A., Vanara, F., 2008. Effects of maize residues on the *Fusarium* spp. infection and deoxynivalenol (DON) contamination of wheat grain. *Crop Prot.* 27, 182-188.
- Mayerle, M., Pancaldi, D., Haidukowski, M., Pascale, M., Ravaglia, S., 2007. Fusariosi e grano tenero: quali sono le varietà più resistenti. *L'Informatore Agrario* 32, 45-49.
- Mesterhazy, A., Bartok, T., Lamper, C., 2003. Influence of wheat cultivar, species of *Fusarium*, and isolate aggressiveness on the efficacy of fungicides for control of *Fusarium* head blight. *Plant Dis.* 87, 1107-1115.

- Mesterházy, A., Bartók, T., Kászonyi, G., Varga, M., Tóth, B., Varga, J., 2005. Common resistance to different *Fusarium* spp. causing Fusarium head blight in wheat. *Eur. J. Plant Pathol.* 112, 267-281.
- McMullen, M., Halley, S., Schatz, B., Meyer, S., Jordahl, J., Ransom, J., 2008. Integrated strategies for Fusarium head blight management in the United States. *Cereal Res. Commun.* 36 (Suppl.B45), 563-568.
- Miller, J.D., Culley, J., Fraser, K., Hubbard, S., Meloche, F., Ouellet, T., Seaman, L., Seifert, K. A., Turkington, K., Voldeng, H., 1998. Effect of tillage practice on Fusarium head blight of wheat. *Can. J. Plant Pathol.* 20, 95-103.
- Neumann, G., Lombaert, G.A., Kotello, S., Fedorowich, N., 2009. Determination of deoxynivalenol in soft wheat by immunoaffinity column cleanup and LC-UV detection: interlaboratory study. *Journal of AOAC International* 92, 181-189.
- Nita, M., Dewolf, E., Madden, L., Paul, P., Shaner, G., Adhikari, T., Ali, S., Stein, J., Osborne L., 2006. Effect of corn residue level, fungicide application and cultivar resistance level on disease incidence and severity of Fusarium head blight and DON concentration. In: *Proc. 2006 National Fusarium Head Blight Forum*, Dec. 10-12, 2006, Research Triangle Park, NC., East Lansing: Michigan State University, Page 49.
- Obst, A., Lepschy, J., Beck, R., Bauer, G., Bechtel, A., 2000. The risk of toxins by *Fusarium graminearum* in wheat – interactions between weather and agronomic factors. *Mycotoxin Res.* 16A, 16-20.
- Palazzini, J.M., Ramirez, M.L., Torres, A.M., Chulze, S.N., 2007. Potential biocontrol agents for Fusarium head blight and deoxynivalenol production in wheat. *Crop Prot.* 26, 1702-1710.
- Parry, D.W., Jenkinson, P., McLeod, L., 1995. Fusarium ear blight (scab) in small grain cereal - Review. *Plant Pathol.* 44, 207-238.
- Paul, P.A., Lipps, P.E., Hershman D.E., McMullen, M.P., Draper, M.A., Madden, L.V., 2008. Efficacy of triazole-based fungicides for Fusarium Head Blight and deoxynivalenol control in wheat: a multivariate meta-analysis. *Phytopathology* 98, 999-1011.
- Pereyra, S.A., Dill-Macky, R., 2008. Colonization of the residues of diverse plant species by *Gibberella zeae* and their contribution to Fusarium Head Blight inoculum. *Plant Dis.* 92, 900-806.
- Pirgozliev, S.R., Edwards, S.G., Hare, M.C., Jenkinson, P., 2003. Strategies for the control of Fusarium head blight in cereals. *Eur. J. Plant Pathol.* 109, 731-742.
- Poels, P., Sztor, E., Cannaert, F., 2006. Fusariosie, elle peut migrer de la semence à l'épi. *Phytoma – La defense des vegetaux*, 593, 9-12.

- Ramirez, M.L., Chulze, S., Magan, N., 2004. Impact of environmental factors and fungicides on growth and deoxynivalenol production by *Fusarium graminearum* isolates from Argentinian wheat. *Crop Prot.* 23, 117-125.
- Ruske, R.E., Gooding, M.J., Jones, S.A., 2003. The effect of adding picoxystrobin, azoxystrobin and nitrogen to a triazole programme on disease control, flag leaf senescence, yield and grain quality of winter wheat. *Crop Prot.* 22, 975-987.
- Schaafsma, A.W., Tamburic-Illinic, L., Miller, J.D., Hooker, D.C., 2001. Agronomic consideration for reducing deoxynivalenol in wheat grain. *Can. J. Plant Pathol.* 23, 279-285.
- Smith-White, J.L., Burgess, L.W., Summerell, B.A., 2004. Sorghum, an intermediate host of the head blight fungus *Gibberella zea*. In: Proc. Third Soilborne Diseases Symposium, Feb. 8-11, 2004, South Australian Research and Development Institute, Adelaide, Pages 81-82.
- Snijders, C.H.A., 2004. Resistance in wheat to *Fusarium* infection and trichothecene formation. *Toxicol. Lett.* 153, 37-46.
- Sutton, J.C., 1982. Epidemiology of wheat head blight and maize ear rot caused by *Fusarium graminearum*. *Can. J. Plant Pathol.* 4, 195-209.
- Tóth, B., Kászonyi, G., Bartók, T., Varga, J., Mesterházy, A., 2008. Common resistance of wheat to members of the *Fusarium graminearum* species complex and *F. culmorum*. *Plant Breeding* 127, 1-8.
- Xu, X.M., 2003. Effects of environmental conditions on the development of *Fusarium* ear blight. *Europ. J. Plant Pathol.* 109, 683-689.
- Zadoks, J.C., Chang, T.T., Konzak, C.F., 1974. A decimal code for the growth stages of cereals. *Weed Res.* 14, 415-421.

Table 1

Main trial information for the field experiments conducted in the 2005-2008 period in 3 sites in North Italy

Growing season	Site	IM	RC	
	Location	Imola (IM)	Riva presso Chieri (TO)	S. Ang
	Geographic coordinates	44° 21' N, 11° 42' E	44° 54' N, 7° 24' E;	45°
	Altitude (m)	21	262	
	Soil ^(a)	Silty-sandy-loamy, Vertic Haplustepts	Loamy, Aquic Frugiudalf	Sandy
2005-2006	Previous crop	Sorghum	Maize	
	Sowing date	22/11/05	28/10/05	
	Date of fungicide application	18/05/06	18/05/06	
	Harvest date	12/07/06	10/07/06	
2006-2007	Previous crop	Sorghum	Maize	
	Sowing date	18/10/06	27/10/06	
	Date of fungicide application	26/04/07	06/05/07	
	Harvest date	21/06/07	28/06/07	
2007-2008	Previous crop	Maize	Maize	
	Sowing date	22/10/07	02/11/07	
	Date of fungicide application	07/05/08	16/05/08	
	Harvest date	07/07/08	15/07/08	

(a) USDA classification

Table. 2

Monthly rainfall and growing degree days (GDD 0s) from March to July 2006-2008 in the research sites.

Year		2006			2007			
Site	Month	Rainfall (mm)	Rainy days (d)	GDD 0s ^a (°C d ⁻¹)	Rainfall (mm)	Rainy days (d)	GDD 0s ^a (°C d ⁻¹)	Rainfall (mm)
IM	March	49	18	256	70	16	317	54
	April	58	18	396	12	5	463	62
	May	54	15	532	63	12	579	70
	June	27	13	641	53	11	650	62
	April - June	188	64	1825	198	44	2008	248
RC	March	17	3	277	29	8	365	20
	April	21	8	430	13	3	509	105
	May	60	9	558	76	14	606	137
	June	19	6	679	107	12	667	97
	April - June	116	26	1944	225	37	2147	359
SL	March	23	5	275	42	8	335	31
	April	65	14	418	13	4	517	77
	May	49	12	565	101	11	603	96
	June	18	5	683	79	12	673	80
	April - June	156	36	1942	235	35	2127	283

^(a) Accumulated growing degree days for each decade using a 0°C base.

Table. 3

Analysis of variance for FHB severity, field experiments conducted at three sites in North Italy over a three years period.

Year	Site	Source of variation	2006			2007			2008		
			P	sem	protected LSD	P	sem	protected LSD	P	sem	protected LSD
IM	Tillage (A)	0.020	0.16		0.012	0.03		0.014	0.18		
	Variety (B)	0.002	0.18		0.003	0.05		0.013	0.28		
	Fungicide (C)	0.031	0.37		0.029	0.07		0.001	0.40		
	A X B	0.003	0.26		0.004	0.07	0.47	0.243	0.40		
	A X C	0.042	0.52		0.065	0.10		0.021	0.56	3.86	
	B X C	0.013	0.23		0.064	0.10		0.284	0.34		
	A X B X C	0.014	0.32	3.35	0.572	0.13		0.937	0.48		
RC	Tillage (A)	0.251	0.82		0.367	1.86		0.002	0.99		
	Variety (B)	< 0.001	0.93		0.208	2.31		< 0.001	1.66		
	Fungicide (C)	< 0.001	0.79		< 0.001	1.79		< 0.001	1.12		
	A X B	0.629	1.32		0.611	3.27		0.038	2.34		
	A X C	0.938	1.12		0.230	2.53		< 0.001	1.59		
	B X C	0.001	1.17	6.37	0.793	2.10		< 0.001	1.40		
	A X B X C	0.923	1.65		0.653	2.96		0.008	1.98	14.89	
SL	Tillage (A)	0.004	0.13		0.018	0.87		0.012	1.72		
	Variety (B)	< 0.001	0.12		0.001	1.55		0.751	2.27		
	Fungicide (C)	< 0.001	0.10		0.015	1.07		0.000	0.67		
	A X B	0.001	0.18	1.21	0.008	2.19		0.155	3.21		
	A X C	< 0.001	0.14	0.98	0.005	1.51		< 0.001	0.94		
	B X C	< 0.001	0.15	1.02	0.127	0.87		0.022	0.90		
	A X B X C	0.536	0.21		0.001	1.23	12.86	0.002	1.27	13.25	

The data reported in the table refer to the level of significance (P) and the standard error of mean (sem).

When interactions between factors are significant, the Fisher's Least Significant Difference (LSD), protected by Bonferroni at $P \leq 0.05$, is reported.

Table. 4

Effect of tillage, variety susceptibility to FHB and triazole fungicide application on FHB severity of winter wheat (%); field experiments conducted at three sites in North Italy over a three years period.

Site	Tillage	Fungicide	2006			2007			2008		
			Variety			Variety			Variety		
			S	MR	Mean	S	MR	Mean	S	MR	Mean
IM	Direct sowing	Untreated	3.51	0.81	2.16	0.80	0.23	0.52	5.55	4.21	4.88
		Fungicide	0.95	0.17	0.56	0.35	0.08	0.22	1.70	0.81	1.26
		Mean	2.23	0.49	1.36	0.58	0.16	0.37	3.63	2.51	3.07
	Ploughing	Untreated	0.31	0.21	0.26	0.17	0.06	0.11	2.73	1.97	2.35
		Fungicide	0.21	0.15	0.18	0.03	0.11	0.07	0.99	0.61	0.80
		Mean	0.26	0.18	0.22	0.10	0.09	0.09	1.86	1.29	1.58
	Tillage mean	Untreated	1.91	0.51	1.21	0.48	0.15	0.31	4.14	3.09	3.62
		Fungicide	0.58	0.16	0.37	0.19	0.10	0.14	1.35	0.71	1.03
		Mean	1.25	0.34		0.34	0.12		2.74	1.90	
RC	Direct sowing	Untreated	10.59	2.28	6.43	14.91	14.38	14.65	55.72	20.09	37.91
		Fungicide	3.35	0.73	2.04	4.36	2.55	3.46	9.39	3.89	6.64
		Mean	6.97	1.51	4.24	9.64	8.47	9.05	32.55	11.99	22.27
	Ploughing	Untreated	9.38	1.53	5.45	13.05	10.28	11.66	34.72	8.66	21.69
		Fungicide	2.12	0.14	1.13	4.44	2.01	3.23	8.07	3.14	5.61
		Mean	5.75	0.84	3.29	8.75	6.14	7.44	21.39	5.90	13.65
	Tillage mean	Untreated	9.98	1.91	5.94	13.98	12.33	13.16	45.22	14.38	29.80
		Fungicide	2.74	0.44	1.59	4.40	2.28	3.34	8.73	3.52	6.12
		Mean	6.36	1.17		9.19	7.31		26.97	8.95	
SL	Direct sowing	Untreated	7.85	3.63	5.74	26.55	7.85	17.20	30.71	37.96	34.34
		Fungicide	3.21	0.77	1.99	14.73	5.31	10.02	13.72	11.00	12.36
		Mean	5.53	2.20	3.87	20.64	6.58	13.61	22.22	24.48	23.35
	Ploughing	Untreated	4.23	1.75	2.99	10.95	6.44	8.70	13.84	8.76	11.30
		Fungicide	1.34	0.43	0.89	6.91	4.55	5.73	5.32	3.69	4.51
		Mean	2.79	1.09	1.94	8.93	5.50	7.22	9.58	6.22	7.90
	Tillage mean	Untreated	6.04	2.69	4.37	18.75	7.15	12.95	22.28	23.36	22.82
		Fungicide	2.27	0.60	1.44	10.82	4.93	7.87	9.52	7.35	8.43
		Mean	4.16	1.65		14.79	6.04		15.90	15.35	

Significance for the differences of the compared means are reported in Table 3.

FHB severity was calculated as the percentage of spikelets per ear with symptoms of disease at the soft dough stages (GS 85).

Tillage: the previous crop was grain sorghum at site IM (years 2006 and 2007) and grain maize at site IM (2008), RC and SL.

Variety: S = susceptible to FHB, MR = moderately resistant to FHB

Fungicide: metconazole was applied at 0.06 kg active ingredient (AI) ha⁻¹ at wheat heading.

Table. 5

Analysis of variance for grain yield, field experiments conducted at three sites in North Italy over a three years period.

Site	Source of variation	2006			2007			2008		
		P	sem	protected LSD	P	sem	protected LSD	P	sem	protected LSD
IM	Tillage (A)	0.062	0.32		0.160	0.20		0.740	0.12	
	Variety (B)	0.039	0.26		0.001	0.22		0.566	0.12	
	Fungicide (C)	0.002	0.19		0.006	0.23		0.003	0.22	
	A X B	0.583	0.37		0.514	0.31		0.621	0.16	
	A X C	0.354	0.27		0.667	0.32		0.234	0.31	
	B X C	0.143	0.38		0.380	0.53		0.204	0.26	
	A X B X C	0.428	0.53		0.864	0.76		0.083	0.37	
	RC	Tillage (A)	0.003	0.17		0.004	0.19		< 0.001	0.10
Variety (B)		0.001	0.26		0.002	0.15		0.282	0.33	
Fungicide (C)		0.009	0.13		0.043	0.17		0.000	0.25	
A X B		0.123	0.37		0.452	0.21		0.160	0.47	
A X C		0.883	0.18		0.786	0.24		0.192	0.36	
B X C		0.710	0.23		0.028	0.19	0.53	0.168	0.29	
A X B X C		0.457	0.32		0.986	0.27		0.980	0.41	
SL		Tillage (A)	0.354	0.25		0.024	0.12		< 0.001	0.05
	Variety (B)	< 0.001	0.11		< 0.001	0.08		0.289	0.09	
	Fungicide (C)	0.011	0.12		0.003	0.18		0.004	0.15	
	A X B	0.048	0.16		0.002	0.12	0.80	0.068	0.12	
	A X C	0.033	0.16		0.664	0.26		0.658	0.21	
	B X C	0.194	0.12		0.124	0.21		0.634	0.21	
	A X B X C	0.038	0.17	1.56	0.153	0.30		0.447	0.29	

The data reported in the table refer to the level of significance (P) and the standard error of mean (sem).

When interactions between factors are significant, the Fisher's Least Significant Difference (LSD), protected by Bonferroni at $P \leq 0.05$, is reported.

Table. 6

Effect of tillage, variety susceptibility to FHB and triazole fungicide application on grain yield of winter wheat ($t\ ha^{-1}$); field experiments conducted at three sites in North Italy over a three years period.

Site	Tillage	Fungicide	2006			2007			2008		
			Variety			Variety			Variety		
			S	MR	Mean	S	MR	Mean	S	MR	Mean
IM	Direct sowing	Untreated	6.77	7.62	7.20	6.65	8.34	7.49	6.90	7.50	7.20
		Fungicide	7.81	8.30	8.06	7.78	9.03	8.40	8.19	7.79	7.99
		Mean	7.29	7.96	7.63	7.21	8.68	7.95	7.55	7.64	7.59
	Ploughing	Untreated	7.81	8.77	8.29	6.36	7.91	7.14	7.08	6.98	7.03
		Fungicide	9.46	9.40	9.43	7.43	8.36	7.90	8.20	8.31	8.25
		Mean	8.64	9.09	8.86	6.89	8.14	7.52	7.64	7.64	7.64
	Tillage mean	Untreated	7.29	8.20	7.75	6.50	8.12	7.31	6.99	7.24	7.11
		Fungicide	8.64	8.85	8.74	7.60	8.70	8.15	8.19	8.05	8.12
		Mean	7.96	8.52	8.24	7.05	8.41		7.59	7.64	
RC	Direct sowing	Untreated	6.24	7.26	6.75	4.92	4.61	4.77	1.86	2.59	2.23
		Fungicide	6.42	7.66	7.04	5.42	4.67	5.05	4.11	4.46	4.29
		Mean	6.33	7.46	6.89	5.17	4.64	4.91	2.99	3.53	3.26
	Ploughing	Untreated	7.63	8.25	7.94	6.15	5.98	6.07	4.40	4.50	4.45
		Fungicide	7.94	8.48	8.21	6.60	5.99	6.29	6.21	5.95	6.08
		Mean	7.78	8.37	8.08	6.38	5.99	6.18	5.31	5.23	5.27
	Tillage mean	Untreated	6.93	7.76	7.34	5.54	5.30	5.42	3.13	3.54	3.34
		Fungicide	7.18	8.07	7.62	6.01	5.33	5.67	5.16	5.21	5.18
		Mean	7.06	7.91		5.77	5.31		4.15	4.38	
SL	Direct sowing	Untreated	5.87	6.49	6.18	4.12	6.04	5.08	3.69	3.75	3.72
		Fungicide	6.22	7.39	6.80	5.44	6.58	6.01	4.41	4.19	4.30
		Mean	6.05	6.94	6.49	4.78	6.31	5.55	4.05	3.97	4.01
	Ploughing	Untreated	6.03	7.46	6.74	5.57	6.25	5.91	6.33	6.53	6.43
		Fungicide	6.22	7.48	6.85	6.39	7.04	6.71	6.98	7.24	7.11
		Mean	6.12	7.47	6.80	5.98	6.64	6.31	6.66	6.88	6.77
	Tillage mean	Untreated	5.95	6.98	6.46	4.84	6.15	5.49	5.01	5.14	5.07
		Fungicide	6.22	7.43	6.83	5.92	6.81	6.36	5.70	5.72	5.71
		Mean	6.08	7.20		5.38	6.48		5.35	5.43	

Significance for the differences of the compared means are reported in Table 5.

Tillage: the previous crop was grain sorghum at site IM (years 2006 and 2007) and grain maize at site IM (2008), RC and SL.

Variety: S = susceptible to FHB, MR = moderately resistant to FHB

Fungicide: metconazole was applied at $0.06\ kg\ active\ ingredient\ (AI)\ ha^{-1}$ at wheat heading.

Table. 7

Analysis of variance for DON content, field experiments conducted at three sites in North Italy over a three years period.

Site	Source of variation	2006			2007			2008		
		P	sem	protected LSD	P	sem	protected LSD	P	sem	protected LSD
IM	Tillage (A)	0.004	56		0.043	23		0.015	54	
	Variety (B)	0.001	112		0.001	19		< 0.001	20	
	Fungicide (C)	0.028	47		0.004	28		0.001	38	
	A X B	0.002	159	1091	0.013	27	184	< 0.001	28	
	A X C	0.091	66		0.028	39	218	0.001	54	
	B X C	0.108	19		0.016	39	219	0.003	38	
	A X B X C	0.521	27		0.204	55		0.005	54	569
RC	Tillage (A)	< 0.001	39		0.001	5		< 0.001	1359	
	Variety (B)	< 0.001	56		0.008	42		< 0.001	1001	
	Fungicide (C)	< 0.001	67		0.001	16		< 0.001	822	
	A X B	< 0.001	79	434	0.091	60		0.428	1416	
	A X C	0.054	95		0.028	22		< 0.001	1163	6354
	B X C	< 0.001	97	528	0.029	22		< 0.001	623	3404
	A X B X C	0.838	137		0.032	32	238	0.819	881	
SL	Tillage (A)	0.018	5		< 0.001	5		< 0.001	120	
	Variety (B)	< 0.001	7		0.001	65		0.001	489	
	Fungicide (C)	0.001	5		0.001	36		0.005	1262	
	A X B	0.002	9		0.044	91	626	0.359	691	
	A X C	0.003	7		0.018	51	352	0.082	1784	
	B X C	0.001	7		0.601	150		0.002	125	
	A X B X C	0.003	11	110	0.606	212		< 0.001	176	1845

The data reported in the table refer to the level of significance (P) and the standard error of mean (sem).

When interactions between factors are significant, the Fisher's Least Significant Difference (LSD), protected by Bonferroni at $P \leq 0.05$, is reported.

Table. 8

Effect of tillage, variety susceptibility to FHB and triazole fungicide application on DON contamination of winter wheat ($\mu\text{g kg}^{-1}$); field experiments conducted at three sites in North Italy over a three years period.

Site	Tillage	Fungicide	2006			2007			2008			
			Variety			Variety			Variety			
			S	MR	Mean	S	MR	Mean	S	MR	Mean	
IM	Direct sowing	Untreated	1708	407	1057	374	96	235	1073	369	721	
		Fungicide	1508	233	871	85	30	57	364	113	238	
		Mean	1608	320	964	230	63	146	718	241	480	
	Ploughing	Untreated	181	37	109	115	n.d.	63	86	37	62	
		Fungicide	116	24	70	n.d.	n.d.	10	38	n.d.	24	
		Mean	149	31	90	63	n.d.	36	62	24	43	
	Tillage mean	Untreated	945	222	583	245	53	149	580	203	391	
		Fungicide	812	129	471	47	20	34	201	61	131	
		Mean	878	175		146	37		390	132		
	RC	Direct sowing	Untreated	2552	211	1381	261	66	163	27223	22338	24781
			Fungicide	1113	27	570	130	38	84	14392	12294	13343
			Mean	1832	119	976	195	52	124	20808	17316	19062
Ploughing		Untreated	1491	21	756	93	47	70	12951	7206	10078	
		Fungicide	255	n.d.	132	64	20	42	5379	2177	3778	
		Mean	873	15	444	78	33	56	9165	4691	6928	
Tillage mean		Untreated	2021	116	1069	177	56	117	20087	14772	17429	
		Fungicide	684	18	351	97	29	63	9886	7235	8561	
		Mean	1353	67		137	43		14986	11004		
SL		Direct sowing	Untreated	167	n.d.	88	1103	547	825	19804	18163	18983
			Fungicide	47	n.d.	28	768	213	490	14885	9135	12010
			Mean	107	n.d.	58	935	380	658	17345	13649	15497
	Ploughing	Untreated	50	n.d.	30	446	71	258	6513	2440	4477	
		Fungicide	29	n.d.	20	225	18	122	2560	673	1616	
		Mean	40	n.d.	25	335	45	190	4537	1556	3046	
	Tillage mean	Untreated	108	n.d.	59	774	309	542	13159	10301	11730	
		Fungicide	38	n.d.	24	496	116	306	8723	4904	6813	
		Mean	73	n.d.		635	212		10941	7603		

Significance for the differences of the compared means are reported in Table 7.

Tillage: the previous crop was grain sorghum at site IM (years 2006 and 2007) and grain maize at site IM (2008), RC and SL.

Variety: S = susceptible to FHB, MR = moderately resistant to FHB

Fungicide: metconazole was applied at 0.06 kg active ingredient (AI) ha^{-1} at wheat heading.

nd: not detected. The detection limit was 20 $\mu\text{g kg}^{-1}$.

1 **Table. 9**
 2 Severity of the effect of several agricultural practices on deoxynivalenol (DON)
 3 contamination in winter wheat grain in natural conditions and on average of the other
 4 experimental factors included.

Country	Factor	Value in			Reference
		Numerator	Denominator	Severity of effect	
Italy	Tillage (after maize or sorghum)	Direct sowing	Ploughing (30 cm)	4.8	Data reported in the present manuscript
	Variety	Highly susceptible	Moderately resistant	5.5	
	Fungicide application	Without	With	2.4	
USA (MN)	Preceding crop	Maize	Soybean	2.0	Dill-Macky and Jones, 2000
	Tillage	Direct sowing	Ploughing (30 cm)	1.4	
Canada (ON) ^(a)	Preceding crop	Maize	Soybean	2.9	Schaafsma et al., 2001
	Tillage ^(b)	Minimum tillage	Ploughing (30 cm)	1.6	
	Variety	Highly susceptible	Moderately resistant	3.7	
Germany ^(a)	Preceding crop	Winter wheat	Sugar beet	4.3	Koch et al., 2006
	Tillage ^(b)	Direct sowing	Ploughing (30 cm)	2.7	
	Variety	Highly susceptible	Moderately resistant	4.3	
Germany	Tillage (after wheat)	Direct sowing	Ploughing (30 cm)	3.4	Koch et al., 2006
	Variety	Highly susceptible	Moderately resistant	5.6	
	Fungicide application	Without	With	2.1	
Italy	Variety	Highly susceptible	Moderately resistant	1.7	Blandino et al., 2006
	Fungicide application	Without	With	2.4	
Argentina	Tillage (after maize)	Direct sowing	Ploughing (30 cm)	1.3	Lori et al., 2009
	Variety	Highly susceptible	Moderately resistant	1.7	


5
 6 ^(a) Average data obtained from commercial wheat farm fields

7 ^(b) From commercial wheat farm fields with different previous crop

8

9

1 **Figure 1.**
 2 Effect of different agronomic scenarios, obtained from the combination of tillage, variety
 3 susceptibility and fungicide application, on the relative DON content.



4
 5 Reductions are expressed in relation to the worst case scenario (direct sowing, S variety, no fungicide =
 6 100% DON content).
 7 The reported data are the average of the relative DON content of 9 field experiments, expressed in relation
 8 to the worst case scenario in each trial.
 9 The simulations of the combined effect of factors were obtained from the additive
 10 computation of the effect of a single factor in relation to the worst case scenario .