

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Calcium signals and FGF-2 induced neurite growth in cultured parasympathetic neurons: spatial localization and mechanisms of activation**

**This is a pre print version of the following article:**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/127631> since 2021-04-10T11:28:40Z

*Published version:*

DOI:10.1007/s00424-013-1257-5

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

*The final publication is available at Springer via <http://dx.doi.org/10.1007/s00424-013-1257-5>*

**Calcium signals and FGF-2 induced neurite growth in cultured parasympathetic neurons:  
spatial localization and mechanisms of activation**

P. Zamburlin<sup>1\*</sup>, F. A. Ruffinatti<sup>1,2\*</sup>, A. Gilardino<sup>1,2</sup>, S. Farcito<sup>1</sup>, M. Parrini<sup>1#</sup>, D. Lovisolo<sup>1,2,3</sup>

<sup>1</sup> Department of Life Sciences and Systems Biology, University of Torino, Torino, I-10123 Italy

<sup>2</sup> NIS Centre of Excellence, University of Torino, Italy

<sup>3</sup> Neuroscience Institute of Torino, University of Torino, Italy

\* these Authors contributed equally to this work

# M. Parrini present address is at the Department of Neuroscience and Brain Technologies, Italian Institute of Technology, Genova, Italy.

**Address correspondence to:**

Davide Lovisolo

Department of Life Sciences and Systems Biology

University of Torino

Via Accademia Albertina 13, 10123 Torino, Italy

**[davide.lovisolo@unito.it](mailto:davide.lovisolo@unito.it)**

**Keywords:**

Calcium channels, neurotrophic factor, FGF-2, neurite growth, wavelet analysis

## Abstract

1  
2  
3  
4 The growth of neuritic processes in developing neurons is tightly controlled by a wide set of  
5 extracellular cues, that act by initiating downstream signalling cascades, where calcium signals play  
6 a major role. Here we analyze the calcium dependence of the neurite growth promoted by basic  
7 Fibroblast Growth Factor (bFGF, or FGF-2) in chick embryonic ciliary ganglion neurons, taking  
8 advantage of dissociated, organotypic and compartmentalized cultures. We report that signals at  
9 both the growth cone and the soma are involved in the promotion of neurite growth by the factor.  
10 Blocking calcium influx through L- and N-type voltage dependent calcium channels and TRPC  
11 channels reduces, while release from intracellular stores does not significantly affect, the growth of  
12 neuritic processes. Simultaneous recordings of calcium signals elicited by FGF-2 at the soma and at  
13 the growth cone show that the factor activates different patterns of responses in the two  
14 compartments: steady and sustained responses at the former, oscillations at the latter. At the soma,  
15 both voltage dependent channel and TRPC blockers strongly affect steady state levels. At the  
16 growth cone, the changes in the oscillatory pattern are more complex; therefore we used a tool  
17 based on wavelet analysis to obtain a quantitative evaluation of the effects of the two classes of  
18 blockers. We report that the oscillatory behaviour at the growth cone is dramatically affected by all  
19 the blockers, pointing to a role for calcium influx through the two classes of channels in the  
20 generation of signals at the leading edge of the elongating neurites.  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Introduction

1  
2  
3  
4  
5 In the developing nervous system, the oriented growth of neuritic processes is a crucial step in the  
6  
7 establishment of the correct neuronal connections; it is a highly regulated process, under the tight  
8  
9 control of a wide array of extracellular cues [12, 29]. Neurotrophic factors play a prominent role in  
10  
11 this context: they can either promote elongation [24, 57, 60] or act as a stop signal and induce  
12  
13 branching (see e.g. Refs. [9, 50]), but above all they act as attractive or repulsive agents, influencing  
14  
15 not only growth but orientation [31, 45, 56]. Calcium signalling is a key player in these processes  
16  
17 [18, 21, 26, 62], and the different responses to extracellular signals may depend on the specific  
18  
19 neuronal type, but also reflect different spatiotemporal patterns of calcium signalling, in turn  
20  
21 supported by different calcium mobilizing mechanisms. The growth cone is the leading edge of the  
22  
23 extending neurite that senses the extracellular environment. Its behaviour and adaptation to the  
24  
25 changing milieu are finely regulated by local changes in  $[Ca^{2+}]_i$  (see e.g. Ref [18]), and, at least in  
26  
27 part, in the expression of membrane and cytoskeletal proteins [32, 36], but the integration and  
28  
29 crosstalk between signalling mechanisms at the periphery and at the soma are not completely  
30  
31 understood. In particular, while evidence of retrograde signalling of transcription factors to the  
32  
33 soma is available (see e.g. Ref. [10]), and a pioneering paper [7] suggested that the presence of the  
34  
35 NGF at the soma was not necessary for neurite elongation, the role of somatic signals has not been  
36  
37 extensively investigated to date.

38  
39 Basic Fibroblast Growth Factor (FGF-2, or bFGF) is one of the best characterized neurotrophic  
40  
41 factors involved in the control of neurite growth and orientation [41, 53]. In a well established  
42  
43 model of peripheral neurons, cultured E7 chick ciliary ganglion neurons, we have shown that it  
44  
45 induces long lasting calcium influx, dependent on the tyrosine kinase activity of FGFRs [13], and  
46  
47 promotes neuritic growth by increasing the mean velocity of elongation of individual neurites [60].  
48  
49 The prototypical receptor for this agonist, FGFR1, is expressed in these neurons both in the somatic  
50  
51 and in the neuritic compartment [16]. The purpose of this paper was to investigate the role of the  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 calcium signals activated by FGF-2 in the promotion of neurite growth, the specific contribution of  
2 signals at the soma and at the growth cone, and the calcium mobilizing mechanisms involved. To  
3  
4 this end, taking advantage of compartmentalized cultures and of calcium imaging, we provide  
5  
6 evidence that while the presence of FGF-2 at the soma plays a minor role, signalling at the growth  
7  
8 cone is the main event responsible for the promotion of neurite outgrowth. On the other hand both  
9  
10 at the growth cone and at the soma, calcium influx is involved in this action, through voltage  
11  
12 dependent and TRPC (Transient Receptor Potential Canonical) channels. At the growth cone, the  
13  
14 factor elicits a peculiar pattern of oscillations of  $[Ca^{2+}]_i$ , that can be strongly reduced by  
15  
16 pharmacological blockers of the two channel families, suggesting a relationship between the onset  
17  
18 of calcium oscillations and the neurite growth promoting action of FGF-2.  
19  
20  
21  
22  
23  
24  
25  
26  
27

## 28 **Methods**

### 29 *Dissociated cultures*

30  
31  
32  
33  
34 Chick ciliary ganglia were dissected from E7 embryos and maintained in a chemically defined N2  
35  
36 medium [5] as previously described [13]. Briefly, ganglia were both enzymatically (0.06% trypsin,  
37  
38 in cation-free phosphate-buffered saline (PBS), for 5 min at 37°C) and mechanically dissociated  
39  
40 and resuspended in N2 medium. Nearly 15,000 cells were plated in the middle area of plastic dishes  
41  
42 or glass coverslips coated with poly-D-lysine (PL; 100 µg/ml) and laminin (LN; 2 µg/cm<sup>2</sup>) in N2  
43  
44 medium, alone or with added FGF-2 (20 ng/ml, Alomone Labs, Israel).  
45  
46  
47  
48  
49  
50  
51  
52

### 53 *Organotypic cultures*

54  
55 Chick ciliary ganglia were obtained from E7 embryos and collected in PBS. Explants were cultured  
56  
57 in rat tail collagen gel as previously reported [60] and plated in 12-multiwell plates previously  
58  
59 treated with PL and LN. Briefly, ganglia were taken one by one in a small drop of PBS with a  
60  
61  
62  
63  
64  
65

1 Pasteur pipette and placed in the middle of an empty well with as little PBS as possible. A 15  $\mu$ l  
2 drop of collagen gel mix was added on the top of each ganglion and kept at 37°C in a CO<sub>2</sub> incubator  
3  
4 for 40 min until the gel had polymerized. Subsequently 1 ml of N2 medium was added to each well,  
5  
6 alone or with added 20 ng/ml FGF-2.  
7  
8  
9

### 10 11 *Immunocytochemistry*

12 For immunocytochemical experiments, dissociated cells were fixed in paraformaldehyde (PFA) 4%  
13  
14 for 20 min at room temperature, whereas CG explants were fixed in methanol at -20°C for 10 min.  
15  
16 For neurite outgrowth assays, both dissociated CG neurons and CG explants were stained with a  
17  
18 monoclonal anti-NF68 antibody to take into account only the contribution of neurons. Chick CG  
19  
20 neurons and CG explants fixed, respectively, after 24 and 48 hr of culture were both incubated  
21  
22 overnight with the anti-NF68 antibody (1:300) at room temperature. On the following day,  
23  
24 dissociated neurons were incubated with an anti-mouse IgG conjugated with the Cy3 fluorophore  
25  
26 (1:1,000). In the case of organotypical cultures, the reaction was developed with a horse  
27  
28 biotinylated antimouse IgG (Vector Laboratories, Burlingame, CA; 1:200) detected with the  
29  
30 peroxidase-conjugated biotin-avidin complex (Vector Laboratories), followed by a brief incubation  
31  
32 with 3,30-diaminobenzidine tetrahydrochloride (DAB; 15 mg/ml) and H<sub>2</sub>O<sub>2</sub> (8  $\mu$ g/ml).  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42

### 43 *Assessment of cell survival*

44 Cell survival assays were performed on dissociated cultures. Cells were plated on 60-mm-  
45  
46 diameter/2-mm-grid culture dishes (Corning Inc., Corning, NY), and two square fields of the grid in  
47  
48 each dish were recorded by means of a CoolSNAP-Pro-color CCD camera (Roper Scientific/  
49  
50 Photometrics, Germany) connected to an inverted microscope (Olympus, IX50). Survival levels  
51  
52 were assessed by counting neurons, in the same field, after 24 and 48 h of culture, and giving the  
53  
54 values as percentages of the counts at 2 h.  
55  
56  
57  
58  
59

60 Three independent experiments, in duplicate, were performed for each individual treatment.  
61  
62  
63  
64  
65

## 1 2 *Assessment of neurite outgrowth*

3  
4 In dissociated cultures, after immunocytochemical staining, neurite outgrowth was assessed by  
5  
6 measuring the total area occupied by neurofilament-positive elements using the image processing  
7  
8 program ImageJ [38] as previously described [60]. Briefly, for dissociated CG cultures, eight to ten  
9  
10 fields ( $1,024 \times 1,024$  pixels, 16-bit gray scale) per culture dish were acquired by means of a  
11  
12 Fluoview 200 laser scanning confocal microscope (Olympus) with a  $50\times$  oil immersion objective, at  
13  
14 an excitation wavelength of 568 nm. Fields were selected by observing the DIC (Differential  
15  
16 Interference Contrast) images and randomly choosing non overlapping fields. Contribution of  
17  
18 neuronal somata was manually subtracted and the analysis was performed by applying to 16-bit  
19  
20 images a threshold in order to obtain binary images (black pixels corresponding to neurites and  
21  
22 white pixels to background). Total number of neurons was evaluated from DIC images of the same  
23  
24 fields analyzed, and these values were used to normalize the total area occupied by black pixels  
25  
26 ( $\mu\text{m}^2$ ). Performing analysis on fluorescent images allowed to exclude the potential contribution of  
27  
28 glial cell processes, without substantially losing any contribution from the neurites, as can be seen  
29  
30 merging a DIC image with a fluorescent one of the same field (Suppl. Fig. 1).

31  
32 As in previous experiments [16] analysis was performed on cultures at 24 h, since for longer times  
33  
34 extensive neurite fasciculation could lead to incorrect evaluation of global neurite occupancy.

35  
36 Experiments were repeated three to four times in duplicate.

37  
38 In the case of CG explants images were acquired in bright field mode with an inverted microscope  
39  
40 Olympus IX50 equipped with a CoolSNAP-Pro CCD camera (Roper Scientific/Photometrics,  
41  
42 Germany) and an image analysis software (Image Pro Plus 4.1 for Windows; Media Cybernetics).

43  
44 After subtraction of the area corresponding to the body of the whole explant, previously  
45  
46 reconstructed from a collage of partial images ( $1,392 \times 1,040$  pixels, 8-bit gray scale), binary  
47  
48 images were obtained again by applying a threshold. The percentage of black square pixels over the  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



1 total area ( $2,000 \times 2,000$  pixels) was used as an index of global neurite growth. As previously  
2 described [16, 60] analysis was performed at 48 h of culture.  
3

4 Data are the average of three to six independent experiments, and five to twelve explants per  
5 condition were analyzed.  
6  
7

### 8 9 10 11 *Compartmentalized cultures*

12 In order to assess the relative contribution of FGF-2 signalling and of the downstream calcium  
13 fluxes at the somata and at the growth cone of the neurites, we performed a set of experiments with  
14 compartmentalized cultures, using Campenot chambers [7, 8] that allow to present FGF-2 (20  
15 ng/ml) and the inhibitors to the cells in one or more compartments. Briefly, the plastic dishes were  
16 coated with 100  $\mu\text{g/ml}$  PL and 2  $\mu\text{g/cm}^2$  LN, and 30,000 cells (obtained from E7 embryos) were  
17 plated in the central part of a three compartment chamber (Tyler Research, Canada), sealed with  
18 silicone grease to the bottom of the dish. To allow cells to adhere, chambers were kept for two  
19 hours at 37°C in a humidified atmosphere containing 5% CO<sub>2</sub>. Afterwards, specific volumes of N2  
20 medium supplemented with B27 (dilution 1:50; Gibco, Invitrogen Corporation, Carlsbad, CA,  
21 USA) were added into both central and side compartments.  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37

38 Some Authors have used the average neurite length or the length of the longest neurites to quantify  
39 neurite growth (see e.g. Ref. [37, 54]). However, considering neurite length may not give the  
40 correct information on the effects of FGF-2 signalling on the global outgrowth of neuritic processes.  
41  
42  
43  
44  
45

46 Others have measured the global axonal density by means of Western Blot assessment of beta-  
47 tubulin levels [47]. We chose a different approach: in analogy with the protocols used to evaluate  
48 global neurite occupancy in dissociated and organotypic cultures, we measured the total area  
49 covered by neurites in each single track, by staining neurites with the anti-neurofilament antibody.  
50  
51  
52  
53  
54

55 Staining and acquisition protocols were the same as for organotypic cultures.  
56  
57

58 Three independent experiments (in duplicate) were performed for each treatment  
59  
60  
61  
62  
63  
64  
65

### *Calcium imaging*

1  
2 [Ca<sup>2+</sup>]<sub>i</sub> was monitored using the Ca<sup>2+</sup> indicator dye FURA-2 acetoxymethylester (FURA-2AM,  
3  
4 Molecular Probes, Inc.). Cells were loaded for 45 min at 37°C with 2 μM FURA-2AM in N2  
5  
6 medium and subsequently washed in Tyrode Standard solution (NaCl 154 mM, KCl 4 mM,  
7  
8 CaCl<sub>2</sub> 2 mM, MgCl<sub>2</sub> 1 mM, HEPES 5 mM, glucose 5.5 mM, pH 7.34 with NaOH). After dye  
9  
10 loading cells were transferred to a perfusion chamber (Bioptechs, USA) connected to a  
11  
12 peristaltic pump and mounted on an inverted fluorescence microscope (Nikon TE-2000-S), a Xenon  
13  
14 lamp illumination system and a CoolSNAP CCD camera (Roper Scientific/Photometrics,  
15  
16 Germany). All experiments were performed at 37°C. A gravity microperfusion system, regulated by  
17  
18 electrovalves, was employed to switch from the standard Tyrode solution to the same medium  
19  
20 containing human recombinant FGF-2 (40 ng/ml) either with or without the inhibitors. Calcium  
21  
22 measurements were performed exciting FURA-2AM for 0.5 s alternatively at 340 nm and 380 nm,  
23  
24 with a dark interval of 1 s, and emission was recorded at 510 nm. Images were visualized on a  
25  
26 computer with the dedicated acquisition software Metafluor (Universal Imaging Corporation, PA).  
27  
28  
29 In experiments performed at 24 h of culture, a 20× objective was used, and ROIs (Region of  
30  
31 Interest) corresponding to the cell bodies were chosen. For simultaneous recordings from the soma  
32  
33 and the growth cone, experiments were performed after 6 h of culture, when neurite extension was  
34  
35 still limited and all the compartments could be observed by means of a 100× objective. One or at  
36  
37 most two cells per dish could be recorded. Since at this short culture time cells could still be  
38  
39 recovering from the dissociation procedure and some perturbation in the membrane could be  
40  
41 expected, all cells were challenged with 40 mM KCl and non responsive cells were discarded.  
42  
43  
44 Raw data were analysed with IGOR Pro software: peak amplitudes of somatic signals are given as  
45  
46  $\Delta R/R$  (increase in 340/380 ratio over the basal value).  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57

### *Wavelet analysis of calcium signals*

58  
59  
60  
61  
62  
63  
64  
65

1 To give a quantitative evaluation of the changes in the oscillatory pattern of calcium signals at the  
2 growth cone following administration of FGF-2 and of the calcium channel blockers, an approach  
3 based on wavelet analysis was employed, using KYM 0.4 software  
4 (<http://sourceforge.net/projects/kym/>) and according to the formulation given in [40]. The original  
5 trace  $f(t)$  was first detrended and denoised (see e.g. Fig. 6E, upper box), then its wavelet transform  
6  $W(a,b) = |a|^{-1/2} \int \psi^*[(t-b)/a] f(t) dt$  was computed, according to equation 2 of [40], as a function of  
7 time  $b$  and scale  $a$  (\* denotes complex conjugation). Morlet function  $\psi(t) \approx \pi^{-1/4} \exp(-0.5 t^2)$   
8  $\exp(ist)$  was used as mother wavelet ( $s$  is a constant parameter;  $s = 5$  in our case; see equations 3,4,5  
9 of [40]). The resulting scaleogram  $|W(t,v)|$  was plotted with an appropriate Lookup Table (LUT),  
10 after having renamed  $b$  as  $t$  and having done the substitution  $v = s/(2\pi a)$ , in order to get a  
11 representation of the signal in the time-frequency domain (see e.g. Fig. 6E, middle box). Then the  
12 index  $J(t) \approx \int |W(t,v)|^2 v dv$  was calculated as formally defined in equation 9 of [40] and plotted as a  
13 function of time (e.g. Fig. 6E, lower box), so providing a reliable estimate of the oscillatory activity  
14 of the signal  $f(t)$ . Finally, for each temporal window of interest (i.e. control, FGF-2 perfusion and  
15 addition of blockers) the average  $J$  ( $\langle J \rangle$ ) was calculated in order to estimate the increase of  
16 oscillatory activity in terms of ratio between activity during the treatments (FGF-2 or FGF-2 plus  
17 blockers) and activity in control conditions (ctrl):  $r_J = \langle J \rangle_{\text{drug}} / \langle J \rangle_{\text{ctrl}}$  (as defined in equation 10 and  
18 11 of [40]). By this way we could compare the average value of  $r_J$  during FGF-2 administration  
19 with its decrease in the presence of the blockers (see e.g. Fig. 6D), similarly to what we did with  
20  $\Delta R/R$  values obtained from somatic signals (see e.g. Fig. 6B).

### 51 *Statistical analysis*

52 All data are expressed as mean  $\pm$  standard error of the mean (SEM). Normality of the residual  
53 distributions has been verified through Shapiro–Wilk normality test; Levene’s test showed  
54 variances to be quite homogeneous over each condition. For these reasons one-way ANOVA and  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 Tukey-corrected post hoc tests were a suitable method for significance assessment. Statistical  
2 analyses of control and experimental groups were performed using SPSS software.  
3  
4

5  
6  
7 If not otherwise specified, all chemicals and drugs were purchased from Sigma Chemical Co. (St.  
8  
9 Louis, MO).  
10

## 11 **Results**

### 12 **Calcium dependence of neurite growth induced by FGF-2**

13  
14  
15  
16  
17  
18  
19  
20  
21  
22 In order to evaluate the role of calcium signals in the promotion of neurite growth induced by FGF-  
23  
24  
25  
26  
27 2, we cultured for 24 h dissociated neurons loaded with the membrane-permeant calcium chelator  
28  
29 BAPTA-AM, at concentrations (1-5  $\mu$ M) that we checked to have no influence on survival in  
30  
31 dissociated cultures (FGF-2:  $86.99 \pm 6.35\%$  at 24 h,  $78.77 \pm 8.86\%$  at 48 h; FGF-2 + BAPTA 1  $\mu$ M:  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000

90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000

1 dependent increase in neurite growth was significantly inhibited in dissociated cultures (where the  
2 inhibition was complete) and in whole ganglia, respectively at 24 h (Figs. 1 and 2B) and 48 h (Figs.  
3 1 and 2C), providing evidence that calcium influx through voltage-dependent channels is a  
4 necessary step in the cascade of events linking FGFR1 activation to promotion of neurite  
5 elongation. Notably, at these concentrations the two blockers have been shown to have no effect on  
6 FGF-2 dependent neuronal survival [13].

7 E7 CG neurons in culture may show spontaneous electrical activity, that is enhanced by acute  
8 perfusion with FGF-2 [13]; in long-term culture with the factor many cells show calcium signals  
9 that are partially abolished by the voltage-dependent Na<sup>+</sup> channel blocker tetrodotoxin, TTX [2].

10 Therefore, we tested if blocking voltage dependent Na<sup>+</sup> channels could affect the promotion of  
11 neurite growth by FGF-2. In dissociated cultures at 24 h, 10 μM TTX had no effect both on FGF2-  
12 induced neurite growth (Figs. 1 and 2D) and neuronal survival ratios (FGF-2: 95.93 ± 2.53% at 24  
13 h, 93.15 ± 2.85% at 48 h; FGF-2 + TTX 10 μM: 98.33 ± 1.61% at 24 h, 98.56 ± 1.95% at 48 h; p >  
14 0.05).

15 It can be concluded that some additional mechanism(s) must be responsible for the FGF-2 induced  
16 depolarization of the membrane potential and subsequent activation of calcium VOCs. At the  
17 growth cone of extending axons, it has been shown that several neurotrophic factors and guidance  
18 molecules can activate calcium permeable cationic channels of the TRPC family [19, 27, 31, 45,  
19 56]; the ensuing depolarization of the membrane potential may be sufficient to recruit voltage  
20 dependent calcium channels, thus further enhancing calcium influx [17, 56]. While no data are  
21 available for a similar role of FGF-2, we have previously shown [1] that the factor can induce a  
22 calcium influx at least partially dependent on TRPC1 activation in endothelial cells. In the light of  
23 these observations, we tested the role of this family of channels by using SKF 96365, a non  
24 selective blocker of TRPC channels.

25 At both 0.5 and 2 μM, SKF 96365 completely abolished the FGF-2 induced enhancement of neurite  
26 growth in dissociated cultures, while not influencing basal growth in control conditions (Figs. 1 and  
27 2C).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

2E); in organotypic cultures, with 2  $\mu$ M SKF the inhibition was again specific and statistically significant, even if not complete (Figs. 1 and 2F). Notably, up to 2  $\mu$ M, the blocker had no effect on FGF-2 dependent neuronal survival ratios (FGF-2:  $98.71 \pm 3.41\%$  at 24 h,  $101.51 \pm 2.90\%$  at 48 h; FGF-2 + SKF 0.5  $\mu$ M:  $104.18 \pm 2.23\%$  at 24 h,  $106.13 \pm 3.31\%$  at 48 h; FGF-2 + SKF 2  $\mu$ M:  $108.29 \pm 3.28\%$  at 24 h,  $110.41 \pm 4.54\%$  at 48 h;  $p > 0.05$ ).

These data provide evidence that activation of TRPC channels is another key step in the signalling cascade downstream of FGF-2 controlling the growth of neuritic processes in cultured embryonic chick ciliary ganglion neurons.

Finally, calcium release from intracellular stores has been shown to play a role in neurite growth and orientation [28, 30]. In dissociated cultures of ciliary ganglion neurons, 10 nM thapsigargin (TG), an agent that irreversibly depletes intracellular stores, had no effect on neuronal survival ratios (FGF-2:  $84.06 \pm 1.93\%$  at 24 h,  $80.30 \pm 3.08\%$  at 48 h; FGF-2 + TG 10 nM:  $85.86 \pm 2.47\%$ ,  $78.90 \pm 3.64\%$  at 48 h;  $p > 0.05$ ) and on neurite outgrowth (Figs. 1 and 2G), both in control conditions and in the presence of FGF-2. Even at the relatively low concentration of 10 nM, in CG neurons at 24 h of culture, TG effectively induced calcium release from intracellular stores. The release was maximal, as a subsequent perfusion with 100 nM TG did not elicit any additional increase in  $[Ca^{2+}]_i$  ( $n = 65$ ; a representative trace is shown in Fig. 2H). When 2 mM of calcium were reintroduced in the extracellular solution, an increase in  $[Ca^{2+}]_i$  could be observed, indicative of the activation of store operated calcium entry, SOCE [49]; remarkably, subsequent stimulation with FGF-2 elicited a further increase in  $[Ca^{2+}]_i$ .

### **Spatial compartmentalization of FGF-2 induced responses**

A major issue, when dealing with the role of neurotrophic factors in growth and orientation of neuritic processes, is the understanding of the spatial specificity and subcellular localization of the

1 signals activated by the extracellular cue. While most data refer to changes in  $[Ca^{2+}]_i$  at the growth  
2 cone (see e.g. Refs. [17, 52]), the role of the signals originated at the somatic compartment has been  
3  
4 little studied. For this reason we employed compartmentalized cultures by using Campenot  
5  
6 chambers [7, 8] (Fig. 3A).  
7

8  
9 Since protocols with Campenot chambers require the neurons to be cultured for several days, and  
10  
11 since the effect of the addition of FGF-2 in the different compartments had to be compared with  
12  
13 control conditions in which the neurons are cultured with the sole presence of adhesion molecules, a  
14  
15 bias due to cellular death could not be excluded. For this reason, all experiments were performed  
16  
17 with the B27 supplement [6] added to the N2 culture medium. In parallel experiments with  
18  
19 dissociated cultures maintained up to 5 days in the presence of B27, with and without 20 ng/ml  
20  
21 FGF-2, survival ratios were not significantly different in the two experimental conditions: in the  
22  
23 presence of 20 ng/ml LN + B27, survival ratios were  $74.74 \pm 9.56\%$  at 2 d,  $45.43 \pm 8.66\%$  at 5 d;  
24  
25 with added FGF-2, the ratios were  $100.29 \pm 5.00\%$  at 2 d and  $42.81 \pm 5.29\%$  at 5 d ( $p > 0.05$ ).  
26  
27  
28

29  
30 In compartmentalized cultures, when FGF-2 was absent in all compartments, after 5 days in culture  
31  
32 virtually no growth of neuritic processes could be detected in the lateral chambers. When the  
33  
34 neurotrophic factor was added to any compartment, outgrowth of neurites in the lateral  
35  
36 compartments could be detected starting from 3 days of culture (Fig. 3B). All subsequent analysis  
37  
38 was performed after 2 additional days, i.e. at 5 days.  
39  
40  
41

42  
43 A rectangular region of  $175 \mu\text{m} \times 3500 \mu\text{m}$  was selected for each track, starting at  $350 \mu\text{m}$  from the  
44  
45 Teflon barrier (Fig. 3B). Results are shown in Fig. 3C, and can thus be summarized:  
46  
47

- 48 - when FGF-2 was present in the somatic compartment, a bilateral growth occurred, with modest  
49  
50 total occupancy in both lateral compartments;
- 51  
52 - when the factor was present in one lateral compartment, growth was significantly enhanced in this  
53  
54 compartment, while a moderate growth could be detected also in the unstimulated compartment: the  
55  
56 amount of the latter was not significantly different from what observed in the previous experimental  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 condition. These data provide a first evidence that FGF-2 is sensed by the growth cones of  
2 elongating neurites;

3  
4 - when FGF-2 was present in both the central and one lateral compartment, global neurite growth in  
5 both compartments was slightly increased, but not significantly different from what observed with  
6 the factor in one lateral compartment only. Therefore, no synergic effect of FGF-2 presence at the  
7 somatic and neuritic compartments could be evidenced.  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18

### 19 **Characterization and localization of the calcium entry pathways controlling neurite growth**

20  
21  
22  
23  
24 Compartmentalized cultures were next used to localize the contributions of the different calcium  
25 mobilizing mechanisms. All subsequent protocols were performed in the presence of 20 ng/ml FGF-  
26 2 in both the central and one lateral chamber (control condition) and neurite growth was measured  
27 in the chamber in which FGF-2 was present; the blockers were added either in the somatic or in the  
28 neuritic chamber or in both.  
29  
30  
31  
32  
33  
34  
35

36 Addition of the voltage dependent calcium channel blockers nifedipine (10  $\mu$ M) and  $\omega$ -conotoxin (2  
37  $\mu$ M) either in the central or in the lateral compartment reduced global neurite growth by about 60-  
38 70% (Fig. 4A); when the blockers were present in both compartments, a comparable effect was  
39 observed (percentages of neurite occupation: CTRL:  $17.64 \pm 1.57\%$ ; Soma:  $5.21 \pm 0.52\%$ ; Neurites:  
40  $6.30 \pm 1.70\%$ ; Soma+Neurites:  $6.17 \pm 1.00\%$ ).  
41  
42  
43  
44  
45  
46  
47  
48

49 The blocker of TRPC channels, SKF 96365 (2  $\mu$ M), caused effects comparable to those induced by  
50 the VOCs blockers (Fig. 4B); the reduction of global neurite growth was between 50 and 60%  
51 (Percentages of neurite occupation: CTRL:  $17.64 \pm 1.57\%$ ; Soma:  $9.12 \pm 1.67\%$ ; Neurites:  $5.79 \pm$   
52  $2.97\%$ ; Soma+Neurites:  $7.67 \pm 1.71\%$ ).  
53  
54  
55  
56  
57

58 In both cases, no cooperative effect could be evidenced when the blockers were present in both the  
59 central and lateral compartments.  
60  
61  
62  
63  
64  
65



## Spatial localization and specificity of FGF-2 induced calcium signals

To investigate the spatial localization of calcium signals elicited by FGF-2 and of the mechanisms involved in their generation, we simultaneously recorded from the soma and from the growth cone of the same cell. For this purpose, recordings were performed after about 6 hours of culture, in order to be able to visualize at high magnification neurons with relatively short neurites and identifiable growth cones (See Fig. 5A). By this approach, 1-2 neurons per dish, with 1-2 neurites each, could be analyzed.

In a previous paper [13] we used both 20 and 40 ng/ml in acute experiments, obtaining identical results. Here we opted for 40 ng/ml, since the perfusion pipette had to be positioned quite far from the cell, due to the critical adhesion at this early time of culture, and the actual concentration sensed by the cells could not be determined. We performed a set of control experiments with 20 ng/ml FGF-2: the percentage of responsive cells was 45% (9 out of 20 cells). In cells that did not respond to 20 ng/ml FGF-2, subsequent perfusion with the higher concentration failed to induce an increase in  $[Ca^{2+}]_i$  (not shown); in responding cells, perfusion with 40 ng/ml FGF-2 did not elicit any additional response (Suppl. Fig. 2).

Subsequently, a total of 102 cells, with 133 neuritic processes, and 15 isolated neurites (whose soma was outside the field of view) was challenged with 40 ng/ml FGF-2. Responses were recorded in 47 somata (46.03%) and in 86 growth cones (58.10%). The increases in  $[Ca^{2+}]_i$  showed a markedly different time course in the two compartments: at the soma the response was in most cases a smooth and gradual increase in  $[Ca^{2+}]_i$ , followed by either a maintained or a slowly decaying plateau (Fig. 5B, left), in accordance with the responses observed at 24 h [13, 61]. In some cases, the plateau was preceded by a transient phase, indicative of a possible contribution of release from intracellular stores, that has been previously [13] observed in a subpopulation of cells. At the growth cone, the factor usually elicited an oscillatory behaviour, in some cases superimposed on a steady increase

1 (Fig. 5B, right). This difference could reflect differences in cell geometry such as the  
2 surface/volume ratio; however, the picture is probably more complex, since in some instances it was  
3 possible to record signals from two growth cones of the same cell (the same cell of Fig. 5A),  
4 showing sharply different time courses of the responses (Fig. 5C, left and right): therefore, the  
5 spatial specificity of the responses to FGF-2 must be dependent, at least in part, from differences in  
6 the calcium mobilizing mechanisms activated in the different compartments.  
7

8  
9  
10  
11  
12  
13  
14 On both cell bodies and growth cones, substitution of the standard Tyrode solution with another one  
15 containing 0.5 mM EGTA and 0 mM calcium completely and reversibly abolished the FGF-2  
16 induced calcium signals ( $n = 4$ ; Fig. 5D, left: soma; right: growth cone), pointing to a crucial and  
17 necessary role of calcium influx from the extracellular medium.  
18

19  
20  
21  
22  
23  
24 In some cells, spontaneous calcium oscillations could be observed at the growth cone (24.8%), and  
25 to a lesser degree (17.9%) at the soma. These cells responded to FGF-2 in percentages comparable  
26 to basally silent cells (Fig. 5E, left: soma; right: growth cone).  
27

28  
29  
30  
31 Finally, we tested the effects of channel blockers on the responses elicited by FGF-2 in the two  
32 subcellular compartments. In a previous paper [13] we reported that blockers of voltage-dependent  
33 calcium channels did not affect the plateau of the somatic response to FGF-2. The observation was  
34 based on data from a limited set of cells ( $n = 8$ ). This issue is of particular relevance, since we have  
35 shown that block of somatic calcium VOCs significantly reduces FGF-2 induced neurite growth.  
36

37  
38  
39  
40  
41 Therefore, we preliminarily re-evaluated this aspect on a larger population of neurons after 24 h in  
42 culture, the same time of the recordings shown in [13].  
43

44  
45  
46  
47  
48 The effects of the combined addition of the L- and N-type calcium channel blockers on the somatic  
49 response to FGF-2 were evaluated in 52 cells: a significant ( $p < 0.01$ ) and reversible reduction of  
50 the plateau response could be observed in 69.2% of cells; the mean response amplitude ( $\Delta R/R$ ) was  
51  $18.57 \pm 2.27\%$  before and  $11.29 \pm 1.55\%$  after administration of the blockers. A typical recording is  
52 shown in the inset of Fig. 6A. Therefore, at 24 h, a component of the plateau response to FGF-2 at  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 the neuronal soma can be ascribed, in the majority of cells investigated, to calcium influx through  
2 voltage-dependent calcium channels.  
3

4 In CG cultures at 6 h, when the VOCs blockers and SKF were presented to the cells prior to  
5 perfusion with the factors, no change in baseline  $[Ca^{2+}]_i$  could be detected (n = 3; not shown).  
6

7 Acute application of the voltage dependent channel blockers  $\omega$ -conotoxin (2  $\mu$ M) and nifedipine (10  
8  $\mu$ M) affected the somatic and growth cone responses to FGF-2 in quite different ways. 15 cells,  
9 with a total of 19 neurites, and 4 isolated neurites were analyzed. At the soma, the effects were  
10 comparable to those observed at 24 h: a partial, but significant, and reversible reduction of the  
11 amplitude of the plateau response could be recorded in 9 out of 15 somata (Fig. 6A);  $\Delta R/R$  before  
12 application of the blockers was  $61.59 \pm 16.35\%$ , and in the presence of the blockers was  $33.77 \pm$   
13  $13.55\%$  (Fig. 6B).  
14

15 In 15 out of 23 growth cones an effect was observed: in some cases the blockers caused a partial  
16 reduction of the amplitude of the response, but in general they affected the oscillatory behaviour in  
17 a more complex way (Fig. 6C). To extract quantitative information about the change in the pattern  
18 of oscillations induced by the blockers, an approach based on wavelet analysis [40] was employed,  
19 as described in the Methods section, taking  $r_j$  as an index of the change in the oscillatory activity.  
20

21 The average value of  $r_j$  during FGF-2 administration was  $22.14 \pm 7.57$  and decreased to  $6.87 \pm 2.74$   
22 in the presence of the blockers (Fig. 6D, 6E).  
23

24 When SKF 96365 was added to the medium during the response to FGF-2, the somatic and growth  
25 cone effects were remarkably similar to those observed with the VOCs blockers. 19 cells with a  
26 total of 21 neurites, and 4 isolated neurites were analyzed. The TRPC blocker reduced the response  
27 at 12 cell somata (Fig. 7A). Average  $\Delta R/R$  at the soma was  $61.89 \pm 9.40\%$  in the presence of FGF-2  
28 and was reduced to  $41.68 \pm 7.79\%$  following addition of the blocker (Fig 7B). In 18 out of 25  
29 growth cones, a change in activity could be detected (Fig. 7C). Wavelet analysis provided evidence  
30 for a reduction in the oscillatory activity following perfusion with SKF 96365: average  $r_j$  was  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

22.90 ± 7.18 in FGF-2 and decreased to 8.47 ± 2.15 after addition of the blocker (Fig. 7D, 7E).

## Discussion

The data presented in this paper provide new insight into the role of calcium signals activated in response to a neurotrophic factor, FGF-2, in the control of neurite growth and elongation, and point to a specific involvement of different subcellular compartments in this process.

Even if FGF-2 has been considered to be a target derived signal [46, 50], in our case no data are available to support this hypothesis; on the other hand, FGF-2 is expressed by ciliary ganglion neurons in dissociated cultures (A.G. and S.F., unpublished observation). Therefore, it is likely that in the CG it acts as a paracrine/autocrine factor, rather than as a target-derived one. It must be recalled that, even if our data are obtained from mixed cultures, we have previously shown that the role of FGF-2 is cell autonomous [61].

Taking advantage of compartmentalized cultures, we show that FGF-2 exerts a growth promoting action on the neurites. The signalling pathways activated by FGF-2 are not spatially restricted to the growth cone: somatic signals by themselves can induce moderate outgrowth, but only when FGF-2 is sensed by growth cones massive growth occurs. No cooperative effect could be detected when the factor was added in both central and one lateral compartment, suggesting that the signalling at the growth cone induces saturating activation of the machinery responsible for neurite elongation.

A key player in this context is calcium. Growth and orientation of neuritic processes have been shown to depend on calcium signals at the growth cone [18, 62], and to be affected by blockers of both voltage dependent calcium channels (see e.g. Ref. [57]) and of cationic channels of the TRPC family (see e.g. Refs. [28, 31, 56]). The available data provide a quite complex picture: activation of calcium VOCs has been associated with increased growth promoted by FGF-2 [57] and other molecules, including neurotransmitters [22, 44] and with the turning of growth cones towards chemoattractive guidance molecules [17]; on the other hand, it has been reported that their block

1 can rescue, and activation inhibit, neurite and axonal growth [39, 42, 51]. Interestingly, calcium  
2 influx through this class of channels seems not to be a universal prerequisite: in *Xenopus* spinal cord  
3 explants, a mixture of calcium VOCs blockers did not affect axonal growth rate [28].  
4

5 TRPC channels are a more recent entry into the neuritic/axonal growth affair, but the findings are  
6  
7 similarly diverging: while the data of their involvement in growth cone turning behaviour point to a  
8  
9 chemoattractant role, both promotion and inhibition of growth rate have been reported. For instance,  
10  
11 conflicting results have been reported for the role of TRPC5 in axonal growth of hippocampal  
12  
13 neurons [11, 19]; it has been shown to promote neurite growth in a neuroblastoma cell line [58] and  
14  
15 to inhibit it in PC12 cells [25]. Remarkably, Jacques-Fricke et al. [28], reported that blocking TRPC  
16  
17 channels reduced neurite growth rate, while blocking TRPV and stretch activated calcium  
18  
19 permeable channels enhanced it. We obtained information on the specific calcium mobilizing  
20  
21 pathways activated by FGF-2 by means of a pharmacological approach. While the specificity of the  
22  
23 VOCs blockers is well established, the case for SKF-96365 is more controversial [3, 33, 43, 48].  
24  
25

26 However, most of the nonspecific effects, in particular block of other members of the TRP  
27  
28 superfamily or of L-type voltage-dependent calcium channels, are detectable at doses almost two  
29  
30 orders of magnitude above 2  $\mu$ M, the higher concentration used in our experiments. Therefore, the  
31  
32 selectivity of the pharmacological tool used in the present work can be considered sufficient to  
33  
34 prove the involvement of TRPC channels in the signals and biological effects downstream the  
35  
36 activated FGF receptor. A more direct approach, such as silencing of the channel proteins, is not  
37  
38 easily performed on these primary cultures, and due to the high redundancy of the different  
39  
40 members and their complex heteromerization patterns, only multiple silencing could give useful and  
41  
42 reliable answers.  
43  
44

45 Both classes of blockers significantly reduced neurite growth induced by FGF-2. Data from  
46  
47 dissociated and organotypic cultures have been confirmed by those on compartmentalized cultures:  
48  
49 when applied either in the central or in one lateral compartment the blockers markedly reduced the  
50  
51 global growth of neuritic processes. The effects of the two classes were quite similar, and did not  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 completely abolish neurite growth (probably underlying the partial contribution of the two  
2 pathways); on the other hand, other calcium mobilizing mechanisms may be activated by FGF-2.  
3  
4 Moreover, the effects were comparable when the blockers were added either in the central  
5  
6 compartment or at the neurites, and no additivity could be evidenced when they were present in  
7  
8 both.  
9

10 Calcium release from intracellular stores has been reported to play a role in this context, too [17,  
11  
12 28]. In our case no effects could be observed in neurons cultured for 24 h following irreversible  
13  
14 depletion of the stores by TG.  
15  
16

17 To investigate the spatial specificity of calcium signalling in neurons challenged with a  
18  
19 neurotrophic factor, we performed simultaneous recordings of FGF-2 induced changes in  $[Ca^{2+}]_i$   
20  
21 from the growth cone and the soma. To perform this challenging protocol, we had to image calcium  
22  
23 signals at high resolution in cells showing relatively short neurites, so that both compartments could  
24  
25 be visualized in the same field, i.e. in neurons at about 6 h of culture. In these experimental  
26  
27 conditions, FGF-2 elicited long lasting calcium signals in a substantial percentage of neurons, with  
28  
29 specific time courses in the two compartments: a rather stable plateau at the soma (comparable to  
30  
31 signals recorded at 24-48 h [13], and trains of oscillations, in most cases superimposed on a steady  
32  
33 increase in  $[Ca^{2+}]_i$ , at the growth cone. By means of a quantitative analytical approach based on  
34  
35 wavelet analysis we show that channel blockers cause a substantial reduction in the peculiar  
36  
37 oscillatory behaviour at the growth cone, where the growth promoting effect of FGF-2 is more  
38  
39 marked.  
40  
41  
42  
43  
44  
45  
46  
47

48 It is of relevance that the information provided by the different experimental protocols was obtained  
49  
50 at quite different culture times, from 6 h in acute experiments to 1-5 days for assessment on neurite  
51  
52 growth: the substantial agreement between observations at different time scales strengthens the  
53  
54 validity of the results.  
55  
56

57 A final, and relevant, issue is: why blocking calcium channels at the soma, where FGF-2 has a  
58  
59 moderate effect on neurite growth, gave the same reduction observed when the blockers were  
60  
61  
62  
63  
64  
65

1 applied to the neurites? The hypothesis of a retrograde signal could help to explain this observation:  
2 when the factor is presented at the growth cone, by some way the information is transferred to the  
3  
4 soma, where recruitment of voltage dependent and independent calcium channels activates the  
5  
6 machinery responsible for neurite outgrowth. Thus, blocking the channels at the somatic  
7  
8 compartment impairs the process, even if the neuritic compartment is not perturbed. Retrograde  
9  
10 signals have been reported by several Authors for the prototypical neurotrophin, NGF [55, 34] but  
11  
12 also for FGFs [4, 15, 35].  
13  
14

15  
16 The balance between local and global calcium signals in developing neurons, and the mechanisms  
17  
18 by which they can propagate from one compartment to another, has been addressed in a few recent  
19  
20 papers. In cortical neurons Hutchins and Kalil [26] have observed local and global calcium signals  
21  
22 in elongating axons and their collaterals, dependent on calcium VOCs activation. Interestingly, no  
23  
24 changes in  $[Ca^{2+}]_i$  were detected at the soma. Guan et al. [20], have shown that extracellular cues,  
25  
26 such as the repellent Slit-1, act on neuronal migration when applied at the growth cone and not at  
27  
28 the soma: the agonist elicits a local calcium increase that retrogradely propagates to the soma.  
29  
30

31  
32 Finally, Yamada et al. [59] found that glutamate did induce axonal retraction in dentate granule  
33  
34 cells when applied at the somatodendritic area but not when presented at the growth cone; the  
35  
36 response was mediated by a calcium wave originating at the soma and propagated anterogradely to  
37  
38 the growth cone, while an increase in  $[Ca^{2+}]_i$  at the latter site was without effect. Therefore, it  
39  
40 appears that different forms of signal propagation, dependent on specific calcium mobilizing  
41  
42 mechanisms, can modulate the motility of the somatic and growth cone compartments.  
43  
44

45  
46 In our case, some form of crosstalk between the soma (or more precisely, the perisomatic  
47  
48 compartment) and the growth cone appears to exist: while the oscillations at the growth cone are  
49  
50 localized signals and are likely involved in the massive growth observed when the factor is present  
51  
52 only at the periphery, the steady component can represent the global signal that activates  
53  
54 downstream neurite growth-promoting mechanisms dependent on perisomatic calcium influx.  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

In conclusion, by taking advantage of a set of different experimental approaches we have provided a coherent, even if not conclusive, picture of the interplay between the calcium influx mechanisms activated by a neurotrophic factor at different subcellular compartments and their involvement in the promotion of neurite growth. The most relevant findings can be summarized as follows:

- FGF-2 promotion of neurite growth in GC neurons is mainly dependent on calcium influx;
- somatic signals play a role in this process, but maximum growth can be observed only with FGF-2 at the growth cone;
- both calcium VOCs and TRPC channels are involved. Blockers of each of the two channel families can completely abolish the oscillations at the growth cone, thus indicating that a complex interplay between voltage dependent and independent calcium channels is necessary for the activation of this peculiar response. On the other hand, the sustained component appears to be a good candidate for the global signal that can transfer information between the periphery and the soma.

### **Acknowledgements**

Funding from the Compagnia di San Paolo to the NIS Centre of Excellence is gratefully acknowledged. We thank Martina Gullà for help in data analysis.

The Authors declare no competing financial interests.



## References

1. Antoniotti S, Lovisolo D, Fiorio Pla A, Munaron L (2002) Expression and functional role of bTRPC1 channels in native endothelial cells. *FEBS Lett* 510:189-195.
2. Ariano P, Erriquez J, Gilardino A, Ferraro M, Lovisolo D, Distasi C (2006) Calcium signals and the in vitro migration of chick ciliary ganglion cells. *Cell Calcium*. 40(1):63-71.
3. Barbiero G, Munaron L, Antoniotti S, Baccino FM, Bonelli G, Lovisolo D (1995) Role of mitogen-induced calcium influx in the control of the cell cycle in Balb-c 3T3 fibroblasts. *Cell Calcium* 18:542-556.
4. Blottner D, Stapf C, Meisinger C, Grothe C (1997) Localization, differential expression and retrograde axonal transport suggest physiological role of FGF-2 in spinal autonomic neurons of the rat. *Eur J Neurosci* 9:368-377.
5. Bottenstein JE (1983) Defined media for dissociated neural cultures. *Curr Methods Cell Neurobiol* 4:107-130.
6. Brewer GI, Torricelli JR, Evege EK, Price PJ (1993) Optimized survival of hippocampal neurons in B27-supplemented Neurobasal, a new serum-free medium combination, *J Neurosci Res*. 35:567-576.
7. Campenot RB (1977) Local control of neurite development by nerve growth factor. *Proc Natl Acad Sci U S A* 74:4516-4519.
8. Campenot RB, Walji AH, Draker DD (1991) Effects of sphingosine, staurosporine, and phorbol ester on neurites of rat sympathetic neurons growing in compartmented cultures. *J Neurosci* 11:1126-1139.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- 9 Catapano LA, Arlotta P, Cage TA, Macklis JD (2004) Stage-specific and opposing roles of BDNF, NT3 and bFGF in differentiation of purified callosal projection neurons toward cellular repair of complex circuitry. *Eur J Neurosci* 19:2421-434.
  - 10 Cox LJ, Hengst U, Gurskaya NG, Lukyanov KA, Jaffrey SR (2008) Intra-axonal translation and retrograde trafficking of CREB promotes neuronal survival. *Nat Cell Biol* 10:149-159.
  11. Davare MA, Fortin DA, Saneyoshi T, Nygaard S, Kaech S, Banker G, Soderling TR, Wayman GA (2009) Transient receptor potential canonical 5 channels activate Ca<sup>2+</sup>/calmodulin kinase Igamma to promote axon formation in hippocampal neurons. *J Neurosci* 29:9794-9808.
  12. Dickson BJ (2002) Molecular mechanisms of axon guidance. *Science* 298:1959-1964.
  13. C, Torre M, Antoniotti S, Munaron L, Lovisolo D (1998) Neuronal survival and calcium influx induced by basic fibroblast growth factor in chick ciliary ganglion neurons. *Eur J Neurosci* 10:2276-2286.
  - 14 Dryer SE (1994) Functional development of the parasympathetic neurons of the avian ciliary ganglion: a classic model system for the study neuronal differentiation and development. *Progr Neurobiol* 43:281–322.
  - 15 Ferguson IA, Johnson EM Jr (1991) Fibroblast growth factor receptor-bearing neurons in the CNS: identification by receptor-mediated retrograde transport. *J Comp Neurol* 313:693-706.
  16. Gilardino A, Farcito S, Zamburlin P, Audisio C, Lovisolo D (2009) Specificity of the second messenger pathways involved in basic fibroblast growth factor-induced survival and neurite growth in chick ciliary ganglion neurons. *J Neurosci Res* 87:2951-2962.
  17. Gomez TM (2005) Channels for pathfinding. *Nature* 434:835-838.

18. Gomez TM, Zheng JQ (2006) The molecular basis for calcium-dependent axon pathfinding. *Nat Rev Neurosci* 7:115-125.
19. Greka A, Navarro B, Oancea E, Duggan A, Clapham DE (2003) TRPC5 is a regulator of hippocampal neurite length and growth cone morphology. *Nat Neurosci* 6:837-845.
20. Guan CB, Xu HT, Jin M, Yuan XB, Poo MM (2007) Long-range Ca<sup>2+</sup> signaling from growth cone to soma mediates reversal of neuronal migration induced by slit-2. *Cell* 129:385-395.
21. Henley J, Poo MM (2004). Guiding neuronal growth cones using Ca<sup>2+</sup> signals. *Trends Cell Biol* 14:320-330.
22. Homma K, Kitamura Y, Ogawa H, Oka K (2006) Serotonin induces the increase in intracellular Ca<sup>2+</sup> that enhances neurite outgrowth in PC12 cells via activation of 5-HT<sub>3</sub> receptors and voltage-gated calcium channels. *J Neurosci Res* 84:316-325.
23. Hong K, Nishiyama M, Henley J, Tessier-Lavigne M, Poo, MM (2000) Calcium signalling in the guidance of nerve growth by netrin-1. *Nature* 403:93–98.
24. Hossain WA, Morest DK (2000) Fibroblast growth factors (FGF-1, FGF-2) promote migration and neurite growth of mouse cochlear ganglion cell in vitro: immunohistochemistry and antibody perturbation. *J Neurosci Res* 62:640–655.
25. Hui H, McHugh D, Hannan M, Zeng F, Xu SZ, Khan SU, Levenson R, Beech DJ, Weiss JL (2006) Calcium-sensing mechanism in TRPC5 channels contributing to retardation of neurite outgrowth. *J Physiol* 572:165-172.
26. Hutchins BI, Kalil K (2008) Differential outgrowth of axons and their branches is regulated by localized calcium transients. *J Neurosci* 28:143-153.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
27. Hutchins BI, Li L, Kalil K (2012) Wnt-induced calcium signaling mediates axon growth and guidance in the developing corpus callosum. *Sci Signal*. 2012 Jan 10;5(206):pt1. doi: 0.1126/scisignal.2002523.
28. Jacques-Fricke BT, Seow Y, Gottlieb PA, Sachs F, Gomez TM (2006) Ca<sup>2+</sup> influx through mechanosensitive channels inhibits neurite outgrowth in opposition to other influx pathways and release from intracellular stores. *J Neurosci* 26:5656-5664.
29. Kolodkin AL, Tessier-Lavigne M (2011) Mechanisms and Molecules of Neuronal Wiring: A Primer. *Cold Spring Harb Perspect Biol* 3 (6) pii: a001727. doi: 10.1101/cshperspect.a001727.
30. Li L, Hutchins BI, Kalil K (2009) Wnt5a induces simultaneous cortical axon outgrowth and repulsive axon guidance through distinct signaling mechanisms. *J Neurosci* 29:5873-5883.
31. Li Y, Jia YC, Cui K, Li N, Zheng ZY, Wang YZ, Yuan XB (2005) Essential role of TRPC channels in the guidance of nerve growth cones by brain-derived neurotrophic factor. *Nature* 434:894-898.
32. Lin AC, Holt CE (2007) Local translation and directional steering in axons. *EMBO J* 26:3729-3736.
33. Merritt JE, Armstrong WP, Benham CD, Hallam TJ, Jacob R, Jaxa-Chamiec A, Leigh BK, McCarthy SA, Moores KE, Rink TJ (1990) SK&F 96365, a novel inhibitor of receptor-mediated calcium entry. *Biochem J* 271:515-522.
34. Mok SA, Campenot RB (2007) A nerve growth factor-induced retrograde survival signal mediated by mechanisms downstream of TrkA. *Neuropharmacology* 52:270-278.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
35. Mufson EJ, Kroin JS, Sendera TJ, Sobreviela T (1999) Distribution and retrograde transport of trophic factors in the central nervous system: functional implications for the treatment of neurodegenerative diseases. *Prog Neurobiol* 57:451-484.
  36. Pertz OC, Wang Y, Yang F, Wang W, Gay LJ, Gristenko MA, Clauss TR, Anderson DJ, Liu T, Auberry KJ, Camp DG 2nd, Smith RD, Klemke RL (2008). Spatial mapping of the neurite and soma proteomes reveals a functional Cdc42/Rac regulatory network. *Proc Natl Acad Sci U S A* 105:1931-1936.
  37. Posse de Chaves E, Vance DE, Campenot RB, Vance JE (1995) Axonal synthesis of phosphatidylcholine is required for normal axonal growth in rat sympathetic neurons. *J Cell Biol* 128:913-918.
  38. Rasband, WS (1997-2012) ImageJ, U. S. National Institutes of Health, Bethesda, Maryland, USA, <http://imagej.nih.gov/ij/>.
  39. Roehm PC, Xu N, Woodson EA, Green SH, Hansen MR (2008) Membrane depolarization inhibits spiral ganglion neurite growth via activation of multiple types of voltage sensitive calcium channels and calpain. *Mol Cell Neurosci* 37:376-387.
  40. Ruffinatti FA, Lovisolo D, Distasi C, Ariano P, Erriquez J, Ferraro M (2011) Calcium signals: analysis in time and frequency domains. *J Neurosci Methods* 199:310-320.
  41. Sánchez-Camacho C, Rodríguez J, Ruiz JM, Trousse F, Bovolenta P (2005) Morphogens as growth cone signalling molecules. *Brain Res Brain Res Rev* 2005 49:242-252.
  42. Sann SB, Xu L, Nishimune H, Sanes JR, Spitzer NC (2008) Neurite outgrowth and in vivo sensory innervation mediated by a Ca(V)2.2-laminin beta 2 stop signal. *J Neurosci* 28:2366-2374.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
43. Schwarz G, Droogmans G, Nilius B (1994) Multiple effects of SK&F 96365 on ionic currents and intracellular calcium in human endothelial cells. *Cell Calcium* 15:45–54.
  44. Schmitz Y, Luccarelli J, Kim M, Wang M, Sulzer D (2009) Glutamate controls growth rate and branching of dopaminergic axons. *J Neurosci* 29:11973-1181.
  45. Shim S, Goh EL, Ge S, Sailor K, Yuan JP, Roderick HL, Bootman MD, Worley PF, Song H, Ming GL (2005) XTRPC1-dependent chemotropic guidance of neuronal growth cones. *Nat Neurosci* 8:730-735.
  46. Shirasaki R, Lewcock JW, Lettieri K, Pfaff SL (2006) FGF as a target-derived chemoattractant for developing motor axons genetically programmed by the LIM code. *Neuron* 50:841-853.
  47. Singh KK, Miller FD (2005) Activity regulates positive and negative neurotrophin-derived signals to determine axon competition. *Neuron* 45:837-845.
  48. Singh A, Hildebrand ME, Garcia E, Snutch TP (2010) The transient receptor potential channel antagonist SKF96365 is a potent blocker of low-voltage-activated T- type calcium channels. *Br J Pharmacol* 160:1464–1475.
  49. Smyth JT, Hwang SY, Tomita T, DeHaven WI, Mercer JC, Putney JW (2010) Activation and regulation of store-operated calcium entry. *J Cell Mol Med* 14:2337-2349.
  50. Szebenyi G, Dent EW, Callaway JL, Seys C, Lueth H, Kalil K (2001) Fibroblast growth factor-2 promotes axon branching of cortical neurons by influencing morphology and behavior of the primary growth cone. *J Neurosci* 21:3932-3941.
  51. Spitzer NC, Lautermilch NJ, Smith RD, Gomez TM (2000) Coding of neuronal differentiation by calcium transients. *Bioessays* 22:811-817.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
52. Tojima T, Hines JH, Henley JR, Kamiguchi H (2011) Second messengers and membrane trafficking direct and organize growth cone steering. *Nat Rev Neurosci* 12:191-203.
  53. Thomas KA (1987) Fibroblast growth factors. *FASEB J* 1:434-440.
  54. Tonge D, Chan K, Zhu N, Panjwani A, Arno M, Lynham S, Ward M, Snape A, Pizzey J (2008) Enhancement of axonal regeneration by in vitro conditioning and its inhibition by cyclopentenone prostaglandins. *J Cell Sci* 12:2565-2577.
  55. Ure DR, Campenot RB (1994) Leukemia inhibitory factor and nerve growth factor are retrogradely transported and processed by cultured rat sympathetic neurons. *Dev Biol* 162:339-347.
  56. Wang GX, Poo MM (2005) Requirement of TRPC channels in netrin-1-induced chemotropic turning of nerve growth cones. *Nature* 434:898-904.
  57. Williams EJ, Furness J, Walsh FS, Doherty P (1994) Characterisation of the second messenger pathway underlying neurite outgrowth stimulated by FGF. *Development* 120:1685-1693.
  58. Wu G, Lu ZH, Obukhov AG, Nowycky MC, Ledeen RW (2007) Induction of calcium influx through TRPC5 channels by cross-linking of GM1 ganglioside associated with alpha5beta1 integrin initiates neurite outgrowth. *J Neurosci* 27:7447-758.
  59. Yamada RX, Sasaki T, Ichikawa J, Koyama R, Matsuki N, Ikegaya Y (2008) Long-range axonal calcium sweep induces axon retraction. *J Neurosci* 28:4613-4618.
  60. Zamburlin P, Gilardino A, Dalmazzo S, Ariano P, Lovisolo D (2006) Temporal dynamics of neurite outgrowth promoted by basic fibroblast growth factor in chick ciliary ganglia. *J Neurosci Res* 84:505-514.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

61. Zamburlin P, Ruffinatti FA, Gilardino A, Farcito S, Lovisolo D (2012) Calcium signals induced by FGF-2 in parasympathetic neurons: Role of second messenger pathways. *Neurosci Lett* 23:30-34.
62. Zheng JQ, Poo MM. (2007) Calcium signaling in neuronal motility. *Annu Rev Cell Dev Biol* 23:375-404.



## Figure legends

1  
2  
3  
4  
5  
6 Figure 1. Evaluation of global neurite growth in dissociated and organotypic cultures.

7  
8 Images of dissociated (upper row) and organotypic (lower row) cultures of E7 chick CG neurons,  
9  
10 respectively at 24 and 48 h of culture, in control conditions (P+LN), with FGF-2 alone and in the  
11  
12 added presence of BAPTA and of the various inhibitors of calcium mobilizing mechanisms.

13  
14 Staining procedures as described in the Methods section. Scale bars: A 5  $\mu\text{m}$ , B 20  $\mu\text{m}$ .

15  
16  
17  
18  
19  
20  
21  
22  
23 Figure 2. Effects of blockers of different calcium mobilizing pathways on neurite growth at 24 h in  
24  
25 dissociated (A, B, D, E, G) and at 48 h in organotypic (C, F) cultures of E7 chick CG neurons.

26  
27 A. In dissociated neurons, incubation with 1-5  $\mu\text{M}$  BAPTA-AM did reduce global neurite growth  
28  
29 dependent on FGF-2 without affecting growth in the sole presence of adhesion molecules. Values of  
30  
31 neurite occupation were: CTRL (P+LN):  $471.61 \pm 16.45 \mu\text{m}^2$ ; FGF-2:  $610.86 \pm 22.79 \mu\text{m}^2$ ; BAPTA  
32  
33 1  $\mu\text{M}$ :  $426.04 \pm 17.41 \mu\text{m}^2$ ; BAPTA 5  $\mu\text{M}$ :  $403.72 \pm 21.99 \mu\text{m}^2$ ; FGF-2 + BAPTA 1  $\mu\text{M}$ :  $492.42 \pm$   
34  
35  $21.18 \mu\text{m}^2$ ; FGF-2 + BAPTA 5  $\mu\text{M}$ :  $415.30 \pm 15.06 \mu\text{m}^2$ . \*\*  $p < 0.01$  vs. CTRL condition; ++  $p <$   
36  
37  $0.01$  vs. FGF-2 condition.

38  
39  
40  
41  
42 B. 2  $\mu\text{M}$   $\omega$ -conotoxin and 10  $\mu\text{M}$  nifedipine (CTX +NIFE) reduced FGF-2 induced neurite growth  
43  
44 in dissociated cultures. Values of neurite occupation were: CTRL:  $621.42 \pm 45.33 \mu\text{m}^2$ ; FGF-2:  
45  
46  $869.30 \pm 86.90 \mu\text{m}^2$ ; CTX+NIFE  $608 \pm 43.66 \mu\text{m}^2$ ; FGF-2 + CTX+NIFE:  $497.17 \pm 25.27 \mu\text{m}^2$ . \*\*  $p$   
47  
48  $< 0.01$  vs. CTRL; ++  $p < 0.01$  vs. FGF-2 condition.

49  
50  
51 C. Comparable effects of the two  $\text{Ca}^{2+}$  VOCs blockers were observed in organotypic cultures.

52  
53 Percentages of neurite occupation were: CTRL:  $14.23 \pm 3.21 \%$ ; FGF-2:  $42.45 \pm 4.39 \%$ ;

54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

D. Incubation with tetrodotoxin (TTX), blocker of Na<sup>+</sup> VOCs, had no significant effect on neurite growth in dissociated cultures. CTRL: 544.85 ± 21.10 μm<sup>2</sup>; FGF-2: 743.72 ± 36.09 μm<sup>2</sup>; TTX: 598 ± 22.36 μm<sup>2</sup>; FGF-2 + TTX: 698.59 ± 22.84 μm<sup>2</sup>. \*\* p < 0.01 vs. CTRL; # p < 0.05 vs. TTX condition.

E. Neurite occupation promoted by FGF-2 in dissociated cultures was reduced by the TRPC channel blocker SKF 96365. CTRL: 623.70 ± 67.81 μm<sup>2</sup>; FGF-2: 918.71 ± 81.93 μm<sup>2</sup>; SKF 0.5 μM: 586.81 ± 50.39 μm<sup>2</sup>; SKF 2 μM: 517.72 ± 43.50 μm<sup>2</sup>; FGF-2 + SKF 0.5 μM: 519.34 ± 27.99 μm<sup>2</sup>; FGF-2 + SKF 2 μM: 575.53 ± 35.95 μm<sup>2</sup>. \*\* p < 0.01 vs CTRL; ++ p < 0.01 vs. FGF-2 condition.

F. Comparable effects of SKF 96365 were observed in organotypic cultures. CTRL: 8.73 ± 1.42 %; FGF-2: 35.91 ± 2.68 %; SKF 0.5 μM: 8.53 ± 1.64 %; SKF 2 μM: 5.13 ± 1.20 %; FGF-2 + SKF 0.5 μM: 35.27 ± 3.71 % ; FGF-2 + SKF 2 μM: 22.13 ± 3.54 %. \*\* p < 0.01 vs CTRL; + p < 0.05 vs FGF-2 condition.

G. Interfering with calcium release from intracellular stores by preincubation of dissociated cultures with 10 nM thapsigargin (TG), an inhibitor of SERCA calcium pumps, had no significant effect on neurite growth. CTRL: 612.45 ± 29.68 μm<sup>2</sup>; FGF-2: 838.40 ± 62.80 μm<sup>2</sup>; TG: 562.29 ± 20.85 μm<sup>2</sup>; FGF-2 + TG: 761.88 ± 32.42 μm<sup>2</sup>. \*\* p < 0.01 vs CTRL; ## p < 0.01 vs TG.

H. 10 nM TG induced saturating calcium release from intracellular stores in CG neurons in dissociated cultures at 24 h, when added to a nominally calcium free solution with 0.5 mM EGTA. Subsequent stimulation with 100 nM TG did not elicit any additional increase in [Ca<sup>2+</sup>]<sub>i</sub>. Reintroduction of 2 mM calcium in the extracellular medium evoked an increase in [Ca<sup>2+</sup>]<sub>i</sub>; further stimulation with 40 ng/ml FGF-2 was still able to evoke an additional increase in [Ca<sup>2+</sup>]<sub>i</sub>.

Figure 3. Compartmentalized cultures: the contribution of somatic and neuritic signalling activated by FGF-2 to global neurite growth.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

A. Schematic drawing of the Campenot chambers used in our experiments.

B. Images of a few tracks in a lateral compartment of a Campenot chamber at 5 days of culture, with neurites, stained with the anti neurofilament antibody, extending along them. In gray is depicted the field selected for analysis. Scale bar: 500  $\mu$ m.

C. Neurite occupancy in the left (L) and right (R) lateral compartments of Campenot chambers when 20 ng/ml FGF-2 was present either at the somatic or at one lateral compartment, or in both. Percentages of occupation were: FGF-2 in the central compartment, L  $2.75 \pm 1.11$  %, R  $2.39 \pm 0.66$  %; FGF-2 in R lateral compartment, L  $0.91 \pm 0.24$  %, R  $12.15 \pm 2.90$  %; FGF-2 in both central and R lateral compartment, L  $2.21 \pm 1.39$  %, R  $17.63 \pm 1.57$  %. \*  $p < 0.05$  and \*\*  $p < 0.01$  vs. L and R with FGF-2 in the central compartment, vs. L with FGF-2 in the R compartment, and vs. L with FGF-2 in both central and R compartment.

Figure 4. Effects of addition of the calcium channel blockers in the different compartments on FGF-2 induced neurite growth.

A. 2  $\mu$ M  $\omega$ -conotoxin and 10  $\mu$ M nifedipine (CTX+NIFE) significantly reduced neurite growth when added in either the central compartment or in the lateral compartment, or in both. CTRL: 20 ng/ml FGF-2 alone. In all protocols, FGF-2 was present in both the central and one lateral compartment. Percentages of neurite occupation were: CTRL:  $17.64 \pm 1.57$  %; Blockers on soma:  $5.21 \pm 0.52$  %; Blockers on neurites:  $6.30 \pm 1.70$  %; Blockers on soma and neurites:  $6.17 \pm 1.00$  %. \*\*  $p < 0.01$  vs. all other conditions.

B. The TRPC channel blocker SKF-96365 exerted effects comparable to those of the VOCs blockers. Percentages of neurite occupation: CTRL:  $17.64 \pm 1.57$  %; SKF on soma:  $9.12 \pm 1.67$  %; SKF on neurites:  $5.79 \pm 2.97$  %; SKF on soma and neurites:  $7.67 \pm 1.71$  %. \*  $p < 0.05$  vs. all other conditions.

1  
2 Figure 5. Calcium signals at the soma and at the growth cone activated by acute application of 40  
3  
4 ng/ml FGF-2 in CG neurons at 6 h of culture.  
5  
6

7 A. Image obtained by Intensity Modulated Display (IMD) mode of a neuron with two neurites. The  
8  
9 neuron was loaded with Fura-2 and challenged with 40 ng/ml FGF-2. Scale bar 5  $\mu\text{m}$ .  
10

11 B. Left: a representative trace showing a somatic long lasting plateau increase in  $[\text{Ca}^{2+}]_i$  in response  
12  
13 to FGF-2. Right: trace from the growth cone of the same cell. Perfusion with the factor induced the  
14  
15 onset of an oscillatory behaviour. Responding somata were 47 of 102 (46.03%); responding growth  
16  
17 cones 86 of 148 (58.10%).  
18  
19  
20

21 C. Recording from the two growth cones of the neuron shown in A. Left: growth cone marked with  
22  
23 +; right: growth cone marked with \*.  
24  
25

26 D. Perfusion with a low calcium extracellular solution reversibly abolished responses to FGF-2 both  
27  
28 at the soma (left) and at the growth cone (right).  $N = 4$ .  
29  
30

31 E. An example of a cell showing spontaneous oscillations of  $[\text{Ca}^{2+}]_i$  both at the soma (left) and at  
32  
33 the growth cone (right). Subsequent perfusion with 40 ng/ml FGF-2 induced a change in  $[\text{Ca}^{2+}]_i$  in  
34  
35 both subcellular compartments. Percentages of cells showing spontaneous oscillations: 17.9 % at  
36  
37 the soma, 24.8 % at the growth cone.  
38  
39  
40  
41  
42  
43  
44  
45

46 Figure 6. Blockers of calcium VOCs affect the FGF-2 induced calcium signals.  
47

48 A. A representative trace showing the reduction of the somatic plateau response to 40 ng/ml FGF-2  
49  
50 by the blockers of N- and L- type calcium channels (2  $\mu\text{M}$   $\omega$ -conotoxin and 10  $\mu\text{M}$  nifedipine).  
51

52 Inset: a representative trace showing the effects of the calcium VOCs blockers on FGF-2 induced  
53  
54 somatic increases in  $[\text{Ca}^{2+}]_i$  at 24 h of culture.  
55  
56  
57

58 B. Average reduction of the plateau response to FGF-2 by the calcium VOCs blockers.  $N = 9$ ; \*  $p <$   
59  
60 0.05; paired-samples t-test.  
61  
62  
63  
64  
65

1  
2 C. Complex effects of the VOCs blockers on the oscillatory response to FGF-2 at the growth cone  
3 of the same neuron.

4  
5 D. Average increase of oscillatory activity relative to basal behaviour,  $r_J$ , during administration of  
6 FGF-2 and after addition of the VOCs blockers.  $N = 15$ ; \*  $p < 0.05$ ; paired-samples t-test.

7  
8  
9 E. Wavelet analysis of the oscillatory activity at the same growth cone as in C in basal conditions,  
10 following administration of 40 ng/ml FGF-2 and subsequent addition of the calcium VOCs  
11 blockers. Upper box: original trace (in blue) and the same signal after a detrending and denoising  
12 procedure (in red). Middle box: plot of the scaleogram representing the wavelet transform of the red  
13 trace in the upper box. Lower box: plot of the activity index  $J$  as a function of time. Vertical red  
14 lines in the upper and lower boxes mark the changes in the extracellular solution indicated in C.  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28

29 Figure 7. Effects of the TRPC blocker, SKF-96365, on FGF-2 induced calcium signals.

30  
31 A. A representative trace showing the reduction of the somatic plateau response to 40 ng/ml FGF-2  
32 by SKF 96365.  
33

34  
35 B. Average reduction of the plateau response to FGF-2 by SKF 96365.  $N = 12$ ; \*\*  $p < 0.01$ ; paired-  
36 samples t-test.  
37  
38

39  
40 C. Complex effects of the TRPC blocker on the oscillatory response to FGF-2 at the growth cone.  
41

42  
43 D. Average increase of oscillatory activity relative to basal behaviour,  $r_J$ , during administration of  
44 FGF-2 and after addition of SKF 96365.  $N = 18$ ; \*  $p < 0.05$ ; paired-samples t-test.  
45

46  
47 E. Wavelet analysis of the oscillatory activity at the same growth cone as in C in basal conditions,  
48 following administration of 40 ng/ml FGF-2 and subsequent addition of SKF 96365. Upper box:  
49 original trace (in blue) and the same signal after a detrending and denoising procedure (in red).  
50  
51 Middle box: plot of the scaleogram representing the wavelet transform of the red trace in the upper  
52 box. Lower box: plot of the activity index  $J$  as a function of time. Vertical red lines in the upper and  
53 lower boxes mark the changes in the extracellular solution indicated in C.  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Figure 1  
[Click here to download high resolution image](#)





Figure 2

[Click here to download high resolution image](#)




Figure3  
[Click here to download high resolution image](#)






Figure4

[Click here to download high resolution image](#)





Figure 5

[Click here to download high resolution image](#)




Figure6  
[Click here to download high resolution image](#)





Figure 7

[Click here to download high resolution image](#)



Supplementary Figure 1

[Click here to download Supplementary Material: Suppl Figure 1.tif](#)

Supplementary Figure2

[Click here to download Supplementary Material: Suppl Figure 2.tif](#)

Supplementary Fig. Legends

[Click here to download Supplementary Material: Suppl.Fig. Legends.doc](#)